



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## The association between use of asthma drugs and the prevalence of hypomineralized permanent first molar teeth in 6 to 8-year old Danish children

Nechifor, Ioan Alin; Boila, Adela Corina; Wogelius, Pia; Haubek, Dorte; Tvedebrink, Torben; Nørgaard, Mette; Poulsen, Sven

*Publication date:*  
2007

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### Citation for published version (APA):

Nechifor, I. A., Boila, A. C., Wogelius, P., Haubek, D., Tvedebrink, T., Nørgaard, M., & Poulsen, S. (2007). *The association between use of asthma drugs and the prevalence of hypomineralized permanent first molar teeth in 6 to 8-year old Danish children*. Poster presented at Forskningens Dag, Aalborg, Denmark.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

# The association between use of asthma drugs and the prevalence of hypomineralized permanent first molar teeth in 6 to 8-year old Danish children

Ioan Alin Nechifor<sup>4</sup>, Adela Corina Boila<sup>4</sup>, Pia Wogelius<sup>1,2</sup>, Dorte Haubek<sup>1</sup>, Torben Tvedebrink<sup>3</sup>, Mette Nørgaard<sup>1</sup>, Sven Poulsen<sup>2</sup>

1. Department of Clinical Epidemiology, Aalborg Hospital Aarhus University Hospital, Denmark.
2. Department of Community Oral Health and Pediatric Dentistry, School of Dentistry, Faculty of Health Sciences, Aarhus University, Denmark.
3. Center for Cardiovascular Research, Aalborg Hospital Aarhus University Hospital, Denmark
4. University of Medicine and Pharmacy, Targu-Mures, Romania.

## Background

Permanent first molar hypomineralization (PFMH) is a common developmental tooth defect, characterized by an insufficient calcium deposit in the enamel. There is a wide range of lesion types but the more serious ones appear clinically as yellow opacities or lesions with loss of tooth substance. The etiology is sparsely elucidated but asthma drugs have been suspected.

## Aim

The aim of this study was to examine the association between redeemed prescriptions of asthma drugs and the prevalence of PFMH among Danish children.

## Methods

We performed a cross-sectional population-based study in two Danish municipalities. All children at age 6, 7, and 8 years were included ( $n=891$ ) and we examined the teeth of 745(83.6%) of them. We restricted the analyses to the 647 children with eruption of all four permanent first molars.

Data on use of asthma drugs from birth until the time of the dental examination were obtained from a prescription-database.

## Results

A positive association was found between use of both inhaled  $\beta_2$  agonists and inhaled corticosteroids (asthma drugs) and risk of hypomineralizations. Among 47 children with prescriptions of asthma drugs before the age of three years 6(12.8%), and 9(19.29%) had lesions with loss of tooth substance, and yellow opacities or loss of tooth substance, respectively. The corresponding figures among 264 children with no prescriptions for either inhaled or per oral asthma drugs were 13(4.9%), and 30(11.4%) (Table 1). The prevalence ratios are presented in Table 2.

## Conclusions

Children with prescription of inhaled asthma drugs before the age of three have an increased risk for hypomineralization with macroscopic loss of enamel. It is uncertain whether this was a consequence of the drugs, the asthma or both.




Photo: Dorte Haubek

Table 1. Number (and %) of children with prescriptions of asthma drugs, according to gender and the type of hypomineralization.

Number of children	Inhaled $\beta_2$ -agonists and corticosteroids		No asthma drug prescriptions
	Anytime (%)	Before 3 <sup>rd</sup> year (%)	Anytime (%)
N	65	47	264
Gender			
Male	36 (55.4)	29 (61.7)	127 (48.1)
Female	29 (44.6)	18 (38.3)	137 (51.9)
Type of hypomineralization			
Lesions with loss of tooth substance	5 (7.7)	6 (12.8)	13 (4.9)
Yellow opacities or lesions with loss of tooth substance	9 (13.9)	9 (19.2)	30 (11.4)

Table 2. Prevalence ratio (with 95% CI) of hypomineralization in molar teeth among children with use of asthma drugs.

Hypomineralization lesions	Prevalence ratio with 95% CI		
	All	Boys	Girls
Yellow opacities or lesions with loss of tooth substance	1.69 (0.86-3.32)	1.29 (0.52-3.21)	2.34 (0.86-6.41)
Lesions with loss of tooth substance	2.59 (1.04-6.48)	2.19 (0.42-11.39)	3.38 (1.16-9.87)

## Acknowledgements

Thanks to the dental staff in Støvring and Nibe municipalities who collected the clinical data to this study. The study received financial support from the "Augustinus Foundation"