



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The influence of eccentric hip abductor strength on contralateral pelvic drop and mediolateral knee movement

Brund, René Børge Korsgaard; Rasmussen, Sten; Nielsen, Rasmus Østergaard; Kersting, Uwe G.; Voigt, Michael

Published in:

International Calgary Running Symposium, ICRS, 14-17 August 2014

Publication date:
2014

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Brund, R. B. K., Rasmussen, S., Nielsen, R. Ø., Kersting, U. G., & Voigt, M. (2014). The influence of eccentric hip abductor strength on contralateral pelvic drop and mediolateral knee movement. In *International Calgary Running Symposium, ICRS, 14-17 August 2014* (pp. 103-104). University of Calgary.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

The influence of eccentric hip abductor strength on contralateral pelvic drop and mediolateral knee movement

René Korsgaard, Sten Rasmussen, Rasmus O. Nielsen, Michael Voigt and Uwe G. Kersting

Affiliations

Background

Running is associated with overuse injuries and **approximately 50% of all recreational and competitive runners experience this type of injuries yearly** (1). The predominant sites of injuries are located in the knee. Hip Abductor weakness is related to knee injuries like Patellofemoral pain and illiotibial band syndromes (2,5,6). This relationship is likely due to the hip abductor muscles is stabilizing the hip and knee by preventing a hip adduction, internal rotation of femur and a knee abduction (3,4). The hip abductors works eccentric during stance and therefore low eccentric hip abductor strength may be related with a higher contralateral pelvic drop (CPD) and mediolateral knee movement (MKM) during stance (2-4). Therefore a weak eccentric hip abductor muscle may be a potential risk of knee injuries.

Objectives

The objective of the study was to elaborate the influence of eccentric hip abductor strength on CPD and MKM during stance.

Hypothesis

Low eccentric hip abductor strength induces the highest CPD and MKM during stance


Design

26 male recreational runners running at least 2 times a week and had at least 2 years of running experience and without any pain for the last 3months. Hip abductor strength, CPD and MKM was measured on both legs and the each leg was considered as independent.

Hip abductor strength and demographics for High vs low Strength group (mean +/-sd)

	Low (n=26 legs)	High (n = 26 legs)	P-value
<i>Eccentric hip abduction</i>			
Strength (nm)	144,42 +/-16,14	200,54 +/-22,95	0,000*
Strength / kg (nm/kg)	1,91+/-0,23	2,55+/-2,55	0,000*
Bodyweigth (kg)	76,08+/-7,77	79,46+/-7,65	0,06
Age (years)	34,12+/-7,97	30,58+/-10,35	0,08
Height (m)	178,00+/-23,54	176,46+/-23,16	0,4

Outcome



Discussion/Perspective

Low eccentric hip abductor strength probably has an influence on CPD and MKM but should be adjusted for weight and maybe static knee alignment and strike pattern as well. Results need power and therefore more subjects are needed.

References

- 1: Ferber R., Hreljac A. and Mkin K. D. K. (2009) suspected mechanisms in the cause of overuse running injuries: A clinical review. *American Orthopaedic Society for Sports Medicine* 1(3), 242-246
- 2: Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreicher N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin. J. Sport Med.* 2000;10(3):169-75.
- 3: Thorborg K, Bandholm T, Petersen J, et al. Hip abduction strength training in the clinical setting: with or without external loading? *Scand. J. Med. Sci. Sports.* 2010;20 Suppl 2:70-7.
- 4: Noehren B, Davis I, Hamill J. ASB clinical biomechanics award winner 2006 prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clin. Biomech. (Bristol, Avon).* 2007;22(9):951-6.
- 5: Waryasz GR, McDermott AY. Patellofemoral pain syndrome (PFPS): a systematic review of anatomy and potential risk factors. *Dyn. Med.* 2008;7:9.
- 6: Jacobsen JS, Thorborg K, Søballe K, Ulrich-Vinther M. Eccentric hip abductor weakness in patients with symptomatic external snapping hip. *Scand. J. Med. Sci. Sports.* 2012;22(6):e140-6.