



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Progesterone, estradiol, arachidonic acid, oxytocin, forskolin and cAMP influence on aquaporin 1 and 5 expression in porcine uterine explants during the mid-luteal phase of the estrous cycle and luteolysis

an in vitro study

Skowronska, Agnieszka; Mlotkowska, Patrycja; Wojciechowicz, Bartosz; Okrasa, Stanislaw; Nielsen, Søren; Skowronski, Mariusz T

Published in:
Reproductive Biology and Endocrinology

DOI (link to publication from Publisher):
[10.1186/s12958-015-0004-5](https://doi.org/10.1186/s12958-015-0004-5)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Skowronska, A., Mlotkowska, P., Wojciechowicz, B., Okrasa, S., Nielsen, S., & Skowronski, M. T. (2015). Progesterone, estradiol, arachidonic acid, oxytocin, forskolin and cAMP influence on aquaporin 1 and 5 expression in porcine uterine explants during the mid-luteal phase of the estrous cycle and luteolysis: an in vitro study. *Reproductive Biology and Endocrinology*, 13, Article 7. <https://doi.org/10.1186/s12958-015-0004-5>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

RESEARCH

Open Access

Progesterone, estradiol, arachidonic acid, oxytocin, forskolin and cAMP influence on aquaporin 1 and 5 expression in porcine uterine explants during the mid-luteal phase of the estrous cycle and luteolysis: an *in vitro* study

Agnieszka Skowronska^{1*}, Patrycja Młotkowska², Bartosz Wojciechowicz², Stanisław Okrasa², Soren Nielsen³ and Mariusz T Skowronski^{2,4}

Abstract

Background: The cell membrane water channel protein, aquaporins (AQPs), regulate cellular water transport and cell volume and play a key role in water homeostasis. Recently, AQPs are considered as important players in the field of reproduction. In previous studies, we have established the presence of AQP1 and 5 in porcine uterus. Their expression at protein level altered in distinct tissues of the female reproductive system depending on the phase of the estrous cycle. However, the regulation of aquaporin genes and proteins expression has not been examined in porcine uterine tissue. Therefore, we have designed an *in vitro* experiment to explain whether steroid hormones, progesterone (P4) and estradiol (E2), and other factors: oxytocin (OT), arachidonic acid (AA; substrate for prostaglandins synthesis) as well as forskolin (FSK; adenylate cyclase activator) and cAMP (second messenger, cyclic adenosine monophosphate) may impact AQPs expression.

Methods: Uterine tissues were collected on Days 10–12 and 14–16 of the estrous cycle representing the mid-luteal phase and luteolysis. Real-time PCR and Western blot analysis were performed to examine the expression of porcine AQP1 and AQP5. Their expression in the uterine explants was also evaluated by immunohistochemistry.

Results: The results indicated that uterine expression of AQP1 and AQP5 potentially remains under control of steroid hormones and AA-derived compounds (e.g. prostaglandins). P₄, E₂, AA, FSK and cAMP cause translocation of AQP5 from apical to the basolateral plasma membrane of the epithelial cells, which might affect the transcellular water movement (through epithelial cells) between uterine lumen and blood vessels. The AC/cAMP pathway is involved in the intracellular signals transduction connected with the regulation of AQPs expression in the pig uterus.

Conclusions: This study documented specific patterns of AQP1 and AQP5 expression in response to P₄, E₂, AA, FSK and cAMP, thereby providing new indirect evidence of their role in maintaining the local fluid balance within the uterus during the mid-luteal phase of the estrous cycle and luteolysis in pigs.

Keywords: Aquaporins, Uterus, Estrous cycle, Pig, *in vitro* study

* Correspondence: agnieszka.skowronska@uwm.edu.pl

¹Department of Human Physiology, Faculty of Medical Sciences, University of Warmia and Mazury in Olsztyn, Olsztyn, Poland

Full list of author information is available at the end of the article

Background

Since the discovery of aquaporins (AQPs), water channel proteins, a high rate of transcellular water flow is believed to be mediated by these specialized protein transporters. Recently, AQPs have become considered to be important players in the field of reproduction, see reviews [1, 2]. Several AQP isoforms are expressed in the female reproductive tissues: ovary, uterus, placenta, amnion and chorion cytotrophoblasts [3–10]. Their specific expression pattern suggests that they participate in water movement between the intraluminal, interstitial and capillary compartments. Further studies have demonstrated that AQPs are also involved in endometrial development, cell migration and invasion [11]. Aquaporin 1 and 5 are water selective and belong to the classical AQP family [12]. AQP1 is a 28-kDa water channel protein expressed in the endothelial and epithelial cells of many tissues, increasing water permeability of the cell membrane. AQP5 has mainly been localized in apical plasma membranes of various secretory glands [13]. The important role of AQP5 in water homeostasis is evidenced by AQP5-null mice which have reduced saliva secretion [14].

The expression of AQPs in uterine tissue was first described by Li et al. [15], who confirmed the presence of AQP1 transcript in the human uterus. Afterwards, Li et al. [16] demonstrated that AQP1 mRNA expression in the rat uterus is up-regulated by estradiol. Accumulating evidence indicated that ovarian steroids can affect the expression of several AQPs in the reproductive system, including the uterus [17, 18]. The presence of AQP1, 2 and 5 has also been studied throughout the estrous cycle in bitches [19]. Very recently, Klein et al. [20] showed uterine mRNA expression of 12 different AQPs subtypes in endometrium of cyclic and pregnant mares. Moreover, it was demonstrated that cAMP is involved in up-regulation of some AQPs in a variety of cell types [9,21–24]. The presence of AQP1 in human endometrial blood vessels indicates its involvement in the regulation of edema, and in the regulation of angiogenesis [25] as well as pathological processes related to ovulatory uterine bleeding in women [26].

Studies on pigs suggest a functional collaboration among diverse AQPs within the uterus during different phases of the estrous cycle and early pregnancy [27]. It has been shown that AQP5 is localized in myometrial and epithelial cells of the uterus, but AQP1 in uterine endometrial and myometrial blood vessels [7, 27]. Their expression at a protein level was also altered in distinct tissues depending on the phase of the estrous cycle and the stages of early pregnancy. However, the regulation of aquaporin genes and protein expression has not been examined in porcine uterine tissue. Therefore, we have designed an *in vitro* experiment to explain whether steroid hormones, progesterone (P_4) and estradiol (E_2), and other factors: oxytocine

(OT), arachidonic acid (AA; substrate for prostaglandins synthesis) as well as forskolin (FSK; adenylate cyclase activator) and cAMP (cyclic adenosine monophosphate; second messenger) may have impact on the AQPs expression. Consequently, the primary aim of this study was; (i) to examine the changes in AQP1 and AQP5 at mRNA and protein levels in porcine uterine explants in the presence of P_4 , E_2 , OT, AA, FSK and cAMP; and then (ii) to describe the effect of tissue exposition duration to the experimental factors on the AQPs expression; (iii) to compare their expression in the uterine explants, representing the mild-luteal phase of the estrous cycle and luteolysis; (iv) to determine the localization of AQP1 and 5 in uterine explants after the treatments. This study provided additional information concerning factors potentially responsible for water homeostasis in porcine uterus during the mid-luteal phase and luteolysis in the pig.

Methods

Experimental animals and collection of uterine tissue

All experiments were performed in accordance with Animal Ethics Committee, University of Warmia and Mazury in Olsztyn, Poland (AEC approval No. 66/2010/DTN). Tissue samples were recovered from mature cross-bred gilts (Large White × Polish Landrace) on Days 10–12 ($n = 5$ per group) during the mid-luteal phase and on Days 14–16 ($n = 5$ per group) of the estrous cycle (the stage of luteolysis). Gilts were observed daily for estrous behavior, and they were used in the study during their third consecutive normal estrous cycle. The animals were slaughtered at a local abattoir on Days 10–12, corresponding to the period with increased plasma progesterone (P_4) concentration, or Days 14–16, the period of decreasing plasma P_4 concentration and luteal regression, of the estrous cycle. Additionally, stage of the cycle was verified by utero-ovarian morphology [28]. Uteri were placed immediately in ice-cold phosphate-buffered saline (PBS) supplemented with 100 IU/ml penicillin (Polfa, Poland) and 100 μ g/ml streptomycin (Polfa, Poland) and transported to the laboratory on ice within 1 to 1.5 h for *in vitro* tissue culture.

Preparation and incubation of uterine slices

Sections of the middle part of uterine horn collected from pigs were opened longitudinally on the mesometrial surface. Uteri were washed three times in sterile PBS then carefully cut into small pieces (400 mg weight) and then washed three times in medium M199 (Sigma, USA). Individual uterine slices were placed in culture vials containing 2 ml Medium 199 supplemented with 0.1% BSA (Sigma), 20 μ g nystatin (Sigma) and 20 μ g gentamicin (Krka, Novo Mesto, Slovenia) and then preincubated *in vitro* under atmosphere of 95% O_2 and 5% CO_2 at 37°C for 18 h. After preincubation, the culture

medium was replaced with fresh medium, and the explants were treated with vehicle (control) or P_4 (10^{-5} M; Sigma), E_2 (10^{-9} M; Sigma), OT (10^{-7} M; Sigma), AA (10^{-5} M; Sigma), FSK (10 μ g/mL; Sigma) and cpt-cAMP analog (200 μ M; Sigma) and incubated for an additional 3 or 24 h. All treatments were performed in triplicates. Furthermore, uterine tissue explants were snap-frozen in liquid nitrogen (for RNA and protein extraction) and stored at -80°C until further use.

Total RNA isolation, cDNA synthesis and quantitative real-time polymerase chain reaction analysis

Total RNA was extracted, using the total RNA Prep Plus kit (A&A Biotechnology, Gdansk Poland) according to the manufacturer's protocol, from uterine explants collected after *in vitro* culture. Total RNA quality and quantity were determined with spectrophotometry (NanoDrop ND-1000, Thermo Scientific, Wilmington, DE, USA). Total RNA samples were transcribed to cDNA using an Enhanced Avian HS RT-PCR Kit (Sigma) and a mix of dNTPs and random hexamers as primers. Real-Time PCR was performed in duplicate for each sample using a 7300 Real-Time PCR system and SYBR[®]Green PCR Master Mix (Life Technologies, Grand Island, NY, USA). Real-Time PCR reaction included 12.5 μ l SYBR Green PCR master mix, 1 μ M forward and reverse primers each and reverse transcribed cDNA (3.5 μ l of diluted RT product) supplemented with water to a volume of 25 μ l. The conditions of the thermal cycling for each gene were: initial denaturation for 10 min at 95°C , denaturation for 15 sec at 95°C , primer annealing for 1 min at 60°C . Specific primers for *AQP1* and *AQP5* (Table 1) were designed with the Primer Express 3.0 software (Life Technologies) and their specificities were confirmed by comparison of their sequences with the sequence of *AQP1* and *AQP5* deposited in a database and calculation of the statistical significance of the match was performed using the Basic Local Alignment Search Tool (BLAST). For the specificity control, non-template controls and dissociation curve analysis of the amplified products were used for each amplification.

Table 1 Primer pairs used in the study

PCR product	Sequence
<i>AQP1</i>	Forward: 5'-CCAGCGAGTTCAAGAAGAAG-3' Reverse: 5'-GCGACACCTTCACGTTATC-3'
<i>AQP5</i>	Forward: 5'-CTATGAGTCCGAGGAGGATT-3' Reverse: 5'-GCTTCGCTGTCATCTGT-3'
<i>18S rRNA</i>	Forward: 5'-GGCTACCACATCCAAGGAAG-3' Reverse: 5'-TCCAATGGATCCTCGCGAA-3'
<i>GADPH</i>	Forward: 5'-GACCTCCACTACATGGTCTA-3' Reverse: 5'-AAGATGGTGATGGCCTTC-3'

The specificity of amplifications was further validated with electrophoresis of the putative amplicons in a 2% agarose gel and, after extraction from gel, automated sequencing using 3730xl DNA Analyzer (Life Technologies). Levels of gene expression were calculated using the $\Delta\Delta$ Ct method and normalized using the geometrical means of reference genes expression levels, glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) and *18S rRNA*.

SDS-PAGE and Western blot

The tissues were placed in ice-cold dissection buffer (0.3 M sucrose, 25 mM imidazol, 1 mM EDTA in ddH₂O, pH 7.2) containing 8.4 μ M leupeptin and 0.4 mM pefabloc [29]. The tissue samples were homogenized using an ultra Turrax T8 homogeniser (IKA Labor Technik, Staufen, Germany) and centrifuged at $4,000 \times g$ for 15 min at 4°C . The supernatant was diluted in SDS buffer contained a final concentration of 62 mM Tris (hydroxymethyl)-aminomethane, 0.1 M sodium dodecyl sulphate (SDS), 8.7% glycerol, 0.09 mM bromophenol blue and 0.04 M dithiothreitol (DTT), pH 6.8. The protein samples were heated for 5 min at 90°C and stored in refrigerator for further analysis. Total protein amounts were determined with spectrophotometry (NanoDrop ND-1000, Thermo Scientific, Wilmington, DE, USA). The samples warmed up to 37°C were loaded into 12.5% polyacrylamide gels and proteins were separated by electrophoresis. The proteins of studied gels were then electro-transferred onto nitrocellulose membranes (Hybond ECL RPN3032D, Amersham Pharmacia Biotech, Little Chalfont, UK) for 1 h at 100 V. The membranes were blocked with 5% milk in PBS-T (80 mM Na₂HPO₄, 20 mM NaH₂PO₄, 100 mM NaCl, pH 7.5 and 0.1% vol/vol Tween 20) for 1 h. After washing, the membranes were incubated overnight at 5°C with anti-AQPs or β -actin antibodies. Thereafter, the membranes were washed and incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG secondary antibody (P448, diluted 1:3,000, Dako A/S, Glostrup, Denmark) in PBS-T for 1 h. After washing with PBS-T, the sites of antibody-antigen reaction were visualized with an enhanced chemiluminescence (ECL) system (Amersham Pharmacia Biotech, Little Chalfont, UK) and exposure to photographic film (Hyperfilm ECL, RPN3103K, Amersham Pharmacia Biotech, Little Chalfont, UK). The results of Western blotting were quantified by densitometric scanning of immunoblots with GelScan for Windows ver. 1.45 software (Kucharczyk, Poland). Data were expressed as a ratio of AQP proteins relative to actin protein in OD units.

Immunohistochemistry

Tissues were fixed by immersion in 4% paraformaldehyde for 24 hr. For preparation of paraffin-embedded tissue

sections (4 μm thickness), the tissues were dehydrated in ethanol followed by xylene and finally embedded in paraffin [27]. The staining was carried out using indirect immunoperoxidase. The sections were dewaxed and rehydrated. For immunoperoxidase labeling, endogenous peroxidase was blocked by 0.5% H_2O_2 in absolute methanol for 10 min at room temperature. To reveal antigens, the sections were submerged in 1 mM Tris solution (pH 9.0) supplemented with 0.5 mM EGTA and heated in a microwave oven. After the treatment, the sections were left for 30 min in the buffer for cooling. Nonspecific binding of IgG was eliminated by incubating the sections in 50 mM NH_4Cl for 30 min, followed by blocking in PBS supplemented with 1% BSA, 0.05% saponin and 0.2% gelatin. The sections were incubated overnight at 4°C with primary antibodies diluted in PBS supplemented with 0.1% BSA and 0.3% Triton X-100. The sections were rinsed with PBS supplemented with 0.1% BSA, 0.05% saponin and 0.2% gelatin, and then incubated with horseradish peroxidase-conjugated secondary antibody (Dako A/S, Glostrup, Denmark). Labeling was visualized by 0.05% 3,3 diaminobenzidine tetrahydrochloride (DAB). The microscopy was carried out using an Olympus light microscope (BX51, Japan).

Primary antibodies

Antibodies to AQP1 and AQP5, used in Western blot analysis and immunocytochemistry, were previously characterized, respectively by Terris et al. [30] and Nielsen et al. [13]. All polyclonal antibodies were affinity-purified (SulfoLink Kit, Pierce, Rockford, IL). Moreover, the anti- β -actin antibody was used (cat. no. A2066; Sigma-Aldrich, St Louis, MO). In our previous study, we demonstrated that anti-AQP1 and anti-AQP5 antibodies preincubated with the immunizing peptide prevented labeling in the pig uterus [7]. In addition, immunoglobulins from non-immunized rabbit were used as a negative control.

Statistical analysis

All numerical data were analyzed by one-way ANOVA and least significant difference (LSD) post hoc test and reported as the means \pm S.E.M. from five separate experiments (pigs), each performed in duplicates. Statistical analyses were performed using the Statistica program (StatSoft Inc., Tulsa, USA). Values for $p < 0.05$ were considered statistically significant.

Results

AQP1 mRNA expression in porcine uterine explants

The control abundance of AQP1 transcript harvested in the uterine tissue from mid-luteal phase (Days 10–12) was about 4-fold and 2-fold higher ($p < 0.05$) than on Days 14–16 (luteolysis) during 3-h and 24-h incubation, respectively (Figure 1A and B). In porcine uterine

explants from Days 10–12 (the mid-luteal phase) of the estrous cycle, AQP1 mRNA expression significantly ($p < 0.05$) decreased after 3-h treatment with E_2 and AA, in comparison to the respective controls. In turn, longer treatment (24 h) with FSK significantly ($p < 0.05$) increased AQP1 mRNA expression, but with P_4 , oxytocin and AA it decreased.

Aquaporin 1 gene expression in porcine uterine tissue explants from Days 14–16 (luteolysis), significantly ($p < 0.05$) increased after both 3- and 24-h treatment with E_2 (Figure 1B). In contrast, AQP1 mRNA expression decreased after 3-h treatments with P_4 and OT ($p < 0.05$). Arachidonic acid, FSK and cAMP did not affect AQP1 mRNA expression during shorter incubation. In turn, an increase in AQP1 mRNA expression was seen after 24-h treatment with FSK and cAMP ($p < 0.05$). A particularly strong increase (~7-fold) in AQP1 mRNA expression was noted after 24-h treatment with FSK.

AQP5 mRNA expression in uterine explants

Quantitative expression of AQP5 mRNA in the uterine explants on Days 10–12 and 14–16 of the estrous cycle is presented on Figure 2A and B, respectively. The content of AQP5 transcript in uterine explants following control incubation was about 2.5 fold lower after 3-h incubation ($p < 0.05$) on Days 10–12 of the estrous cycle in comparison to Days 14–16. Such a difference was not observed in the case of longer incubation (24 h). During the mid-luteal phase of the cycle, markedly lower amounts of AQP5 transcript were noted after treatments with P_4 , E_2 , oxytocin, AA, FSK and cAMP for 3 h as compared to control value ($p < 0.05$) and remained lower after 24 hours ($p < 0.05$) except for treatment with cAMP.

Treatment with E_2 for 3 h of the uterine explants representing luteolysis (Days 14–16) of the estrous cycle significantly increased ($p < 0.05$) AQP5 mRNA expression ($p < 0.05$) but OT and FSK treatments significantly decreased it. Incubation with P_4 , E_2 , OT and AA did not result in any detectable change in porcine uterine explants AQP5 gene expression after 24-hours incubation. Treatment with cAMP for 24 h significantly decreased AQP5 gene expression. Conversely, incubation with FSK after 24 h, significantly increased AQP5 gene expression.

Protein content of AQP1 in uterine explants

The effects of P_4 , E_2 , oxytocin, AA, FSK and cAMP on AQP1 protein expression in uterine tissue explants harvested during the mid-luteal phase (Days 10–12) and luteolysis (Days 14–16) are shown in Figure 3A and B. The protein expression of AQP1 in porcine uterine explants increased ($p < 0.05$) after treatment with P_4 after 3- and 24-h incubation. Estradiol treatment for 3 h did not significantly affect AQP1 protein expression in the tissue from mid-luteal phase, however, after longer




Figure 1 *AQP1* mRNA expression in porcine uterine explants. The expression of *AQP1* mRNA in the pig uterine tissue harvested on (A) Days 10–12 ($n = 5$) and (B) Days 14–16 ($n = 5$) of the estrous cycle after treatment with progesterone (P_4 ; 10^{-5} M), estradiol (E_2 ; 10^{-9} M), oxytocin (OT; 10^{-7} M), arachidonic acid (AA; 10^{-5} M), forskolin (FSK; $10 \mu\text{g/mL}$) and cyclic-AMP (cAMP; $200 \mu\text{M}$) for 3 and 24 h. The data are presented as the mean values \pm SEM of *AQP1* expression in relation to expression of *GAPDH* and *18 sRNA*. Different letters (a,b,c,d) indicate significant differences ($p < 0.05$) between each treatment and respective control for 3- (a,b) or 24-h (c,d) incubations.

incubation (24 h) E_2 stimulated its expression ($p < 0.05$). *AQP1* protein expression significantly increased after 3-h treatment with FSK and cAMP. In turn, treatment with OT and AA did not influence *AQP1* expression. During luteolysis, the content of *AQP1* was increased (from 2.5- to 3-fold; $p < 0.05$) by P_4 , E_2 , AA, FSK and cAMP after 3-h incubation (Figure 3A). After 24-h incubation, only P_4 and E_2 significantly stimulated *AQP1* expression ($p < 0.05$), similar to the levels noted after 3-h incubation (Figure 3A). It is noteworthy that the responses of *AQP1* to E_2 , AA and forskolin, after short incubation, appeared to be higher on Days 14–16 than on Days 10–12 of the estrous cycle.

Protein content of *AQP5* in uterine explants

AQP5 expression evaluated by Western blot analysis of in porcine uterus explants from the mid-luteal phase and the luteolysis is shown in Figure 4A and B. Bands of *AQP5* protein product of the expected size (28 kDa) were clearly detected in both stages of the cycle. During the mid-luteal phase, uterine *AQP5* protein expression

significantly ($p < 0.05$) increased after 3-h and 24-h of treatment with P_4 , E_2 , FSK and cAMP. During luteolysis, a high increase (~ 3.0 -fold) in *AQP5* protein expression in the tissue was observed in response to P_4 , E_2 , FSK and cAMP after both 3-h and 24-h incubations ($p < 0.05$). *AQP5* protein content in the uterine tissue was only elevated (~ 2.5 -fold) by 3-h treatment with AA ($p < 0.05$). In turn, treatment with OT did not influence *AQP5* expression in the studied tissue. It is noteworthy that *AQP5* protein response to some of the studied factors appeared to be higher on Days 14–16 than on Days 10–12 of the estrous cycle. In particular, during the luteolysis a higher *AQP5* protein expression was observed after 3-h treatment with P_4 , E_2 , FSK and cAMP and remained elevated for 24 hours in comparison to the mid-luteal phase.

Immunohistochemical localization of *AQP1* and *AQP5* in uterine tissue explants

In the tissue sections of the pig uterine explants representing the mid-luteal phase and luteolysis, immunoperoxidase labeling for *AQP1* was associated with uterine




Figure 2 *AQP5* mRNA expression in porcine uterine explants. The expression of *AQP5* mRNA in the pig uterine tissue harvested on (A) Days 10–12 ($n = 5$) and (B) Days 14–16 ($n = 5$) of the estrous cycle after treatment with progesterone (P_4 ; 10^{-5} M), estradiol (E_2 ; 10^{-9} M), oxytocin (OT; 10^{-7} M), arachidonic acid (AA; 10^{-5} M), forskolin (FSK; $10 \mu\text{g/mL}$) and cyclic-AMP (cAMP; $200 \mu\text{M}$) for 3 and 24 h. The data are presented as the mean values \pm SEM of *AQP5* expression in relation to expression of *GAPDH* and *18 sRNA*. Different letters (a,b,c,d) indicate significant differences ($p < 0.05$) between each treatment and respective control for 3- (a,b) or 24-h (c,d) incubations.




Figure 3 Protein content of AQP1 in uterine explants. Semi-quantitative Western blot analysis of AQP1 in homogenates of the pig uterine explants from (A) Days 10–12 (n = 5) and (B) Days 14–16 (n = 5) of the estrous cycle (each lane represents a sample from 1 pig), after treatment with progesterone (P₄; 10⁻⁵ M), estradiol (E₂; 10⁻⁹ M), oxytocin (OT; 10⁻⁷ M), arachidonic acid (AA; 10⁻⁵ M), forskolin (FSK; 10 μg/mL), and cyclic-AMP (cAMP; 200 μM). Densitometric analysis of the immunoblots was performed and different small letters (a, b,c,d) indicate significant differences (p < 0.05) between each treatment and respective control for 3- (a,b) or 24-h (c,d) incubations. Different large letters (A, B, C, D) indicate significant differences between the same treatments for 3- (A,B) or 24-h (C,D) incubations for different periods (Days 10–12 and 14–16) of the estrous cycle (p < 0.05).




Figure 4 Protein content of AQP5 in uterine explants. Semi-quantitative Western blot analysis of AQP5 in homogenates of the pig uterine explants from (A) Days 10–12 (n = 5) and (B) Days 14–16 (n = 5) of the estrous cycle (each lane represents a sample from 1 pig), after treatment with progesterone (P₄; 10⁻⁵ M), estradiol (E₂; 10⁻⁹ M), oxytocin (OT; 10⁻⁷ M), arachidonic acid (AA; 10⁻⁵ M), forskolin (FSK; 10 μg/mL) and cyclic-AMP (cAMP; 200 μM). Different small letters (a, b, c, d) indicate significant differences (p < 0.05) between each treatment and respective control for 3- (a,b) or 24-h (c,d) incubations. Different large letters (A, B, C, D) indicate significant differences between the same treatments for 3- (A,B) or 24-h (C,D) incubations for different periods (Days 10–12 and 14–16) of the estrous cycle (p < 0.05).

endothelial cells (Figure 5A and B). Both apical and basal plasma membranes exhibited stable AQP1 labeling -without intracellular changes in localization - after 3-h and 24-h incubation with all examined factors (Figure 5A and B). However, expression of AQP1 increased after 3- and 24-h tissue exposition to P_4 and E_2 on Days 10–12 and 14–16 as well as after 3-h incubation with FSK and cAMP on Days 10–12, and AA, FSK and cAMP on Days 14–16 of the estrous cycle. As a positive control, AQP1 labeling was seen in the apical and basolateral plasma membrane in the proximal tubule cells of the pig kidney (Figure 5Y).

In turn, in sections of the explants treated with examined factors, immunoperoxidase staining for AQP5 was associated with uterine epithelial (Figure 5C) and smooth muscle cells (Figure 5U and W). Moreover, AQP5 labeling in the uterine epithelial cells was more intensive in the apical plasma membrane after 3- and 24-h treatment with OT, AA on Days 10–12 and OT as well as 24-h treatment with AA on Days 14–16 than those observed in respective controls. Following 3- and 24-h treatments of the tissue with P_4 , E_2 , FSK and cAMP on Days 10–12 and 14–16 of the estrous cycle, prominent AQP5 labeling was seen in both the apical and basolateral plasma membranes of the epithelial cells (Figure 5D-S). A similar pattern of AQP5 labeling was seen after 3-h treatment with AA on Days 14–16 of the estrous cycle (Figure 5T). In the smooth muscle cells, in contrast to the epithelial cells, changes in AQP5 localization within the cell membranes were not observed in response to applied treatments (Figure 5U-W). As a positive control, AQP5 antibody noticeably stained (Figure 5Z) the apical plasma membrane of the type I pulmonary epithelial cells of the pig.

Discussion

In the present study, we have demonstrated that two isoforms of AQP5 (*1* and *5*) mRNAs are expressed in the porcine uterine explants in both studied periods. In our previous study, performed with the use of immunohistochemistry and Western blot, AQP1 and 5 were clearly detected in untreated uterine tissue on Days 10–12 and 14–16 of the estrous cycle. In cyclic gilts, endometrial and myometrial expression of AQP1 did not change significantly, in turn endometrial AQP5 protein expression was significantly higher on Days 14–16 than on Days 10–12 of the cycle [7]. In the present study, we aimed to further investigate, using an *in vitro* system, whether these AQPs are regulated by steroid hormones, cAMP, forskolin, arachidonic acid or oxytocin in porcine uterine tissue at mRNA and protein levels.

In the present study, steroid hormones (P_4 and E_2) differentially influenced the expression of AQP1 and 5 in porcine uterine tissue under *in vitro* conditions. *AQP1* gene expression was down-regulated by E_2 treatment for

3-h during the mid-luteal phase. Furthermore, P_4 inhibited *AQP1* mRNA gene expression during luteolysis, after short incubation (3 h). Steroid hormones (P_4 and E_2) increased AQP1 protein expression during both studied stages of the cycle and incubations. Moreover, E_2 more effectively stimulated AQP1 protein expression on Days 14–16 than on Days 10–12 of the cycle. In turn, *AQP5* gene expression was also down-regulated after treatments with E_2 and P_4 (3 h and 24 h) during the mid-luteal phase, but up-regulated by E_2 during luteolysis (3 h). As in the case of *AQP1*, longer incubation with steroids down-regulated *AQP5* mRNA expression during the mid-luteal phase. Interestingly, the basic (control) level of *AQP1* transcript was higher (after 3- and 24-h incubations) in the mid-luteal phase (2.07 ± 0.61 and 1.03 ± 0.09 , respectively) than during luteolysis, but in the case of *AQP5* (after 3-h incubation) the relationship was inverted (1.59 ± 0.35 and 4.48 ± 0.71 , respectively). At the protein level, the changes of AQP1 and 5 expression in response to E_2 and P_4 , were not in full agreement with those noted for mRNAs.

Our findings concerning the E_2 effect on AQP expression are comparable to those reported by Li et al. [16] and Kobayashi et al. [31], who studied this steroid action on the uterine expression of AQP1 in rat and AQP5 in mouse, respectively. In addition, Kobayashi et al. [31] revealed the presence of functional estrogen response element in AQP5 promoter regions which suggests the possibility of direct action of estrogens on AQP5 expression. Richard et al. [5] found increased *AQP1* mRNA expression in mice myometrium in response to estrogen, but AQP5 expression was induced by estrogen only in progesterone-primed uteri. In several studies, the role of P_4 alone or in combination with estrogens in controlling AQP expression in uterus has been confirmed. Lindsay and Murphy [6] reported increased expression of AQP5 in uterine epithelial cells by progesterone alone and in combination with estrogen. The same authors [18] also noted progesterone-dependent expression of AQP5 and AQP1 in the rat uterus; AQP5 in glandular epithelium and AQP1 in the inner circular layer of myometrium. Aralla et al. [19] demonstrated coincidently elevated expression of AQP5 in the apical plasma membrane of uterine epithelial cells with increased concentrations of P_4 in plasma. Furthermore, in the mouse uterus exogenous estrogen strongly up-regulated the expression of AQP2, without any effect on AQP5 [17]. Very recently, the presence of mRNA for all AQPs (*AQP0* to *12*) has been shown in equine endometrium, while the Western blot analysis confirmed protein expression of AQP0, 2 and 5 [20]. To the contrary, progesterone treatment of anoestrus mares did not enhance the expression of AQPs, indicating that factors other than progesterone or some factors in addition to progesterone are required for




Figure 5 (See legend on next page.)

(See figure on previous page.)

Figure 5 Immunohistochemical localization of AQP1 and AQP5 in uterine tissue explants. Immunoperoxidase staining of AQP1 in paraffin-embedded sections of the uterine explants from pigs. Anti-AQP1 antibody labels endothelial cells of the uterine explants (arrows). Both apical and basal plasma membranes exhibited stable AQP1 labeling (**A/** an example of the staining for the control and **B/** from the E_2 -treated uterine explants for 24-h on Days 14–16 of the cycle). Immunoperoxidase labeling of AQP1 in the pig kidney cortex (**Y/** positive control). The labeling is seen in both of the apical and basolateral plasma membrane in proximal tubule cells. AQP5 antibody stains epithelial cells of the uterine explants (arrows). The labeling is seen only in the apical plasma membranes of the epithelial cells (**C/** an example of the staining for the control on Days 14–16 of the cycle). Continuously, 3- and 24-h treatments on Days 10–12 and 14–16 of the estrous cycle of the tissue explants with P_4 (**D-E and F-G**), E_2 (**H-I and J-K**), FSK (**L-M and N-O**) and cAMP (**P-Q and R-S**), respectively, prominent AQP5 labeling is seen in both the apical and basolateral plasma membranes of the epithelial cells. A similar pattern of AQP5 labeling is also seen after 3-h treatment with AA on Days 14–16 of the estrous cycle (**T**). AQP5 antibody also stains smooth muscle cells (**U/** an example of the staining for the control and **W/** from the E_2 -treated uterine explants for 24-h on Days 14–16 of the cycle). The anti-AQP5 labels apical membrane of type I pulmonary epithelial cells of the pig (**Z/** positive control). No staining was observed with using non-immune immunoglobulins (negative controls: **X1** – AQP1 and **X5** – AQP5 controls). L – lumen. Bar = 50 μ m.

the up-regulation of certain AQP subtypes. The above observations are not entirely consistent, the discrepancies seem to result from different animal models and/or experimental protocols applied in the study. For example, Jablonski et al. [17] used ovariectomized and hormonally treated mice, but in the present study porcine uterine explants were used. In turn, the lack of a full relationship – observed in the present study – between the concentration of gene transcripts and respective proteins may result from differentiated stability of mRNAs and/or proteins as well as from independent regulation of transcription, posttranscriptional processes or translation and functioning feedbacks, *i.e.* high protein concentration may suppress mRNA synthesis. Moreover, steroids themselves may alter the stability of mRNA [32]. It is also noteworthy that the processes of transcription and translation are not equally efficient.

The uterus is a target organ for ovarian steroid hormones and undergoes marked changes during the estrous cycle, including: tissue expansion (by 40–60%) and an increase in uterine gland activity during luteal phase [33], hyperemia [34] as well as increased capillary permeability [33]. These changes require increased synthesis of AQPs. In the present study, as documented by Western blot analysis, P_4 and E_2 appeared to be effective in controlling AQP1 and AQP5 expression in pigs, suggesting their crucial role in this regulation during the mid-luteal phase of the estrous cycle and luteolysis.

Arachidonic acid is metabolized in the uterus to prostaglandins (PGE_2 and $PGF_{2\alpha}$), which are involved in many reproductive activities including luteolysis, maternal recognition of pregnancy, endometrial gene expression and conceptus development [35–37]. Prostaglandin synthesis is thus dependent upon accessibility of AA [38] and the activity of enzymes involved in its metabolism [39]. In our studies, its potential effect on AQP1 and 5 expression in uterine tissue of cyclic pigs was tested. The expression of AQP1 and 5 mRNAs were similarly down-regulated by AA during the mid-luteal phase (after 3- and 24-h exposition). Nevertheless, the expression of both AQPs at the protein level was stimulated only during luteolysis in

response to shorter exposition to AA. This well corresponds with physiological situation, since $PGF_{2\alpha}$ is released by endometrium (epithelial and stromal cells) [40] in a pulsatile manner to cause corpus luteum regression [41]. Thus, our data indirectly indicate that prostaglandins may exert regulatory effects on AQP1 and 5 in uterine tissue. Studies performed by Zelenina et al. [42] have confirmed the interaction of PGE_2 with AVP in the regulation of AQP2 in the rat renal medulla. Nevertheless, the potential involvement of prostaglandins in the regulation of AQPs in the pig uterus requires explanation in further experiments.

Oxytocin is one of the key hormones implicated in controlling the uterine functions. Among others, it affects phosphoinositide hydrolysis [43], expression of COX-2 in uterine tissue [37] and regulates $PGF_{2\alpha}$ secretion [43]. The main reason for investigating the effect of oxytocin on AQP1 and AQP5 expression was that pig endometrium secretes oxytocin [44] and possesses its receptors [45]. In the uterus, the concentration of OT receptor in the endometrium changes during the estrous cycle. It has been reported that in the middle luteal phase of the cycle, the density of OT receptor increases in the endometrium and myometrium [46]. In the present experiments, OT decreased AQP1 and AQP5 mRNA expression (after 3- and 24-h incubations), without visible changes in protein content. In the pig during luteolytic period, oxytocin is responsible for pulsatile secretion of $PGF_{2\alpha}$ and contractions of myometrium [47]. It might be hypothesized that under physiological conditions, inhibitory action of OT on AQP1 and 5 uterine expression at a transcriptional level is connected with remodeling of endometrial tissues taking place at the end of the luteal phase [33]. Very recently, Ducza et al. [48] demonstrated increased AQP2 mRNA, but reduced AQP5 mRNA expression by OT in the rat uterus on Day 18 of pregnancy. Since the role of oxytocin, similar to PGs, in the regulation of AQPs expression and uterine fluid balance, so far is not sufficiently defined, therefore further work is needed to better delineate it.

Previous studies confirmed that intracellular signaling pathway consisting of adenylate cyclase (AC) and cAMP

may be involved in the regulation of AQP expression [21–23]. In the present study, the effects of cAMP and forskolin (AC activator) on AQP1 and 5 in the uterine tissue were tested. The treatment with cAMP did not cause striking changes in expression of studied AQP mRNAs in the uterine tissue; only increased *AQP1* expression on Days 14–16 (after 24 h) and decreased *AQP5* on Days 10–12 (after 3 h) and Days 14–16 (after 24 h) of the estrous cycle. In a different experimental model, Wang et al. [22] failed to see any changes in the expression of *AQP1*, 8 or 9 genes in human amnion-derived WISH cells after *in vitro* treatment with cAMP analog. In our studies, the opposite reaction of *AQP5* mRNA versus protein in response to cAMP is complicated. This discrepancy may result from the regulatory mechanism functioning at the transcription and translation levels, as discussed earlier and requires an explanation in further studies. The effects of forskolin on *AQP1* and 5 mRNAs were very similar to those observed in response to cAMP; i.e. stimulation of *AQP1* and mostly inhibition of *AQP5* (except stimulation on Day 14–16 after 24 h incubation). At the protein level, the expression of AQP1 and AQP5 was up-regulated by forskolin, as in response to cAMP. Studies performed with the use of different cells/tissues [9, 22, 23, 49] or cell lines [21, 23, 24] also confirmed the stimulatory effect of cAMP and/or forskolin on the expression of AQP1 [24], AQP5 [23] and AQP3, 8 and 9 [9, 21, 23, 49]. Furthermore, AQP1 appeared to be up-regulated by arginine vasopressin and cAMP analogue in trophoblast cells [24]. On the basis of our study, it might be concluded that the AC/cAMP pathway participates in the regulation of AQP5 expression in the uterine tissue. However, the engagement of the AC/cAMP pathway, particularly in the regulation of AQP1 expression in this tissue during the estrous cycle remains to be elucidated.

The changes in cellular localization of AQP1 and AQP5 in response to the studied factors, visualized by immunohistochemistry, are particularly interesting (Figure 5). In the uterine tissue, localization of AQP1 was predominantly associated with apical and basal membranes of endothelial cells, but AQP5 with apical membranes of epithelial cells. It is noteworthy that steroid hormones (P_4 and E_2), cAMP and forskolin caused an emergence of AQP5 in basolateral membrane of the epithelial cells during both studied periods. It might be thus hypothesized, that these changes are connected with potentially bidirectional transcellular water movement through uterine epithelial cells. Garcia et al. [50] indicated that cAMP may induce insertion of AQP8 within intracellular vesicular structures and translocation to plasma membranes in the rat hepatocytes. Further experiments are necessary to further explain the role of AQPs in uterine water balance in cyclic gilts.

Conclusions

The present studies delineated the specific AQP1 and AQP5 expression patterns in response to P_4 , E_2 , AA, FSK and cAMP and indirectly provided novel evidence for the role of AQP1 and AQP5 in maintaining local fluid balance within the uterus during mid-luteal phase of the estrous cycle and luteolysis in pigs. They revealed that: (1) steroid hormones (P_4 and E_2), oxytocin and presumably AA metabolites (prostaglandins) exhibit potential for regulation of AQP1 and 5 expression in porcine uterine tissue during the luteal phase; (2) AQPs responses to the studied factors depended on the stage of luteal phase and duration of their action; (3) P_4 , E_2 , AA, forskolin and cAMP cause translocation of AQP5 from apical to the basolateral plasma membrane of the epithelial cells, which might affect the transcellular water movement (through epithelial cells) between uterine lumen and blood vessels; (4) the AC/cAMP pathway is involved in the intracellular signals transduction connected with the regulation of AQPs expression in the pig uterus.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MTS and AS conceived and designed the experiments. AS, PM and MTS performed the experiments. SN, BW, AS, MTS and SO contributed reagents/materials/analysis tools. AS wrote this manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was support by Grant Number N N308 5848 40 awarded to MTS from the Polish Ministry of Science and Higher Education. This article was edited by a distinguished professional service.

Author details

¹Department of Human Physiology, Faculty of Medical Sciences, University of Warmia and Mazury in Olsztyn, Olsztyn, Poland. ²Department of Animal Physiology, Faculty of Biology and Biotechnology, University of Warmia and Mazury in Olsztyn, Olsztyn, Poland. ³Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark. ⁴Institute of Veterinary, Poznań University of Life Sciences, Poznań, Poland.

Received: 19 August 2014 Accepted: 7 February 2015

Published online: 18 February 2015

References

- Zhang D, Tan YJ, Qu F, Sheng JZ, Huang HF. Functions of water channels in male and female reproductive systems. *Mol Asp Medic*. 2012;33:676–90.
- Hua Y, Jiang W, Zhang W, Shen Q, Chen M, Zhu X. Expression and significance of aquaporins during pregnancy. *Front Biosc*. 2013;18:1373–83.
- Meng QX, Gao HJ, Xu CM, Dong MY, Sheng X, Sheng J-Z, et al. Reduced expression and function of aquaporin-3 in mouse metaphase-II oocytes induced by controlled ovarian hyperstimulation were associated with subsequent low fertilization rate. *Cell Physiol Biochem*. 2008;21:123–8.
- McConnell NA, Yunus RS, Gross SA, Bost KL, Clemens MG, Hughes Jr FM. Water permeability of an ovarian antral follicle is predominantly transcellular and mediated by aquaporins. *Endocrinology*. 2002;143:2905–12.
- Richard C, Gao JU, Brown N, Reese J. Aquaporin water channel genes are differentially expressed and regulated by ovarian steroids during the periimplantation period in the mouse. *Endocrinology*. 2003;144:1533–41.
- Lindsay LA, Murphy CR. Redistribution of aquaporins in uterine epithelial cells at the time of implantation in the rat. *Acta Histochem*. 2004;106:299–307.

7. Skowronski MT. Distribution and quantitative changes in amounts of aquaporin 1, 5 and 9 in the pig uterus during the estrous cycle and early pregnancy. *Reprod Biol Endocrinol*. 2010;8:109.
8. Damiano AE, Zotta E, Goldstein J, Reisin I, Ibarra C. Water channel proteins AQP3 and AQP9 are present in syncytiotrophoblast of human term placenta. *Placenta*. 2001;22:776–81.
9. Wang S, Amidi F, Beall M, Gui L, Ross MG. Aquaporin 3 expression in human fetal membranes and its up-regulation by cyclic adenosine monophosphate in amnion epithelial cell culture. *J Soc Gynecol Investig*. 2006;13:181–5.
10. Zhu XQ, Jiang SS, Zhu XJ, Zou SW, Wang Y, Hu YC. Expression of aquaporin 1 and aquaporin 3 in fetal membranes and placenta in human term pregnancies with oligohydramnios. *Placenta*. 2009;30:670–6.
11. Zou LB, Zhang RJ, Tan YJ, Ding GI, Shi S, Zhang D, et al. Identification of estrogen response element in the aquaporin -2 gene that mediates estrogen-induced cell migration and invasion in human endometrial carcinoma. *J Clin Endocrinol Metab*. 2011;96:E1399–408.
12. Agre P, King L, Yasui M, Guggino WB, Ottersen OP, Fujiyoshi Y, et al. Aquaporin water channels – from atomic structure to clinical medicine. *J Physiol*. 2002;542:3–16.
13. Nielsen S, King LS, Christensen BM, Agre P. Aquaporins in complex tissue. II. Subcellular distribution in respiratory and glandular tissues of rat. *Am J Physiol*. 1997;273:C1549–61.
14. Verkman AS. More than just water channels: unexpected cellular roles of aquaporins. *J Cell Sci*. 2005;118:3225–32.
15. Li X, Yu H, Koide SS. The water channel gene in human uterus. *Biochem Mol Biol Int*. 1994;32:371–7.
16. Li XJ, Yu HM, Koide SS. Regulation of water channel gene (AQP-CHIP) expression by estradiol and androstadiol in rat uterus. *Yao Xue Xue Bao*. 1997;32:586–92.
17. Jablonski EM, McConnell NA, Hughes Jr FM, Huet-Hudson YM. Estrogen regulation of aquaporins in the mouse uterus: potential roles in uterine water movement. *Biol Reprod*. 2003;69:1481–7.
18. Lindsay LA, Murphy CR. Redistribution of aquaporins 1 and 5 in the rat uterus is dependent on progesterone: a study with light and electron microscopy. *Reproduction*. 2006;131:369–78.
19. Aralla M, Borromeo V, Gropetti D, Secchi C, Cremonesi F, Arrighi S. A collaboration of aquaporins handles water transport in relation to the estrous cycle in the bitch uterus. *Theriogenology*. 2009;72:310–21.
20. Klein C, Troedsson MHT, Rutlant J. Expression of aquaporin water channels in equine endometrium is differentially regulated during the oestrous cycle and early pregnancy. *Reprod Dom Anim*. 2013;48:529–37.
21. Wang S, Chen J, Au KT, Ross MG. Expression of aquaporin 8 and its up-regulation by cyclic adenosine monophosphate in human WISH cells. *Am J Obstet Gynecol*. 2003;188:997–1001.
22. Wang S, Amidi F, Yin S, Beall M, Ross MG. Cyclic adenosine monophosphate regulation of aquaporin gene expression in human amnion epithelia. *Reprod Sci*. 2007;14:234–40.
23. Yang F, Kawedia JD, Menon AG. Cyclic AMP regulates aquaporin 5 expression at both transcriptional and post-transcriptional levels through a protein kinase A pathway. *J Biol Chem*. 2003;278:32173–80.
24. Belkacemi L, Beall MH, Magee TR, Pourtemour M, Ross MG. AQP1 gene expression is up-regulated by arginine vasopressin and cyclic AMP agonist in trophoblast cells. *Life Sci*. 2008;82:1272–80.
25. Hildenbrand A, Lalitkumar L, Nielsen S, Gemzell-Danielsson K, Stavreus-Evers A. Expression of aquaporin 2 in human endometrium. *Reprod Endocrinol*. 2006;86:1452–8.
26. Feng C, Sun CC, Wang TT, He RH, Sheng JZ, Huang HF. Decreased expression of endometrial vessel AQP1 and endometrial epithelium AQP2 related to an ovulatory uterine bleeding in premenopausal women. *Menopause*. 2011;15:648–54.
27. Skowronski MT, Kwon TH, Nielsen S. Immunolocalization of aquaporin 1, 5, and 9 in the female pig reproductive system. *J Histochem Cytochem*. 2009;57:61–7.
28. Akins EL, Morrisette MC. Gross ovarian changes during the estrous cycle of swine. *Am J Vet Res*. 1968;29:1953–7.
29. Skowronski MT, Lebeck J, Rojek A, Praetorius J, Fuchtbauer EM, Frokiaer J, et al. AQP7 is localized in capillaries of adipose tissue, cardiac and striated muscle: implications in glycerol metabolism. *Am J Physiol Renal Physiol*. 2007;292:F956–65.
30. Terris J, Ecelbarger CA, Nielsen S, Knepper MA. Long-term regulation of four renal aquaporins in rats. *Am J Physiol*. 1996;271:F414–22.
31. Kobayashi M, Takahashi E, Miyagawa S, Watanabe H, Iguchi T. Chromatin immunoprecipitation-mediated target identification proved aquaporin 5 is regulated directly by estrogen in the uterus. *Genes Cells*. 2006;11:1133–43.
32. Ing NH. Steroid hormones regulate gene expression posttranscriptionally by altering the stabilities of messenger RNAs. *Biol Reprod*. 2005;72:1290–6.
33. Krzymowski T, Stefanczyk-Krzybowska S. The role of endometrium in endocrine regulation of the animal oestrous cycle. *Reprod Dom Anim*. 2008;43:80–91.
34. Finlay TH, Katz J, Kirsh L, Levitz M, Nathoo SA, Seiler S. Estrogen-stimulated uptake of plasminogen by the mouse uterus. *Endocrinology*. 1983;112:856–61.
35. Norman SJ, Poyser N. Effects of inhibitors of arachidonic acid turnover on the production of prostaglandins by the guinea-pig uterus. *J Reprod Fertil*. 2000;118:181–6.
36. Hertelendy F, Zakar T. Prostaglandins and the myometrium and cervix. *Prostaglandin Leukotriens Essential Fatty Acids*. 2004;70:207–22.
37. Franczak A, Kotwica G, Kurowicka B, Oponowicz A, Waclawek-Potocka J, Petroff BK. Expression of enzymes of cyclooxygenase pathway and secretion of prostaglandin E₂ and F_{2α} by porcine myometrium during luteolysis and early pregnancy. *Theriogenology*. 2006;66:1049–56.
38. Irvine RF. How is the level of free arachidonic acid controlled in cells? *Biochem J*. 1982;204:3–16.
39. Blitek A, Waclawik A, Kaczmarek MM, Stadejek T, Pejask Z, Ziecik AJ. Expression of cyclooxygenase-1 and -2 in the porcine endometrium during the oestrous cycle and early pregnancy. *Reprod Dom Anim*. 2006;41:251–7.
40. Blitek A, Ziecik A. Prostaglandins F_{2α} and E₂ secretion by porcine epithelial and stromal endometrial cells on different days of the oestrus cycle. *Reprod Dom Anim*. 2004;39:340–6.
41. Gleeson AR, Thorburn GD, Cox RI. Prostaglandin F concentrations in the utero-ovarian venous plasma of the sow during the late luteal phase of the oestrous cycle. *Prostaglandins*. 1974;5:521–9.
42. Zelenina M, Christensen BM, Palmer J, Nairn AC, Nielsen S, Aperia A. Prostaglandin E₂ interaction with AVP: effects on AQP2 phosphorylation and distribution. *Am J Physiol Renal Physiol*. 2000;278:F388–94.
43. Uzumcu M, Braileanu GT, Carnahan KG, Ludwig TE, Mirando MA. Oxytocin – stimulated phosphoinositide hydrolysis and prostaglandin F secretion by luminal epithelial, glandular epithelial and stromal cells from pig endometrium. I. Response of cyclic pig on day 16 postestrus. *Biol Reprod*. 1998;59:1259–65.
44. Trout WE, Smith GW, Gentry PC, Galvin JM, Keisler DH. Oxytocin secretion by the endometrium of the pig during maternal recognition of pregnancy. *Biol Reprod*. 1995;52 Suppl 1:189. abstract.
45. Whiteaker SS, Mirando MA, Becker WC, Hostetler CE. Detection of functional oxytocin receptors on endometrium of pigs. *Biol Reprod*. 1994;51:92–8.
46. Okano A, Okuda K, Takahashi M, Schams D. Oxytocin receptors in porcine endometrium during the estrous cycle and early pregnancy. *Animal Reprod Sci*. 1996;41:61–70.
47. Gilbert CL, Jenkins, Wathes DC. Pulsatile release of oxytocin into the circulation of the ewe during oestrous, mating and the early luteal phase. *J Reprod Fertil*. 1991;91:337–46.
48. Ducza E, Seres AB, Hajagos-Toth J, Falkay G, Gaspar R. Oxytocin regulates the expression of aquaporin 5 in the late-pregnant rat uterus. *Mol Reprod Dev*. 2014;81:524–30.
49. Wang S, Kallichanda N, Song W, Ramirez B, Ross MG. Expression of aquaporin 8 in human placenta and chorioamniotic membranes: evidence of molecular mechanism for intramembranous amniotic fluid resorption. *Am J Obstet Gynecol*. 2001;185:1226–31.
50. Garcia F, Kierbel A, Larocca MC, Gradilone SA, Splinter P, LaRusso NF, et al. The water channel aquaporin-8 is mainly intracellular in rat hepatocytes and its plasma membrane insertion is stimulated by cyclic AMP. *J Biol Chem*. 2001;276:12147–52.