



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Therapeutic drug quantification using targeted mass spectrometry

Meyer, Michael Kruse; Andersen, Marlene; Stausbo , Troels Vindbæk ; Bennike, Tue Bjerg; Andersen, Grethe Neumann; Stensballe, Allan

*Publication date:*  
2016

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Meyer, M. K., Andersen, M., Stausbo , T. V., Bennike, T. B., Andersen, G. N., & Stensballe, A. (2016). *Therapeutic drug quantification using targeted mass spectrometry*. Abstract from 8th MaxQuant Summer School on Computational Mass Spectrometry-based Proteomics, Oxford, United Kingdom.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Therapeutic Drug Quantification using Targeted Mass Spectrometry

Michael Kruse Meyer<sup>1,2</sup>, Marlene Andersen<sup>1</sup>, Troels Vindbæk Stausbo<sup>2</sup>, Tue Bjerg Bennike<sup>3</sup>, Grethe Neumann Andersen<sup>1</sup> & Allan Stensballe<sup>2</sup>

## Author Affiliations:

1 Department of Rheumatology, North Denmark Regional Hospital, Denmark, and Center for Clinical Science, Aalborg University, Denmark

2. Department of Health Science and Technology, Aalborg University, Denmark

3. Departments of Pathology, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts, USA

\* To whom correspondence should be addressed:

Michael Kruse Meyer, Department of Rheumatology, North Denmark Regional Hospital, Denmark. Bispensgade 37 – 9800 Hjoerring, Denmark. Email: michael.o@rn.dk, Phone: +45 20 31 39 03

MaxQuant Summer School 8th

**Abstract** (2000)

## Keywords

Biological Treatment, disease modifying anti-rheumatic treatment (DMARD), b biological DMARD (bDMARD), parallel reaction monitoring

## Introduction:

Rheumatic diseases involve auto-immune, and inflammatory diseases, in which the inflammatory activity degrades cartilage, and eventually destroys the joints. We have the past five years, systematically reduced treatment frequency and dosage using empirical clinical data and patient

outcome. Previous, determination of the optimal treatment dosage and frequency for biological drugs was based on clinical symptoms and patient questionnaire feedback, with no truly objective biochemical evaluations. Hence, we seek to quantify the biological drug concentrations in the patients to establish a more objective dosing strategy. Previous studies have utilized immuno-based assays, which are sensitive, but limited to availability of good antibodies. They also require high volumes of sample, and are not readily multiplexed. In this study, we developed a parallel reaction monitoring (PRM) based method on four selected biological drugs on a QExactive HF.

### **Method:**

A standard method for ELISA preparation was optimized and scaled down for nUPLC-MS/MS. Four biological drugs and serum samples from patients were immunoprecipitated, and characterized on a Q Exactive high field mass spectrometer. Based on the results, a target list of proteotypic peptides and a spectral database was constructed, and used to establish a PRM method in Skyline, which was utilized to reanalyze the samples.

### **Results:**

We have successfully established a method able to quantify three of the four biological drugs. Interestingly, we show that the PRM method from one commonly administered biological drug, infliximab, also works on the biosimilar version of infliximab with no modifications made to the method used above.

### **Conclusion:**

We have established the foundation for accurately determining the concentration of several biological drugs routinely used in clinical settings. The methods can provide objective measurements to help establish the correct treatment dosage and frequency.