Aalborg Universitet



## Investigation of a 12-weeks community-based training intervention in children with obesity

Importance of intensity and exercise modality

Eggertsen, Charlotte Nørkjær; Bech Pedersen, Jeppe; Hagstrøm, Søren; Vestergaard, Esben Thyssen; Larsen, Ryan Godsk; Simonsen, Morten Bilde

Publication date: 2023

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA):

Eggertsen, C. N., Bech Pedersen, J., Hagstrøm, S., Vestergaard, E. T., Larsen, R. G., & Simonsen, M. B. (2023). *Investigation of a 12-weeks community-based training intervention in children with obesity: Importance of intensity and exercise modality*. Abstract from 22nd Annual Meeting of The International Society of Behavioral Nutrition and Physical Activity, Sweden.

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
  You may freely distribute the URL identifying the publication in the public portal -

Take down policy If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

## Investigation of a 12-weeks community-based training intervention in children with obesity: Importance of intensity and exercise modality.

**Dr. Charlotte Eggertsen**<sup>1,2</sup>, Mr. Jeppe Bech Pedersen<sup>4</sup>, Prof. Søren Hagstrøm<sup>1,2</sup>, Dr. Esben Thyssen Vestergaard<sup>3</sup>, Associate Professor Ryan Godsk Larsen<sup>4</sup>, Associate Professor Morten Bilde Simonsen<sup>5</sup> <sup>1</sup>Department of Pediatrics and Steno Diabetes Center North Denmark, Aalborg University Hospital, Aalborg, Denmark, <sup>2</sup>Department of Clinical Medicine, Aalborg University Hospital, Aalborg, Denmark, <sup>3</sup>Department of Pediatrics, Regional Hospital Randers, Randers, Denmark, <sup>4</sup>Department of Health Science and Technology, Aalborg University, Aalborg, Denmark, <sup>5</sup>Department of Materials and Production, Aalborg University, Aalborg, Denmark

Poster Session #2, June 16, 2023, 10:50 AM - 12:00 PM

SIG - Primary Choice: L. Assessment and Methodologies in Behavioral Nutrition and Physical Activity

**Purpose** Childhood obesity is a major public health concern. Physical activity as part of a multidisciplinary lifestyle intervention is a keystone in treating obesity. High-intensity interval-training (HIIT) has shown several physiological benefits, including weight reduction and cardiovascular health. Few studies have monitored the intensity during HIIT or compared different types of exercises. This study is part of a randomized controlled project investigating the effect of HIIT in a group of children with obesity, and focuses on the training intensity of different activities in a community-based training setting.

**Methods** Sixty-three children with obesity were enrolled in a 12-weeks training intervention, with three weekly sessions. Each session included 4x4 minutes of HIIT focusing on one of three specific exercise modalities; Ball games, CrossFit, or running. Heart rate (HR) was measured to quantify time spent in high (>85% of max HR) intensity activity. In nine of the HIIT sessions, enjoyment (Physical Activity Enjoyment Scale) and perceived exertion (Borg CR-10 scale) were recorded. A three-way (sex, modality, intensity) mixed ANOVA was used to compare the HR data. Friedman and Wilcoxon Signed-rank tests (post hoc) were used to compare enjoyment and perceived exertion between types of activity.

**Results** The participants' median age was 12.3 years, with a body-mass index of  $28.84 \pm 4.04$ . Ball games elicited less time in the high-intensity zone ( $4.96\pm5.75$  minutes) than CrossFit ( $7.73\pm7.25$  minutes, P = 0.010) and running ( $6.36\pm5.80$  minutes, P = 0.025). No gender interaction was observed (p=0.995). The enjoyment of physical activity was rated high, on average 4.5 for all activities (scale from 1(lowest) – 5(highest)), but no significant difference was found. The perceived exertion was rated lower by adolescents compared to children in ball games ( $5.1\pm1$ ) compared to CrossFit ( $6\pm1$ , P = 0.017) and running ( $6\pm1$ , P = 0.004). Furthermore, with all participants combined, ball games were perceived as less exhausting than CrossFit (P = 0.002) and running (P = 0.003).

**Conclusions** The participants had high enjoyment ratings during all three modalities. CrossFit and running seem favorable to elicit more time in the high-intensity zone compared to ball games. This should be considered in future training interventions.