



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Wolf genetic diversity compared across Europe using the yardstick method

Jan, Maja; Stronen, Astrid Vik; Boljte, Barbara; Černe, Rok; Huber, Đuro; Iosif, Ruben; Kljun, Franc; Konec, Marjeta; Kos, Ivan; Krofel, Miha; Kusak, Josip; Luštrik, Roman; Majić Skrbinišek, Aleksandra; Promberger–Füerpass, Barbara; Potočnik, Hubert; Rigg, Robin; Trontelj, Peter; Skrbinišek, Tomaž

Published in:
Scientific Reports

DOI (link to publication from Publisher):
[10.1038/s41598-023-40834-x](https://doi.org/10.1038/s41598-023-40834-x)

Creative Commons License
CC BY 4.0

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Jan, M., Stronen, A. V., Boljte, B., Černe, R., Huber, Đ., Iosif, R., Kljun, F., Konec, M., Kos, I., Krofel, M., Kusak, J., Luštrik, R., Majić Skrbinišek, A., Promberger–Füerpass, B., Potočnik, H., Rigg, R., Trontelj, P., & Skrbinišek, T. (2023). Wolf genetic diversity compared across Europe using the yardstick method. *Scientific Reports*, 13, Article 13727. <https://doi.org/10.1038/s41598-023-40834-x>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -



OPEN Wolf genetic diversity compared across Europe using the yardstick method

Maja Jan¹✉, Astrid Vik Stronen^{1,2,3}, Barbara Boljte^{1,2}, Rok Černe⁴, Đuro Huber⁵, Ruben Iosif⁶, Franc Kljun¹, Marjeta Konec^{1,2}, Ivan Kos¹, Miha Krofel¹, Josip Kusak⁵, Roman Luštrik⁷, Aleksandra Majić Skrbinšek^{1,2}, Barbara Promberger-Fürerpass⁶, Hubert Potočník¹, Robin Rigg^{1,8}, Peter Trontelj¹ & Tomaž Skrbinšek^{1,2}

Integrating data across studies with traditional microsatellite genetic markers requires careful calibration and represents an obstacle for investigation of wide-ranging species where populations require transboundary management. We used the “yardstick” method to compare results published across Europe since 2002 and new wolf (*Canis lupus*) genetic profiles from the Carpathian Mountains in Central Europe and the Dinaric Mountains in Southeastern Europe, with the latter as our reference population. We compared each population with Dinaric wolves, considering only shared markers (range 4–17). For each population, we calculated standard genetic diversity indices plus calibrated heterozygosity (*Hec*) and allelic richness (*Ac*). *Hec* and *Ac* in Dinaric (0.704 and 9.394) and Carpathian wolves (0.695 and 7.023) were comparable to those observed in other large and mid-sized European populations, but smaller than those of northeastern Europe. Major discrepancies in marker choices among some studies made comparisons more difficult. However, the yardstick method, including the new measures of *Hec* and *Ac*, provided a direct comparison of genetic diversity values among wolf populations and an intuitive interpretation of the results. The yardstick method thus permitted the integration of diverse sources of publicly available microsatellite data for spatiotemporal genetic monitoring of evolutionary potential.

Comparison of results from traditional microsatellite genetic markers across studies requires calibration and thus complicates broad-scale studies¹. This has negative consequences for evolutionary research and conservation management of wide-ranging species where many populations range across multiple international boundaries^{2–4} and single studies rarely include entire populations. Rapid environmental changes and associated range shifts in many wild species further augment the need to understand their population genetic structure and evolutionary potential across wider scales (e.g., Ref.⁵). New genomic techniques are increasingly permitting broad-scale investigation also for non-model organisms, including marine and terrestrial taxa (e.g., Ref.^{6,7}). However, many genetic monitoring programs still use microsatellite markers, which offer important temporal perspectives in cases where DNA sources no longer exist, but legacy data are available in the form of microsatellite profiles⁸. Given the comparative strength of microsatellite markers for detecting changes across limited temporal and spatial scales⁹, more efforts are needed to mitigate the problems that researchers and conservation managers encounter when seeking to integrate these valuable datasets. Efforts toward standardizing non-invasive genetic methods would also greatly facilitate the study of dispersal, and in turn help discern biological differences in dispersal parameters from variability owing to differences in methodology (reviewed in Ref.¹⁰).

To help alleviate these constraints, a “yardstick” method was proposed for comparison of genetic diversity indices among populations and applied to brown bears (*Ursus arctos*) as a relevant example¹¹. The method calibrates genetic diversity indices to a common denominator, the reference or “yardstick” population, allowing comparisons to be made. The authors suggested that this method may be even more useful for species with

¹Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia. ²DivjaLabs d.o.o., Aljaževa ulica 35a, 1000 Ljubljana, Slovenia. ³Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, 9220 Aalborg, Denmark. ⁴Slovenia Forest Service, Večna pot 2, 1000 Ljubljana, Slovenia. ⁵Faculty of Veterinary Medicine, University of Zagreb, Vjekoslava Heinzelova 55, 10000 Zagreb, Croatia. ⁶Foundation Conservation Carpathia, 27 Calea Feldioarei, 500471 Braşov, Romania. ⁷Genialis Inc, Vojkova cesta 63, 1000 Ljubljana, Slovenia. ⁸Slovak Wildlife Society, Belanská 574/6, P.O. Box 72, Liptovský Hrádok 033 01, Slovakia. ✉email: mjelencic@gmail.com

more diverse (i.e., less overlapping) sets of genetic markers used in different studies, as is the case with the grey wolf (*Canis lupus*)³. The yardstick method could facilitate comparison of existing data from populations with diverse demographic histories¹¹, including habitat fragmentation, bottlenecks, and founder effects. Notably, allelic diversity has been proposed as a better predictor and candidate proxy than heterozygosity of evolutionary potential¹². Recent works maintain that genetic diversity is vital for evolutionary fitness (reviewed in Ref.¹³) and that within-species genetic diversity merits wider recognition in international conservation planning¹⁴. The yardstick method's inclusion of allelic richness could thus provide a flexible and cost-effective measure for genetic monitoring, research, and applied conservation management.

The wolf is a wide-ranging species where discrepancies in genetic markers among studies limit comparisons at relevant spatial scales in Europe³. A review of wolf population genetics across Europe found genetic diversity to be high in northeastern Europe and decreasing toward the southwest¹⁵, and the authors reported that observed and expected heterozygosity showed a clear spatial trend whereas allelic richness did not. They also noted the potential of the yardstick method but were unable to use this approach to analyze the wolf microsatellite data across the continent because of too few (≤ 3) overlapping loci.

In recent years, new genetic data on wolves has become available, allowing comparison of findings across regional scales, based on a larger number of common markers. These new data include populations in Central and Southeastern Europe where relatively abundant and persistent populations of large carnivores² are thought to represent vital genetic diversity for long-term preservation of evolutionary potential. Whereas wolves in many parts of Europe are well-studied at various spatio-temporal scales, those in Central and Southeastern Europe have received less attention (Supplemental Tables S1 and S2) and have high priority for conservation¹⁶.

Analyses across genetic markers—including neutral and potentially adaptive genetic variation—show that wolves from the Carpathian Mountains in Central Europe are separated from wolves farther north^{17,18}, although there is evidence of recent gene flow into the Carpathians¹⁹. The Carpathian population spans the Czech Republic, Slovakia, Poland, Hungary, Ukraine, Serbia, and Romania and is among the largest in Europe²⁰, estimated to number c. 3900–4700 individuals¹⁶.

Recent population genetic analyses of microsatellite markers showed spatial structure within Slovakia that may have been influenced by environmental and anthropogenic factors¹⁹. Wolves were almost extirpated twice from Slovakia during the twentieth century but recovered and have since become relatively abundant in the central and eastern regions^{21,22}. Moreover, further research is needed to evaluate connectivity across the region. A recent genome-wide analysis found signs of weak genetic structure between wolves in Romania and the Carpathian Mountain region of Ukraine²³. Additional analyses with microsatellite markers could help clarify the extent of genetic exchange across short spatio-temporal scales⁹ and whether there are signs of recent resistance to gene flow.

The Carpathian Mountain wolves were reported to be genetically distinct from the Dinaric-Balkan population^{18,24,25}. The latter spans ten countries, from the Italian–Slovenian border in the west, across the Dinaric Mountains, to the Balkan Peninsula and Rhodope Mountains in Greece and Bulgaria in the southeast, with an estimated size of c. 5000–5500 individuals^{16,26}. Wolves in the northwestern Dinaric Mountains (NW Dinaric) were subject to severe persecution in the past and were almost extirpated during the eighteenth and nineteenth centuries. The population started to recover in the second half of the twentieth century^{27,28} and is currently expanding its range into the Alps, where contact with the Alpine population has already been established^{29,30}. Earlier findings from Croatia and Bosnia & Herzegovina have reported findings of genetic structure based on microsatellite markers^{29,31} and although such differentiation was not observed in Serbia³², there was evidence of structure based on mitochondrial DNA³³, suggesting the need for broader-scale analyses.

The main objectives of this study were to (1) place new findings on wolf genetic diversity in the Carpathian and Dinaric regions into a broader context at the European level using previously published studies and (2) apply the yardstick method to a model species where conservation managers require a better understanding of genetic diversity at the continental scale. We also extended the yardstick method with calculation of calibrated heterozygosity and allelic richness estimates, enhancing the interpretability of the results. Finally, we discuss the relevance of these measures for conservation management of wolves and other wide-ranging species.

Materials and methods

Sampling. We included 345 wolf profiles from tissue and noninvasive samples from the Dinaric Mountains (Slovenia, $n = 145$ and Croatia, $n = 148$) and the Carpathian Mountains (Romania, $n = 32$ and Slovakia $n = 20$; Supplemental Fig. S1), after removal of individuals showing any sign of possible hybridization with dogs. For details about sampling, laboratory processing, genotyping and error checking, assessment of hybridization, and identification of wolves, see Supplementary Note S1, Supplemental Tables S3, and S4. We obtained genotypes for 34 microsatellite markers for Dinaric wolves, and a subset of 21 markers for Carpathian wolves. We henceforth refer to wolves in Slovenia and the Gorski kotar region of northwestern Croatia—our reference population—as the northwestern Dinaric (NW Dinaric) population. The broader Dinaric Mountain study area also comprised the Croatian regions of Lika and Dalmatia further south (Supplemental Fig. S1).

Estimation of genetic diversity, reference wolf population and comparison of populations. For data handling, calculation of genetic diversity indices—observed heterozygosity (H_o), expected heterozygosity (H_e) and allelic diversity (A)—and estimation of genotyping error rates, we used the package *adegenet* v.2.1.8³⁴ in R v.4.2.2³⁵. We used the R package *pegas* v. 1.0-1³⁶ to estimate departures from Hardy–Weinberg (HW) equilibrium and corrected for multiple testing with the Holm–Bonferroni multiple test correction with $\alpha = 0.05$ ³⁷.

We collected published genetic diversity data based on microsatellites for other European gray wolf populations through literature review (Supplemental Tables S1 and S2). Where diversity indices were published per locus and, at times, per population (for studies that reported population genetic structure), we compared allelic richness using the yardstick reference population method¹¹. For the purpose of this study, we use ‘population’ to refer to sampling areas and/or population genetic clusters reported in the original studies, although it is important to note that sampling may have been defined by national and regional jurisdictions and does not always correspond with broad-scale studies, where genome-wide analyses have found several transboundary units comprising multiple countries (e.g., Ref.^{18,38,39}). We performed pairwise comparison whereby each population of interest was compared with a reference population, considering only the genetic markers shared between the two studies. Differences in sample size were corrected through resampling with replacement multiple times (~1000) from the larger sample size to the sample size of the smaller data set. For the resulting subsamples, genetic diversity indices with their standard errors were calculated (A , He and Ho). Heterozygosity ratio (Her) and allelic richness ratio (Art) indices were calculated with the corresponding reference population values in the denominator¹¹. As an extension of the method described in the original paper, we multiplied the Her and Art ratio indices with heterozygosity or, respectively, allelic richness of the reference population, obtaining directly interpretable calibrated expected heterozygosity (Hec) and allelic richness (Ac) values. As the values of these parameters were on the same scale as the original heterozygosity and allelic richness, this allowed both direct comparison between otherwise incompatible datasets as well as a more intuitive interpretation of the obtained values. As the relationship between observed and expected heterozygosity is better estimated in the original data (more loci, no resampling), we maintained the relationship between these two parameters in each population by calculating the calibrated observed heterozygosity (Hoc) from Hec and the ratio between observed (Ho) and expected (He) heterozygosity in the original data as $Hoc = Hec * (Ho/He)$.

The NW Dinaric population with a relatively large sample size and locus set was used as the reference population (Supplemental Table S2, Supplemental Fig. S1, Supplemental Note S1). To avoid a possible sampling intensity bias in Slovenia and Croatia, we randomly selected 76 out of 145 individuals from Slovenia to match the number of wolves sampled in Gorski kotar (58), considering that the estimated number of wolf packs living in each area is comparable⁴⁰. The final reference population included 134 individuals. Analyses were performed in R using the “resampleddiversity” package¹¹ (also available on GitHub (<https://github.com/romunov/resampleddiversity>)) and implemented in the function `divRatio` of the `diveRsim` package⁴¹). For each comparison we ran 1000 random subsamples. When using the yardstick method to calculate diversity indices for populations with sample sizes larger than the reference population, we compared the published indices directly (i.e., without resampling) using the loci in common. In these cases, the number of samples was large in both populations and the relative sample sizes were very similar, and any bias should therefore be minimal. Figures were created using base R functions and the R package `ggplot2`⁴². Maps were created using QGIS 3.24.0⁴³.

Results

Dinaric and Carpathian wolves. For Dinaric wolves, locus FH2004 indicated the presence of null alleles (estimated frequency 0.10) and was therefore excluded (Supplemental Note S2, Supplemental Table S3). Genetic diversity values were comparable among the NW Dinaric, Lika, Dalmatia, and Romanian populations whereas Slovakian wolves, where we had the smallest sample size, showed somewhat lower values (Table 1; Fig. 1; Supplemental Table S2).

Comparison of genetic diversity of Dinaric and Carpathian wolves with other European populations. For the studies we examined, the number of microsatellite loci ranged from 10⁴⁴ to 42⁴⁵ (Supplemental Table S2; Fig. 2). The minimum number of loci shared between our reference population and other populations was four, for wolves from the Iberian Peninsula and Russia⁴⁶. We also included data from other wolf studies from these areas that shared a higher number of loci with the reference population. For wolves from the Iberian Peninsula, we used data from Ref.⁴⁵, and for wolves in NW Russian we used results from Ref.^{47,48}. The maximum number of loci shared between our reference population and another wolf population was 17, for Iberian wolves⁴⁵.

Heterozygosity and allelic richness were generally higher in populations in N and NE Europe (European Russia, the Baltic States, Finland) (Fig. 3). In many cases He and Ho seem far apart (Fig. 1), suggesting a departure from HW equilibrium and possible population-level processes driving this departure. Examples are Iberia^{45,46,49}, Poland¹⁷, and Bulgaria²⁴, where the authors had noted known or probable underlying genetic structure. In general, allelic richness ratio and heterozygosity were lowest for Scandinavian wolves in 1991–2001 ($Art = 0.685$) and highest for wolves in European Russia ($Art = 2.252$).

Calibrated heterozygosity and allelic richness in Dinaric wolves were higher than those found in Scandinavian, Italian and Iberian wolves. In Carpathian wolves, both parameters were higher in Romania than in Slovakia, and comparable to those of Dinaric wolves. Our results from the Dinaric and Carpathian regions showed lower values for heterozygosity and allelic richness than those observed in NE Europe.

Discussion

We assessed the genetic diversity of wolves in the Dinaric and Carpathian regions in Europe and examined our findings in a wider context at the continental level. The yardstick method¹¹ allowed us to compare our results with genetic diversity parameters published in various studies from across Europe, which included samples originating from 1829 to 2018 (Supplemental Table S2). For Dinaric and Carpathian wolves we observed comparable calibrated heterozygosity and allelic richness, and genetic diversity values were, as expected, lower than in the large populations of northeastern Europe (Ref.¹⁵ and Supplemental Table S2). Within the Carpathians, the




Figure 1. Observed (H_o) versus expected (H_e) heterozygosity and calibrated allelic richness (A_c) with wolves in NW Dinaric (Slovenia and Gorski kotar, Croatia) used as the reference population. Dashed diagonal line presents the Hardy–Weinberg (HW) equilibrium of $H_o = H_e$, and the positions of the points indicate the direction of deviation from HWE. The graph on the right is a close-up of the marked region in the first image. Some studies did not report values for all parameters as noted for Supplemental Table S2.

Populations	<i>N</i>	<i>A</i> (s.e.)	<i>He</i> (s.e.)	<i>Ho</i> (s.e.)
NW Dinaric (Slovenia and Gorski kotar, HR)	134	6.62 (0.32)	0.673 (0.021)	0.661 (0.025)
Lika (HR)	34	5.91 (0.27)	0.687 (0.018)	0.699 (0.023)
Dalmatia (HR)	56	6.47 (0.27)	0.702 (0.016)	0.680 (0.020)
Slovakia	20	4.86 (0.22)	0.640 (0.031)	0.650 (0.039)
Romania	32	5.71 (0.33)	0.690 (0.025)	0.700 (0.039)

Table 1. Genetic diversity measures for Dinaric (NW Dinaric, Lika, Dalmatia) and Carpathian (Slovakia and Romania) wolves. *N* number of individuals, *A* allelic diversity, *He* expected heterozygosity, *Ho* observed heterozygosity, with standard error (s.e.).

larger population segment in Romania also exhibited higher values than wolves in Slovakia. The meta-analysis provided a valuable continental-level overview of patterns of genetic diversity of this species, enabling a better understanding of the conservation status and ongoing recovery process of wolves in Europe.

Meta-analysis of genetic diversity in European wolves. The observed distribution of genetic diversity in European wolves fits well with the general picture found in earlier studies^{15,38,39,44,46,50–52}. Interestingly, none of the populations (Supplemental Table S2) showed exceedingly high or low heterozygosity despite some small populations have been isolated. This contrasts with the situation found in brown bears where major differences in heterozygosity were observed¹¹. On the other hand, the differences in calibrated allelic richness (*Ac*) among wolf populations are considerable and, in many cases, correspond better with known population histories than the heterozygosity estimates. Ref.¹⁵ noted that allelic richness was distributed relatively evenly across Europe, showing only a weak spatial signal. In our study, *Ac* nonetheless shows a general increase from southwest to north-east, which corresponds with the distribution of genetic diversity across European populations observed by Ref.¹⁵. As allelic richness seems to be a good indicator of severe, short bottlenecks⁵³, and may more effectively reflect a population’s long-term evolutionary potential^{12,54}, temporal data on allelic richness can provide an effective means of monitoring smaller and relatively isolated populations identified as vulnerable (Ref.¹⁵ and references therein).

There is a general pattern of deviations from HW equilibrium in most reported wolf studies (Fig. 1). In some cases, this is clearly caused by the Wahlund effect⁵⁵, but deviations occur at a relatively small spatial scale for a highly mobile mammal^{29,30,49,56}. A simple explanation is that because many populations survived the persecution of previous centuries in very small numbers, high genetic drift caused by low effective population sizes leads to detectable genetic differentiation between populations, or even between population fragments within the




Figure 2. Comparison of original and calibrated heterozygosity and allelic richness values for European wolf populations. N = number of samples, loc/used = number of loci in the population/number of loci used in comparisons. Lines indicate the confidence interval of each estimate ($\pm 1.96 \times \text{s.e.}$).




Figure 3. Distribution of wolves in Europe (based on Kaczensky et al.²⁶) and genetic diversity indices reported in various studies calibrated using the “yardstick” method¹¹. Overall, both calibrated heterozygosity and calibrated allelic richness showed a general increase from south-west to north-east, similar to that observed by Hindrikson et al.¹⁵. Studies that reported parameter values are included—see Supplemental Table S2.

same metapopulation, also at relatively small spatial scales. Founder effects during recolonization have resulted in persistent genetic drift⁵⁷. Moreover, some areas were colonized from populations that were quite recently established by a limited number of founders, such as wolves in Denmark⁵⁸, and a “double” founder effect could thus be affecting wolves in such areas.

However, in other cases there are clear deviations from HW equilibrium where a hidden substructure would be unexpected given the spatial and landscape characteristics of the area⁵⁹. Such deviations seem more frequent in populations that are geographically close to other genetically differentiated wolf populations, than in divergent populations separated by larger geographic distances, as observed in our study. This general pattern of deviations from HW equilibrium at the continental scale may suggest that, as wolves in Europe continue to recover, gene flow may be starting between previously isolated populations. However, the detectable effects of gene flow usually follow clear and predictable patterns. At first, direct immigrants (i.e., dispersers) are expected to cause the Wahlund effect⁵⁵ resulting in excess homozygotes and a positive *Fis*⁶⁰, which may be misinterpreted as a sign of assortative breeding if no other analysis is done. This can be an explanation for populations that are above the diagonal dashed line in Fig. 1 where there is no reasonable expectation of genetic substructure (which would cause the same effect). Subsequent reproduction of these dispersers with resident wolves would cause an excess of heterozygotes (“isolate breaking”⁵⁵), shifting the deviation from HW equilibrium in the other direction and causing negative *Fis*, which can also be misinterpreted as a sign of inbreeding avoidance. This may be the case for the populations below the dashed diagonal line in Fig. 1. Whereas these interesting processes require further research at the continental level and need to be interpreted using other knowledge and data about specific populations, an increase of gene flow could be seen as a positive development from the conservation perspective.

Utility of the yardstick method for analyses of European wolves. Originally, the yardstick method calculates the ratios between the genetic diversity indices in target versus reference populations, but they are not always intuitive to interpret. We extended the method to calculate calibrated heterozygosity and allelic richness, which enables a much more intuitive interpretation. Moreover, we compared the “original” genetic diversity values obtained in the reviewed studies with the calibrated values (Fig. 2). The differences were small in studies with large sample sizes and large number of loci but became pronounced in small studies with a limited number of loci (Table S2). In many cases it would be impossible to meaningfully compare these studies with other studies of the same species. Using the yardstick method such comparisons became possible, but the confidence intervals around such estimates were quite large.

If sample size is small, estimates of some parameters, particularly allelic diversity, can become severely biased, and Ref.¹¹ used bootstrap resampling to account for this issue when demonstrating the yardstick method. In our study the sample size for the reference (NW Dinaric) population was lower than in nine of the populations used for comparison. These nine studies all had relatively large sample sizes, and we therefore compared the published *He* and *Ho* for the shared loci directly (i.e., without resampling).

One challenge for our investigation was the availability of data from the published studies. In some cases, we were unable to calculate the calibrated allelic richness or observed heterozygosity ratio with the yardstick method for some of the reported population clusters, because the authors did not report locus-level allelic diversity for each separate unit. Neither were the temporal extents, sampling periods, or sampling areas clearly defined in all cases, and the results should therefore be interpreted with caution. For example, we noticed quite a difference between the genetic diversity observed in Slovakia in our study (study SK2) and that of Ref.²⁴ (study SK1). Our sampling area was focused on northern Slovakia, and we included only 20 individuals, whereas Ref.²⁴ had a sample of 194.

We genotyped 33 microsatellites, and only Ref.⁴⁵ among the compared studies reported analyzing more markers (42 loci). However, the numbers of overlapping markers available for comparison between the reference and other populations varied widely, ranging from 21 to only four loci. The marker panels differed considerably among studies, reflecting the high diversity of markers available for canids, and nearly every study included here had its own marker panel. Conversely, in brown bears, where the number of markers routinely investigated is smaller, more than half of the compared studies had eight loci available for comparison with the reference population¹¹. Whereas the high diversity of canid markers presents some challenges and complicates the comparison of populations that have few overlapping markers with the reference population, it also highlights the yardstick method’s utility in our objective of comparing wolves across Europe.

The yardstick method can inform conservation management for wide-ranging species where populations encompass multiple jurisdictions, such as large carnivores in the Carpathian and Dinaric-Balkan regions where populations of wolves, bears, and lynx (*Lynx lynx*) extend across multiple countries (Ref.^{2,61}). The same holds for many ungulate species, such as red deer (*Cervus elaphus*) and wild boar (*Sus scrofa*) that are broadly distributed across Europe^{62,63}. As with wolves in parts of our study area⁶⁴, red deer, wild boar and other ungulate species in Europe have been affected by hybridization in parts of their range (reviewed in Ref.⁴). Such events confound analyses of genetic structure and variability, and make it difficult to distinguish natural patterns of gene flow and diversity from those influenced by human activities⁶². Broad-scale investigations that permit direct comparison of results across geographic regions therefore remain an important priority⁴.

Genetic diversity of wolves in the Dinaric and Carpathian regions. The genetic diversity of wolves in the Dinaric Mountains reflects the population’s demographic history. Although it is comparable to many other recovering wolf populations in Europe, even the largest of these populations share the same history of persecution and recent recovery²⁰ and may also have suffered population fragmentation and reduction of genetic diversity. Observed deviations from HW equilibrium are not unexpected and are probably due to the Wahlund effect as the population seems to be divided into two or three subpopulations²⁹. Substructure at this scale may be

a legacy of past persecution when wolves often survived in small and isolated population fragments^{29,51,65}, where genetic drift quickly created a strong signal of population structure. Even relatively large populations, including Dinaric wolves, have therefore lost a considerable part of the original genetic diversity within each isolated fragment. There are indications, however, that the current recovery is starting to create gene flow between some population fragments^{19,29,30,51,66,67}, which should help dissolve or at least reduce some of the existing population structure and augment genetic diversity.

Within the Carpathian Mountains, wolves in Romania seem to have higher genetic diversity than wolves in Slovakia, which is expected as they represent a major part (70%) of the Carpathian population⁶¹. Romanian wolves also appear relatively well-connected with wolves in the Carpathian Mountains of western Ukraine²³. Wolves in Slovakia seem to have lower genetic diversity than their Dinaric counterparts, but still higher than the Scandinavian (study SCAN2) and Italian (study IT2) populations that are known to have gone through severe bottlenecks. Recent findings also suggest a dynamic situation with signs of recent gene flow between the Carpathian Mountains and wolves farther north and west^{19,67}, which requires further monitoring and investigation.

Implications for conservation and management. Although we observed considerable differences among wolf populations in Europe, even at finer spatial scales, these differences may gradually dissolve as recovering wolf populations become increasingly reconnected. This process is already seen in the Apennine Mountains²⁹, in the Alps and Dinaric Mountains^{29,30}, in this study, and possibly in the Carpathians Mountains¹⁹, and Poland⁶⁷. Hence, this may be a wider phenomenon connected with the ongoing continent-wide recovery of wolves²⁰.

The newly presented measures for calibrated heterozygosity and allelic richness allow a direct comparison of genetic diversity values among populations and a more intuitive interpretation of the results. However, it is still crucial to interpret the results carefully, including deviations from HW equilibrium, as this may indicate the presence of cryptic population structure⁵⁹. For certain populations showing structure across relatively short geographic distances, natural selection has also been suggested as a possible contributing factor^{18,30,51,68} although it is difficult to see how this could have occurred in recent times. Given that most of these populations have been considerably reduced and their effective population size was (and probably still is) low, selection would need to be extremely strong not to be overpowered by genetic drift⁶⁹, making adaptive evolutionary changes in such populations unlikely. However, this selection could have occurred in the past, in which case some of the currently observed population structure in wolves might not be a result of human activity, but possibly a result of natural selection (e.g., Ref.^{70,71}). These issues have consequences for applied conservation of wild species, for example, whether some populations should be conserved in situ and admixture with other populations encouraged only via natural dispersal. These questions require further genomic research that can evaluate both neutral and functional genetic variation.

Gene flow from populations adapted to warmer and drier conditions could become increasingly important given rapid environmental changes⁷², and it would therefore seem farsighted to preserve the genetic diversity currently found in habitats where organisms are more likely to have experienced such selective pressures. The role of adaptive potential at range edges is important also for highly mobile species⁷³, and future research could help clarify the extent to which allelic richness may function as an indicator of adaptive potential, where the yardstick method could be informative.

Much conservation effort is focused at the species level, including the backbone of European biodiversity conservation policy, the Habitats Directive (Council Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora). Extinction rates for populations, however, are estimated to be three to eight times higher than those for species⁷⁴. On the other hand, practical conservation in Europe is frequently done at the level of administrative units—countries, or even provinces—causing further “administrative fragmentation” of conservation management. To promote conservation, restore genetic diversity, and facilitate increased gene flow in wolves and other large carnivores, conservation managers should move towards population-level actions⁷⁵. This process should be supported by the scientific community by shifting the focus from national-level research towards broader studies coordinated at the level of biogeographic regions (e.g., <https://www.lifewolfalps.eu/>) or even the entire continent. For highly mobile species such as large carnivores, broad-scale approaches likely present the best option to reduce human-wildlife conflicts and ensure their sustainable future in the human-dominated landscapes of modern-day Europe. Tracking allelic richness across the spatial scales relevant for gene flow in wolves and other highly mobile mammals could thus provide a cost-effective measure for genetic monitoring, especially given its capacity to connect past and present data across multiple jurisdictions.

Data availability

Most of the data on wolf genetic diversity that we refer to in our analysis come from previously published studies, cited in Supplemental Table S2 and Supplemental Note S3. The new data we have presented from Dinaric and Carpathian wolves are available in the Repository of the University of Ljubljana (<https://repozitorij.uni-lj.si/IzpisGradiva.php?id=148315>).

Received: 19 January 2023; Accepted: 17 August 2023

Published online: 22 August 2023

References

1. Presson, A. P., Sobel, E., Lange, K. & Papp, J. C. Merging microsatellite data. *J. Comput. Biol.* **16**, 1131–1147 (2006).
2. Kaczensky, P. et al. Status, management and distribution of large carnivores—Bear, lynx, wolf & wolverine—in Europe. Part 1—Europe Summaries. Report: 1–72. A Large Carnivore Initiative for Europe Report prepared for the European Commission (2013).

3. de Groot, G. A. *et al.* Decades of population genetic research reveal the need for harmonization of molecular markers: The grey wolf *Canis lupus* as a case study. *Mammal Rev.* **46**, 44–59 (2006).
4. Iacolina, L., Corlatti, L., Bužan, E., Safner, T. & Šprem, N. Hybridisation in European ungulates: An overview of the current status, causes, and consequences. *Mammal Rev.* **49**, 45–59 (2019).
5. Rutkowski, R. *et al.* A European concern? Genetic structure and expansion of golden jackals (*Canis aureus*) in Europe and the Caucasus. *PLoS ONE* **10**(11), e0141236 (2015).
6. Le Moan, A., Gagnaire, P.-A. & Bonhomme, F. Parallel genetic divergence among coastal–marine ecotype pairs of European anchovy explained by differential introgression after secondary contact. *Mol. Ecol.* **25**, 3187–3202 (2016).
7. McDevitt, A. D. *et al.* Next-generation phylogeography resolves post-glacial colonization patterns in a widespread carnivore, the red fox (*Vulpes vulpes*), Europe. *Mol. Ecol.* **31**, 993–1006 (2022).
8. Stronen, A. V. *et al.* Wolf-dog admixture highlights the need for methodological standards and multidisciplinary cooperation for effective governance of wild x domestic hybrids. *Biol. Conserv.* **266**, 109467 (2022).
9. Putman, A. I. & Carbone, I. Challenges in analysis and interpretation of microsatellite data for population genetic studies. *Ecol. Evol.* **4**, 4399–4428 (2014).
10. Morales-González, A., Fernández-Gil, A., Quevedo, M. & Revilla, E. Patterns and determinants of dispersal in grey wolves (*Canis lupus*). *Biol. Rev.* **97**, 466–480 (2022).
11. Skrbinišek, T. *et al.* Using a reference population yardstick to calibrate and compare genetic diversity reported in different studies: an example from the brown bear. *Heredity* **109**, 299–305 (2012).
12. Allendorf, F. W. Genetic drift and the loss of alleles versus heterozygosity. *Zoo Biol.* **5**, 181–190 (1986).
13. DeWoody, J. A., Harder, A. M., Mathus, S. & Willoughby, J. R. The long-standing significance of genetic diversity in conservation. *Mol. Ecol.* **30**, 4147–4154 (2021).
14. Hoban, S. *et al.* Genetic diversity targets and indicators in the CBD post-2020 Global Biodiversity Framework must be improved. *Biol. Conserv.* **248**, 108654 (2020).
15. Hindrikson, M. *et al.* Wolf population genetics in Europe: A systematic review, meta-analysis and suggestions for conservation and management. *Biol. Rev.* **92**, 1601–1629 (2017).
16. Boitani, L. *et al.* *Assessment of the conservation status of the Wolf (Canis lupus) in Europe*. Report for the Convention on the Conservation of European Wildlife and Natural Habitats Standing Committee (2022). Prepared by the Large Carnivore Initiative for Europe. https://lciepub.nina.no/pdf/638036032684557257_LCIE%20CoE%20Wolf%20status%20report%202022.pdf.
17. Czarnomska, S. D. *et al.* Concordant mitochondrial and microsatellite DNA structuring between Polish lowland and Carpathian Mountain wolves. *Conserv. Genet.* **14**, 573–588 (2013).
18. Stronen, A. V. *et al.* Genome-wide analyses suggest parallel selection for universal traits may eclipse local environmental selection in a highly mobile carnivore. *Ecol. Evol.* **5**, 4410–4425 (2015).
19. Hulva, P. *et al.* Wolves at the crossroad: Fission–fusion range biogeography in the Western Carpathians and Central Europe. *Divers. Distrib.* **24**, 179–192 (2018).
20. Chapron, G. *et al.* Recovery of large carnivores in Europe’s modern human-dominated landscapes. *Science* **346**, 1517–1519 (2014).
21. Kutal, M. & Rigg, R. (eds). Perspectives of wolves in Central Europe. In *Proceedings from the Conference Held on 9th April 2008 in Malenovice, Beskydy Mts., Czech Republic*. Hnutí Duha, Olomouc 48 (2008).
22. Boitani, L. Wolf conservation and recovery. In *Wolves: Behaviour, Ecology, and Conservation* (eds Mech, L. D. & Boitani, L.) 317–340 (The University of Chicago Press, 2003).
23. Ericson, H. E. *et al.* Genome-wide profiles indicate wolf population connectivity within the eastern Carpathian Mountains. *Genetica* **148**, 33–39 (2020).
24. Bakan, J., Lavadinová, V., Popović, Z. & Paule, L. Genetic differentiation of grey wolf population (*Canis lupus* L.) from Balkan and Carpathians. *Balkan J. Wildl. Res.* **1**, 87–93 (2014).
25. Rigg, R., Skrbinišek, T. & Linnell, J. *Engaging hunters and other stakeholders in a pilot study of wolves in Slovakia using non-invasive genetic sampling*. Report to DG Environment, European Commission. Contract no. 07.0307/2013/654446/SER/B (2014).
26. Kaczensky, P. *et al.* *Distribution of large carnivores in Europe 2012–2016: Distribution maps for brown bear, Eurasian lynx, grey wolf, and wolverine*. Dryad, Dataset. <https://doi.org/10.5061/dryad.pc866t1p3> (2021)
27. Adamič, M., Jerina, K., Zafran, J. & Marinčič, A. *Izhodišča za oblikovanje strategije ohranitvenega upravljanja s populacijo volka (Canis lupus) v Sloveniji*. Biotehniška fakulteta, Ljubljana (2004).
28. Štrbenac, A. *et al.* *Plan upravljanja vukom u Hrvatskoj*. Državni zavod za zaštitu prirode, Zagreb (2005). https://mingor.gov.hr/UserDocsImages/NASLOVNE%20FOTOGRAFIJE%20I%20KORI%20C%20A%20TENI%20LOGOTIPOVI/doc/plan_upravljanja_vukom_u_republici_hrvatskoj_za_razdoblje_od_2010_do_2015.pdf.
29. Fabbri, E. *et al.* Genetic structure of expanding wolf (*Canis lupus*) populations in Italy and Croatia, and the early steps of the recolonization of the Eastern Alps. *Mamm. Biol.* **79**, 138–148 (2014).
30. Ražen, N. *et al.* Long-distance dispersal connects Dinaric-Balkan and Alpine grey wolf (*Canis lupus*) populations. *Eur. J. Wildl. Res.* **62**, 137–142 (2016).
31. Šnjegota, D., Stefanović, M., Veličković, N., Čirović, D. & Djan, M. Genetic characterization of grey wolves (*Canis lupus* L. 1758) from Bosnia and Herzegovina: Implications for conservation. *Conserv. Genet.* **19**, 755–760 (2018).
32. Djan, M. *et al.* Genetic variability and population structure of grey wolf (*Canis lupus*) in Serbia. *Russ. J. Genet.* **52**, 821–827 (2016).
33. Djan, M. *et al.* Genetic diversity and structuring of the grey wolf population from the Central Balkans based on mitochondrial DNA variation. *Mamm. Biol.* **79**, 277–282 (2014).
34. Jombart, T. Adegnet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**, 1403–1405 (2008).
35. R Core Team. (2020). *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, 2020), <https://www.R-project.org/>.
36. Paradis, E. Pegas: An R package for population genetics with an integrated-modular approach. *Bioinformatics* **26**, 419–420 (2010).
37. Holm, S. A simple sequentially rejective multiple test procedure. *Scand. J. Stat.* **6**, 65–70 (1979).
38. vonHoldt, B. M. *et al.* A genome-wide perspective on the evolutionary history of enigmatic wolf-like canids. *Genome Res.* **21**, 1294–1305 (2011).
39. Pilot, M. *et al.* Genome-wide signatures of population bottlenecks and diversifying selection in European wolves. *Heredity* **112**, 428–442 (2014).
40. Potočnik, H. *et al.* *Projektno poročilo za Akcijo C1(LIFE08 NAT/SLO/000244 SloWolf): Spremljanje stanja populacije volka v Sloveniji (3). 1., 2. in 3. sezona—2010/11, 2011/12 in 2012/13 (2014). Projekt LIFE+ SloWolf* (Ljubljana, 2014). https://www.volkovi.si/wp-content/uploads/2014/10/porocilo_c1_koncno.pdf
41. Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W. & Prodöhl, P. A. diveRsity: An R package for the estimation of population genetics parameters and their associated errors. *Methods Ecol. Evol.* **4**, 782–788 (2013).
42. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2016).
43. QGIS. *QGIS Geographic Information System*. Open Source Geospatial Foundation Project (2022). <http://qgis.org>.
44. Aspi, J., Roininen, E., Ruokonen, M., Kojola, I. & Vilà, C. Genetic diversity, population structure, effective population size and demographic history of the Finnish wolf population. *Mol. Ecol.* **15**, 1561–1576 (2006).
45. Godinho, R. *et al.* Genetic evidence for multiple events of hybridization between wolves and domestic dogs in the Iberian Peninsula. *Mol. Ecol.* **20**, 5154–5166 (2011).

46. Sastre, N. *et al.* Signatures of demographic bottlenecks in European wolf populations. *Conserv. Genet.* **12**, 701–712 (2010).
47. Aspi, J. *et al.* Genetic structure of the northwestern Russian wolf populations and gene flow between Russia and Finland. *Conserv. Genet.* **10**, 815–826 (2009).
48. Jansson, E., Ruokonen, M., Kojola, I. & Aspi, J. Rise and fall of a wolf population: Genetic diversity and structure during recovery, rapid expansion and drastic decline. *Mol. Ecol.* **21**, 5178–5193 (2012).
49. Silva, P. *et al.* Cryptic population structure reveals low dispersal in Iberian wolves. *Sci. Rep.* **8**, 14108 (2018).
50. Lucchini, V., Galov, A. & Randi, E. Evidence of genetic distinction and long-term population decline in wolves (*Canis lupus*) in the Italian Apennines. *Mol. Ecol.* **13**, 523–536 (2004).
51. Moura, A. E. *et al.* Unregulated hunting and genetic recovery from a severe population decline: The cautionary case of Bulgarian wolves. *Conserv. Genet.* **15**, 405–417 (2014).
52. Pilot, M. *et al.* Genetic variability of the grey wolf *Canis lupus* in the Caucasus in comparison with Europe and the Middle East: Distinct or intermediary population?. *PLoS ONE* **9**(4), e93828 (2014).
53. Leberg, P. L. Estimating allelic richness: Effects of sample size and bottlenecks. *Mol. Ecol.* **11**, 2445–2449 (2008).
54. Petit, R. J., El Mousadik, A. & Pons, O. Identifying populations for conservation on the basis of genetic markers. *Conserv. Biol.* **12**, 844–855 (1998).
55. Wahlund, S. Zusammensetzung von populationen und korrelationserscheinungen vom standpunkt der vererbungslehre aus betrachtet. *Hereditas* **11**, 65–106 (1928).
56. Wabakken, P. *et al.* Multistage, long-range natal dispersal by a global positioning system–collared Scandinavian wolf. *J. Wildl. Manag.* **71**, 1631–1634 (2007).
57. Lesniak, I. *et al.* Population expansion and individual age affect endoparasite richness and diversity in a recolonising large carnivore population. *Sci. Rep.* **7**, 41730 (2017).
58. Andersen, L. W. *et al.* Long-distance dispersal of a wolf, *Canis lupus*, in northwestern Europe. *Mamm. Res.* **60**, 163–168 (2015).
59. Hindrikson, M. *et al.* (2013) Spatial genetic analyses reveal cryptic population structure and migration patterns in a continuously harvested grey wolf (*Canis lupus*) population in North-Eastern Europe. *PLoS ONE* **8**(9), e75765 (2013).
60. Wright, S. The genetical structure of populations. *Ann. Eugen.* **15**, 323–354 (1951).
61. Boitani, L. *et al.* Key actions for large carnivore populations in Europe. Institute of Applied Ecology, Rome, Italy. Report to DG Environment, European Commission, Bruxelles (2015). Contract no. 07.0307/2013/654446/SER/B3.
62. Niedziałkowska, M. *et al.* Molecular biogeography of red deer *Cervus elaphus* from eastern Europe: Insights from mitochondrial DNA sequences. *Acta Theriol.* **56**(1), 1–12 (2011).
63. Veličković, N. *et al.* Demographic history, current expansion and future management challenges of wild boar populations in the Balkans and Europe. *Heredity* **117**, 348–357 (2016).
64. Kusak, J. *et al.* Wolf-dog hybridization in Croatia. *Vet. Arhiv* **88**, 375–395 (2018).
65. Fabbri, E. *et al.* From the Apennines to the Alps: Colonization genetics of the naturally expanding Italian wolf (*Canis lupus*) population. *Mol. Ecol.* **16**, 1661–1671 (2007).
66. Karamanlidis, A. A. *et al.* Wolf population genetics at the south-eastern edge of their European range. *Mamm. Biol.* **81**, 506–510 (2016).
67. Szweczyk, M. *et al.* Dynamic range expansion leads to establishment of a new, genetically distinct wolf population in Central Europe. *Sci. Rep.* **9**, 19003 (2019).
68. Pilot, M. *et al.* Ecological factors influence population genetic structure of European grey wolves. *Mol. Ecol.* **15**, 4533–4553 (2006).
69. Allendorf, F. W. & Luikart, G. *Conservation and the Genetics of Populations* (Wiley, 2009).
70. Schweizer, R. M. *et al.* Genetic subdivision and candidate genes under selection in North American grey wolves. *Mol. Ecol.* **25**, 380–402 (2016).
71. Werhahn, G. *et al.* The unique genetic adaptation of the Himalayan wolf to high-altitudes and consequences for conservation. *Glob. Ecol. Conserv.* **16**, e00455 (2018).
72. Exposito-Alonso, M. *et al.* Genomic basis and evolutionary potential for extreme drought adaptation in *Arabidopsis thaliana*. *Nat. Ecol. Evol.* **2**, 352–358 (2018).
73. Koen, E. L., Bowman, J., Murray, D. L. & Wilson, P. J. Climate change reduced genetic diversity of Canada lynx at the trailing range edge. *Ecography* **37**, 754–762 (2014).
74. Hughes, J. B., Daily, G. C. & Ehrlich, P. R. Population diversity: Its extent and extinction. *Science* **278**, 689–692 (1997).
75. Linnell, J.D.C., Salvatori, V. & Boitani, L. *Guidelines for population level management plans for large carnivores in Europe*. A Large Carnivore Initiative for Europe report prepared for the European Commission (contract 070501/2005/424162/MAR/B2) Large carnivore initiative for Europe, Rome (2008). https://ec.europa.eu/environment/nature/conservation/species/carnivores/pdf/guidelines_for_population_level_management.pdf

Acknowledgements

The work in Slovenia was co-financed and supported by the Ministry of the Environment and Spatial Planning of the Republic of Slovenia, by the Slovenian Research and Innovation Agency through the Research Programs P1-0184 and P4-0059, and by the European Commission through projects LIFE SLOWOLF (LIFE08 NAT/SLO/000244) and LIFE WOLFALPS EU (LIFE18 NAT/IT/000972). We would like to thank Slovenia Forest Service personnel for help with sample collection and the Dinaricum Society and volunteers for help in collecting noninvasive wolf samples in the scope of several yearly wolf monitoring sessions in Slovenia. The work in Croatia was funded by Bernd Thies Foundation, UK Wolf Conservation Trust, EURONATUR, Croatian Agency for Nature and Environment, “Sjeverni Velebit” National Park, “Velebit” Nature Park and “Plitvice Lakes” National Park. We thank all members of the Large Carnivores Emergency Team of Croatia, game wardens and all others who helped in the sample collection. In Slovakia, the work was financed by the European Commission (contract no. 07.0307/2013/654446/SER/B.3) through the Institute of Applied Ecology, Rome. We thank the Slovak Wildlife Society, the Slovak Hunting Association, Forests of the Slovak Republic state enterprise, the State Nature Conservancy, the University of Ljubljana, the Wolves and Humans Foundation and the IUCN/SSC Large Carnivore Initiative for Europe. For the work in Romania, we thank rangers and volunteers from Foundation Conservation Carpathia, Piatra Craiului National Park and Bârsa Hunting Association for help with sample collection. Romanian wolves were sampled with support from the OAK Foundation grant number OCAY-11-136, and from the European Commission through the Operational Programme ‘Environment’, grant number SMIS 102086.

Author contributions

M.J., B.B., R.Č., Đ.H., R.I., F.K., I.K., M.Kr., J.K., B.P.F., H.P., R.R., T.S. organized sampling and helped in organizing the sample collection. M.J., B.B. and M.Ko. carried out the laboratory analysis and organized the data. M.J.,

R.L. and T.S. ran analysis. M.J. wrote the original draft. T.S. prepared the figures and the map. M.J., A.V.S. and T.S. wrote and edited the manuscript. R.R. provided language review. P.T. provided consultation and support. A.M.S., T.S., H.P., I.K., M.Kr., B.P.F. and R.R. contributed essential parts of the results via ancillary research projects. All authors contributed to writing through reviewing and commenting on the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-40834-x>.

Correspondence and requests for materials should be addressed to M.J.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023