



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Ecophysiology of novel core phylotypes in activated sludge wastewater treatment plants with nutrient removal

McIlroy, Simon Jon; Awata, Takanori; Nierychlo, Marta; Saunders, Aaron Marc; Albertsen, Mads; Szyszka, Anna; Starnawski, Piotr Marian; Kindaichi, Tomonori; Nielsen, Jeppe Lund; Nielsen, Per Halkjær

Publication date:
2014

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

McIlroy, S. J., Awata, T., Nierychlo, M., Saunders, A. M., Albertsen, M., Szyszka, A., Starnawski, P. M., Kindaichi, T., Nielsen, J. L., & Nielsen, P. H. (2014). *Ecophysiology of novel core phylotypes in activated sludge wastewater treatment plants with nutrient removal*. Poster presented at 15th International Symposium on Microbial Ecology, Seoul, Korea, Republic of.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Ecophysiology of novel core phylotypes in activated sludge wastewater treatment plants with nutrient removal

Simon Jon McIlroy¹, Takanori Awata^{1,2}, Marta Nierychlo¹, Aaron Marc Saunders¹, Mads Albertsen¹, Anna Szyszka¹, Piotr Starnawski¹, Tomonori Kindaichi², Jeppe Lund Nielsen¹ & Per Halkjær Nielsen¹

¹ Center for Microbial Communities, Section of Biotechnology, Aalborg University, Aalborg, Denmark
² Department of Civil Engineering, Graduate School of Engineering, Hiroshima University, Higashihiroshima, Japan



Introduction

An understanding of the ecology of activated sludge (AS) biological nutrient removal (BNR) wastewater treatment plants requires detailed knowledge of the community composition and metabolic activities of individual members. Recent 16S rRNA gene amplicon surveys of full-scale systems in Denmark indicate a core set of bacterial genera. These core genera are suggested to be responsible for the bulk of nutrient transformations underpinning the functions of these plants. While we know the basic *in situ* activities of some of these genera, there is little to no information for the majority, and thus no indication as to their relevance to the ecology of these systems.


Aim

To characterise the ecophysiology of core genera in full-scale BNR plants with *in situ* based methods.

Methods


Stable isotope probing

Stable isotope probing was applied under nitrite reducing conditions with a fully ¹³C labelled complex media to assist in the identification of putative active core denitrifiers.



MAR-FISH

Eight core phylotypes were selected and fluorescent *in situ* hybridisation (FISH) was applied in combination with microautoradiography (MAR) to assess the ability of these organisms for key metabolic activities. New FISH probes were optimised for 7 of the phylotypes.



Results




Fig. 1. FISH images of phylotypes in full-scale AS. Scale bars = 20 µm.

Table 1. Phylogeny of selected core genera

Classification				
Phylum	Class	Order	Family	Genus
Proteobacteria;	β-proteobacteria	Burkholderiales	Comamonadaceae	<i>Rhodoferax</i>
Proteobacteria	β-proteobacteria	Rhodocyclales	Rhodocyclaceae	<i>Sulfuritalea</i>
Proteobacteria	β-proteobacteria	Rhodocyclales	Rhodocyclaceae	<i>Dechloromonas</i>
Proteobacteria	β-proteobacteria	SC-1-84	SC-1-84	<i>A21b</i>
Proteobacteria	γ-proteobacteria	Xanthomonadales	Xanthomonadaceae	<i>Thermomonas</i>
Proteobacteria	δ-proteobacteria	Myxococcales	Haliangiaceae	<i>Haliangium</i>
Acidobacteria	Holophagae	Sub-group 10	43F-1404R	<i>AKAB1</i>
Chloroflexi	Caldilineae	Caldilineales	CCD21	<i>B45</i>

Table 2. Summary of *in situ* characterisation of selected phylotypes

	<i>Rhodoferax</i>	<i>Sulfuritalea</i>	<i>Dechloromonas</i>	<i>A21b</i> (β-proteobacteria)	<i>Thermomonas</i>	<i>Haliangium</i>	<i>AKAB1</i> (Holophagae)	<i>B45</i> (Caldilineae)
Metabolic feature								
Substrate uptake								
Pyruvate	+	+	+	+	N.D.	N.D.	-	-
Acetate	+	+	+	-	+	-	-	-
Propionate	+	±	N.D.	-	N.D.	N.D.	-	-
Butyrate	+	-	N.D.	-	N.D.	N.D.	-	-
Oleate	-	-	-	-	N.D.	N.D.	-	-
Amino acids	+	+	+	-	+	+	-	-
Glucose	±	±	-	-	N.D.	N.D.	-	+
Glycerol	±	-	N.D.	-	N.D.	N.D.	-	-
Ethanol	-	-	-	-	N.D.	N.D.	-	-
N-acetylglucosamine	-	-	N.D.	-	N.D.	N.D.	-	-
Anaerobic carbon uptake	±	+	+	+	±	-	-	+
Denitrification	+	+	+	-	+	+	-	N.D.
Autotrophy (incl. oxidation of: NH ₄ ⁺ ; NO ₂ ⁻ ; H ₂ ; S ₂ O ₃ ²⁻)	N.D.	-	N.D.	-	N.D.	N.D.	-	N.D.
Polyphosphate storage	-	N.D.	±	N.D.	N.D.	N.D.	N.D.	-

+ = positive; - = negative; ± = varied; N.D. = Not determined.




Fig. 2. Boxplot of the 50 most abundant OTUs (clustered at 94 %) based on 16S rRNA gene amplicon sequencing surveys of 13 full-scale BNR plants (Saunders *et al.* submitted). Selected genera studied here are in bold. Dots denote outliers. OTU labels are the lowest assigned taxonomic rank (x-axis). Genera for which *in situ* ecophysiological information is available are colour coded by their defining traits (acknowledging that each genera may fit into to multiple categories).

Denitrifier, Ammonia oxidiser, Saprophyte, Unresolved, Fermenter, Lipid accumulator, GAO, PAO

Conclusions

Members of the *Rhodoferax*, *Sulfuritalea*, *Thermomonas* and *Haliangium* appear to be core denitrifiers. Members of the novel genera selected have more specialised physiologies requiring further investigation. Future work will focus on characterisation of additional ‘core’ genera, including genomic based approaches. The ecophysiology profiles generated will be made available on the ‘Microbial Database of Activated Sludge’ (MiDAS) field guide website (midasfieldguide.org); a public resource for the benefit of individuals interested in AS.



sjm@bio.aau.dk



www.cmc.aau.dk



Environmental Biotechnology
Aalborg University