



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The Quality of Prescribing for Psychiatric Patients

Sørensen, Ann Lykkegaard; Nielsen, Lars Peter; Poulsen, Birgitte Klindt; Lisby, Marianne; Mainz, Jan

Published in:
Value in Health

Publication date:
2014

Document Version
Tidlig version også kaldet pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Sørensen, A. L., Nielsen, L. P., Poulsen, B. K., Lisby, M., & Mainz, J. (2014). The Quality of Prescribing for Psychiatric Patients. *Value in Health*, 17(7).

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

The Quality of Prescribing for Psychiatric Patients

Soerensen AL^{1,2}, Nielsen LP^{3,4}, Poulsen BK³, Lisby M^{3,5}, Mainz J^{6,7}

¹Danish Center for Healthcare Improvements, Faculty of Social Sciences and Faculty of Health Sciences, Aalborg University, Denmark; ²University College of Northern Denmark; ³Department of Clinical Pharmacology, Aarhus University Hospital, Aarhus, Denmark; ⁴Institute of Biomedicine, Aarhus University, Aarhus; ⁵Centre of Emergency Medicine Research, Aarhus University Hospital & Aarhus University, Aarhus, Denmark; ⁶Aalborg University Hospital, Psychiatry and ⁷Aalborg University, Aalborg; Denmark

OBJECTIVES: Prescribing for adult psychiatric patients is often highly complex due to the nature of psychiatric conditions, but also due to somatic comorbidity. Therefore, the aim of this study was to identify prevalence and types of potential inappropriate prescribing (PIP), assess the severity of potential clinical consequences and identify possible predictive factors of PIP.

METHODS: The study was designed as a prospective study of PIP using medication reviews. Patients who were admitted during a 4 month period (August 2013 - November 2013) to a psychiatric university hospital were included (n=219). The medication reviews, including an assessment of potential severity, were carried out by clinical pharmacologists after admission and after the attending physician had seen the patient. Frequencies and categories of PIP were analyzed in absolute numbers and as percentages. Severity of PIP was assessed using four categories. Logistic regression analysis was used to identify possible predictive factors of PIP.

RESULTS: The proportion of patients with one or more PIPs was 123/219(56%). “Interaction between drugs” was the most common category for potentially serious and potentially fatal PIPs with 49/123(40%) and 32/45(71%), respectively. Of 32 identified potentially fatal drug-drug interactions, 15/32(47%) involved two or more antipsychotic drugs and 12/32(37%) involved antipsychotic drugs in combination with antidepressants. The remaining 5/32(16%) potentially fatal drug-drug interactions involved propranolol, erythromycin, simvastatin and promethazine. After adjusting for age, gender, alcohol/substance abuse, number of prescriptions, number of somatic diagnoses and level of kidney function, only polypharmacy (>5 prescriptions) increased the odds for a PIP significantly; OR=4,82(95% CI:2.33-9.98), p<0.0001.

CONCLUSIONS: PIP is frequent and might have serious or fatal consequences. Special attention should be given to drug-drug interactions involving antipsychotics and antidepressants but also somatic medications and polypharmacy threatens medication safety. There is a pressing need to improve the quality in prescribing for psychiatric patients.