

## An inequality of rearrangements on the unit circle

Draghici, Cristina

*Publication date:*  
2005

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Draghici, C. (2005). *An inequality of rearrangements on the unit circle*. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2005-31

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

AALBORG UNIVERSITY

**An inequality of rearrangements  
on the unit circle**

by

Cristina Draghici

R-2005-31


October 2005

DEPARTMENT OF MATHEMATICAL SCIENCES  
AALBORG UNIVERSITY

Fredrik Bajers Vej 7 G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 96 35 80 80 ■ Telefax: +45 98 15 81 29

URL: <http://www.math.aau.dk>



# AN INEQUALITY OF REARRANGEMENTS ON THE UNIT CIRCLE

CRISTINA DRAGHICI

ABSTRACT. We prove that the integral of the product of two functions over a symmetric set in  $\mathbb{S}^1 \times \mathbb{S}^1$ , defined as  $E = \{(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma_1(x), \sigma_2(y)) \leq \alpha\}$ , where  $\sigma_1, \sigma_2$  are diffeomorphisms of  $\mathbb{S}^1$  with certain properties and  $d$  is the geodesic distance on  $\mathbb{S}^1$ , increases when we pass to their symmetric decreasing rearrangement. We also give a characterization of these diffeomorphisms  $\sigma_1, \sigma_2$  for which the rearrangement inequality holds. As a consequence, we obtain the result for the integral of the function  $\Psi(f(x), g(y))$  ( $\Psi$  a supermodular function) with a kernel given as  $k[d(\sigma_1(x), \sigma_2(y))]$ , with  $k$  decreasing.

## 1. INTRODUCTION

On a measure space  $(X, \mu)$ , the Hardy-Littlewood inequality asserts [4]:

$$\int_X f(x)g(x) d\mu(x) \leq \int_0^{\mu(X)} f^*(t)g^*(t) dt,$$

where  $f^*$  and  $g^*$  are the decreasing rearrangements of  $f$  and  $g$ , respectively. In what follows,  $X = \mathbb{S}^1$ , or  $X = [-\pi, \pi]$ , and the above inequality can be written as:

$$(1.1) \quad \int_{-\pi}^{\pi} f(x)g(x) dx \leq \int_{-\pi}^{\pi} f^\sharp(x)g^\sharp(x) dx,$$

with  $f^\sharp, g^\sharp$  the symmetric decreasing rearrangements of  $f$  and  $g$ , given by  $f^\sharp(x) = f^*(2|x|)$  and  $g^\sharp(x) = g^*(2|x|)$ .

These inequalities can be proved using *the layer-cake formula* [10]: Every measurable function  $f : X \rightarrow \mathbb{R}_+$  can be written as an integral of the characteristic function of its level sets:

$$(1.2) \quad f(x) = \int_0^\infty \chi_{\{f>t\}}(x) dt.$$

A more general rearrangement inequality on  $X = \mathbb{R}^n$  is the Riesz-Sobolev inequality:

$$(1.3) \quad \int_{\mathbb{R}^{2n}} f(x)g(y)h(x-y) dx dy \leq \int_{\mathbb{R}^{2n}} f^\sharp(x)g^\sharp(y)h^\sharp(x-y) dx dy,$$

where  $f, g, h$  are non-negative functions which vanish at infinity in a weak sense. The case  $n = 1$  is due to Riesz in 1930 (see [12]), and the case  $n > 1$  is due to Sobolev in 1938 (see [13]). The proof can be found in the book by Hardy, Littlewood, Pólya [9] which sets the beginning of the systematic study of rearrangement inequalities. A more general version of this inequality in  $\mathbb{R}^n$ , involving  $n$  functions can be found in [5].

---

1991 *Mathematics Subject Classification.* 26D15, 28A25, 52A10.

*Key words and phrases.* symmetrization, rearrangement, circle integral inequality, convolution.

The equivalent of (1.3) for three non-negative functions on the unit circle was proved by Baernstein [1]:

$$(1.4) \quad \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} f(e^{i\phi})g(e^{i\theta})h(e^{i(\phi-\theta)}) d\theta d\phi \leq \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} f^{\sharp}(e^{i\phi})g^{\sharp}(e^{i\theta})h^{\sharp}(e^{i(\phi-\theta)}) d\theta d\phi.$$

The proof of this inequality uses a variational principle applied to the convolution of characteristic functions of sets which does not seem to generalize in higher dimensions.

The Riesz-Sobolev inequality (1.3) is equivalent to the Brunn-Minkowski inequality from convex geometry [8, 11, 7] which states that if  $K$  and  $L$  are measurable sets in  $\mathbb{R}^n$ , then their Minkowski (pointwise) sum  $K + L$  is related to the measure of the sets  $K$  and  $L$  by

$$V(K + L)^{1/n} \geq V(K)^{1/n} + V(L)^{1/n},$$

where  $V$  denotes the  $n$ -dimensional volume. An analog of this inequality for  $\mathbb{S}^n$  is not known, and, since the proof of rearrangement inequalities in  $\mathbb{R}^n$  require it, an analog of the Riesz-Sobolev inequality (1.3) is not known in  $\mathbb{S}^n$ , for  $n > 1$ .

However, a partial result in  $\mathbb{S}^n$  was proved by Baernstein and Taylor in [2]. They considered a version of the Riesz-Sobolev inequality where one of the functions is symmetric decreasing. They showed that, if  $h = K$  is already symmetric decreasing then

$$\int_{\mathbb{S}^n} \int_{\mathbb{S}^n} f(x)g(y)K(x \cdot y) d\sigma(x)d\sigma(y) \leq \int_{\mathbb{S}^n} \int_{\mathbb{S}^n} f^{\sharp}(x)g^{\sharp}(y)K(x \cdot y) d\sigma(x)d\sigma(y),$$

where  $d\sigma$  is the surface measure on the unit sphere  $\mathbb{S}^n$  in  $\mathbb{R}^{n+1}$ ,  $x \cdot y$  is the usual inner product and  $K(t)$  is an increasing function on  $[-1, 1]$ . Since  $x \cdot y = \cos \alpha$ , where  $\alpha$  is the angle between the vectors  $x$  and  $y$ , we can write  $K(x \cdot y) = k(d(x, y))$ , with  $k$  decreasing. Here  $d(x, y)$  is the great circle (geodesic) distance between  $x$  and  $y$ . Their proof is based on the polarization technique. They showed first that the inequality holds for the polarizations of  $f$  and  $g$  in any hyperplane and then they passed to the limit for the general case. They were led to this version of the Riesz-Sobolev inequality while trying to generalize a 2-dimensional result stating that  $u$  is subharmonic implies its star function is also subharmonic.

In this paper we are interested in the case  $n = 1$  of this inequality with  $K$  replaced by the characteristic function of a symmetric set which does not depend on the distance between two points, but rather on the distance between their images under two diffeomorphisms  $\sigma_1, \sigma_2$  of  $\mathbb{S}^1$ . We will also obtain a characterization of these diffeomorphisms for which the inequality holds. With the set  $E$  defined as

$$E = \{(x, y) : d(\sigma_1(x), \sigma_2(y)) \leq \alpha\},$$

we will show that

$$(1.5) \quad \int_E f(x)g(y) dx dy \leq \int_E f^{\sharp}(x)g^{\sharp}(y) dx dy,$$

for every  $\alpha > 0$ . This result implies *the main result* of this paper, Theorem 3.6:

$$\begin{aligned} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f(x), g(y))k[d(\sigma_1(x), \sigma_2(y))] dx dy \\ \leq \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f^{\sharp}(x), g^{\sharp}(y))k[d(\sigma_1(x), \sigma_2(y))] dx dy, \end{aligned}$$

with  $k$  decreasing and  $\Psi$  the distribution function of a measure  $\mu$ .

The paper is organized as follows: We will first prove (1.5) for  $f$  and  $g$  replaced by characteristic functions  $\chi_A$ ,  $\chi_B$ , and  $\sigma_2$  the identity. Then we will deduce the result (1.5) mentioned above, and we will show that we can replace the product  $f(x)g(y)$  by a function  $\Psi(f(x), g(y))$  and that we can replace  $\chi_E$  by a decreasing function of the distance between  $\sigma_1(x)$  and  $\sigma_2(y)$ , yielding Theorem 3.6.

## 2. PRELIMINARIES

Recall that a function  $f : I \rightarrow \mathbb{R}$ , defined on an interval  $I \subset \mathbb{R}$ , is called convex if, for every  $0 < \lambda < 1$  and every  $a, b \in I$ , the following inequality holds:

$$f(\lambda a + (1 - \lambda)b) \leq \lambda f(a) + (1 - \lambda)f(b).$$

A convex function is differentiable almost everywhere on  $I$  and its derivative is increasing.

We denote by  $\mathbb{S}^1$  the unit circle in  $\mathbb{R}^2$ , i.e.,  $\mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$ , and by  $\mathbb{S}_+^1$  the upper half unit circle,

$$\mathbb{S}_+^1 = \{e^{i\theta} : 0 \leq \theta \leq \pi\}.$$

**Definition 2.1.** A function  $\sigma : \mathbb{S}_+^1 \rightarrow \mathbb{S}_+^1$  is called convex if the function  $\sigma_1 : [0, \pi] \rightarrow [0, \pi]$ , defined as :

$$\sigma(e^{i\theta}) = e^{i\sigma_1(\theta)}, \quad 0 \leq \theta \leq \pi,$$

is convex on  $[0, \pi]$ .

Let  $f : \mathbb{S}^1 \rightarrow \mathbb{R}_+$  be a non-negative measurable function. We define its *distribution function*:

$$\lambda_f(t) = |\{f > t\}|, \quad t \in [0, \infty),$$

where  $\{f > t\} := \{z \in \mathbb{S}^1 : f(z) > t\}$  denote the level sets of  $f$ , and  $|A|$  is the linear measure on  $\mathbb{S}^1$  of  $A$ . Functions which have the same distribution function are called *equimeasurable*.

We define the *symmetric decreasing rearrangement* of  $f$  to be the function  $f^\sharp : \mathbb{S}^1 \rightarrow \mathbb{R}_+$ , given by:

$$f^\sharp(z) = \inf\{t : \lambda_f(t) \leq 2d(1, z)\},$$

where  $d(1, z)$  is the geodesic distance on  $\mathbb{S}^1$  between  $z$  and 1.

It is clear that  $f^\sharp(z) = f^\sharp(\bar{z})$  and that  $f^\sharp$  decreases as  $d(1, z)$  increases. Also,  $f$  and  $f^\sharp$  are equimeasurable.

If we write  $z = e^{i\theta}$ ,  $-\pi \leq \theta < \pi$ , then  $d(1, z) = d(1, e^{i\theta}) = |\theta|$ , and we can think of  $f$  as a function of  $\theta$  via the relation

$$\tilde{f}(\theta) = f(e^{i\theta}).$$

For  $\tilde{f} : [-\pi, \pi] \rightarrow \mathbb{R}_+$ , one defines its symmetric decreasing rearrangement as:

$$\tilde{f}^\sharp(\theta) = \inf\{t : \lambda_{\tilde{f}}(t) \leq 2|\theta|\},$$

where, as before,  $\lambda_{\tilde{f}}(t) = |\{\tilde{f} > t\}|$ , and thus, there is a one-to-one correspondence between  $f^\sharp$  and  $\tilde{f}^\sharp$ , given by

$$\tilde{f}^\sharp(\theta) = f^\sharp(e^{i\theta}).$$

Whenever necessary, we will think of a function  $f$  defined on  $\mathbb{S}^1$  as a function on  $[-\pi, \pi]$ . If  $f = \chi_A$  is the characteristic function of a measurable set  $A \subset \mathbb{S}^1$ , then

$f^\sharp = \chi_{A^\sharp}$ , where  $A^\sharp$  is the open interval on the unit circle centered at 1, having the same linear measure as  $A$ .

Next, we introduce the Hardy-Littlewood-Pólya preorder relation  $\prec$  for non-negative functions defined on the interval  $[-\pi, \pi]$ . We say that (see [3, 4]):

$$f \prec F \quad \text{iff} \quad \int_{-t}^t f^\sharp(s) ds \leq \int_{-t}^t F^\sharp(s) ds, \quad \text{for all } 0 \leq t \leq \pi.$$

This is equivalent to

$$\int_{-\pi}^{\pi} f^\sharp(s) h^\sharp(s) ds \leq \int_{-\pi}^{\pi} F^\sharp(s) h^\sharp(s) ds,$$

for every positive symmetric decreasing function  $h^\sharp$  defined on  $[-\pi, \pi]$ . To see this, write  $h^\sharp(s) = \int_0^\infty \chi_{\{h^\sharp > t\}}(s) dt$  (this is the layer cake formula (1.2)), and, using Fubini's formula and the fact that  $\{h^\sharp > t\} = (-l(t), l(t))$  is a symmetric interval,

$$\begin{aligned} \int_{-\pi}^{\pi} f^\sharp(s) h^\sharp(s) ds &= \int_0^\infty \left[ \int_{-l(t)}^{l(t)} f^\sharp(s) ds \right] dt \\ &\leq \int_0^\infty \left[ \int_{-l(t)}^{l(t)} F^\sharp(s) ds \right] dt = \int_{-\pi}^{\pi} F^\sharp(s) h^\sharp(s) ds. \end{aligned}$$

Yet another equivalent characterization is:

$$f \prec F \Leftrightarrow \int_E f(s) ds \leq \int_E F(s) ds, \quad \text{for every } E \subset [-\pi, \pi].$$

The next result is well-known and it follows from the proof of the equality case in the Hardy-Littlewood inequality, presented by Lieb and Loss in [10, pp.82]. We will include a proof here for consistency.

**Lemma 2.2.** *Let  $f : [-\pi, \pi] \rightarrow \mathbb{R}_+$  be a measurable function such that*

$$(2.1) \quad \int_{-t}^t f(x) dx \geq \int_{-t}^t f^\sharp(x) dx, \quad \text{for every } 0 \leq t \leq \pi.$$

*Then  $f = f^\sharp$  a.e. on  $[-\pi, \pi]$ .*

*Proof.* From (1.1) applied to  $\chi_{(-t,t)}$  and  $f$ , it follows that we must have equality in (2.1), i.e.,

$$(2.2) \quad \int_{-t}^t f(x) dx = \int_{-t}^t f^\sharp(x) dx.$$

We will use the layer-cake formula to write  $f(x) = \int_0^\infty \chi_{\{f > s\}}(x) ds$ , and similarly for  $f^\sharp(x)$ .

Using (1.1), we obtain:

$$(2.3) \quad \int_{-t}^t \chi_{\{f > s\}}(x) dx \leq \int_{-t}^t \chi_{\{f^\sharp > s\}}(x) dx, \quad \text{for every } s \geq 0.$$

Fubini's theorem and (2.2) imply that:

$$\begin{aligned} \int_{-t}^t f(x) dx &= \int_0^\infty \left[ \int_{-t}^t \chi_{\{f > s\}}(x) dx \right] ds \\ &= \int_0^\infty \left[ \int_{-t}^t \chi_{\{f^\sharp > s\}}(x) dx \right] ds = \int_{-t}^t f^\sharp(x) dx. \end{aligned}$$

From this equality and (2.3) it follows that, for a fixed  $t$ , there exists a set of measure zero  $S_t$ , such that

$$\int_{-t}^t \chi_{\{f>s\}}(x) dx = \int_{-t}^t \chi_{\{f^\#>s\}}(x) dx, \quad \text{for every } s \in (0, \infty) \setminus S_t.$$

Next, we choose  $T_N$  a countable dense set in  $[0, \pi]$  and we denote by  $S_{T_N} = \cup_{t \in T_N} S_t$ . Then:

$$(2.4) \quad \int_{-t}^t \chi_{\{f>s\}}(x) dx = \int_{-t}^t \chi_{\{f^\#>s\}}(x) dx, \quad \text{for every } t \in T_N \text{ and } s \in (0, \infty) \setminus S_{T_N}.$$

Since for every fixed  $s$ ,  $t \rightarrow \int_{-t}^t \chi_{\{f>s\}}(x) dx$  is a continuous function of  $t$ , in fact (2.4) holds for every  $0 \leq t \leq \pi$ . Thus,

$$\int_{-t}^t \chi_{\{f>s\}}(x) dx = \int_{-t}^t \chi_{\{f^\#>s\}}(x) dx, \quad \text{for all } 0 \leq t \leq \pi \text{ and a.e. } s \in (0, \infty).$$

Now, let  $t$  be such that  $\{f^\# > s\} = (-t, t)$ . Then, it follows that  $\{f > s\} = (-t, t) = \{f^\# > s\}$  a.e., and thus,  $f = f^\#$  by the layer cake formula.  $\square$

The following result shows that  $\int_{-t}^t f^\#(x) dx$  is attained as a supremum. A proof can be found in [4, Theorem 7.5, pp.82].

**Theorem 2.3.** (*J. V. Ryff*) *For every measurable function  $f$  as in Lemma 2.2, there exists a measure preserving transformation  $T$  such that  $f = f^\# \circ T$ . This guarantees, for every  $t$ , the existence of a set  $A \subset [-\pi, \pi]$  of measure  $2t$  such that  $\int_A f(x) dx = \int_{-t}^t f^\#(x) dx$ .*

### 3. MAIN RESULTS: INEQUALITIES ON THE CIRCLE

**Notation.** As before,  $d$  is the geodesic distance, also called the arclength, on the unit circle  $\mathbb{S}^1$ . We have:

$$(3.1) \quad d(u, v) = d(u\bar{v}, 1), \quad \text{for all } u, v \in \mathbb{S}^1,$$

where  $\bar{v}$  denotes the complex conjugate of  $v$ .

We define, for  $\alpha > 0$ , the function:

$$\chi_\alpha(u, v) = \begin{cases} 1, & \text{if } d(u, v) \leq \alpha, \\ 0, & \text{otherwise} \end{cases}$$

and we observe that  $\chi_\alpha(u, v) = \chi_\alpha(u\bar{v}, 1)$ , by (3.1).

We introduce a new function, which we call again  $\chi_\alpha : \mathbb{S}^1 \rightarrow \mathbb{R}_+$ , given by  $\chi_\alpha(z) := \chi_\alpha(z, 1)$ , which is the characteristic function of the closed interval on  $\mathbb{S}^1$  of linear length  $2\alpha$ , centered at 1.

We will make use, in what follows, of the relation:

$$(3.2) \quad \chi_\alpha(u\bar{v}) = \chi_\alpha(u, v), \quad \text{for all } u, v \in \mathbb{S}^1.$$

Given two positive measurable functions  $f, g : \mathbb{S}^1 \rightarrow \mathbb{R}_+$ , their convolution,  $f * g$ , is defined to be the function:

$$\begin{aligned} (f * g)(z_0) &= \int_{\mathbb{S}^1} f(z_0 \bar{z}) g(z) dz \\ &= \int_{-\pi}^{\pi} f(e^{i(\theta_0 - \theta)}) g(e^{i\theta}) d\theta, \end{aligned}$$

with  $z_0 = e^{i\theta_0}$  and  $dz$  represents the arclength element on  $\mathbb{S}^1$ , usually denoted by  $|dz|$ .

Given three positive functions  $f, g, h$  defined on  $\mathbb{S}^1$ , we can write

$$(3.3) \quad \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} f(e^{i(\theta-t)})g(e^{it})h(e^{i\theta}) dt d\theta = (f * g * h^-)(1),$$

where  $h^-(z) = h(\bar{z})$ , i.e.,  $h^-(e^{i\theta}) = h(e^{-i\theta})$ .

**Theorem 3.1.** *Let  $\sigma : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  be a  $C^1$  diffeomorphism such that  $\sigma(1) = 1$  and  $\sigma(-1) = -1$ . Additionally, we assume that  $\sigma(\mathbb{S}_+^1) \subseteq \mathbb{S}_+^1$  and  $\sigma(\mathbb{S}_-^1) \subseteq \mathbb{S}_-^1$ . Let  $d$  be the geodesic distance on the unit circle,  $\alpha$  be a positive real number, and we define the set  $E = \{(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma(x), y) \leq \alpha\}$ . For  $A, B \subset \mathbb{S}^1$  measurable sets, let*

$$I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_A(x) \chi_B(y) \chi_E(x, y) dx dy.$$

*Then, for any  $A, B$  measurable subsets of  $\mathbb{S}^1$ , and  $\alpha > 0$ ,*

$$(3.4) \quad I_\alpha(A, B) \leq I_\alpha(A^\sharp, B^\sharp),$$

*if and only if,  $\sigma$  is symmetric (i.e.  $\overline{\sigma(z)} = \sigma(\bar{z})$ , for every  $z \in \mathbb{S}^1$ ) and convex on  $\mathbb{S}_+^1$ .*

*Proof. Sufficiency.* We define  $\sigma_1 : [-\pi, \pi) \rightarrow [-\pi, \pi)$  by  $e^{\sigma_1(\theta)} := \sigma(e^{i\theta})$  and we assume that  $\sigma_1$  is convex on  $(0, \pi)$ . Using change of variables,  $(\sigma(x), y) = (u, v)$ , the integral  $I_\alpha$  becomes:

$$I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_{\sigma(A)}(u) \chi_B(v) \chi_\alpha(u, v) (\sigma^{-1})'(u) du dv.$$

With  $\chi_\alpha(u, v) = \chi_\alpha(u\bar{v})$ , as in (3.2), the above expression becomes:

$$(3.5) \quad I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_{\sigma(A)}(u) \chi_B(v) \chi_\alpha(u\bar{v}) \psi(u) du dv,$$

where  $\psi(e^{i\theta}) = \tau_1'(\theta)$  and  $\tau_1$  is defined by  $\sigma^{-1}(e^{i\theta}) = e^{i\tau_1(\theta)}$ , and is the inverse of  $\sigma_1$ .

Thus, we can write using convolution and (3.3):

$$I_\alpha(A, B) = [(\chi_{\sigma(A)} \cdot \psi) * \chi_\alpha * \chi_B^-](1),$$

where we used the fact that  $\chi_\alpha$  is a symmetric function.

It was proved in [1] (see also (1.4)) by Baernstein that, for any three positive measurable functions  $f, g, h$  on  $\mathbb{S}^1$ , the following inequality holds:

$$(3.6) \quad (f * g * h^-)(1) \leq (f^\sharp * g^\sharp * h^\sharp)(1).$$

One can replace  $h^-$  in the inequality above by  $h$  since they are equimeasurable functions. Thus, based on (3.6) and the fact that  $\chi_\alpha$  is symmetric decreasing, we conclude that:

$$(3.7) \quad I_\alpha(A, B) \leq [(\chi_{\sigma(A)} \cdot \psi)^\sharp * \chi_\alpha * \chi_{B^\sharp}](1).$$

**Fact:** If  $F$  is a positive symmetric decreasing function and if  $f \prec F$  in the sense of Hardy-Littlewood-Pólya (i.e.  $\sup_{|G|=2\theta} \int_G f \leq \int_{-\theta}^\theta F$ ), then  $f^\sharp$  in inequality (3.6) can be

replaced by  $F$ . Indeed,  $f \prec F$  is equivalent to  $\int_{\mathbb{S}^1} f^\sharp(z) g^\sharp(z) dz \leq \int_{\mathbb{S}^1} F(z) g^\sharp(z) dz$ ,



for all positive symmetric decreasing functions  $g^\sharp$ . Now, since  $g^\sharp * h^\sharp$  is symmetric decreasing and since the convolution  $(f^\sharp * g^\sharp * h^\sharp)(1)$  can be written as the integral of the product  $f^\sharp(z)(g^\sharp * h^\sharp)(z)$ , we conclude that:

$$(f^\sharp * g^\sharp * h^\sharp)(1) \leq (F * g^\sharp * h^\sharp)(1).$$

Therefore, using (3.7) and the Fact, we can prove (3.4) if we show that  $\chi_{\sigma(A)}\psi \prec \chi_{\sigma(A^\sharp)}\psi$ , i.e.

$$(3.8) \quad \int_E \chi_{\sigma(A)}\psi \leq \int_{E^\sharp} \chi_{\sigma(A^\sharp)}\psi.$$

Let  $E' = \sigma^{-1}(E)$ , and  $E'' = \sigma^{-1}(E^\sharp)$ . With these notations, inequality (3.8) becomes:

$$\int_{A \cap E'} dx \leq \int_{A^\sharp \cap E''} dx,$$

or equivalently,  $|A \cap E'| \leq |A^\sharp \cap E''|$ , which is true if  $|E'| \leq |E''|$ , since  $E''$  is symmetric. Since  $\psi$  is symmetric decreasing, we have that  $\int_E \psi(u)du \leq \int_{E^\sharp} \psi(u)du$ , which is equivalent to  $\int_{\sigma^{-1}(E)} dx \leq \int_{\sigma^{-1}(E^\sharp)} dx$ , using change of variables. The latter inequality simply states that  $|E'| \leq |E''|$ , and the proof of the sufficiency is now complete.

*Necessity.* Dividing (3.5) by  $2\alpha$ , and letting  $\alpha$  tend to zero, we obtain:

$$I_0(A, B) = \int_{\mathbb{S}^1} \chi_{\sigma(A)}(u) \chi_B(u) \psi(u) du,$$

and inequality (3.4) implies that:

$$(3.9) \quad I_0(A, B) \leq I_0(A^\sharp, B^\sharp).$$

With the notation  $\tau = \sigma^{-1}$ ,  $\psi$  the Jacobian of  $\tau$ , and  $x = \tau(u)$ ,  $I_0$  becomes:

$$(3.10) \quad I_0(A, B) = \int_{\mathbb{S}^1} \chi_A(x) \chi_{\tau(B)}(x) dx = |A \cap \tau(B)|.$$

First, we will show that the symmetry condition is necessary. Suppose  $\tau$  is not symmetric. Then, there exists a point  $x = e^{i\theta}$  in  $\mathbb{S}_+^1$  such that  $\tau(x) \neq \tau(\bar{x})$ . If we consider  $A = \tau(\{e^{it} : |t| < \theta\})$  and  $B = \{e^{it} : |t| < \theta\}$ , then we have:  $|A \cap \tau(B)| = |\tau(B)| > |A^\sharp \cap \tau(B^\sharp)|$ , since  $\tau(B^\sharp)$  is not symmetric and  $|A| = |\tau(B)|$ . But this contradicts (3.9) and therefore (3.4).

Suppose now that  $\tau_1$  is symmetric, but not concave (or, equivalently,  $\sigma_1$  is symmetric, but  $\sigma_1$  is not convex on  $(0, \pi)$ ). Then, there exist  $e^{ib}, e^{ic} \in \mathbb{S}_+^1$  with  $b, c \in (0, \pi)$  such that:

$$(3.11) \quad \frac{\tau_1(b) + \tau_1(c)}{2} > \tau_1\left(\frac{b+c}{2}\right).$$

Without loss of generality we can assume that  $b > c$  and let us denote by  $a = \frac{b+c}{2}$ . Letting  $B = \{e^{it} : -c < t < b\}$ , it follows that  $B^\sharp = \{e^{it} : -a < t < a\}$ . We calculate  $|\tau(B)| = \tau_1(b) - \tau_1(-c) = \tau_1(b) + \tau_1(c)$  and  $|\tau(B^\sharp)| = 2\tau_1(a)$ .

From (3.11) we obtain that  $|\tau(B)| > |\tau(B^\sharp)|$  which shows that  $I_0(\mathbb{S}^1, B) > I_0(\mathbb{S}^1, B^\sharp)$  and contradicts (3.4). Therefore,  $\tau$  must also be concave.  $\square$

**Theorem 3.2.** *Suppose we have two functions  $\sigma_1, \sigma_2$  satisfying the conditions of  $\sigma$  in Theorem 3.1 and define  $E = \{(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma_1(x), \sigma_2(y)) \leq \alpha\}$ , for  $\alpha \in \mathbb{R}_+$ . Let*

$$I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_A(x) \chi_B(y) \chi_E(x, y) dx dy.$$

*Then, for any  $A, B$  subsets of  $\mathbb{S}^1$  and  $\alpha > 0$ ,*

$$(3.12) \quad I_\alpha(A, B) \leq I_\alpha(A^\sharp, B^\sharp),$$

*if and only if  $\sigma_1, \sigma_2$  are symmetric and convex on  $\mathbb{S}_+^1$ .*

*Proof. Sufficiency.* Very similar to Theorem 3.1. Using change of variables,  $(\sigma_1(x), \sigma_2(y)) = (u, v)$ , the integral becomes:

$$I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_{\sigma_1(A)}(u) \chi_{\sigma_2(B)}(v) \chi_\alpha(u\bar{v}) \psi_1(u) \psi_2(v) du dv,$$

where  $\psi_1, \psi_2$  are defined similarly to  $\psi$  in Theorem 3.1 (see (3.5)). Using convolution, this integral can be written as:

$$I_\alpha(A, B) = [(\chi_{\sigma_1(A)} \cdot \psi_1) * \chi_\alpha * (\chi_{\sigma_2(B)} \cdot \psi_2)^-](1).$$

We have already proven that  $\chi_{\sigma_1(A)} \psi_1 \prec \chi_{\sigma_1(A^\sharp)} \psi_1$  and  $\chi_{\sigma_2(B)} \psi_2 \prec \chi_{\sigma_2(B^\sharp)} \psi_2$ , from which it follows that  $I_\alpha(A, B) \leq I_\alpha(A^\sharp, B^\sharp)$ .

*Necessity.* Using change of variable  $v = \sigma_2(y)$ ,  $I_\alpha$  becomes:

$$I_\alpha(A, B) = \int_{\mathbb{S}^1 \times \mathbb{S}^1} \chi_A(x) \chi_{\{(x, v) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma_1(x), v) \leq \alpha\}} \chi_{\sigma_2(B)}(v) \psi_2(v) dx dv.$$

Dividing by  $\alpha$  and letting  $\alpha \rightarrow 0$ , we obtain:

$$I_0(A, B) = \int_{\mathbb{S}^1} \chi_A(x) \chi_{\sigma_2(B)}(\sigma_1(x)) \psi_2(\sigma_1(x)) dx.$$

Inequality (3.12) of the theorem implies the following inequality:

$$(3.13) \quad I_0(A, B) \leq I_0(A^\sharp, B^\sharp),$$

for all subsets  $A$  and  $B$  of  $\mathbb{S}^1$ .

Now let  $B = \mathbb{S}^1$  in the above identity. Then:

$$I_0(A, \mathbb{S}^1) = \int_{\mathbb{S}^1} \chi_A(x) \psi_2(\sigma_1(x)) dx \leq \int_{\mathbb{S}^1} \chi_{A^\sharp}(x) \psi_2(\sigma_1(x)) dx,$$

or equivalently,

$$\int_A \psi_2(\sigma_1(x)) dx \leq \int_{A^\sharp} \psi_2(\sigma_1(x)) dx,$$

for every measurable set  $A \subset \mathbb{S}^1$ . Since the inequality is true for every measurable set  $A$ , we conclude by Lemma 2.2 and Theorem 2.3 that  $\psi_2 \circ \sigma_1$  is symmetric (i.e.,  $\psi_2(\sigma_1(z)) = \psi_2(\sigma_1(\bar{z}))$ ) and decreasing, which implies that  $\psi_2$  is decreasing on  $\mathbb{S}_+^1$ . Likewise,  $\psi_1 \circ \sigma_2$  is symmetric and decreasing on  $\mathbb{S}_+^1$ , implying that  $\psi_1$  is decreasing on  $\mathbb{S}_+^1$ . Thus,  $\sigma_1^{-1}$  and  $\sigma_2^{-1}$  are concave on  $\mathbb{S}_+^1$  and therefore,  $\sigma_1$  and  $\sigma_2$  are convex on  $\mathbb{S}_+^1$ .

Next, we denote by  $\tau = \sigma_1^{-1} \circ \sigma_2$ . With this notation,  $I_0$  becomes:

$$\begin{aligned} I_0(A, B) &= \int_{\mathbb{S}^1} \chi_A(x) \chi_{\sigma_2(B)}(\sigma_1(x)) [\psi_2 \circ \sigma_1](x) dx \\ &= \int_{\mathbb{S}^1} \chi_A(x) \chi_{\tau(B)}(x) [\psi_2 \circ \sigma_1](x) dx = \int_{A \cap \tau(B)} [\psi_2 \circ \sigma_1](x) dx. \end{aligned}$$

We will show that  $\tau$  is symmetric, i.e.,  $\tau(\bar{x}) = \overline{\tau(x)}$ , for every  $x \in \mathbb{S}^1$ . Suppose this is not the case. Then there exists  $x = e^{i\theta}$ , with  $\theta \in (0, \pi)$ , such that  $\overline{\tau(x)} \neq \tau(\bar{x})$ . Let  $B = \{e^{it} : |t| < \theta\} = B^\sharp$  and  $A = \tau(B) \neq A^\sharp$ . Then, we have that  $A^\sharp \cap \tau(B^\sharp) \subset A \cap \tau(B) = A$  and  $|A \cap \tau(B)| > |A^\sharp \cap \tau(B^\sharp)|$ . Since  $\psi_2 \circ \sigma_1$  is positive, it follows that  $I_0(A, B) > I_0(A^\sharp, B^\sharp)$ , which contradicts (3.13). Thus,  $\sigma_1^{-1} \circ \sigma_2$  is symmetric. We have shown before that  $\psi_1 \circ \sigma_2$  is also symmetric.

*Claim:*  $\sigma_1^{-1} \circ \sigma_2$  and  $\psi_1 \circ \sigma_2$  symmetric imply  $\sigma_2$  is symmetric.

*Proof of claim:* We define  $f_2$  on the interval  $[-\pi, \pi]$  as follows:

$$\sigma_2(e^{i\theta}) = e^{if_2(\theta)}.$$

Since  $\psi_1 \circ \sigma_2$  is symmetric and  $[\psi_1 \circ \sigma_2](e^{i\theta}) = \psi_1(e^{if_2(\theta)}) = \tau_1'(f_2(\theta))$ , as in (3.5), it follows that  $\tau_1' \circ f_2$  is even.

Since  $[\sigma_1^{-1} \circ \sigma_2](e^{i\theta}) = e^{i\tau_1(f_2(\theta))}$  is symmetric, it follows that  $\tau_1 \circ f_2$  is odd.

Now,  $(\tau_1 \circ f_2)' = (\tau_1' \circ f_2) \cdot f_2'$  is even and  $\tau_1' \circ f_2$  is also even (as we have previously shown) and nonzero, so that  $f_2'$  is even and thus  $f_2$  is odd. Therefore  $\sigma_2$  is symmetric and the proof of the claim is now complete.

Following exactly the same steps, we can show that  $\sigma_1$  is symmetric. We have shown that  $\sigma_1, \sigma_2$  are symmetric and convex on  $\mathbb{S}_+^1$ .  $\square$

**Corollary 3.3.** *With  $\sigma$ ,  $\alpha$  and  $E = \{(x, y) \in \mathbb{S}^1 : d(\sigma(x), y) \leq \alpha\}$ , as in Theorem 3.1, we have the following result: For every  $f, g : \mathbb{S}^1 \rightarrow \mathbb{R}_+$  positive measurable functions, and every  $\alpha > 0$ ,*

$$(3.14) \quad \int_E f(x)g(y) dx dy \leq \int_E f^\sharp(x)g^\sharp(y) dx dy,$$

*if and only if,  $\sigma$  is symmetric, and convex on  $\mathbb{S}_+^1$ .*

To sketch the proof, we write  $f$  and  $g$  as the integrals of their level sets, using the layer-cake representation formula (1.2):

$$\begin{aligned} f(x) &= \int_0^\infty \chi_{\{f>t\}}(x) dt \quad \text{and} \\ g(y) &= \int_0^\infty \chi_{\{g>t\}}(y) dt, \end{aligned}$$

and we notice that  $\{f > t\}^\sharp = \{f^\sharp > t\}$  and  $\{g > t\}^\sharp = \{g^\sharp > t\}$  so that inequality (3.14) reduces to the case where  $f$  and  $g$  are characteristic functions, and thus, Theorem 3.1 applies.

**Corollary 3.4.** *Let  $\sigma_1, \sigma_2$  and  $E = \{(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma_1(x), \sigma_2(y)) \leq \alpha\}$  be as in Theorem 3.2. For every  $f, g : \mathbb{S}^1 \rightarrow \mathbb{R}_+$  positive measurable functions, and every  $\alpha > 0$ ,*

$$(3.15) \quad \int_E f(x)g(y) dx dy \leq \int_E f^\sharp(x)g^\sharp(y) dx dy,$$

*if and only if,  $\sigma_1$  and  $\sigma_2$  are symmetric, and convex on  $\mathbb{S}_+^1$ .*

The proof of Corollary 3.4 is indeed very similar to the proof of Corollary 3.3, in which one represents  $f$  and  $g$  as integrals of the characteristic functions of their level sets.

The next theorem is a generalization of the previous results, where one replaces the product by a function  $\Psi$  defined as follows:

$\Psi : \mathbb{R}_+^2 \rightarrow \mathbb{R}$  vanishes on the boundary of  $\mathbb{R}_+^2$ , i.e.,  $\Psi|_{\{x_1=0\}} = \Psi|_{\{x_2=0\}} = 0$ , and

$$\Psi(x_1, x_2) + \Psi(y_1, y_2) \leq \Psi(x_1 \wedge x_2, y_1 \wedge y_2) + \Psi(x_1 \vee x_2, y_1 \vee y_2).$$

If  $\Psi$  is twice continuously differentiable, then the above inequality is equivalent to  $\partial_{12}\Psi \geq 0$ .

Crowe, Zweibel and Rosenbloom [6] noticed that a continuous such  $\Psi$  is the distribution function of a Borel measure  $\mu$  on  $\mathbb{R}_+^2$ , i.e.,

$$(3.16) \quad \Psi(s, t) = \mu([0, s] \times [0, t]),$$

and using Fubini's theorem:

$$(3.17) \quad \int \Psi(f(x), g(y)) dx dy = \int_{\mathbb{R}_+^2} \left[ \int \chi_{\{f>s\}}(x) \chi_{\{g>t\}}(y) dx dy \right] d\mu(s, t).$$

We are now ready to state our next result.

**Theorem 3.5.** *With  $\sigma_1$ ,  $\sigma_2$  and  $E = \{(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1 : d(\sigma_1(x), \sigma_2(y)) \leq \alpha\}$  as in Theorem 3.2, and  $\Psi$  the distribution function of a Borel measure  $\mu$  on  $\mathbb{R}_+^2$  as in (3.16), the following inequality holds for every  $\alpha > 0$ :*

$$\int_E \Psi(f(x), g(y)) dx dy \leq \int_E \Psi(f^\#(x), g^\#(y)) dx dy,$$

*if and only if,  $\sigma_1$  and  $\sigma_2$  are symmetric on  $\mathbb{S}^1$ , and convex on  $\mathbb{S}_+^1$ .*

Again, we can reduce  $\Psi(f(x), g(y))$  to a product of characteristic functions, using (3.17), and the result follows from Theorem 3.2.

The next theorem shows that we can replace the characteristic function of the set  $E$  by a decreasing function of the distance between  $\sigma_1(x)$  and  $\sigma_2(y)$ , call it  $k[d(\sigma_1(x), \sigma_2(y))]$ .

**Theorem 3.6.** *Let  $\sigma_1$ ,  $\sigma_2$  be as in Theorem 3.2 and let  $k : [0, \infty) \rightarrow [0, \infty)$  be a decreasing function, and  $\Psi$  the distribution function of a Borel measure  $\mu$  on  $\mathbb{R}_+^2$  as in (3.16). Then, the following inequality holds for every decreasing function  $k$ ,*

$$\begin{aligned} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f(x), g(y)) k[d(\sigma_1(x), \sigma_2(y))] dx dy \\ \leq \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f^\#(x), g^\#(y)) k[d(\sigma_1(x), \sigma_2(y))] dx dy, \end{aligned}$$

*if and only if,  $\sigma_1$  and  $\sigma_2$  are symmetric on  $\mathbb{S}^1$ , and convex on  $\mathbb{S}_+^1$ .*

*Proof.* Using (1.2), we can write:

$$k(\tau) = \int_0^\infty \chi_{\{k>t\}}(\tau) dt = \int_0^\infty \chi_{[0, l(t)]}(\tau) dt,$$

and substituting  $d(\sigma_1(x), \sigma_2(y))$  for  $\tau$  in the above formula, we have

$$(3.18) \quad k[d(\sigma_1(x), \sigma_2(y))] = \int_0^\infty \chi_{[0, l(t)]}[d(\sigma_1(x), \sigma_2(y))] dt.$$

We define the set  $E_{l(t)}$  as follows:

$$E_{l(t)} = \{(x, y) : d(\sigma_1(x), \sigma_2(y)) \leq l(t)\}.$$

Then

$$\chi_{[0, l(t)]}[d(\sigma_1(x), \sigma_2(y))] = 1 \Leftrightarrow (x, y) \in E_{l(t)}.$$

Using this fact, (3.18), Fubini's theorem and Theorem 3.5 we obtain the conclusion of Theorem 3.6 by:

$$\begin{aligned} & \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f(x), g(y)) k[d(\sigma_1(x), \sigma_2(y))] dx dy \\ &= \int_0^\infty \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f(x), g(y)) \chi_{E_{l(t)}}(x, y) dx dy dt \\ &\leq \int_0^\infty \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f^\sharp(x), g^\sharp(y)) \chi_{E_{l(t)}}(x, y) dx dy dt \\ &= \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \Psi(f^\sharp(x), g^\sharp(y)) k[d(\sigma_1(x), \sigma_2(y))] dx dy. \end{aligned}$$

□

## REFERENCES

- [1] A. Baernstein, II. Convolution and rearrangement on the circle. *Complex Variables Theory Appl.*, 12(1-4):33–37, 1989.
- [2] A. Baernstein, II and B. A. Taylor. Spherical rearrangements, subharmonic functions, and  $*$ -functions in  $n$ -space. *Duke Math. J.*, 43(2):245–268, 1976.
- [3] A. Baernstein II. A unified approach to symmetrization. *Partial Differential Equations of Elliptic Type*, eds. A. Alvino et al, *Symposia Mathematica*, Cambridge Univ. Press, 35:47–91, 1995.
- [4] C. Bennett and R. Sharpley. *Interpolation of operators*. Academic Press Inc., 1988.
- [5] H. J. Brascamp, E. H. Lieb, and J. M. Luttinger. A general rearrangement inequality for multiple integrals. *J. Funct. Anal.*, 17:227–237, 1974.
- [6] J. A. Crowe, J. A. Zweibel, and P. C. Rosenbloom. Rearrangements of functions. *J. Funct. Anal.*, 66(3):432–438, 1986.
- [7] H. Federer. *Geometric measure theory*. Springer-Verlag New York Inc., New York, 1969.
- [8] H. Hadwiger and D. Ohmann. Brunn-Minkowskischer Satz und Isoperimetrie. *Math. Z.*, 66:1–8, 1956.
- [9] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1988. Reprint of the 1952 edition.
- [10] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [11] R. Osserman. The isoperimetric inequality. *Bull. Amer. Math. Soc.*, 84(6):1182–1238, 1978.
- [12] F. Riesz. Sur une inégalité intégrale. *J. London Math. Soc.*, 5:162–168, 1930.
- [13] S. L. Sobolev. On a theorem of functional analysis. *Mat. Sb. (N.S.)*, 4:471–497, 1938. English transl., *Amer. Math. Soc. Transl.*, 2(34):29–68, 1963.

DEPARTMENT OF MATHEMATICAL SCIENCES, AALBORG UNIVERSITY, FR. BAJERS VEJ 7G,  
DK-9220 AALBORG EAST, DENMARK

*E-mail address*: cristina@math.aau.dk