



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The consequence of combined pain and stress on work ability in female laboratory technicians

Jay, Kenneth; Friberg, Maria Kristine; Sjøgaard, Gisela; Jakobsen, Markus Due; Sundstrup, Emil; Petersen, Mikkel Brandt; Andersen, Lars Louis

Published in:

International Journal of Environmental Research and Public Health

DOI (link to publication from Publisher):

[10.3390/ijerph121215024](https://doi.org/10.3390/ijerph121215024)

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Jay, K., Friberg, M. K., Sjøgaard, G., Jakobsen, M. D., Sundstrup, E., Brandt, M., & Andersen, L. L. (2015). The consequence of combined pain and stress on work ability in female laboratory technicians: a cross-sectional study. *International Journal of Environmental Research and Public Health*, 12(12), 15834-15842. <https://doi.org/10.3390/ijerph121215024>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Article

The Consequence of Combined Pain and Stress on Work Ability in Female Laboratory Technicians: A Cross-Sectional Study

Kenneth Jay ^{1,2,*}, Maria Kristine Friberg ², Gisela Sjøgaard ¹, Markus Due Jakobsen ², Emil Sundstrup ², Mikkel Brandt ^{2,3} and Lars Louis Andersen ^{2,3}

Received: 20 October 2015; Accepted: 8 December 2015; Published: 11 December 2015

Academic Editor: Cary Cooper

¹ Department of Physical Activity and Health, Institute for Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense 5230, Denmark; gsjogaard@health.sdu.dk

² National Research Centre for the Working Environment, Copenhagen 2100, Denmark; mkf@nrcwe.dk (M.K.F.); mdj@nrcwe.dk (M.D.J.); esu@nrcwe.dk (E.S.); mbp@nrcwe.dk (M.B.); lla@nrcwe.dk (L.L.A.)

³ Physical Activity and Human Performance group, SMI Department of Health Science and Technology, Aalborg University, Aalborg 9220, Denmark

* Correspondence: kennethjay@icloud.com; Tel.: +45-39-165-200 (ext. 123); Fax: +45-39-165-200

Abstract: Musculoskeletal pain and stress-related disorders are leading causes of impaired work ability, sickness absences and disability pensions. However, knowledge about the combined detrimental effect of pain and stress on work ability is lacking. This study investigates the association between pain in the neck-shoulders, perceived stress, and work ability. In a cross-sectional survey at a large pharmaceutical company in Denmark 473 female laboratory technicians replied to questions about stress (Perceived Stress Scale), musculoskeletal pain intensity (scale 0–10) of the neck and shoulders, and work ability (Work Ability Index). General linear models tested the association between variables. In the multi-adjusted model, stress ($p < 0.001$) and pain ($p < 0.001$) had independent main effects on the work ability index score, and there was no significant stress by pain interaction ($p = 0.32$). Work ability decreased gradually with both increased stress and pain. Workers with low stress and low pain had the highest Work Ability Index score (44.6 (95% CI 43.9–45.3)) and workers with high stress and high pain had the lowest score (32.7 (95% CI 30.6–34.9)). This cross-sectional study indicates that increased stress and musculoskeletal pain are independently associated with lower work ability in female laboratory technicians.

Keywords: pain stress relationship; behavior; social factors; fear-avoidance; biopsychosocial; learned helplessness; resources and demands

1. Introduction

In the working population reduced work ability caused by physical or mental disabilities is highly prevalent and is the leading cause of long-term sick leave [1–3] staff turnover and early retirement [4,5]. Reduced work ability is not only costly for the individual, but also for the organization and society due to lost productivity and compensation costs [3]. Work ability is the result of the interaction between the employee and his or her work. Consequently, work ability can be described as the balance between the employee's resources and the work demands [6]. The employee contributes to his or her work ability via health and functional abilities, knowledge, skills, attitude and motivation. The workplace influences work ability through organizational factors—through leadership, management issues, work demands and social factors [7]. On the basis of comprehensive

clinical assessments and statistical analyses, Tuomi and colleagues identified a short set of questions ultimately resulting in a score indicating the employee's work ability, aptly named the Work Ability Index (WAI) [8]. Ilmarinen and Tuomi have shown that people with high WAI scores have a lower risk for early retirement and a higher quality of life—even after retirement [6,7,9]. Furthermore, studies using the WAI have also presented data suggesting that it is possible to sustainably improve work ability—even at older age—provided the right measures are taken [6,7,9]. The WAI has been associated with musculoskeletal pain, chronic disease, work productivity, sickness, untimely retirement, as well as all-cause mortality [6,8]. In occupational groups where employees are exposed to repetitive, monotonous and/or forceful exertion, compromised body positions and/or insufficient recovery, there is an elevated risk of both impaired work ability and chronic musculoskeletal disorders in the long term [10].

Research on the relationship between stress and musculoskeletal pain has previously largely focused on back pain and multisite pain but only to a limited extent on neck-shoulder pain specifically. Oberlinner *et al.* demonstrated, by conducting a comprehensive survey combining questionnaire data and medical examinations in one division of a major chemical company in Germany, that although occupational stressors were perceived differently, there was no difference in the prevalence of back pain between different working groups within the company [11]. Conversely, a 2000 review by Davis and Haeney [12] on the relationship between psychosocial work characteristics and low back pain established that while it is problematic to argue for causal inferences, it appears that psychosocial characteristics are related to at least some lower back pain outcomes. Furthermore, employees' reactions to psychosocial work characteristics (e.g., job dissatisfaction and job stress) are more consistently related to lower back pain than the psychosocial work characteristics themselves. In conjunction with a systematic review by Hoogendroom *et al.* on the psychosocial factors at work and in private life as risk factors for back pain, there is confirmation for an effect of work-related psychosocial factors on the perception of physical well-being [13]. Congruent with this, Lindegaard *et al.* reported a combination of frequent pain and stress constituted the highest risk for a decrease in work performance evaluated by a single item from the WAI questionnaire and conclude that workplace interventions should focus on promoting musculoskeletal well-being and encompass both individual and organizational interventions to minimize the risk of increased work-related stress [14]. Furthermore, Ganster *et al.* [15], and Cohen *et al.* [16,17] concluded that there are three main approaches to the definition and study of work stress. In brief, the first approach conceptualizes stress as a characteristic of the environment that affects the individual. The second approach defines stress as the psychological reaction (psychological, physiological and behavioral) on environmental stimuli. The third and most prevalent approach views stress as the interaction between environmental characteristics and the subjective reaction to these characteristics [15]. In the present study, we used the third definition as we defined psychological stress as a process by which environmental stimuli initiate cognitive and physiological reactions that ultimately can affect wellbeing. Psychological stress occurs when an individual perceives that environmental demands exceed his or her adaptive capacity [16]. The aim of the present study was to investigate the association between pain in the neck and shoulders, stress and work ability measured by the WAI.

2. Experimental Section

2.1. Study Design

This study is an explorative analysis of baseline data obtained during a worksite intervention trial previously described by our research team [18]. Data for this study were collected during the spring of 2014.

2.2. Ethics

Ethical approval was obtained from The Danish National Committee on Biomedical Research Ethics (The local ethical committee of Frederiksberg and Copenhagen; H-3-2010-062) as part of the research program “Implementation of physical exercise at the workplace (IRMA)”. The trial “Implementation of physical exercise at the Workplace (IRMA09)—Laboratory technicians” was registered in the ClinicalTrials.gov register (NCT02047669) prior to participant enrolment. All experimental conditions conformed to The Declaration of Helsinki.

2.3. Participants

Out of 756 laboratory technicians at a large pharmaceutical company in Denmark, 539 completed questionnaires on musculoskeletal pain, perceived level of stress and work ability. Of these, 473 were women and included in the analysis. Table 1 shows participant demographics of relevant data. All eligible participants were informed about the purpose and content of the study. Table 1 shows participant characteristics of relevant data.

Table 1. Descriptive characteristics of the female laboratory technicians included in the analysis.

Descriptive Characteristics	Mean (S.D.)
Number of included observations	473
Age, years	46.0 (9.5)
Body mass index(kg·m ⁻²)	24.2 (3.7)
Smokers, percentage	1.9 (0.3)
Seniority, years	20.9 (10.8)
Number of working hours per week	36.9 (4.3)
Neck-shoulder pain intensity (0–10 VAS-scale)	2.6 (2.4)
The Perceived Stress Scale score (0–40)	12.3 (6.4)
Work Ability Index score (7–49)	41.8 (4.9)

2.4. Work Ability

The WAI consists of seven indicators providing a score between 7 and 49 (higher is better) that gauge occupational well-being by providing subjective estimations on work ability. This includes capability of employees to perform their work tasks with respect to demands, health, as well as mental resources [6,19]. Further, the WAI includes the dimensions of the determinants (e.g., health) and outcomes (sick leave and functional limitations) of work ability. Currently, the WAI has been translated into 26 languages and is used in numerous countries throughout the world [20]. The validity of WAI is well documented [21] and the test-retest reliability have been found to be a stable measure [22]. For reference, we have classified the sum score into the following three categories of work ability: Poor (score 7–27), Moderate (score 28–36), Good/Excellent (score 36–49).

2.5. Pain

We asked the participants to rate pain intensity in the neck and shoulder on a modified VAS scale (0–10). For reference, “0” is “no pain” and “10” is “worst imaginable pain”. The neck and shoulder were defined by drawings from the Nordic Questionnaire [23] and an average pain score of the two regions was subsequently calculated and used in the statistical analysis. We defined the following cut points for the level of pain: (0 < low pain ≤ 2), (2 < moderate pain ≤ 5), and (high pain > 5) [24–27].

2.6. Stress

The Perceived Stress Scale (PSS), developed by Cohen, Kamarck, and Mermelstein is a comprehensive stress questionnaire and was designed to measure “the degree to which individuals appraise situations in their lives as stressful”. Items evaluate the degree to which people find that

life is unpredictable, uncontrollable, or overloaded [28]. These three aspects have been confirmed as vital elements of the experience of stress and provide a thorough insight into the degree of learned helplessness experienced by the individual [29–33]. The Perceived Stress Scale includes questions intended to evaluate the current level of stress experienced by the subject. The PSS-10 is an abbreviated version of the scale, consisting of only 10 items (the full version has 14 items), administered in only a few minutes, and easily scored. Because the perceived stress scale assesses general beliefs about perceived stress without providing subjects with a list of specific life events, scores are not biased by event content or by differential recall of previous life experiences. In brief, each item on the PSS-10 questionnaire is rated on a 5-point Likert scale ranging from “never” (0) to “almost always” (4). Positively worded items are reverse scored, and the ratings are summed, with higher scores indicating more perceived stress. The PSS-10 score is obtained by reversing the scores on the four positive items: For example, 0 = 4, 1 = 3, 2 = 2, *etc.* and then summing across all 10 items. A score of 13 is considered average and stress scores of more than 20 indicate high stress [28]. For reference, we divided the scoring into three categories with the following cut-off points: (low stress ≤ 10), (10 < moderate stress ≤ 20) and (high stress > 20). Examples of questions from the PSS-10 questionnaire include: (i) “In the past month, how often have you been angry because of things that happened that were outside of your control?”, (ii) “In the past month, how often have you felt that things were going your way?” and (iii) “In the past month, how often have you felt unable to control the important things in your life?” [28].

2.7. Statistics

We performed all statistical analyses using the SAS statistical software for Windows (SAS Institute, Cary, NC, USA). Using the General Linear Modelling procedure (proc GLM), we analyzed the following relationships between stress, pain and WAI. (1) Stress to pain with neck-shoulder pain as the dependent variable; (2) stress to WAI with WAI as the dependent variable; (3) pain to WAI with WAI as the dependent variable and (4) stress + pain, and stress by pain interaction to WAI with WAI as the dependent variable. All analyses were controlled for age. To assess simple associations, we also used the CORR procedure to determine Pearson’s correlation coefficients. An alpha level of <0.05 was accepted as statistically significant. Results are reported as least square means (95% confidence limits).

3. Results

Stress and neck-shoulder pain were positively correlated with a Pearson’s correlation coefficient of 0.29 ($p < 0.0001$). Additionally, stress was independently correlated to WAI with a coefficient of -0.51 ($p < 0.0001$) indicating a lower WAI score with increased stress. Similarly, chronic neck-shoulder pain also correlated to WAI with a coefficient of -0.36 ($p < 0.0001$) indicating a lower WAI score with increased neck-shoulder pain. Figure 1 shows a stress + pain plot with WAI as the dependent outcome variable.

In the multi-adjusted model, both stress ($p < 0.001$) and pain ($p < 0.001$) had independent main effects on WAI, and there was no significant stress by pain interaction ($p = 0.32$). WAI decreased gradually with both increased stress and pain. Workers with low stress and low pain ($n = 133$) had the highest work ability (44.6 (95% CI 43.9–45.3)) and workers with high stress and high pain ($n = 14$) had the lowest (32.7 (95% CI 30.6–34.9)). WAI is therefore dependent on the magnitude of both stress and neck-shoulder pain in an additive fashion. In this population, WAI score decreased 26.6% from low stress + low pain to high stress + high pain.




Figure 1. Least square means stress + pain plot with WAI as the dependent outcome variable. As stress and neck-shoulder pain increase, the work ability of female laboratory technicians decreases. Developing chronic musculoskeletal pain while experiencing high stress returns the lowest work ability score.

4. Discussion

Our study showed that increased stress and musculoskeletal pain were independently associated with lower work ability in female laboratory technicians.

4.1. Stress and Pain

Stress and neck-shoulder pain were associated with work ability in an additive fashion. This result supports that psychosocial factors interact with somatic problems and is congruent with studies reporting a correlation between back pain and psychological stress [11,12,34,35]. As most research on stress and musculoskeletal pain has focused on back pain and multisite pain, this study contributes to the field by indicating an association specifically between neck-shoulder pain and perceived stress. Due to the cross-sectional nature of the present study it is not possible to infer about causality between stress and neck-shoulder pain, but research suggests that stress and pain are positively associated [11]. Workers with neck-shoulder pain may be more likely to perceive themselves as more stressed, as pain potentially limits normal social interactions and work ability [36]. Possibly, perceived stress intensifies or causes neck-shoulder pain through the physical response when subjected to psychological stress. Psychological stress activates the sympathetic nervous system, which releases noradrenaline from widely distributed synapses and adrenaline from the adrenal medulla. The secretion of these hormones primes the body for action causing the muscles tense up, blood vessels to constrict and blood pressure to increase [37–41].

4.2. Stress, Pain and Work Ability

To our knowledge, only a few studies have examined the interaction between musculoskeletal pain and perceived stress on work ability [11,14] making the present study the first study to show the combined effect of neck-shoulder pain specifically, and perceived stress on work ability. Our study showed that stress and pain did not interact in relation to work ability, but that the effect was additive. The result implicate that an intervention strategy aimed at reducing both neck-shoulder pain and perceived stress may be beneficial to prevent reduced work ability and is congruent with other studies attenuating multifactorial intervention strategies [42,43]. Several randomized controlled trials have reported positive reductions in chronic pain in the upper extremity following both strenuous and non-strenuous resistance training using both weights and elastic bands [44–50],

thus providing insight into factors positively affecting work ability and factors reducing both chronic and non-chronic musculoskeletal pain. For instance, in a recent wide-ranging study Kettunen *et al.* showed that low- to moderate physical activity coupled with improvements in cardiorespiratory fitness, are closely associated with long-term (12 months) improved work ability in subjects with low-, moderate- and good work ability at baseline [7]. The researchers measured cardiorespiratory fitness and WAI at baseline, 4 months, 8 months and 12 months in 338 participants and found 2%–3% increases in WAI, at 4 months, again at 8 months and again at 12 months in the exercise training group, whereas the control group experienced a 2% decrease in WAI score [7]. Although the study by Kettunen and colleagues did not include participants with chronic pain, it shows that physical fitness interventions can have a positive effect on work ability. However, as work ability appears to be comprised of a complex set of interactions between psychological, environmental, social and biological factors, single-component interventions are limited as they do not target all aspects of the biopsychosocial model in relation to work ability. Arguably, interventions aimed at promoting work ability do not only have to be multifaceted but must also be implemented on an organizational level [51,52]. Psychological or physiotherapeutic treatment outside of work or in work hours with no involvement of the organization may not be ideal for improving work ability. Involvement of the organization may imply social and emotional support in conjunction with acceptance from the organizational leader. Furthermore, organizational involvement can include a restructuring of work factors that contribute to the employee's pain and stress. As both psychological stress and neck-shoulder pain are highly prevalent [53,54], future research should investigate the effect of multifaceted interventions implemented on an organizational level and targeted specifically at neck-shoulder pain and stress to promote work ability.

4.3. Limitations

This study demonstrates the combined effect of neck-shoulder pain specifically, and perceived stress on work ability. However, some limitations exist. First, the cross-sectional design does not permit examination of causal relationships. Thus, prospective studies should assess the combined effect of stress and pain on deterioration of work ability. On the other hand, high stress and pain may reflect that the worker has already experienced deterioration of work ability. Nevertheless, the results are still of interest, as they give an insight into factors associated with work ability, which can be used in the development of preventive intervention strategies. Second, sample size is relatively small limiting statistical power and the reliability of results. Third, self-reported data are a limitation as they may be influenced by subjective factors. Fourth, given the demographic characteristics of this sample (Danish female laboratory technicians) generalizability remains to be determined. The presented results may not be generalizable to male employees, as gender may modify the examined relations. Conversely, using a homogenous sample consisting of female laboratory technicians is also a noteworthy strength as it limits bias from socioeconomic confounding.

5. Conclusions

This cross-sectional study confirms a necessity to look at both stress and musculoskeletal pain as combined co-factors when implementing interventional strategies to promote work ability.

Acknowledgments: The authors would like to thank the management of the pharmaceutical company for supplying contact information to all employees and allowing their laboratory technicians to answer the questionnaire during working hours.

Author Contributions: Kenneth Jay and Lars L. Andersen designed the questionnaire with help from Markus Due Jakobsen, Emil Sundstrup and Mikkel Brandt. Kenneth Jay and Lars Louis Andersen collected the data. Kenneth Jay and Maria Kristine Friberg wrote the manuscript with valuable input and critique from Gisela Sjogaard, Markus Due Jakobsen, Emil Sundstrup, Mikkel Brandt and Lars Louis Andersen. Lars Louis Andersen performed the statistical calculations and interpreted the results together with Kenneth Jay. All authors were involved in critically reviewing and approving the manuscript before submission.

Conflicts of Interest: The authors declare no conflict of interest. The study received financial support by a commercial organization. The commercial organization does not profit in any way or form from the results of this study.

References

1. Nygård, C.-H.; Arola, H.; Siukola, A.; Savinainen, M.; Luukkaala, T.; Taskinen, H.; Virtanen, P. Perceived work ability and certified sickness absence among workers in a food industry. *Int. Congr. Ser.* **2005**, *1280*, 296–300. [[CrossRef](#)]
2. Kujala, V.; Tammelin, T.; Remes, J.; Vammavaara, E.; Laitinen, J. Work ability index of young employees and their sickness absence during the following year. *Scand. J. Work Environ. Health* **2006**, *32*, 75–84. [[CrossRef](#)] [[PubMed](#)]
3. Sell, L.; Bültmann, U.; Rugulies, R.; Villadsen, E.; Faber, A.; Søgaard, K. Predicting long-term sickness absence and early retirement pension from self-reported work ability. *Int. Arch. Occup. Environ. Health* **2009**, *82*, 1133–1138. [[CrossRef](#)] [[PubMed](#)]
4. Hopsu, L.; Leppänen, A.; Ranta, R.; Louhevaara, V. Perceived work ability and individual characteristics as predictors for early exit from working life in professional cleaners. *Int. Congr. Ser.* **2005**, *1280*, 84–88. [[CrossRef](#)]
5. Sjögren-Rönkä, T.; Ojanen, M.T.; Leskinen, E.K.; Tmustalampi, S.; Mälkiä, E.A. Physical and psychosocial prerequisites of functioning in relation to work ability and general subjective well-being among office workers. *Scand. J. Work Environ. Health* **2002**, *28*, 184–190. [[CrossRef](#)] [[PubMed](#)]
6. Ilmarinen, J.; Tuomi, K.; Klockars, M. Changes in the work ability of active employees as measured by the work ability index over an 11-year period. *Scand. J. Work Environ. Health* **1997**, *23*, 49–57. [[PubMed](#)]
7. Kettunen, O.; Vuorimaa, T.; Vasankari, T. 12-mo intervention of physical exercise improved work ability, especially in subjects with low baseline work ability. *Int. J. Environ. Res. Public Health* **2014**, *11*, 3859–3869. [[CrossRef](#)] [[PubMed](#)]
8. Tuomi, K.; Ilmarinen, J.; Jahkola, A.; Katajarinne, L. *Work Ability Index*; Finnish Institute of Occupational Health: Helsinki, Finland, 1997.
9. Holtermann, A.; Jørgensen, M.B.; Gram, B.; Christensen, J.R.; Faber, A.; Overgaard, K.; Ektor-Andersen, J.; Mortensen, O.S.; Sjøgaard, G.; Søgaard, K. Worksite interventions for preventing physical deterioration among employees in job-groups with high physical work demands: Background, design and conceptual model of FINALE. *BMC Public Health* **2010**, *10*. [[CrossRef](#)] [[PubMed](#)]
10. Sundstrup, E.; Jakobsen, M.D.; Andersen, C.H.; Jay, K.; Persson, R.; Aagaard, P.; Andersen, L.L. Participatory ergonomic intervention *versus* strength training on chronic pain and work disability in slaughterhouse workers: Study protocol for a single-blind, randomized controlled trial. *BMC Musculoskelet Disord.* **2013**, *14*. [[CrossRef](#)] [[PubMed](#)]
11. Oberlinner, C.; Yong, M.; Nasterlack, M.; Pluto, R.-P.; Lang, S. Combined effect of back pain and stress on work ability. *Occup. Med.* **2015**, *65*, 147–153. [[CrossRef](#)] [[PubMed](#)]
12. Davis, K.G.; Heaney, C.A. The relationship between psychosocial work characteristics and low back pain: Underlying methodological issues. *Clin. Biomech.* **2000**, *15*, 389–406. [[CrossRef](#)]
13. Hoogendoorn, W.E.; van Poppel, M.N.; Bongers, P.M.; Koes, B.W.; Bouter, L.M. Systematic review of psychosocial factors at work and private life as risk factors for back pain. *Spine* **2000**, *25*, 2114–2125. [[CrossRef](#)] [[PubMed](#)]
14. Lindegård, A.; Larsman, P.; Hadzibajramovic, E.; Ahlborg, G. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int. Arch. Occup. Environ. Health* **2014**, *87*, 373–379. [[CrossRef](#)] [[PubMed](#)]
15. Ganster, D.C.; Rosen, C.C. Work stress and employee health: A multidisciplinary review. *J. Manag.* **2013**, *39*, 1085–1122. [[CrossRef](#)]
16. Cohen, S.; Janicki-Deverts, D.; Miller, G.E. Psychological stress and disease. *JAMA* **2007**, *298*, 1685–1687. [[CrossRef](#)] [[PubMed](#)]
17. Cohen, S.; Kessler, R.C.; Gordon, L.U. *Measuring Stress: A Guide for Health and Social Scientists*; Oxford University Press: Oxford, UK, 1997.

18. Jay, K.; Brandt, M.; Hansen, K.; Sundstrup, E.; Jakobsen, M.D.; Schraefel, M.C.; Sjogaard, G.; Andersen, L.L. Effect of individually tailored biopsychosocial workplace interventions on chronic musculoskeletal pain and stress among laboratory technicians: Randomized controlled trial. *Pain Physician* **2015**, *18*, 459–471. [[CrossRef](#)] [[PubMed](#)]
19. Feldt, T.; Hyvönen, K.; Mäkikangas, A.; Kinnunen, U.; Kokko, K. Development trajectories of Finnish managers' work ability over a 10-year follow-up period. *Scand. J. Work Environ. Health* **2009**, *35*, 37–47. [[CrossRef](#)] [[PubMed](#)]
20. Ilmarinen, J. The work ability index (WAI). *Occup. Med.* **2006**, *57*, 160–160. [[CrossRef](#)]
21. Ilmarinen, J. Work ability—A comprehensive concept for occupational health research and prevention. *Scand. J. Work Environ. Health* **2009**, *35*, 1–5. [[CrossRef](#)] [[PubMed](#)]
22. De Zwart, B.C.H. Test-retest reliability of the work ability index questionnaire. *Occup. Med.* **2002**, *52*, 177–181. [[CrossRef](#)]
23. Kuorinka, I.; Jonsson, B.; Kilbom, A.; Vinterberg, H.; Biering-Sørensen, F.; Andersson, G.; Jørgensen, K. Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl. Ergon.* **1987**, *18*, 233–237. [[CrossRef](#)]
24. Cohen, S.; Kamarck, T.; Mermelstein, R. A global measure of perceived stress. *J. Health Soc. Behav.* **1983**, *24*, 385–396. [[CrossRef](#)] [[PubMed](#)]
25. Kryger, A.I.; Andersen, J.H.; Lassen, C.F.; Brandt, L.P.; Vilstrup, I.; Overgaard, E.; Thomsen, J.F.; Mikkelsen, S. Does computer use pose an occupational hazard for forearm pain: From the NUDATA study. *Occup. Environ. Med.* **2003**, *60*. [[CrossRef](#)]
26. Kaergaard, A.; Andersen, J.H.; Rasmussen, K.; Mikkelsen, S. Identification of neck-shoulder disorders in a 1 year follow-up study: Validation of a questionnaire-based method. *Pain* **2000**, *86*, 305–310. [[CrossRef](#)]
27. Von Korff, M.; Ormel, J.; Keefe, F.J.; Dworkin, S.F. Grading the severity of chronic pain. *Pain* **1992**, *50*, 133–149. [[CrossRef](#)]
28. Andersen, L.L.; Clausen, T.; Burr, H.; Holtermann, A. Threshold of musculoskeletal pain intensity for increased risk of long-term sickness absence among female healthcare workers in eldercare. *PLoS ONE* **2012**, *7*. [[CrossRef](#)] [[PubMed](#)]
29. Averill, J.R. Personal control over aversive stimuli and its relationship to stress. *Psychol. Bull.* **1973**, *80*. [[CrossRef](#)]
30. Baum, A.; Singer, J.E.; Baum, C.S. Stress and the environment. *J. Soc. Issues* **1981**, *37*, 4–35. [[CrossRef](#)]
31. Crandall, R.; Perrewe, P.L. *Occupational Stress: A Handbook*; CRC Press: Boca Raton, FL, USA, 1995.
32. Lazarus, R.S. *Fifty Years of the Research and Theory of R.S. Lazarus: An Analysis of Historical and Perennial Issues*; Lawrence Erlbaum Associates: Mahwah, NJ, USA, 2013.
33. Seligman, M.E.P. Learned helplessness. *Annu. Rev. Med.* **1972**, *23*, 407–412. [[CrossRef](#)] [[PubMed](#)]
34. Bonzini, M.; Bertu, L.; Veronesi, G.; Conti, M.; Coggon, D.; Ferrario, M.M. Is musculoskeletal pain a consequence or a cause of occupational stress? A longitudinal study. *Int. Arch. Occup. Environ. Health* **2014**, *88*, 607–612. [[CrossRef](#)] [[PubMed](#)]
35. Hurwitz, E.L.; Morgenstern, H.; Chiao, C. Effects of recreational physical activity and back exercises on low back pain and psychological distress: Findings from the UCLA Low Back Pain Study. *Am. J. Public Heal.* **2005**, *95*, 1817–1824. [[CrossRef](#)] [[PubMed](#)]
36. Chapman, C.R.; Tuckett, R.P.; Song, C.W. Pain and stress in a systems perspective: Reciprocal neural, endocrine, and immune interactions. *J. Pain* **2008**, *9*, 122–145. [[CrossRef](#)] [[PubMed](#)]
37. Gunnar, M.; Quevedo, K. The neurobiology of stress and development. *Annu. Rev. Psychol.* **2007**, *58*, 145–173. [[CrossRef](#)] [[PubMed](#)]
38. McEwen, B.S. Brain on stress: How the social environment gets under the skin. *Proc. Natl. Acad. Sci. USA.* **2012**, *109*, 17180–17185. [[CrossRef](#)] [[PubMed](#)]
39. Hellhammer, D.H.; Wüst, S.; Kudielka, B.M. Salivary cortisol as a biomarker in stress research. *Psychoneuroendocrinology* **2009**, *34*, 163–171. [[CrossRef](#)] [[PubMed](#)]
40. De Kloet, E.R.; Joëls, M.; Holsboer, F. Stress and the brain: From adaptation to disease. *Nat. Rev. Neurosci.* **2005**, *6*, 463–475. [[CrossRef](#)] [[PubMed](#)]
41. Bozovic, D.; Racic, M.; Ivkovic, N. Salivary cortisol levels as a biological marker of stress reaction. *Med. Arch.* **2013**, *67*, 374–377. [[CrossRef](#)] [[PubMed](#)]

42. De Boer, A.G.E.M.; Burdorf, A.; van Duivenbooden, C.; Frings-Dresen, M.H.W. The effect of individual counselling and education on work ability and disability pension: A prospective intervention study in the construction industry. *Occup. Environ. Med.* **2007**, *64*, 792–797. [[CrossRef](#)] [[PubMed](#)]
43. Nurminen, E.; Malmivaara, A.; Ilmarinen, J.; Ylöstalo, P.; Mutanen, P.; Ahonen, G.; Aro, T. Effectiveness of a worksite exercise program with respect to perceived work ability and sick leaves among women with physical work. *Scand. J. Work Environ. Health* **2002**, *28*, 85–93. [[CrossRef](#)]
44. Zebis, M.K.; Andersen, L.L.; Pedersen, M.T.; Mortensen, P.; Andersen, C.H.; Pedersen, M.M.; Boysen, M.; Roessler, K.K.; Hannerz, H.; Mortensen, O.S.; *et al.* Implementation of neck/shoulder exercises for pain relief among industrial workers: A randomized controlled trial. *BMC Musculoskelet Disord.* **2011**, *12*. [[CrossRef](#)] [[PubMed](#)]
45. Carpes, F.P.; Reinehr, F.B.; Mota, C.B. Effects of a program for trunk strength and stability on pain, low back and pelvis kinematics, and body balance: A pilot study. *J. Bodyw. Mov. Ther.* **2008**, *12*, 22–30. [[CrossRef](#)] [[PubMed](#)]
46. Freitas-Swerts, F.C.T.; de Robazzi, M.L.; Do, C.C. The effects of compensatory workplace exercises to reduce work-related stress and musculoskeletal pain. *Rev. Lat. Am. Enfermagem* **2014**, *22*, 629–636. [[CrossRef](#)] [[PubMed](#)]
47. Andersen, L.L.; Zebis, M.K.; Pedersen, M.T.; Roessler, K.K.; Andersen, C.H.; Pedersen, M.M.; Feveile, H.; Mortensen, O.S.; Sjøgaard, G. Protocol for work place adjusted intelligent physical exercise reducing musculoskeletal pain in shoulder and neck (VIMS): A cluster randomized controlled trial. *BMC Musculoskelet Disord.* **2010**, *173*. [[CrossRef](#)] [[PubMed](#)]
48. Rasmussen, C.D.N.; Holtermann, A.; Mortensen, O.S.; Søgaard, K.; Jørgensen, M.B. Prevention of low back pain and its consequences among nurses' aides in elderly care: A stepped-wedge multi-faceted cluster-randomized controlled trial. *BMC Public Health* **2013**, *13*. [[CrossRef](#)] [[PubMed](#)]
49. Andersen, L.L.; Christensen, K.B.; Holtermann, A.; Poulsen, O.M.; Sjøgaard, G.; Pedersen, M.T.; Hansen, E.A. Effect of physical exercise interventions on musculoskeletal pain in all body regions among office workers: A one-year randomized controlled trial. *Man Ther.* **2010**, *15*, 100–104. [[CrossRef](#)] [[PubMed](#)]
50. Viljanen, M.; Malmivaara, A.; Uitti, J.; Rinne, M.; Palmroos, P.; Laippala, P. Effectiveness of dynamic muscle training, relaxation training, or ordinary activity for chronic neck pain: Randomised controlled trial. *BMJ* **2003**, *327*. [[CrossRef](#)] [[PubMed](#)]
51. Noordik, E.; Nieuwenhuijsen, K.; Varekamp, I.; van der Klink, J.J.; van Dijk, F.J. Exploring the return-to-work process for workers partially returned to work and partially on long-term sick leave due to common mental disorders: A qualitative study. *Disabil. Rehabil.* **2011**, *33*, 1625–1635. [[CrossRef](#)] [[PubMed](#)]
52. Andersen, M.F.; Nielsen, K.M.; Brinkmann, S. Meta-synthesis of qualitative research on return to work among employees with common mental disorders. *Scand. J. Work Environ. Health* **2012**, *38*, 93–104. [[CrossRef](#)] [[PubMed](#)]
53. Andersen, L.L.; Andersen, C.H.; Skotte, J.H.; Suetta, C.; Søgaard, K.; Saltin, B.; Sjøgaard, G. High-intensity strength training improves function of chronically painful muscles: Case-control and RCT studies. *Biomed. Res. Int.* **2014**, *2014*. [[CrossRef](#)] [[PubMed](#)]
54. Jay, K.; Brandt, M.; Sundstrup, E.; Schraefel, M.C.; Jakobsen, M.D.; Sjøgaard, G.; Andersen, L.L. Effect of individually tailored biopsychosocial workplace interventions on chronic musculoskeletal pain, stress and work ability among laboratory technicians: Randomized controlled trial protocol. *BMC Musculoskelet Disord.* **2014**, *15*. [[CrossRef](#)] [[PubMed](#)]



© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).