



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Genomic insights into members of the candidate phylum Hyd24-12 common in mesophilic anaerobic digesters

Kirkegaard, Rasmus Hansen; Dueholm, Morten Simonsen; McIlroy, Simon Jon; Nierychlo, Marta; Karst, Søren Michael; Albertsen, Mads; Nielsen, Per Halkjær

*Published in:*  
I S M E Journal

*DOI (link to publication from Publisher):*  
[10.1038/ismej.2016.43](https://doi.org/10.1038/ismej.2016.43)

*Creative Commons License*  
CC BY-NC-SA 4.0

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Kirkegaard, R. H., Dueholm, M. S., McIlroy, S. J., Nierychlo, M., Karst, S. M., Albertsen, M., & Nielsen, P. H. (2016). Genomic insights into members of the candidate phylum Hyd24-12 common in mesophilic anaerobic digesters. *I S M E Journal*, 10(10), 2352-2364. <https://doi.org/10.1038/ismej.2016.43>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## ORIGINAL ARTICLE

# Genomic insights into members of the candidate phylum Hyd24-12 common in mesophilic anaerobic digesters

Rasmus Hansen Kirkegaard<sup>1</sup>, Morten Simonsen Dueholm<sup>1</sup>, Simon Jon McIlroy, Marta Nierychlo, Søren Michael Karst, Mads Albertsen and Per Halkjær Nielsen  
Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark

**Members of the candidate phylum Hyd24-12 are globally distributed, but no genomic information or knowledge about their morphology, physiology or ecology is available. In this study, members of the Hyd24-12 lineage were shown to be present and abundant in full-scale mesophilic anaerobic digesters at Danish wastewater treatment facilities. In some samples, a member of the Hyd24-12 lineage was one of the most abundant genus-level bacterial taxa, accounting for up to 8% of the bacterial biomass. Three closely related and near-complete genomes were retrieved using metagenome sequencing of full-scale anaerobic digesters. Genome annotation and metabolic reconstruction showed that they are Gram-negative bacteria likely involved in acidogenesis, producing acetate and hydrogen from fermentation of sugars, and may play a role in the cycling of sulphur in the digesters. Fluorescence *in situ* hybridization revealed single rod-shaped cells dispersed within the flocs. The genomic information forms a foundation for a more detailed understanding of their role in anaerobic digestion and provides the first insight into a hitherto undescribed branch in the tree of life.**

The ISME Journal advance online publication, 8 April 2016; doi:10.1038/ismej.2016.43

## Introduction

Production of methane by anaerobic digestion (AD) is widely used to convert organic waste into biogas and forms an important part of the transition from fossil fuel to sustainable energy production. The AD process is divided into four sequential steps that are performed by specialized microbes: hydrolysis, fermentation (acidogenesis), acetogenesis (dehydrogenation) and methanogenesis (acetoclastic or hydrogenotrophic) (Angenent *et al.*, 2004). Hence, the overall function, stability and efficiency of the AD process are dependent on tightly coupled synergistic activities of the complex microbial communities (Schink, 1997; Weiland, 2010). However, the microbial communities in AD are still poorly understood, and relatively little is known about their diversity and function (Chouari *et al.*, 2005; Werner *et al.*, 2011; Sundberg *et al.*, 2013; De Vrieze *et al.*, 2015). In addition, most of the microorganisms have no pure culture representatives, and, given the

synergistic interactions of members of the community, a reductionist approach to understand the ecology of the system is not possible (Kaeberlein *et al.*, 2002; Fuhrman *et al.*, 2015).

The AD environment also harbours extensive diversity of previously uncharacterized bacterial phyla, often known only by their 16S rRNA gene sequence, making it an ideal environment for the study of novel bacterial lineages (Guermazi *et al.*, 2008; Pelletier *et al.*, 2008; Limam *et al.*, 2014; Sekiguchi *et al.*, 2015). New developments in single-cell genomics and metagenomics have in recent years provided a glimpse into the ecology and evolution of many novel candidate phyla (Dinis *et al.*, 2011; Albertsen *et al.*, 2013; Rinke *et al.*, 2013; Brown *et al.*, 2015; Nobu *et al.*, 2015; Sekiguchi *et al.*, 2015). The genomes have enabled construction of metabolic models that attempt to explain the physiology of these organisms in detail. The genome-based models form the basis of more extensive investigations, such as *in situ* single-cell characterization, metatranscriptomics and proteomics (Koch *et al.*, 2014).

In this study, extensive 16S rRNA gene amplicon sequencing was used to screen anaerobic digesters for the presence of members of the Hyd24-12 lineage, which remains one of the few known candidate phyla for which no genomic information is available with nothing known about their morphology, physiology or ecology (Rinke *et al.*, 2013). Selected

Correspondence: PH Nielsen, Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, Aalborg 9220, Denmark.  
E-mail: phn@bio.aau.dk

<sup>1</sup>These two authors contributed equally to this work.

Received 6 October 2015; revised 15 February 2016; accepted 22 February 2016

samples were subjected to metagenome sequencing and used for retrieval of three near-complete genomes of Hyd24-12 through differential coverage binning. The genomes were used for detailed metabolic reconstruction and design of oligonucleotide probes for the first *in situ* visualization of these hitherto unrecognized players in AD.

## Materials and methods

### *Sample collection and storage*

A total of 306 biomass samples were obtained from 29 anaerobic digesters at 17 Danish wastewater treatment facilities (see Supplementary Table S1). Most digesters were mesophilic (22), whereas 7 were thermophilic. A volume of 50 ml was sampled, homogenized and stored as 2 ml aliquots at  $-80^{\circ}\text{C}$  for DNA extraction.

For fluorescence *in situ* hybridization (FISH) analyses, diluted biomass samples (1:4 in  $1\times$  phosphate-buffered saline) were fixed with 4% (w/v) paraformaldehyde and stored in 50% (v/v) ethanol/ $1\times$  phosphate-buffered saline solution at  $-20^{\circ}\text{C}$ , as previously described by Daims *et al.* (2005).

### *DNA extraction*

DNA was extracted from anaerobic digester sludge using the FastDNA Spin kit for soil (MP Biomedicals, Santa Ana, CA, USA), following the standard protocol except for four times increased bead beating duration and a sludge input volume of 50  $\mu\text{l}$ . These digester-sample-specific modifications to the protocol were found to provide the best trade-off between DNA yield/biomass and DNA integrity (Supplementary Figure S4).

### *Community profiling with 16S rRNA gene amplicon sequencing*

Bacterial community profiling was carried out as recommended by Albertsen *et al.* (2015). The bacterial primers used were 27F (AGAGTTTGAT CCTGGCTCAG) (Lane, 1991) and 534R (ATTACC GCGGCTGCTGG) (Muyzer *et al.*, 1993), which amplify a DNA fragment of  $\sim 500$  bp of the 16S rRNA gene (variable V1–V3 region). PCR amplification was performed using  $1\times$  Platinum High fidelity buffer, 400  $\mu\text{M}$  dNTP, 1.5 mM  $\text{MgSO}_4$ , 2 mU Platinum Taq DNA Polymerase High Fidelity, 5  $\mu\text{M}$  illumina barcoded V1–V3 adaptor mix (see Supplementary Data 1), and 10 ng template DNA. PCR conditions were  $95^{\circ}\text{C}$  for 2 min, 30 cycles of  $95^{\circ}\text{C}$  for 20 s,  $56^{\circ}\text{C}$  for 30 s,  $72^{\circ}\text{C}$  for 60 s, and a final step of elongation at  $72^{\circ}\text{C}$  for 5 min. PCR products were purified using Agencourt AmpureXP (Beckman Coulter, Brea, CA, USA) with a ratio of 0.8 bead solution/PCR solution. Barcoded amplicons were pooled and paired-end sequenced on the Illumina MiSeq platform (v3 chemistry,  $2\times 300$  bp). The paired-end reads were trimmed using trimmomatic (v. 0.32) (Bolger *et al.*,

2014) and then merged using FLASH (v. 1.2.11) (Magoč and Salzberg, 2011). The reads were screened for potential PhiX contamination using USEARCH (v. 7.0.1090) (Edgar, 2010). The reads were clustered into operational taxonomic units (OTUs, sequence identity  $\geq 97\%$ ) using USEARCH and subsequently classified using the RDP classifier (Wang *et al.*, 2007) with the MiDAS database (v. 1.20) (McIlroy *et al.*, 2015). Further processing was carried out in the R environment (v. 3.1.2) using the ampvis package (Albertsen *et al.*, 2015) (v. 1.24.0), which wraps a number of packages including the phyloseq package (v. 1.8.2) (McMurdie and Holmes, 2013), ggplot2 (v. 1.0.1), reshape2 (v. 1.4.1) (Wickham, 2007), dplyr (v. 0.4.2), vegan (v. 2.3-0), knitr (v. 1.10.5), Biostings (v. 2.36.1), data.table (v. 1.9.4), DESeq2 (v. 1.8.1) (Love *et al.*, 2014), ggdendro (v. 0.1–15) and stringr (v. 1.0.0). The samples were subsampled to an even depth of 10 000 reads per sample, and the fraction of reads classified as Hyd24-12 was obtained. The survey data are available at the SRA with the accession IDs ERS861217–ERS861224.

### *In silico analysis of Hyd24-12 source locations*

The Genbank IDs of the sequences classified as Hyd24-12 in SILVA (v. 121, 1982 sequences in total) (Quast *et al.*, 2013) were used to download the corresponding Genbank files. The fields ‘isolation source’ and ‘PUBMED’ were extracted to classify the sequences as originating from either engineered or natural systems.

### *Metagenome sequencing, assembly and binning*

Illumina TruSeq DNA PCR free libraries were prepared for DNA extracts from three of the mesophilic digesters (Supplementary Table S1) according to the manufacturer’s protocol and paired-end sequenced on the Illumina HiSeq 2000 platform ( $2\times 150$  bp) and Illumina MiSeq platform (v3 chemistry,  $2\times 300$  bp). The metagenomic assembly and binning process was carried out as described by Albertsen *et al.* (2013) and detailed at ‘madsalbertsen.github.io/mmgenome/’. Unmerged reads were quality-trimmed and filtered using default settings in CLC Genomics Workbench (v. 7.5.1; CLC Bio, Aarhus, Denmark). The metagenomic reads were assembled separately for each plant using default settings in CLC Genomics Workbench. Reads were mapped to the assemblies using default settings in CLC Genomics Workbench. The assemblies and mappings were exported as .fasta and .sam files, respectively. The exported files and the mmgenome workflow script ‘data.generation.2.1.0.sh’ were used to generate the files necessary for the binning process. 16S rRNA gene sequences were extracted from the assemblies using ‘rRNA.sh’ and classified using the SINA Alignment service (SILVA v 121) (Pruesse *et al.*, 2012); essential genes were called

using Prodigal (Hyatt *et al.*, 2010). Binning was carried out using differential coverage binning in the R environment (v. 3.1.2, R Core Team, 2016) using the R package ‘mmgenome’ (github.com/MadsAlbertsen/mmgenome v. 0.4.1) (Albertsen *et al.*, 2013). The genome bins were checked for completeness, essential single copy genes and coverage distribution using CheckM (v. 0.9.7) (Parks *et al.*, 2015) and the metrics in the mmgenome package. Average nucleotide identity between the genome bins was calculated using JSpecies (Richter and Rosselló-Móra, 2009), and CRISPR arrays were identified with CRT (v. 1.1) (Bland *et al.*, 2007). The genome sequence data have been submitted to DDBJ/EMBL/GenBank databases under accession numbers LKHB000000000, LKHC000000000 and LKHD000000000.

### Genome sequence-based phylogenetic analysis

The genomes were placed within the reference genome tree of CheckM (Parks *et al.*, 2015) (v. 0.9.7) and subsequently visualized in ARB (Ludwig *et al.*, 2004).

### Phylogeny of the 16S rRNA gene and FISH probe design

Phylogenetic analysis and FISH probe design were performed with the ARB software package (Ludwig *et al.*, 2004). Potential probes were assessed *in silico* with the mathFISH software (Yilmaz *et al.*, 2011) for hybridization efficiencies of target and potentially weak, non-target matches (Yilmaz *et al.*, 2011). Unlabelled helper probes (Fuchs *et al.*, 2000) were designed for calculated inaccessible regions. Unlabelled competitor probes were designed for single-base mismatched non-target sequences (Manz *et al.*, 1992). The Ribosomal Database Project (RDP) PROBE MATCH function (Cole *et al.*, 2009) was used to identify non-target sequences with indels (McIlroy *et al.*, 2011). Probe validation and optimization were based on generated formamide dissociation curves (Daims *et al.*, 2005), where average relative fluorescent intensities, of at least 50 cells calculated with ImageJ software (National Institutes of Health, New York, NY, USA), were measured for varied hybridization buffer formamide concentrations in increments of 5% (v/v) over a range of 5–50% (v/v) (data not shown). Where available, weak base mismatch non-target axenic cultures were used for probe optimization, otherwise full-scale anaerobic digester sludge was used (Table 1).

### FISH

FISH was performed essentially as described by Daims *et al.* (2005). Probes were applied, with recommended competitors and helpers, at the stringency conditions given in Table 1 or their original publications. The NON-EUB nonsense probe was used as a negative hybridization control (Wallner *et al.*, 1993). Oligonucleotide probes were

**Table 1** FISH probes designed and optimized in this study

Probe	<i>Escherichia coli</i> pos.	Target group	Coverage <sup>a</sup>	Non-target hits	Sequence (5'–3')	Validated against	Required <sup>b</sup> [FA]%
Hyd24-12_468	468–488	Hyd24-12 clade B-1AC	5/6	0	GTA ACG TCA GGC AAT GGA CAT	Damhusaen full-scale AD	NA
Hyd24-12_468_h1	450–467	Helper for Hyd24-12_468	NA	NA	ATT GCT CCC ACT GCT GTT C	Damhusaen full-scale AD	Yes
Hyd24-12_468_h2	489–506	Helper for Hyd24-12_468	NA	NA	CCG GAG CTT TCT CTG GGG	Damhusaen full-scale AD	Yes
Hyd24-12_659	659–675	Hyd24-12 clade B-1AC	6/6	2	TCC GCA CAC CTC TCC CG	Damhusaen full-scale AD	NA
Hyd24-12_659_c1	659–675	Competitor for Hyd24-12_659	NA	NA	TCC ACA CAC CTC TCC CG	<i>Phenyllobacterium falsum</i> DSM18556	Yes
Hyd24-12_659_c2	659–675	Competitor for Hyd24-12_659	NA	NA	TCC GTA CAC CTC TCC CG	ND	NA
Hyd24-12_659_c3	659–675	Competitor for Hyd24-12_659	NA	NA	TCC GCT CAC CTC TCC CG	<i>Moricaulis maris</i> DSM4734	Yes
Hyd24-12_659_c4	659–675	Competitor for Hyd24-12_659	NA	NA	TCC GCA CTC CTC TCC CG	ND	ND
Hyd24-12_731	731–748	Hyd24-12 clade zEL51	15/17 <sup>d</sup>	0	ACT TGG CCA GTA TGC CGC	NA	ND
Hyd24-12_842	842–858	Hyd24-12 clades zEL51, Hyd-32 & B9.b18	17/31 <sup>d</sup>	0	CGG CTC TGA AGG GGT TGA	NA	ND
Hyd24-12_842_c1	842–858	Competitor for Hyd24-12_842	NA	NA	CGG CAC TGA AGG GGT TGA	NA	ND
Hyd24-12_842_h1	820–841	Helper for Hyd24-12_842	NA	NA	GAC CYC CAA AAC CYA GTR CCC A	NA	ND
Hyd24-12_842_h2	860–881	Helper for Hyd24-12_842	NA	NA	CGG GGY ACT TAA YGC GTT ARC T	NA	ND

Abbreviations: NA, not applicable; ND, not determined.

<sup>a</sup>Coverage of groups as defined in MIDAS Database, Release 1.21 (McIlroy *et al.*, 2015) (see Figure 2). Values given as group hits/ group totals.

<sup>b</sup>Requirement of accessory probes: competitor probes assessed on ability of the mismatch alone to prevent non-target binding at the recommended formamide concentration (note: it is recommended that un-validated competitor probes are included); helper probes assessed on ability to improve the fluorescence signal of the associated labelled probe.

<sup>c</sup>Determined optimal formamide concentration for use in FISH hybridizations.

<sup>d</sup>Together, these probes cover 26/31 sequences of the Hyd24-12 clades zEL51, Hyd-32 and B9.b18.



labelled on both the 3' and 5' ends with either 5(6)-carboxyfluorescein-*N*-hydroxysuccinimide ester (FLUOS) or with the sulphoindocyanine dyes (Cy3 and Cy5) (DOPE-FISH) (Stoecker *et al.*, 2010). Microscopic analysis was performed with an Axioskop epifluorescence microscope (Carl Zeiss, Oberkochen, Germany).

### Genome analysis


Genome annotation was performed in the 'MicroScope' annotation pipeline (Vallenet *et al.*, 2009, 2013). Automatic annotations were manually curated for all genes described using the integrated bioinformatics tools and the proposed annotation rules, which include an amino acid identity of at least 40% to classify homologues and an identity of at least 25% with the support of conserved domains to determine putative homologues (Vallenet *et al.*, 2009, 2013). The set of bioinformatics tools includes BlastP (Altschul *et al.*, 1990) homology searches against the full non-redundant protein sequence databank UniProt (Uniprot Consortium, 2014) and against the well-annotated model organisms *Escherichia coli* K-12 and *Bacillus subtilis* 168 (Vallenet *et al.*, 2013), enzymatic classifications based on COG (Tatusov *et al.*, 2003), InterPro (Mitchell *et al.*, 2015), FIGfam (Meyer *et al.*, 2009) and PRIAM (Claudel-Renard *et al.*, 2003) profiles, and prediction of protein localization using the TMHMM (Sonnhammer *et al.*, 1998), SignalP (Bendtsen *et al.*, 2004) and PSORTb (Gardy *et al.*, 2005) tools. Synteny maps (i.e. conservation of local gene order) were used to validate the annotation of genes located within conserved operons (Vallenet *et al.*, 2009). Metabolic pathways were subsequently identified

with the assistance of the integrated MicroCyc (Vallenet *et al.*, 2009) and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases (Kanehisa *et al.*, 2014).

## Results and discussion

### Survey of 16S rRNA genes of Hyd24-12 in anaerobic digesters

The survey of 22 full-scale mesophilic and 7 thermophilic anaerobic digesters from 17 Danish wastewater treatment plants over 3 years revealed that members of the Hyd24-12 lineage were stably present in most mesophilic but no thermophilic anaerobic digesters (Figure 1). In most mesophilic digesters, they were among the five most abundant bacterial OTUs and constituted around 1–3% and, in some cases, up to 8.2% of all sequenced bacterial reads (see Supplementary Figure S1). No 16S rRNA gene sequences from Hyd24-12 were detected in the incoming surplus sludge from the activated sludge treatment plants, which demonstrates that these bacteria were actively growing in the digesters. The other abundant bacterial phyla in the mesophilic digesters were Actinobacteria, Firmicutes, Chloroflexi, Synergistetes and Bacteroidetes (Figure 1). The best (LCA) classification is shown in Figure 1, but the lack of closely related organisms in the databases and a curated taxonomy hampers taxonomic classification for a number of the most abundant OTUs. In general, the abundance stability of these top genera was high, and that may be due to relatively similar growth conditions for all digesters: feed was primary sludge and surplus activated sludge, temperature in the interval 34–37 °C, pH 7.1–8.2 and total ammonium 0.57–1.1 g N/l (see Supplementary Table S1).



**Figure 1** Heatmap of the 25 most abundant bacterial OTUs in mesophilic digesters at wastewater treatment plants along with their abundance in thermophilic digesters at 17 wastewater treatment plants. The OTU classified as belonging to the Hyd24-12 candidate phylum (purple) was detected exclusively in mesophilic reactors. Classification levels presented are phylum, class, order, family and genus and are separated by a semicolon. The field is empty where no classification at a given level could be provided. The abundance profiles show mean abundances for plants with more reactors (1–4 reactors at each WWTP) and 2–97 samples for each plant over 4 years (Supplementary Table S1). The OTUs are sorted on the basis of the mean abundance across the mesophilic samples.

*In silico* analysis of 16S rRNA gene sequences within Hyd24-12 of SILVA (Quast *et al.*, 2013) from other surveys confirmed that members of Hyd24-12 are widespread in anaerobic environments. The sequences originate from 48 separate studies, with engineered systems such as anaerobic bioreactors accounting for 10 studies, and natural systems such as marine sediments, microbial mats in hydrogen, methane-rich waters and mud volcanoes accounting for 38 studies (see Supplementary References). Furthermore, the 48 studies show that members of Hyd24-12 are globally dispersed (Supplementary Figure S3 and Supplementary Table S2) and are potentially important in many microbial ecosystems besides ADs (Mills *et al.*, 2005; Harris *et al.*, 2012). Some of the surveys of full-scale anaerobic digesters detected some Hyd24-12 sequences (e.g., De Vrieze *et al.*, 2015), while others did not (Sundberg *et al.*, 2013). This was likely because they used the RDP database, where Hyd24-12 sequences are classified as ‘unclassified bacteria’.

#### Recovering genomic information from Hyd24-12

Three full-scale anaerobic digesters were sampled for metagenomic analyses. To ensure differential abundance of microorganisms needed to bin genomes based on coverage profiles (Albertsen *et al.*, 2013), biomass samples were either taken from the sludge and foam layer of reactors or from the same reactor weeks apart. More than 50 gigabases of metagenomic data were generated, and population genomes were recovered by differential coverage binning (Albertsen *et al.*, 2013) from each of the three plants (Table 2). The three population genomes were ~2.2 Mbp with a GC content of ~64%, and the completeness of the genomes were estimated by CheckM (Parks *et al.*, 2015) to be between 86% and 91% with less than 2.2% estimated contamination (Table 2). However, the level of completeness may be underestimated, given that members of the Hyd24-12 are distantly related to other characterized

organisms, and the genes used in the marker sets might be too divergent or simply not present (Rinke *et al.*, 2013; Brown *et al.*, 2015; Sekiguchi *et al.*, 2015). The three genomes each contained a single rRNA operon and shared identical 16S rRNA gene sequences, which suggests that they belong to the same species (Yarza *et al.*, 2014). The JSpecies program determined that these three genomes shared between 99.8% and 99.9% average nucleotide identity (ANIb), supporting the close taxonomic relationship observed from the 16S rRNA gene analysis (Kim *et al.*, 2014). In order to further evaluate the similarity between the strains, the raw metagenome reads from each digester were mapped to the assembled Hyd24-12 genomes obtained from the other two digesters. Complete coverage of all genomes with the metagenome reads from the other digester revealed that the Hyd24-12 genomes were almost identical. This also indicates, along with the high ANIb, that the genomes are more complete than estimated in Table 2 by CheckM. Indeed, the data suggested that the three strains might actually be variants of the same strain with single-nucleotide polymorphisms only. This is very interesting as the digesters were from different parts of Denmark without any exchange of sludge or feed. This could indicate that they are highly adapted to the specific AD environment in this type of mesophilic digesters.


#### Hyd24-12 phylogeny, FISH probe design and morphology

The 16S rRNA genes obtained have a sequence identity of 86% with the original clone Hyd24-12 sequence (AJ535232) (Knittel *et al.*, 2003) and classify to the Hyd24-12-lineage (Figure 2a). Additional phylogenetic analyses, based on the genome sequence, placed the Hyd24-12 genomes within the Fibrobacteres-Chlorobi-Bacteroidetes superphylum (Figure 3). The Hyd24-12 genomes are distantly related to all currently available genomes, supporting its status as a novel phylum.

**Table 2** Genome statistics for the three Hyd24-12 genomes

Genome bin identifier	Dam_1	Ran_1	Vib_1
Source digester	Damhusaaen	Randers	Viborg
Closest env. 16S clone	6E6_cons (EF688250)	6E6_cons (EF688250)	6E6_cons (EF688250)
No. of contigs	247	168	224
Total length	2 182 231	2 013 453	2 188 467
Longest contig	90 522	171 008	113 480
N50 (contigs)	27 448	38 589	27 030
GC (%)	63.8	64.2	63.7
Genome completeness (%) <sup>a</sup>	86.1	90.7	90.7
Genome contamination (%) <sup>a</sup>	2.17	1.09	2.17
No. of tRNA genes	44	41	48
rRNA genes found in genome	5S, 16S, 23S	5S, 16S, 23S	5S, 16S
No. of CDS	2336	1995	2349
No. of CRISPR array	2 (33 repeats in total)	0	0
Coding density (%)	91.8	93.4	92.8

<sup>a</sup>Estimates provided by CheckM.



**Figure 2** Phylogenetic analysis and design of FISH probes. **(a)** Maximum-likelihood (PhyML) 16S rRNA gene phylogenetic tree of target groups and selected related sequences (all > 1200 bp). Phylogenetic classification is taken from the MiDAS database (Release 1.21), which is a version of the SILVA database (Release 119 NR99) (Quast *et al.*, 2013) curated for activated sludge sequences (McIlroy *et al.*, 2015). Clades of the Hyd24-12 lineage are shown in brackets. Probes covering clades are shown in red. The scale bar represents substitutions per nucleotide base. Bootstrap values from 100 re-samplings are indicated for branches when > 50%. **(b)** Composite FISH micrographs of the B-1AC clade members in the Randers anaerobic digester sludge. B-1AC cells appear red (Hyd24-12\_659, Cy3), other bacterial cells appear blue (EUBmix probe set (Amann *et al.*, 1990; Daims *et al.*, 1999), Cy5) and archaeal cells appear green (ARCH915 (Stahl and Amann, 1991), FLUOS).

Several FISH probes were designed to target different clades within the phylum. In the MiDAS taxonomy (v. 1.21) (McIlroy *et al.*, 2015), a version of the SILVA taxonomy (Quast *et al.*, 2013) that is curated for activated sludge-related organisms, the Hyd24-12 lineage is delineated into four clades, designated B-1AC, zEL51, Hyd-32 and B9.18. The Hyd24-12\_468 and Hyd24-12\_659 probes were designed to cover the B-1AC clade, which includes the Hyd24-12 genome sequences obtained in this study (Figure 2). The former probe covers almost all the B-1AC sequences, with the closest non-target sequence match having three internal base mismatches. The Hyd24-12\_659 probe is less specific, having one perfectly matched non-target sequence and several with mismatches not covered by the

competitor probes. Overlap in the coverage of these two probes, labelled with different fluorochromes, allows greater confidence in their specificity. A suitable probe to cover the entire Hyd24-12 lineage was not found. However, the Hyd24-12\_731 and Hyd24-12\_842 probes provide good coverage of the other sequences in the phylum (see Table 1). As sequences covered by these additional probes were not detected in the full-scale anaerobic digesters studied here, optimization and assessment of these probes were not pursued.


When applied to several full-scale anaerobic digester sludge samples, the Hyd24-12\_468 and Hyd24-12\_659 probes hybridized to small rods, approx.  $2 \times 0.4 \mu\text{m}$  in size, dispersed through the flocs (see Figure 2b). Good overlap was observed for



these probes, supporting their specificity. Of the two probes, a much higher signal was observed for the Hyd24-12\_659 probe. There was no observed overlap between the signal of two Hyd24-12 probes and the universal bacterial EUBmix probe set (see Figure 2b), which is supported by the absence of the target site for the probes of the latter in the Hyd24-12 sequences. Quantitative FISH was very difficult to carry out in the digesters due to high levels of background fluorescence. Instead, abundance estimates were carried out for the domains Bacteria, Archaea, Eukarya, and the Hyd24-12 lineage, based on read mapping from the PCR free metagenomes to the 16S rRNA genes of the MiDAS database. It showed that Archaea constituted 4–9% of the reads in sludge samples and 7–13% in foam samples. Reads from the Hyd24-12 lineage constituted 0.4–3.5% in the different samples (Supplementary Table S3).

### Morphology and motility

The rod shape morphology of B-1AC clade organisms observed by FISH is supported by *mreBCD* and *mrdAB* operons in the Hyd24-12 genomes (see Supplementary Data 2). These operons encode proteins involved in the formation of membrane-bound actin filaments, which are essential for the biogenesis of rod-shape stabilizing peptidoglycans



**Figure 3** Phylogenetic position of the Hyd24-12 genomes in the reference genome tree generated by CheckM. The CheckM tree is inferred from the concatenation of 43 conserved marker genes and incorporates 2052 finished and 3604 draft genomes from the IMG database (Parks *et al.*, 2015).

along the lateral cell wall of rod-shaped bacteria (Kruse *et al.*, 2003, 2005; Osborn and Rothfield, 2007; Bendezú and de Boer, 2008).

The cell envelope characteristics of genome-sequenced bacteria can be determined based on PFAM protein families that are substantially enriched or depleted in archetypical monoderm lineages relative to archetypical diderm lineages (Albertsen *et al.*, 2013). A search for such protein families in the Hyd24-12 genomes revealed an archetypical diderm cell envelope with lipopolysaccharides (see Supplementary Figure S2).


None of the Hyd24-12 genomes encode any flagella-related proteins, suggesting limited motility. However, genes associated with type IV pili were identified using the PilFind algorithm (see Supplementary Data 2) (Imam *et al.*, 2011). These pili enable the bacteria to generate surface-associated twitching motility. This allows them to move effectively through environments that contain shear-thinning viscoelastic fluids, such as the extracellular polymeric substances of biofilms (Conrad *et al.*, 2011; Jin *et al.*, 2011). In addition to motility, type IV pili play a role in the attachment to living and non-living surfaces, including those of other bacteria (Giltner *et al.*, 2012).

No genes associated with spore formation were detected in the Hyd24-12 genomes. This suggests that the Hyd24-12 genomes investigated represent non-sporulating bacteria.

### Energy metabolism

The three genomes do not contain any genes for respiration with oxygen, nitrate/nitrite or Fe(III) and seem primarily to have a fermentative metabolism. However, the genomes indicate that the organisms may be able to use elemental sulphur as an electron acceptor, see below. The Hyd24-12 genomes encode a complete glycolysis pathway, along with the non-oxidative branch of the pentose phosphate pathway (Figure 4 and see Supplementary Data 2). This allows Hyd24-12 to potentially catabolize a wide range of hexoses and pentoses to pyruvate, thereby providing the cell with energy in the form of ATP and reducing equivalents in the form of NADH (Stincone *et al.*, 2014). The sugars are probably obtained from the environment through a major facilitator superfamily transporter at the expense of the proton motive force (Madej, 2014; Wisedchaisri *et al.*, 2014). The transporter does not share similarity (>30%) with any experimentally validated transporters, and it is therefore impossible to infer a specific substrate preference. It is known that primary sludge and activated sludge fed into the digesters contain many different polysaccharides (Raunkjaer *et al.*, 1994; Frølund *et al.*, 1996). No genes encoding for extracellular glycosylases were identified, which might indicate that Hyd24-12 is reliant on the hydrolytic action of other organisms present within the anaerobic digesters.





**Figure 4** Metabolic model of Hyd24-12 species in mesophilic anaerobic digesters, based on the annotated genome sequences. Selected metabolic pathways important for the bacteria in the anaerobic digestion process are highlighted. Numbers correspond to annotated genes in Supplementary Data 1.

Hyd24-12 encodes for the complete pathway for glycogen biosynthesis and catabolism (Figure 4 and see Supplementary Data 2) (Preiss *et al.*, 1983; Wilson *et al.*, 2010). Hence, glycogen may serve as a carbon and energy storage which can be utilized to mitigate fluctuations in substrate availability. The Hyd24-12 genomes did not encode for pathways for other storage compounds such as trehalose or polyhydroxyalkanoates.

There are limited catabolic options for the pyruvate formed, for example, by glycolysis. The tricarboxylic acid cycle of Hyd24-12 is incomplete (8 of 10 key enzymes are missing) and probably non-functional. However, pyruvate can be converted into acetyl-CoA by a pyruvate ferredoxin oxidoreductase, providing additional reducing equivalents in the form of reduced ferredoxin (Figure 4 and see Supplementary Data 2) (Menon and Ragsdale, 1997). Acetyl-CoA can then be converted into acetate by the action of phosphate acetyltransferase and acetate kinase, thus providing the bacterium with additional ATP (Latimer and Ferry, 1993; Mai and Adams, 1996).

All three Hyd24-12 genomes also encode for two aldehyde ferredoxin oxidoreductases (Figure 4 and see Supplementary Data 2). These may be used to oxidize formaldehyde and acetaldehyde to formate and acetate, respectively, providing the cell with

energy in the form of additional reduced ferredoxin (Mukund and Adams, 1991). However, the enzyme may also be used in the reverse reaction to regenerate oxidized ferredoxin. The presence of a membrane-embedded, energy-conserving hydrogenase allows the cell to establish a proton motive force, based on the energy-rich reduced ferredoxin, which reduces  $H^+$  to  $H_2$  in the process (Strittmatter *et al.*, 2009). The energy stored in the proton motive force may then be harvested through an ATP synthase to yield ATP.

High concentrations of  $H_2$  inhibit glycolysis and acidogenesis due to thermodynamic considerations (Huang *et al.*, 2015). Hyd24-12 therefore needs a way to remove excess  $H_2$ . This can be achieved by syntrophic association with other microorganisms, or internally by the action of a cytosolic hydrogenase, which couples the oxidation of  $H_2$  with the reduction of  $NAD^+$  (Figure 4 and see Supplementary Data 2). Alternatively, Hyd24-12 may employ a sulfhydrogenase to couple the oxidation of  $H_2$  to  $H^+$  with the reduction of elemental sulphur ( $S^0$ ) or polysulphide to hydrogen sulphide ( $H_2S$ ) as is seen for *Pyrococcus furiosus* (Mukund and Adams, 1991). The genomes do not indicate a potential for sulphate reduction. Elemental sulphur is continuously produced in the digesters because activated sludge fed into the digesters contains oxidized iron (Fe(III)), which in the presence of sulphide produces  $S^0$  and

black iron sulphide (FeS) (Rasmussen and Nielsen, 1996; Nielsen *et al.*, 2005; Omri *et al.*, 2011). Sulphide is a normal compound in digesters and is produced from amino acids and reduction of sulphate. Notably, other studies have also detected members of the Hyd24-12 phylum in sulphur-rich environments such as hydrothermal vents, sulphur-rich springs and sediments (Elshahed *et al.*, 2003; Schauer *et al.*, 2011; Pjevac *et al.*, 2014). Thus, Hyd24-12 related organisms potentially play a role in sulphur transformations in digesters and other environments. Such a role requires further investigation.

The Hyd24-12 genomes do not contain the genes required for fatty acid  $\beta$ -oxidation or for the catabolism of amino acids. Sugars are therefore considered the primary energy source of the Hyd24-12 in anaerobic digesters.

Whereas Hyd24-12 is able to take up carbon in the form of amino acids, carbohydrates, etc., it is unable to carry out fixation of CO<sub>2</sub> as such genes are missing.

#### Amino acid and nitrogen metabolism

Based on the genome annotations, Hyd24-12 is only predicted to be able to synthesize few amino acids (glycine, serine, cysteine, threonine, asparagine, aspartate, glutamate and glutamine). Accordingly, Hyd24-12 might rely on amino acids present within the environment. As most amino acids are found as proteins, which cannot be taken up by the bacterium, Hyd24-12 needs a way to degrade these polymers, and this is achieved by the action of multiple extracellular proteases encoded in the genome, which are likely secreted in a Sec- or Tat-dependent mechanism (Natale *et al.*, 2008) (see Supplementary Data 2). The cells may subsequently import the amino acids using ABC-transporters encoded in the genome. Owing to the lack of experimentally validated homologues from closely related species, it is not possible to predict the substrate specificity of these transporters. A reduced capacity of microorganisms for synthesizing amino acids is known from strict symbionts and, recently, also from a number of candidate phyla with very small genomes (<1 Mbp) (Brown *et al.*, 2015). However, the relatively large size of the Hyd24-12 genomes (~2.2 Mbp) and their dispersed growth in the anaerobic sludge suggest that they are not strict symbionts.

Hyd24-12 does not have the necessary pathways for fixation of nitrogen. The nitrogen metabolism of Hyd24-12 is generally limited. Amino acids may also represent a source of nitrogen. However, nitrogen can also be obtained from ammonium assimilation via the glutamine synthetase/glutamate synthase pathways (Bravo and Mora, 1988).

#### Oxidative stress protection

The three Hyd24-12 genomes each contains a gene cluster encoding for a superoxide reductase, nitric oxide reductase and ferroxidase. These genes are

probably involved in resistance against oxidative stress, and may allow the bacteria to survive in the presence of oxygen. However, 16S rRNA gene sequences from Hyd24-12 have only been observed in oxygen-depleted environments.

#### Ecological significance and concluding remarks

This study applied metagenomic sequencing to obtain genomes from the candidate phylum Hyd24-12 and provides the first morphological and physiological information for the lineage. Members of the phylum were shown to be very abundant and stably present in mesophilic anaerobic digesters, occasionally accounting for the most abundant OTU in the samples, but absent in thermophilic reactors. This indicates that they are likely to play a substantial role in the ecology of mesophilic AD systems at wastewater treatment plants fed with primary sludge and surplus activated sludge. Metabolic reconstruction based on the genomic information showed that members of Hyd24-12 are likely to be fermenters relying on simple sugars. In addition, they may also use elemental sulphur as an electron acceptor, thus forming part of the microbial cycling of sulphur in anaerobic systems and partly responsible for production of hydrogen sulphide. Sulphide is unwanted in the biogas due to toxicity and corrosion (Syed *et al.*, 2006), but will also provide more elemental sulphur by reacting with incoming Fe(III). In that case, members of Hyd24-12 may compete with the methanogens for organics. An *in silico* investigation of environmental 16S rRNA gene surveys suggests that members of the phylum are present in anaerobic environments, often associated with sulphurous compounds and methane production, such as sediment mats and anaerobic bioreactors. The fact that the genomes are auxotrophic for several amino acids and lacking putative secreted glycoside hydrolases also indicates a strict reliance on other organisms for nutrients. The genomes generated in this study provide the foundation for future detailed analyses of members of the phylum, such as metatranscriptomics and metaproteomics. The design of FISH probes for the phylum also revealed their morphology and spatial arrangement in anaerobic digesters and will also facilitate future *in situ* investigations of the phylum in digesters and other environments.

Phylogenetic and genomic analyses of the three Hyd24-12 genomes classified them as a single species within a novel phylum located within the Fibrobacteres-Chlorobi-Bacteroidetes superphylum.

We propose the following taxonomic names for the novel genus and species of Hyd24-12:

- '*Candidatus Fermentibacter*' gen. nov.
- '*Candidatus Fermentibacter daniensis*' gen. et sp. nov.

Based on this, we propose the following names for the phylum, class, order, and family:

- ‘*Candidatus Fermentibacteria*’ phyl. nov.
- ‘*Candidatus Fermentibacteria*’ classis nov.
- ‘*Candidatus Fermentibacterales*’ ord. nov.
- ‘*Candidatus Fermentibacteraceae*’ fam. nov.

**Etymology.** *Fermentibacter* (Fer.men.ti.bac'ter. M.L. n. ferment -um to ferment, Gr. dim. n. bakterion a small rod, M.L. neut. n. Fermentibacter a small fermenting rod-shaped bacterium). *Fermentibacter daniensis* (da.ni. ensis. M.L. fem. adj. daniensis, pertaining to Dania, the Medieval Latin name for the country of Denmark, where the species was first discovered).

## Conflict of Interest

The authors declare no conflict of interest.

## Acknowledgements

This study was supported by the Villum Foundation, the Grundfos Foundation, Aalborg University and Innovation Fund Denmark (NomiGas project).

## References

- Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. (2013). Genome sequences of rare, uncultured bacteria obtained by differential coverage binning of multiple metagenomes. *Nat Biotechnol* **31**: 533–538.
- Albertsen M, Karst SM, Ziegler AS, Kirkegaard RH, Nielsen PH. (2015). Back to basics—the influence of DNA extraction and primer choice on phylogenetic analysis of activated sludge communities. *PLoS One* **10**: e0132783.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403–410.
- Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. (1990). Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl Environ Microbiol* **56**: 1919–1925.
- Angenent LT, Karim K, Al-Dahhan MH, Wrenn BA, Domínguez-Espinosa R. (2004). Production of bioenergy and biochemicals from industrial and agricultural wastewater. *Trends Biotechnol* **22**: 477–485.
- Bendežú FO, de Boer PAJ. (2008). Conditional lethality, division defects, membrane involution, and endocytosis in *mre* and *mrd* shape mutants of *Escherichia coli*. *J Bacteriol* **190**: 1792–1811.
- Bendtsen JD, Nielsen H, von Heijne G, Brunak S. (2004). Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* **340**: 783–795.
- Bland C, Ramsey TL, Sabree F, Lowe M, Brown K, Kypides NC et al. (2007). CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics* **8**: 209.
- Bolger AM, Lohse M, Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.
- Bravo A, Mora J. (1988). Ammonium assimilation in *Rhizobium phaseoli* by the glutamine synthetase-glutamate synthase pathway. *J Bacteriol* **170**: 980–984.
- Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A et al. (2015). Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523**: 208–211.
- Chouari R, Le Paslier D, Dauga C, Daegelen P, Weissenbach J, Sghir A. (2005). Novel major bacterial candidate division within a municipal anaerobic sludge digester. *Appl Environ Microbiol* **71**: 2145–2153.
- Claudel-Renard C, Chevalet C, Faraut T, Kahn D. (2003). Enzyme-specific profiles for genome annotation: PRIAM. *Nucleic Acids Res* **31**: 6633–6639.
- Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ et al. (2009). The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res* **37**: D141–D145.
- Conrad JC, Gibiansky ML, Jin F, Gordon VD, Motto DA, Mathewson MA et al. (2011). Flagella and pili-mediated near-surface single-cell motility mechanisms in *P. aeruginosa*. *Biophys J* **100**: 1608–1616.
- Daims H, Brühl A, Amann R, Schleifer KH, Wagner M. (1999). The domain-specific probe EUB338 is insufficient for the detection of all bacteria: development and evaluation of a more comprehensive probe set. *Syst Appl Microbiol* **22**: 434–444.
- Daims H, Stoecker K, Wagner M. (2005). Fluorescence in situ hybridization for the detection of prokaryotes. *Mol Microb Ecol* **213**: 239.
- De Vrieze J, Saunders AM, He Y, Fang J, Nielsen PH, Verstraete W et al. (2015). Ammonia and temperature determine potential clustering in the anaerobic digestion microbiome. *Water Res* **75**: 312–323.
- Dinis JM, Barton DE, Ghadiri J, Surendar D, Reddy K, Velasquez F et al. (2011). In search of an uncultured human-associated TM7 bacterium in the environment. *PLoS One* **6**: 1–8.
- Edgar RC. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460–2461.
- Elshahed MS, Senko JM, Najjar FZ, Kenton SM, Roe BA, Dewers TA et al. (2003). Bacterial diversity and sulfur cycling in a mesophilic sulfide-rich spring. *Appl Environ Microbiol* **69**: 5609–5621.
- Frolund B, Palmgren R, Keiding K, Nielsen PH. (1996). Extraction of extracellular polymers from activated sludge using a cation exchange resin. *Water Res* **30**: 1749–1758.
- Fuchs BM, Glockner FO, Wulf J, Amann R. (2000). Unlabeled helper oligonucleotides increase the *in situ* accessibility to 16S rRNA of fluorescently labeled oligonucleotide probes. *Appl Environ Microbiol* **66**: 3603–3607.
- Fuhrman JA, Cram JA, Needham DM. (2015). Marine microbial community dynamics and their ecological interpretation. *Nat Rev Microbiol* **13**: 133–146.
- Gardy JL, Laird MR, Chen F, Rey S, Walsh CJ, Ester M et al. (2005). PSORTb v.2.0: expanded prediction of bacterial protein subcellular localization and insights gained



- from comparative proteome analysis. *Bioinformatics* **21**: 617–623.
- Giltner CL, Nguyen Y, Burrows LL. (2012). Type IV pilin proteins: versatile molecular modules. *Microbiol Mol Biol Rev* **76**: 740–772.
- Guermazi S, Daegelen P, Dauga C, Rivière D, Bouchez T, Godon JJ *et al.* (2008). Discovery and characterization of a new bacterial candidate division by an anaerobic sludge digester metagenomic approach. *Environ Microbiol* **10**: 2111–2123.
- Harris JK, Gregory Caporaso J, Walker JJ, Spear JR, Gold NJ, Robertson CE *et al.* (2012). Phylogenetic stratigraphy in the Guerrero Negro hypersaline microbial mat. *ISME J* **7**: 50–60.
- Huang W, Wang Z, Zhou Y, Ng WJ. (2015). The role of hydrogenotrophic methanogens in an acidogenic reactor. *Chemosphere* **140**: 40–46.
- Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. (2010). Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**: 119.
- Imam S, Chen Z, Roos DS, Pohlschröder M. (2011). Identification of surprisingly diverse type IV pili, across a broad range of Gram-positive bacteria. *PLoS One* **6**: e28919.
- Jin F, Conrad JC, Gibiansky ML, Wong GCL. (2011). Bacteria use type-IV pili to slingshot on surfaces. *Proc Natl Acad Sci USA* **108**: 12617–12622.
- Kaerberlein T, Lewis K, Epstein SS. (2002). Isolating ‘uncultivable’ microorganisms in pure culture in a simulated natural environment. *Science* **296**: 1127–1129.
- Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. (2014). Data, information, knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res* **42**: 199–205.
- Kim M, Oh H-SS, Park S-CC, Chun J. (2014). Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* **64**: 346–351.
- Knittel K, Boetius A, Lemke A, Eilers H, Lochte K, Pfannkuche O *et al.* (2003). Activity, distribution, and diversity of sulfate reducers and other bacteria in sediments above gas hydrate (Cascadia Margin, Oregon). *Geomicrobiol J* **20**: 269–294.
- Koch H, Galushko A, Albertsen M, Schintlmeister A, Gruber-Dorninger C, Lucker S *et al.* (2014). Growth of nitrite-oxidizing bacteria by aerobic hydrogen oxidation. *Science* **345**: 1052–1054.
- Kruse T, Bork-Jensen J, Gerdes K. (2005). The morphogenetic MreBCD proteins of *Escherichia coli* form an essential membrane-bound complex. *Mol Microbiol* **55**: 78–89.
- Kruse T, Møller-Jensen J, Løbner-Olesen A, Gerdes K. (2003). Dysfunctional MreB inhibits chromosome segregation in *Escherichia coli*. *EMBO J* **22**: 5283–5292.
- Lane DJ. (1991). 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M (eds). *Nucleic Acid Techniques in Bacterial Systematics* John Wiley and Sons: West Sussex.
- Latimer MT, Ferry JG. (1993). Cloning, sequence analysis, and hyperexpression of the genes encoding phosphotransacetylase and acetate kinase from *Methanosarcina thermophila*. *J Bacteriol* **175**: 6822–6829.
- Limam RD, Chouari R, Mazéas L, Wu T, Di, Li T, Grossin-Debattista J *et al.* (2014). Members of the uncultured bacterial candidate division WWE1 are implicated in anaerobic digestion of cellulose. *MicrobiologyOpen* **3**: 157–167.
- Love MI, Huber W, Anders S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**: 550.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar A *et al.* (2004). ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–1371.
- Madej MG. (2014). Function, structure and evolution of the major facilitator superfamily: the LacY manifesto. *Adv Biol* **2014**: 20.
- Magoč T, Salzberg SL. (2011). FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **27**: 2957–2963.
- Mai X, Adams MW. (1996). Purification and characterization of two reversible and ADP-dependent acetyl coenzyme A synthetases from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Bacteriol* **178**: 5897–5903.
- Manz W, Amann R, Ludwig W, Wagner M, Schleifer K-H. (1992). Phylogenetic oligodeoxynucleotide probes for the major subclasses of Proteobacteria: problems and solutions. *Syst Appl Microbiol* **15**: 593–600.
- McIlroy SJ, Saunders AM, Albertsen M, Nierychlo M, McIlroy B, Hansen AA *et al.* (2015). MiDAS: the field guide to the microbes of activated sludge. *Database* **2015**: bav062.
- McIlroy SJ, Tillett D, Petrovski S, Seviour RJ. (2011). Non-target sites with single nucleotide insertions or deletions are frequently found in 16S rRNA sequences and can lead to false positives in fluorescence *in situ* hybridization (FISH). *Environ Microbiol* **13**: 33–47.
- McMurdie PJ, Holmes S. (2013). phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* **8**: e61217.
- Menon S, Ragsdale SW. (1997). Mechanism of the clostridium thermoaceticum pyruvate: ferredoxin oxidoreductase: evidence for the common catalytic intermediacy of the hydroxyethylthiamine pyrophosphate radical. *Biochemistry* **36**: 8484–8494.
- Meyer F, Overbeek R, Rodriguez A. (2009). FIGfams: yet another set of protein families. *Nucleic Acids Res* **37**: 6643–6654.
- Mills HJ, Martinez RJ, Story S, Sobecky PA. (2005). Characterization of microbial community structure in Gulf of Mexico gas hydrates: comparative analysis of DNA- and RNA-derived clone libraries. *Appl Environ Microbiol* **71**: 3235–3247.
- Mitchell A, Chang H-YH-Y, Daugherty L, Fraser M, Hunter S, Lopez R *et al.* (2015). The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Res* **43**: D213–D221.
- Mukund S, Adams MWW. (1991). The novel tungsten-iron-sulfur protein of the hyperthermophilic archaeobacterium, *Pyrococcus furiosus*, is an aldehyde ferredoxin oxidoreductase: evidence for its participation in a unique glycolytic pathway. *J Biol Chem* **266**: 14208–14216.
- Muyzer G, De Waal EC, Uitterlinden AG. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* **59**: 695–700.
- Natale P, Brüser T, Driessen AJM. (2008). Sec- and Tat-mediated protein secretion across the bacterial



- cytoplasmic membrane—distinct translocases and mechanisms. *Biochim Biophys Acta* **1778**: 1735–1756.
- Nielsen AH, Lens P, Vollertsen J, Hvitved-Jacobsen T. (2005). Sulfide-iron interactions in domestic wastewater from a gravity sewer. *Water Res* **39**: 2747–2755.
- Nobu MK, Narihiro T, Rinke C, Kamagata Y, Tringe SG, Woyke T *et al.* (2015). Microbial dark matter ecogenomics reveals complex synergistic networks in a methanogenic bioreactor. *ISME J* **9**: 1710–1722.
- Omri I, Bouallagui H, Aouidi F, Godon JJ, Hamdi M. (2011). H<sub>2</sub>S gas biological removal efficiency and bacterial community diversity in biofilter treating wastewater odor. *Bioresour Technol* **102**: 10202–10209.
- Osborn MJ, Rothfield LJ. (2007). Cell shape determination in *Escherichia coli*. *Curr Opin Microbiol* **10**: 606–610.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. (2015). CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* **25**: 1043–1055.
- Pelletier E, Kreimeyer A, Bocs S, Rouy Z, Gyapay G, Chouari R *et al.* (2008). ‘*Candidatus* Cloacamonas acidaminovorans’: genome sequence reconstruction provides a first glimpse of a new bacterial division. *J Bacteriol* **190**: 2572–2579.
- Pjevac P, Kamysny A, Dykema S, Musmann M. (2014). Microbial consumption of zero-valence sulfur in marine benthic habitats. *Environ Microbiol* **16**: 3416–3430.
- Preiss J, Yung S, Baecker PA. (1983). Regulation of bacterial glycogen synthesis. *Mol Cell Biochem* **57**: 61–80.
- Pruesse E, Peplies J, Glöckner FO. (2012). SINA: accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* **28**: 1823–1829.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P *et al.* (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* **41**: 590–596.
- R Core Team. (2016). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing: Vienna, Austria. Available from <https://www.R-project.org/>.
- Rasmussen H, Nielsen PH. (1996). Iron reduction in activated sludge measured with different extraction techniques. *Water Res* **30**: 551–558.
- Raunkjaer K, Hvitved-Jacobsen T, Nielsen PH. (1994). Measurement of pools of protein, carbohydrate and lipid in domestic wastewater. *Water Res* **28**: 251–262.
- Richter M, Rosselló-Móra R. (2009). Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* **106**: 19126–19131.
- Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng J-F *et al.* (2013). Insights into the phylogeny and coding potential of microbial dark matter. *Nature* **499**: 431–437.
- Schauer R, Røy H, Augustin N, Gennerich H-H, Peters M, Wenzhoefer F *et al.* (2011). Bacterial sulfur cycling shapes microbial communities in surface sediments of an ultramafic hydrothermal vent field. *Environ Microbiol* **13**: 2633–2648.
- Schink B. (1997). Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol Rev* **61**: 262–280.
- Sekiguchi Y, Ohashi A, Parks DH, Yamauchi T, Tyson GW, Hugenholtz P. (2015). First genomic insights into members of a candidate bacterial phylum responsible for wastewater bulking. *Peer J* **3**: e740.
- Sonnhammer EL, von Heijne G, Krogh A. (1998). A hidden Markov model for predicting transmembrane helices in protein sequences. *Proc Int Conf Intell Syst Mol Biol* **6**: 175–182.
- Stahl DA, Amann R. (1991). Development and application of nucleic acid probes in bacterial systematics. In Stackebrandt E, Goodfellow M (eds), *Nucleic Acid Techniques in Bacterial Systematics*. John Wiley & Sons Ltd. Chichester: England, pp 205–248.
- Stincone A, Prigione A, Cramer T, Wamelink MMC, Campbell K, Cheung E *et al.* (2014). The return of metabolism: biochemistry and physiology of the pentose phosphate pathway. *Biol Rev Camb Philos Soc* **90**: 927–963.
- Stoecker K, Dorninger C, Daims H, Wagner M. (2010). Double labeling of oligonucleotide probes for fluorescence *in situ* hybridization (DOPE-FISH) improves signal intensity and increases rRNA accessibility. *Appl Environ Microbiol* **76**: 922–926.
- Strittmatter AW, Liesegang H, Rabus R, Decker I, Amann J, Andres S *et al.* (2009). Genome sequence of *Desulfobacterium autotrophicum* HRM2, a marine sulfate reducer oxidizing organic carbon completely to carbon dioxide. *Environ Microbiol* **11**: 1038–1055.
- Sundberg C, Al-Soud W a, Larsson M, Alm E, Yekta SS, Svensson BH *et al.* (2013). 454 Pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas digesters. *FEMS Microbiol Ecol* **85**: 612–626.
- Syed M, Soreanu G, Falletta P, Beland M. (2006). Removal of hydrogen sulfide from gas streams using biological processes—a review. *Can Biosyst Eng* **48**: 2.1–2.14.
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin E V *et al.* (2003). The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**: 41.
- Uniprot Consortium. (2014). Activities at the Universal Protein Resource (UniProt). *Nucleic Acids Res* **42**: D191–D198.
- Vallenet D, Belda E, Calteau A, Cruveiller S, Engelen S, Lajus A *et al.* (2013). MicroScope—an integrated microbial resource for the curation and comparative analysis of genomic and metabolic data. *Nucleic Acids Res* **41**: D636–D647.
- Vallenet D, Engelen S, Mornico D, Cruveiller S, Fleury L, Lajus A *et al.* (2009). MicroScope: a platform for microbial genome annotation and comparative genomics. *Database (Oxford)* **2009**: bap021.
- Wallner G, Amann R, Beisker W. (1993). Optimizing fluorescent *in situ* hybridization with rRNA targeted oligonucleotide probes for flow cytometric identification of microorganisms. *Cytometry* **14**: 136–143.
- Wang Q, Garrity GM, Tiedje JM, Cole JR. (2007). Naïve Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73**: 5261–5267.
- Weiland P. (2010). Biogas production: current state and perspectives. *Appl Microbiol Biotechnol* **85**: 849–860.
- Werner JJ, Knights D, Garcia ML, Scalfone NB, Smith S, Yarasheski K *et al.* (2011). Bacterial community structures are unique and resilient in full-scale bioenergy systems. *Proc Natl Acad Sci USA* **108**: 4158–4163.
- Wickham H. (2007). Reshaping data with the {reshape} package. *J Stat Softw* **21**: 1–20.

- Wilson WA, Roach PJ, Montero M, Baroja-Fernández E, Muñoz FJ, Eydallin G *et al.* (2010). Regulation of glycogen metabolism in yeast and bacteria. *FEMS Microbiol Rev* **34**: 952–985.
- Wisedchaisri G, Park M-S, Iadanza MG, Zheng H, Gonen T. (2014). Proton-coupled sugar transport in the prototypical major facilitator superfamily protein XylE. *Nat Commun* **5**: 4521.
- Yarza P, Yilmaz P, Priesse E, Glöckner FO, Ludwig W, Schleifer K-H *et al.* (2014). Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat Rev Microbiol* **12**: 635–645.
- Yilmaz LS, Parnerkar S, Noguera DR. (2011). MathFISH, a web tool that uses thermodynamics-based mathematical models for *in silico* evaluation of oligonucleotide

probes for fluorescence in situ hybridization. *Appl Environ Microbiol* **77**: 1118–1122.



**This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>**

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)