

Pointwise estimates of pseudo-differential operators

Johnsen, Jon

Publication date:
2010

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Johnsen, J. (2010). *Pointwise estimates of pseudo-differential operators*. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2010-12

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

AALBORG UNIVERSITY

**Pointwise estimates of
pseudo-differential operators**

by

Jon Johnsen

R-2010-12

October 2010

DEPARTMENT OF MATHEMATICAL SCIENCES
AALBORG UNIVERSITY

Fredrik Bajers Vej 7 G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 99 40 80 80 ■ Telefax: +45 98 15 81 29

URL: <http://www.math.aau.dk>



POINTWISE ESTIMATES OF PSEUDO-DIFFERENTIAL OPERATORS

JON JOHNSEN

ABSTRACT. As a new technique it is shown how general pseudo-differential operators can be estimated at arbitrary points in Euclidean space when acting on functions u with compact spectra. The estimate is a factorisation inequality, in which one factor is the Peetre–Fefferman–Stein maximal function of u , whilst the other is a symbol factor carrying the whole information on the symbol. The symbol factor is estimated in terms of the spectral radius of u , so that the framework is well suited for Littlewood–Paley analysis. It is also shown how it gives easy access to results on polynomial bounds and estimates in L_p , including a new result for type 1, 1-operators that they are always bounded on L_p -functions with compact spectra.

1. INTRODUCTION

The aim of this note is to show how one can estimate a pseudo-differential operator at an arbitrary point $x \in \mathbb{R}^n$ — and to explore the convenient consequences.

The central theme is to show for a general symbol $a(x, \eta)$, with the associated operator $a(x, D)u(x) = (2\pi)^{-n} \int e^{ix \cdot \eta} a(x, \eta) \hat{u}(\eta) d\eta$, that for $u \in \mathcal{F}^{-1} \mathcal{E}'(\mathbb{R}^n)$,

$$|a(x, D)u(x)| \leq c \cdot u^*(x) \quad \text{for every } x \in \mathbb{R}^n. \quad (1.1)$$

Here u^* denotes the Peetre–Fefferman–Stein maximal function

$$u^*(x) = u^*(N, R; x) = \sup_y \frac{|u(x - y)|}{(1 + |Ry|)^N} = \sup_y \frac{|u(y)|}{(1 + R|x - y|)^N}, \quad (1.2)$$

where $N > 0$, $R > 0$ are parameters; R so large that $x \in \text{supp } \hat{u}$ implies $|x| \leq R$.

One obvious advantage of proving (1.1) in terms of (1.2) is the *immediate* L_p -estimate

$$\int |a(x, D)u(x)|^p dx \leq c^p \int |u^*(x)|^p dx \leq c^p C_p \|u\|_p^p, \quad 1 \leq p \leq \infty, \quad (1.3)$$

where the last step is to invoke the *maximal inequality*

$$\int_{\mathbb{R}^n} |u^*(x)|^p dx \leq C_p \int_{\mathbb{R}^n} |u(x)|^p dx, \quad u \in L_p \cap \mathcal{F}^{-1} \mathcal{E}'. \quad (1.4)$$

This estimate of the non-linear map $u \mapsto u^*$ has for $Np > n$ been known since 1975 from a work of Peetre [Pee75], who estimated $u^*(x)$ by the Hardy–Littlewood maximal function in order to invoke the L_p -boundedness of the latter. A significantly simpler proof is given below.

2000 *Mathematics Subject Classification.* 35S05, 47G30.

Key words and phrases. Pointwise estimates, factorisation inequality, maximal function, symbol factor.

Supported by the Danish Council for Independent Research, Natural Sciences (Grant No. 09-065927).

It is remarkable that little attention has been paid over the decades to pointwise estimates like (1.1) — in comparison Peetre’s proof of (1.4) quickly got a central role in the theory of function spaces; cf [Tri83, 1.4.1]. However, to the author’s knowledge, there has only been a similar attempt by Marschall, who in his thesis [Mar85] suggested to estimate $a(x, D)u(x)$ in terms of Mu ; this was followed up in a series of papers, eg [Mar91, Mar95, Mar96], where the technique was used to derive boundedness under weak assumptions in spaces based on L_p (functions and symbols subject to Besov and Lizorkin–Triebel conditions).

In the present paper the point of view is quite different. First of all because u^* is rather easier to treat and work with than Mu . Secondly, the aim is to explain how pointwise estimates in terms of u^* will simplify well-known topics such as L_p -estimates and Littlewood–Paley analysis of $a(x, D)$.

So as a main result here, (1.1) is also shown to be straightforward to obtain; cf Theorem 4.1–4.5 below. Indeed, the constant c in (1.1) is just an upper bound for the *symbol factor* $F_a(x)$, which is a continuous, bounded function carrying the entire information of the symbol in the *factorisation inequality*

$$|a(x, D)u(x)| \leq F_a(x)u^*(x), \quad u \in \mathcal{F}^{-1}\mathcal{E}'(\mathbb{R}^n). \quad (1.5)$$

As $F_a(x)$ only depends vaguely on u (cf Section 4), this gives a somewhat surprising decoupling.

The inequality is well suited for Littlewood–Paley analysis of $a(x, D)$ as described in Section 5. The set-up there has recently been exploited by the author [Joh10] in proofs of fundamental results for pseudo-differential operators of type 1, 1; this is briefly reviewed in Section 6, where also (1.3) is given as a new theorem for type 1, 1-operators.

2. THE PEETRE–FEFFERMAN–STEIN MAXIMAL FUNCTION

This section explores the definition of $u^*(x)$ in (1.2), in lack of a reference, and gives a straightforward proof of the maximal inequality (1.4).

For the reader’s sake a few easy facts are recalled first. To show that $u^*(x)$ is a ‘slowly’ varying function, note that

$$\frac{|u(x-z)|}{(1+|Rz|)^N} = \frac{|u(y-(z+y-x))|}{(1+R|z+y-x|)^N} \cdot \frac{(1+R|z+y-x|)^N}{(1+|Rz|)^N}, \quad (2.1)$$

so the inequality $1+|x+y| \leq 1+|x|+|y|+|x||y| = (1+|x|)(1+|y|)$ gives

$$u^*(x) \leq u^*(y)(1+R|x-y|)^N. \quad (2.2)$$

Therefore $u^*(x)$ is finite at every $x \in \mathbb{R}^n$ if it is so at one point y . So either $u^*(x) = \infty$ on the entire \mathbb{R}^n , or (2.2) implies that $u^*(x)$ is continuous on \mathbb{R}^n , ie $u^* \in C(\mathbb{R}^n)$.

Finiteness is for large N implied by the (often imposed) assumption that $u \in \mathcal{S}'(\mathbb{R}^n)$ should have its spectrum in the closed ball $\bar{B}(0, R)$ of radius R , ie

$$\text{supp } \mathcal{F}u \subset \bar{B}(0, R). \quad (2.3)$$

Indeed, then $|u(x)| \leq c_R(1+R|x|)^m$ by the Paley–Wiener–Schwartz theorem, when m is the order of \hat{u} . So $N \geq m$ gives $u^*(N, R; 0) \leq c_R$, hence $u^*(N, R; x) < \infty$ for all $x \in \mathbb{R}^n$.

In any case it is clear that $u \mapsto u^*$ is subadditive, ie $(u + v)^* \leq u^* + v^*$, whence

$$|u^*(N, R; x) - v^*(N, R; x)| \leq (u - v)^*(N, R; x). \quad (2.4)$$

Hence $u \mapsto u^*$ is Lipschitz continuous on $L_\infty(\mathbb{R}^n)$ with constant 1, as it is a shrinking map there, ie $\|u^*\|_\infty \leq \|u\|_\infty$. With respect to the Hölder seminorm

$$|u|_\sigma := \sup_{x \neq y} |u(x) - u(y)| / |x - y|^\sigma, \quad 0 < \sigma < 1, \quad (2.5)$$

it is also a shrinking map, for (2.4) gives that

$$|u^*(x + h) - u^*(x)| \leq \sup_{\mathbb{R}^n} \frac{|u(x + h + \cdot) - u(x + \cdot)|}{(1 + R|\cdot|)^N} \leq |u|_\sigma |h|^\sigma. \quad (2.6)$$

Therefore $|u^*|_\sigma \leq |u|_\sigma$ as claimed. In particular one has

Proposition 2.1. *The map $u \mapsto u^*$ is for all $N, R > 0$ a shrinking map on the Hölder space $C^\sigma(\mathbb{R}^n)$, $0 < \sigma < 1$, defined by finiteness of the norm $|u|_\sigma^* = \sup |u| + |u|_\sigma$.*

A main case is when u is in $L_p(\mathbb{R}^n)$, $1 \leq p \leq \infty$. For $p < \infty$ one has $u^* \equiv \infty$ for u equal to $e^{|x|}$ times the characteristic function of $\bigcup_{k \in \mathbb{N}} B(ke_1, e^{-(k+1)2p})$. Such growth is impossible on the subspace of functions fulfilling the spectral condition (2.3), so this is imposed henceforth.

As an a priori analysis of this case, the Nikolskiĭ–Plancherel–Polya inequality implies that $u \in L_p \cap L_\infty$, for it states that if $u \in L_p$ and (2.3) holds, then

$$\|u\|_r \leq c R^{\frac{n}{p} - \frac{n}{r}} \|u\|_p \quad \text{for } p < r \leq \infty. \quad (2.7)$$

For its proof one can take an auxiliary function $\psi \in \mathcal{S}(\mathbb{R}^n)$ so that $\mathcal{F}\psi(\xi) = 0$ for $|\xi| \geq 2$ and $\mathcal{F}\psi(\xi) = 1$ around $B(0, 1)$, for then $u = R^n \psi(R \cdot) * u$, and (2.7) follows from this identity at once by the Hausdorff–Young inequality $\|f * g\|_r \leq \|f\|_p \|g\|_q$, where $\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{r}$; hereby $c = \|\psi\|_q$, that only depends on p, r and n .

To complete the picture, (2.7) extends as it stands to the range $0 < p < r \leq \infty$, provided u is given in $L_p \cap \mathcal{S}'(\mathbb{R}^n)$ with $\text{supp } \mathcal{F}u \in \overline{B}(0, R)$; cf [Tri83, 1.4.1(ii)]. The direct treatment in [JS07] shows that one can take $c = \|\psi\|_\infty^{\frac{1}{p} - \frac{1}{r}}$ for $0 < p \leq 1$. (For $0 < p < 1$, the set $L_p \cap \mathcal{S}'$ itself consists of the $u \in L_1^{\text{loc}} \cap \mathcal{S}'$ fulfilling $\int_{\mathbb{R}^n} |u|^p dx < \infty$, that *per se* requires stricter smallness than L_1 for $|x| \rightarrow \infty$ but gives a global condition on the singularities in the possibly non-compact region where $|u(x)| > 1$.)

By (2.7), pointwise estimates of $u^*(x)$ hold for L_p -functions with compact spectra:

Lemma 2.2. *For every $u \in L_p \cap \mathcal{S}'(\mathbb{R}^n)$, $0 < p \leq \infty$ with $\text{supp } \mathcal{F}u \subset \overline{B}(0, R)$, it holds true on \mathbb{R}^n that*

$$|u(x)| \leq u^*(N, R; x) \leq \|u\|_\infty < \infty \quad \text{for every } N > 0. \quad (2.8)$$

Proof. With $r = \infty$ in (2.7) it follows that $\|u\|_\infty$ is finite; and it dominates $u^*(x)$ as stated, by the definition of u^* in (1.2). Taking $y = 0$ there yields $|u(x)| \leq u^*(x)$. \square

Note that $\mathcal{F}L_p \subset \mathcal{D}'^k$ for the least integer $k > \frac{n}{2} - \frac{n}{p}$ if $p > 2$, so the Paley–Wiener–Schwartz theorem would give the poor condition $N \geq [\frac{n}{2} - \frac{n}{p}] + 1$ for finiteness of u^* .

Example 2.3. As is well known, u^* is useful (when finite) for pointwise control of convolutions, since eg the assumptions $\varphi \in \mathcal{S}$, $u \in \mathcal{F}^{-1}\mathcal{E}'$ clearly give

$$|\varphi * u(x)| \leq \int (1 + R|y|)^N |\varphi(y)| \frac{|u(x-y)|}{(1 + R|y|)^N} dy \leq cu^*(N, R; x). \quad (2.9)$$

Example 2.4. Conversely $u^*(x)$ may be controlled by convolving $|u|$ with

$$f_N(z) = (1 + |z|)^{-N}; \quad (2.10)$$

and cases with $N > n$ are particularly simple as one has

$$u^*(N, R; x) \leq C_N R^n f_N(R \cdot) * |u|(x). \quad (2.11)$$

Indeed, when $u \in L_p$, $1 \leq p \leq \infty$ with $\text{supp } \mathcal{F}u \subset \overline{B}(0, R)$ the compact spectrum of u can be exploited by taking ψ as after (2.7) above, which gives $u = R^n \psi(R \cdot) * u$. Thence

$$\begin{aligned} |u(y)|(1 + R|x-y|)^{-N} &\leq \int \frac{R^n |\psi(R(y-z))u(z)|}{(1 + R|x-y|)^N} dz \\ &\leq \int (1 + R|y-z|)^N |\psi(R(y-z))| \frac{R^n |u(z)|}{(1 + R|x-z|)^N} dz \end{aligned} \quad (2.12)$$

by using $(1 + R|x-y|)(1 + R|y-z|) \geq (1 + R|x-z|)$ in the denominator. This gives

$$u^*(x) \leq C_N \int \frac{R^n |u(z)|}{(1 + R|x-z|)^N} dz, \quad (2.13)$$

where $C_N := \sup(1 + |v|)^N |\psi(v)| < \infty$ because $\psi \in \mathcal{S}$. This shows the claim in (2.11).

As an addendum to Example 2.4, a basic estimate gives in (2.11) that for $p \leq 1$

$$\|u^*\|_p \leq C_N \|R^n f_N(R \cdot) * |u|\|_p \leq C_N \int (1 + |z|)^{-N} dz \cdot \|u\|_p. \quad (2.14)$$

So for $N > n$ this short remark proves a special case of the maximal inequality (1.4).

However, $N > n$ is far from an optimal assumption for (1.4). But a few changes give the improvement $N > n/p$; and also every $p \in]0, \infty]$ can be treated using the Nikolskiĭ–Plancherel–Polya inequality (2.7).

The idea is to utilise the powerful pointwise estimate in (2.11), where eg both sides can be integrated over \mathbb{R}^n (unlike (2.8)). But first it is generalised thus:

Proposition 2.5. *If $u \in \mathcal{S}'$, $\text{supp } \mathcal{F}u \subset \overline{B}(0, R)$ and $N, p \in]0, \infty[$ are arbitrary, then*

$$u^*(N, R; x) \leq C_{n, N, p} \left(\int R^n \frac{|u(x-z)|^p}{(1 + R|z|)^{Np}} dz \right)^{\frac{1}{p}} = C_{n, N, p} (R^n f_N^p(R \cdot) * |u|^p(x))^{\frac{1}{p}} \quad (2.15)$$

for a constant $C_{n, N, p}$ depending only on n, N and p .

Proof. As above $u(x) = R^n \psi(R \cdot) * u = \langle u, R^n \psi(R(x - \cdot)) \rangle$, which can be written as an integral since $(1 + |y|)^{-k} u(y)$ is in $L_1(\mathbb{R}^n)$ for a large k ; therefore (2.12) holds. Suppose now that the right-hand side of (2.15) is finite.

For $1 \leq p < \infty$ one can simply use Hölder's inequality for $p + p' = p'p$ in the passage from (2.12) to (2.13); then $C_{n, N, p} = (\int (1 + |z|)^{Np'} |\psi(z)|^{p'} dz)^{1/p'}$ gives (2.15).

If $0 < p \leq 1$ the L_1 -norm with respect to z in (2.12) can be estimated by the L_p -norm, according to (2.7), for the Fourier transform of $z \mapsto \psi(R(y-z))u(z)$ is supported by $\bar{B}(0, 3R)$. Invoking the specific constant in (2.7) and proceeding as before, this gives

$$u^*(x)^p \leq \int \sup_{y \in \mathbb{R}^n} \frac{\|\psi\|_\infty^{(\frac{1}{p}-1)p} (3R)^{(\frac{n}{p}-n)p} C_N^p R^{np} |u(z)|^p}{(1+R|x-y|)^{Np} (1+R|y-z|)^{Np}} dz \leq C_{n,N,p}^p \int \frac{R^n |u(x-z)|^p}{(1+R|z|)^{Np}} dz, \quad (2.16)$$

where C_N is as in (2.13) and now $C_{n,N,p} = C_N 3^{n/p} \|\psi\|_\infty^{(1-p)/p}$. \square

These elementary considerations give a short proof, in the style of (2.14), of the following important theorem on the L_p -boundedness of the maximal operator $u \mapsto u^*$.

Theorem 2.6. *When $0 < p \leq \infty$ and $N > n/p$, then there is a constant $C'_{n,N,p} > 0$ such that the maximal function $u^*(N, R; x)$ in (1.2) fulfils*

$$\|u^*(N, R; \cdot)\|_p \leq C'_{n,N,p} \|u\|_p \quad (2.17)$$

for every $u \in L_p(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n)$ in the closed subspace with $\text{supp } \mathcal{F}u \subset \bar{B}(0, R)$. On this subspace there is Lipschitz continuity

$$\|u^*(N, R; \cdot) - v^*(N, R; \cdot)\|_p \leq C'_{n,N,p} \|u - v\|_p. \quad (2.18)$$

Proof. Lemma 2.2 yields that u^* is finite and consequently continuous as noted after (2.2), hence measurable. The case $p = \infty$ then follows at once from the lemma. For $0 < p < \infty$ one can integrate both sides of (2.15), which by Fubini's theorem yields

$$\int |u^*(x)|^p dx \leq C_{n,N,p}^p \iint \frac{|u(x-z/R)|^p}{(1+|z|)^{Np}} dz dx = C^p \int |u(x)|^p dx \quad (2.19)$$

for $C^p = C_{n,N,p}^p \int (1+|z|)^{-Np} dz$. Since $Np > n$ this gives $u^* \in L_p$ and (2.17). Now the Lipschitz property follows by integration on both sides of (2.4). \square

Among the further properties there is a Bernstein inequality for u^* , which states that the maximal function of u controls that of the derivatives $\partial^\alpha u$.

Proposition 2.7. *If $\text{supp } \mathcal{F}u \subset \bar{B}(0, R)$ one has $(\partial^\alpha u)^*(N, R; x) \leq C_N^{(\alpha)} R^{|\alpha|} u^*(N, R; x)$.*

While this is known (cf [Tri83, 1.3.1] for $R = 1$), it is natural to give the short proof here. Writing $u(x-y)(1+R|y|)^{-N}$ in terms of the convolution $R^n \psi(R \cdot) * u$, cf Example 2.4, it is straightforward to see by applying ∂_x^α that for $N' > 0$,

$$(\partial^\alpha u)^*(N, R; x) \leq \sup(1+|\cdot|)^{N'} |\partial^\alpha \psi| \sup_y \int \frac{R^{n+|\alpha|} |u(z)|}{(1+R|y|)^N (1+R|x-y-z|)^{N'}} dz. \quad (2.20)$$

For $N' = N + n + 1$ a simple estimate of the denominator, cf Example 2.4, now shows Proposition 2.7 with the constant $C_N^{(\alpha)} = \sup(1+|\cdot|)^{N+n+1} |\partial^\alpha \psi| \int (1+|z|)^{-n-1} dz$.

Remark 2.8. The maximal function u^* was introduced by Peetre [Pee75], inspired by the non-tangential maximal function used by Fefferman and Stein a few years earlier [FS72]. It has been widely used in the theory of Besov and Lizorkin–Triebel spaces, cf [Tri83, Tri92, RS96], where the boundedness in Theorem 2.6 has been a main tool since the 1970’s; cf [Tri83, 1.4.1]. Usually its proof has been based on an estimate in terms of the Hardy–Littlewood maximal function, $M_ru(x) = \sup_{\rho} (\rho^{-n} \int_{|y| < \rho} |u(x+y)|^r dy)^{1/r}$, ie for $\text{supp } \mathcal{F}u \subset \overline{B}(0, R)$,

$$u^*(N, R; x) \leq cM_ru(x), \quad N \geq n/r. \quad (2.21)$$

When $N > n/r$ this results from Proposition 2.5 by splitting the integral ($p = r$) in regions with $2^k \leq |z| \leq 2^{k+1}$. (For $N = n/r$ it was shown by Triebel, cf [Tri83, 1.3.1 ff], by combining an inequality for u^* , $(\partial_j u)^*$ and Mu , due to Peetre [Pee75], with the Bernstein inequality for u^* ; cf Proposition 2.7.) This allowed the inequality $\|M_ru\|_p \leq c\|u\|_p$ to be invoked for $p > r$. The present proofs of Proposition 2.5 and Theorem 2.6 are rather simpler.

3. PREPARATIONS

Notation and notions from distribution theory are the same as in Hörmander’s book [Hör85], unless otherwise mentioned. Eg $[t]$ denotes the largest integer $k \leq t$ for $t \in \mathbb{R}$. The Fourier transformation $\mathcal{F}u(\xi) = \int e^{-ix \cdot \xi} u(x) dx$ is occasionally written as $\mathcal{F}_{x \rightarrow \xi} u(x, y)$ when u depends on further variables y .

As mentioned in the introduction the paper deals with operators given by

$$a(x, D)u = \text{OP}(a)u(x) = (2\pi)^{-n} \int e^{ix \cdot \eta} a(x, \eta) \mathcal{F}u(\eta) d\eta, \quad u \in \mathcal{S}(\mathbb{R}^n). \quad (3.1)$$

Hereby the symbol $a(x, \eta)$ is C^∞ on $\mathbb{R}^n \times \mathbb{R}^n$ and is taken to fulfil the Hörmander condition of order $d \in \mathbb{R}$, ie for all multiindices $\alpha, \beta \in \mathbb{N}_0^n$ there is a constant $C_{\alpha, \beta} > 0$ such that

$$|D_\eta^\alpha D_x^\beta a(x, \eta)| \leq C_{\alpha, \beta} (1 + |\eta|)^{d - \rho|\alpha| + \delta|\beta|}. \quad (3.2)$$

The space of such symbols is denoted by $S_{\rho, \delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$ or $S_{\rho, \delta}^d$; and $S^{-\infty} := \bigcap S_{\rho, \delta}^d$.

The parameters $\rho, \delta \in [0, 1]$ are assumed to fulfil $\delta < \rho$ so that $a(x, D)$ by duality has a continuous extension $a(x, D): \mathcal{S}'(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$. (Type 1, 1-operators, ie $\delta = 1 = \rho$, are considered in Section 6 below.) If desired the reader may specialise to the classical case $\rho = 1, \delta = 0$.

Together with $a(x, D)$ one has the distribution kernel $K(x, y) = \mathcal{F}_{\eta \rightarrow z}^{-1} a(x, \eta)|_{z=x-y}$, that in the usual way is seen to be C^∞ for $x \neq y$ also for $a \in S_{1, 1}^d$. It fulfils

$$\langle a(x, D)u, \varphi \rangle = \langle K, \varphi \otimes u \rangle \quad \text{for all } u, \varphi \in \mathcal{S}. \quad (3.3)$$

As preparations, two special cases are considered: if $u = v + v'$ is any splitting of u with $v \in \mathcal{S}$ and $v' \in \mathcal{F}^{-1} \mathcal{E}'$ then

$$a(x, D)u = a(x, D)v + \text{OP}(a(1 \otimes \chi))v', \quad (3.4)$$

whereby $a(1 \otimes \chi)(x, \eta) = a(x, \eta)\chi(\eta)$ and $\chi \in C_0^\infty(\mathbb{R}^n)$ is chosen so that $\chi = 1$ holds in a neighbourhood of $\text{supp } \mathcal{F}v'$, or just on a neighbourhood of the smaller set

$$\bigcup_{x \in \mathbb{R}^n} \text{supp } a(x, \cdot) \mathcal{F}v'(\cdot). \quad (3.5)$$

Indeed, by linearity on the left-hand side of (3.4) the identity results, for the term $a(x, D)v'$ equals $\text{OP}(a(1 \otimes \chi))v'$ if $v' \in \mathcal{F}^{-1}C_0^\infty(\mathbb{R}^n)$ that extends to $v' \in \mathcal{F}^{-1}\mathcal{E}'$ by mollification of $\mathcal{F}v'$ since $a(1 \otimes \chi) \in S^{-\infty}$.

Moreover, for every auxiliary function $\psi \in C_0^\infty(\mathbb{R}^n)$ equal to 1 in a neighbourhood of the origin, continuity of the adjoint operation $a \mapsto e^{iD_x \cdot D_\eta} \bar{a}$ yields

$$a(x, D)u = \lim_{m \rightarrow \infty} \text{OP}(\psi(2^{-m}D_x)a(x, \eta)\psi(2^{-m}\eta))u. \quad (3.6)$$

4. POINTWISE ESTIMATES

This section develops a flexible framework for discussion of pseudo-differential operators. These are only for convenience restricted to the classes recalled in Section 3.

4.1. The factorisation inequality. In the derivation of (1.5), hence (1.1), the simple result below introduces $u^*(x)$ as a fundamental tool and is therefore given as a theorem.

Formally the idea is to proceed as in Example 2.3, cf (2.9), now departing from

$$a(x, D)u(x) = \int K(x, x-y)u(x-y)dy. \quad (4.1)$$

This leads to the *factorisation inequality* (4.2) below, where the dependence on $a(x, \eta)$ is taken out in the symbol factor F_a , also called the “ a -factor”. This is essentially a weighted L_1 -norm of the distribution kernel. (The estimate shows that the case of an operator is not much worse than that of $\varphi * u$ in Example 2.3.)

Theorem 4.1. *Let $a \in S_{\rho, \delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$ for $0 \leq \delta < \rho \leq 1$. When $u \in \mathcal{S}'(\mathbb{R}^n)$, $\text{supp } \hat{u} \subset \bar{B}(0, R)$, then one has the following pointwise estimate for all $x \in \mathbb{R}^n$:*

$$|a(x, D)u(x)| \leq F_a(N, R; x) \cdot u^*(N, R; x). \quad (4.2)$$

Hereby u^* is as in (1.2) while F_a is bounded and continuous for $x \in \mathbb{R}^n$ and is given in terms of an auxiliary function $\chi \in C_0^\infty(\mathbb{R}^n)$ equal to 1 on a neighbourhood of $\text{supp } \hat{u}$ as

$$F_a(N, R; x) = \int_{\mathbb{R}^n} (1 + R|y|)^N |\mathcal{F}^{-1}(a(x, \cdot)\chi(\cdot))| dy. \quad (4.3)$$

The inequality (4.2) holds for $N > 0$, and remains true if $\chi = 1$ on $\bigcup_{x \in \mathbb{R}^n} \text{supp } a(x, \cdot) \hat{u}(\cdot)$.

Proof. Using formula (3.4) with $v' = u$, and (3.5) for the last statement,

$$a(x, D)u(x) = \text{OP}(a(1 \otimes \chi))u = \langle u, \mathcal{F}_{\eta \rightarrow y}(\frac{e^{ix \cdot \eta}}{(2\pi)^n} a(x, \eta)\chi(\eta)) \rangle \quad (4.4)$$

for the last rewriting is evident from (3.1) if $u \in \mathcal{F}^{-1}C_0^\infty$ and follows for general $u \in \mathcal{F}^{-1}\mathcal{E}'$ by mollification of $\mathcal{F}u$, since $a(1 \otimes \chi)$ is in $S^{-\infty}$.

Now $a(x, \eta)\chi(\eta)$ is in $C_0^\infty(\mathbb{R}^n)$ for fixed x , so $y \mapsto \mathcal{F}_{\eta \rightarrow y}^{-1}(a(1 \otimes \chi))(x, x-y)$ decays rapidly while $u(y)$ grows polynomially by the Paley–Wiener–Schwartz theorem. Therefore the above scalar product on $\mathcal{S}' \times \mathcal{S}$ is an integral, so by the change of variables $y \mapsto x-y$,

$$\begin{aligned} |a(x, D)u(x)| &= \left| \int u(x-y) \mathcal{F}_{\eta \rightarrow y}^{-1}(a(1 \otimes \chi))(x, y) dy \right| \\ &\leq \sup_{z \in \mathbb{R}^n} \frac{|u(x-z)|}{(1+R|z|)^N} \int (1+R|y|)^N |\mathcal{F}_{\eta \rightarrow y}^{-1}(a(1 \otimes \chi))(x, y)| dy \\ &= u^*(x) F_a(x), \end{aligned} \quad (4.5)$$

according to the definition of $u^*(x)$ in (1.2) and that of $F_a(x)$ in (4.3).

That $x \mapsto F_a(x)$ is bounded follows by insertion of $1 = (1+|y|^2)^{N'}(1+|y|^2)^{-N'}$ for $N' > (N+n)/2$ since $\mathcal{F}_{\eta \rightarrow y}^{-1}((1-\Delta_\eta)^{N'}[a(x, \eta)\chi(\eta)])$ is bounded with respect to (x, y) because of the compact support of χ . These estimates also yield continuity of the symbol factor $F_a(x)$. \square

Disregarding the spectral radius R and N , (4.2) may be written concisely as

$$|a(x, D)u(x)| \leq F_a(x) \cdot u^*(x). \quad (4.6)$$

It is noteworthy that the entire influence of the symbol lies in the a -factor $F_a(x)$, while u itself is mainly felt in $u^*(x)$. It is only in a vague way, ie through N and R , that u contributes to $F_a(x)$, so the factorisation inequality is rather convenient.

The theorem is also valid more generally; eg Section 6 gives an extension to symbols of type 1, 1 (extensions to other general symbols can undoubtedly be worked out when needed). To give a version for functions without compact spectrum, $\mathcal{O}_M(\mathbb{R}^n)$ will as usual stand for the space of slowly increasing functions, ie the $f \in C^\infty(\mathbb{R}^n)$ satisfying the estimates

$$|D^\alpha f(x)| \leq C_\alpha (1+|x|)^{N_\alpha}. \quad (4.7)$$

Analogously to the argument after (2.3), $f^*(N, R; \cdot)$ is finite for $N \geq N_{(0, \dots, 0)}$, any $R > 0$. There is a factorisation inequality for such functions, at the expense of a sum over its derivatives:

Theorem 4.2. *When $a(x, \eta)$ is in $S_{\rho, \delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$, $1 \leq \delta < \rho \leq 1$, and $u \in \mathcal{O}_M(\mathbb{R}^n)$ while $N' > (d+n)/2$ is a non-negative integer, then one has for $N, R > 0$ that*

$$|a(x, D)u(x)| \leq c F_a(N, R; x) \sum_{|\alpha| \leq 2N'} (D^\alpha u)^*(N, R; x), \quad (4.8)$$

where F_a is defined by (4.3) for $\chi(\eta) = (1+|\eta|^2)^{-N'}$ and again is in $C(\mathbb{R}^n) \cap L_\infty(\mathbb{R}^n)$.

Proof. That F_a is in $C(\mathbb{R}^n) \cap L_\infty(\mathbb{R}^n)$ can be seen as above, for $a(x, \eta)\chi(\eta) \in S_{\rho, \delta}^{d-2N'}$ is integrable with respect to η . When $a \in S^{-\infty}$ and $u \in \mathcal{S}$,

$$a(x, D)u(x) = \sum_{j=0}^{N'} \binom{N'}{j} \int (1-\Delta)^j u(y) \mathcal{F}_{\eta \rightarrow y} \left(\frac{e^{ix \cdot \eta}}{(2\pi)^n} a(x, \eta) \chi(\eta) \right) dy. \quad (4.9)$$

By continuity this extends to all $u \in \mathcal{S}'$, in particular to $u \in \mathcal{O}_M$; and then to all $a \in S_{\rho,\delta}^d$ since $(1 + |y|)^{-N''} (1 - \Delta)^j u(y)$ is in L_1 for a large N'' . As each term is a function of x ,

$$|a(x, D)u(x)| \leq F_a(N, R; x) \sum \binom{N'}{j} ((1 - \Delta)u)^*(N, R; x). \quad (4.10)$$

Since $(1 - \Delta)^j u = \sum_{|\alpha| \leq 2j} c_{\alpha,j} D^\alpha u$, subadditivity of the maximal operator gives the rest. \square

As a first consequence of the factorisation inequalities, when $u \in \mathcal{O}_M$ then $a(x, D)u$ is of polynomial growth by (4.8), and continuous by (4.9), for $F_a(x)$ is bounded and $D^\alpha u \in \mathcal{O}_M$ so $(D^\alpha u)^*(x)$ has such growth for N sufficiently large; cf (2.2). This applies to the commutator $[D^\beta, a(x, D)]$, in eg $\text{OP}(S_{\rho,\delta}^{d+|\beta|})$, so also $D^\beta a(x, D)u$ has polynomial growth; which proves

Corollary 4.3. $a(x, D)$ is a map $\mathcal{O}_M(\mathbb{R}^n) \rightarrow \mathcal{O}_M(\mathbb{R}^n)$ when $a \in S_{\rho,\delta}^d$, $0 \leq \delta < \rho \leq 1$.

While this is known for $\rho = 1$, $\delta = 0$ from eg [SR91, Cor. 3.8], the above version for $\delta < \rho$ is rather more direct.

Secondly, one may now obtain the L_p -estimate mentioned in the introduction.

Corollary 4.4. For each $a \in S_{\rho,\delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$, $0 \leq \delta < \rho \leq 1$, and $p \in]0, \infty]$ there is to every $R \geq 1$, $N > n/p$ a constant $C(N, R)$ such that

$$\|a(x, D)u\|_p \leq C(N, R) \|u\|_p \quad (4.11)$$

whenever $u \in L_p(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n)$, fulfils $\text{supp } \hat{u} \subset \bar{B}(0, R)$.

Proof. By taking L_p -norms on both sides of the factorisation inequality, (4.11) results at once with $C(N, R) = C_{n,N,p} \sup_x |F_a(N; R; x)|$; this is finite according to Theorem 4.1. \square

Since the spectral condition on u implies $u \in C^\infty$, it is hardly surprising that the L_p -result above is valid for arbitrary orders $d \in \mathbb{R}$. In fact it may, say for $1 < p < \infty$, $(\rho, \delta) = (1, 0)$, be proved simply by observing that $a(x, D)$ has the same action on u as some $b(x, D) \in \text{OP}(S^{-\infty})$ so that boundedness of $b(x, D)$ on L_p gives the rest.

It is noteworthy, however, that the existing proofs of L_p -boundedness use fundamental parts of real analysis, eg Marcinkiewicz interpolation and the Calderon–Zygmund lemma. In contrast to this, pointwise estimates lead straightforwardly to Corollary 4.4. This evident efficacy is also clear from the easy extension to the full range $0 < p \leq \infty$ and to type 1, 1-operators in Section 6.

4.2. Estimates of the symbol factor. To utilise Theorem 4.1 it is of course vital to control F_a . This leads directly to integrals conditions on a , similarly to the Mihlin–Hörmander theorem.

Theorem 4.5. Assume $a(x, \eta)$ is in $S_{\rho,\delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$, $0 \leq \delta < \rho \leq 1$, and let $F_a(N, R; x)$ be given by (4.3) for parameters $R, N > 0$, with the auxiliary function taken as $\chi = \psi(R^{-1} \cdot)$ for $\psi \in C_0^\infty(\mathbb{R}^n)$ equalling 1 in (the closure of) an open set. Then

$$0 \leq F_a(x) \leq c_{n,k} \sum_{|\alpha| \leq k} \left(\int_{R \text{supp } \psi} |R^{|\alpha|} D_\eta^\alpha a(x, \eta)|^2 \frac{d\eta}{R^n} \right)^{1/2} \quad (4.12)$$

for all $x \in \mathbb{R}^n$, when k is the least integer satisfying $k > N + n/2$.

First it is convenient to recall that, for $z \in \mathbb{R}^n$ and $k \in \mathbb{N}$, an expansion yields

$$(1 + |z|)^k \leq \sum_{j=0}^k \binom{k}{j} (|z_1| + \cdots + |z_n|)^j = \sum_{|\alpha| \leq k} C_{k,\alpha} |z^\alpha|. \quad (4.13)$$

Proof. The idea is to pass to the L_2 -norm in (4.3) using Cauchy–Schwarz’ inequality and that $(\int \mathbb{R}^n (1 + |Ry|)^{-n-\varepsilon} dy)^{1/2} < \infty$ for $\varepsilon > 0$. Thus, if ε is so small that $k \geq N + (n + \varepsilon)/2$,

$$F_a(N, R; x) \leq c_n R^{-n/2} \left(\int (1 + |Ry|)^{2k} |\mathcal{F}_{\eta \rightarrow y}^{-1}[a(x, \cdot) \psi(R^{-1} \cdot)]|^2 dy \right)^{1/2}. \quad (4.14)$$

Applying (4.13) to $z = Ry$ and intertwining the resulting polynomials $(Ry)^\alpha$ with the inverse Fourier transformation, it is seen that for fixed $x \in \mathbb{R}^n$,

$$\begin{aligned} F_a(x) &\leq c_n R^{-n/2} \sum_{|\alpha| \leq k} C_{k,\alpha} \left(\int |\mathcal{F}_{\eta \rightarrow y}^{-1}[(iR\partial_\eta)^\alpha a(1 \otimes \psi(R^{-1} \cdot))](x, y)|^2 dy \right)^{1/2} \\ &\leq c_n R^{-n/2} \sum_{|\alpha+\beta| \leq k} c_{k,\alpha,\beta} \left(\int_{R \text{ supp } \psi} (R^{|\alpha+\beta|} |D_\eta^\alpha a(x, \eta)| |D^\beta(\psi(R^{-1}\eta))|)^2 d\eta \right)^{1/2}. \end{aligned} \quad (4.15)$$

Since $D^\beta(\psi(R^{-1} \cdot)) = R^{-|\beta|} (D^\beta \psi)(R^{-1} \cdot)$ is bounded, the result follows. \square

Remark 4.6. As an alternative to the estimate $|a(x, D)u(x)| \leq F_a(x)u^*(x)$, it deserves to be mentioned that other useful properties can be obtained in a similar fashion: by defining an a -factor in terms of an L_2 -norm, ie

$$\tilde{F}_a(N, R; x)^2 = \int_{\mathbb{R}^n} (1 + |Ry|)^{2N} |\mathcal{F}_{\eta \rightarrow y}^{-1}(a(x, \cdot) \chi(\cdot))|^2 dy, \quad (4.16)$$

the Cauchy–Schwarz inequality gives

$$|a(x, D)u(x)| \leq \tilde{F}_a(N, R; x) \left(\int_{\mathbb{R}^n} \frac{|u(x-y)|^2}{(1 + |Ry|)^{2N}} dy \right)^{1/2} \leq c \tilde{F}_a(N, R; x) u^*(\varepsilon, R; x) \quad (4.17)$$

where $c = (\int (1 + |Ry|)^{-2(N-\varepsilon)} dy)^{1/2} < \infty$ whenever $N > n/2 + \varepsilon$ for some $\varepsilon > 0$.

For one thing $\tilde{F}_a^2 \in C^\infty(\mathbb{R}^n)$, with bounded derivatives of any order. Secondly, this gives a version of Theorem 4.5 where only estimates with $|\alpha| \leq [n/2] + 1$ is required, as in the Mihlin–Hörmander theorem. But it would not be feasible in general to replace $u^*(N, R; x)$ by $u^*(\varepsilon, R; x)$ for small ε as above, so $\tilde{F}_a(x)$ is only mentioned in this remark.

Although it is a well-known exercise to control (4.12) in terms of symbol seminorms, it is important to control the behaviour with respect to R and to verify that it improves when $a(x, \cdot) \hat{u}(\cdot)$ is supported in a corona. Therefore the special case in (4.19) below is included:

Corollary 4.7. Assume $a \in S_{1,\delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$, $0 \leq \delta < 1$, and let N , R and ψ have the same meaning as in Theorem 4.5. When $R \geq 1$ and $k > N + n/2$, $k \in \mathbb{N}$, then there is a seminorm p on $S_{1,1}^d$ and some $c_k > 0$ independent of R such that

$$0 \leq F_a(x) \leq c_k p(a) R^{\max(d,k)} \quad \text{for all } x \in \mathbb{R}^n. \quad (4.18)$$

Moreover, if $\text{supp } \psi$ is contained in a corona

$$\{\eta \mid \theta_0 \leq |\eta| \leq \Theta_0\}, \quad (4.19)$$

and $\psi(\eta) = 1$ holds for $\theta_1 \leq |\eta| \leq \Theta_1$, whereby $0 \neq \theta_0 < \theta_1 < \Theta_1 < \Theta_0$, then

$$0 \leq F_a(x) \leq c'_k R^d p(a) \quad \text{for all } x \in \mathbb{R}^n, \quad (4.20)$$

with $c'_k = c_k \max(1, \theta_0^{d-k}, \theta_0^d)$.

Remark 4.8. For general $\rho \in]0, 1]$, the corresponding asymptotics of $F_a(x)$ for $R \rightarrow \infty$ will be $\mathcal{O}(R^{\max(d+(1-\rho)k, k)})$ and $\mathcal{O}(R^{d+(1-\rho)k})$, respectively. Details are left out for simplicity's sake.

Proof. Setting $p_{\alpha, \beta}(a) = \sup(1 + |\eta|)^{-d+|\alpha|-\delta|\beta|} |D_x^\beta D_\eta^\alpha a(x, \eta)|$ and continuing from the proof of Theorem 4.5, the change of variables $\eta = R\zeta$ gives

$$F_a(x) \leq c \sum_{|\alpha| \leq k} p_{\alpha, 0}(a) \left(\int_{\text{supp } \psi} |(1 + |R\zeta|)^{d-|\alpha|} R^{|\alpha|} d\zeta \right)^{\frac{1}{2}} \leq C'_{n,k} R^{\max(d,k)} \sum_{|\alpha| \leq k} p_{\alpha, 0}(a). \quad (4.21)$$

In fact $d \geq k \geq |\alpha|$ gives $R^{|\alpha|} (1 + R|\zeta|)^{d-|\alpha|} \leq R^d (1 + |\zeta|)^{d-|\alpha|}$ since $R \geq 1$; whilst if $d < k$ the crude estimate $R^{|\alpha|} (1 + R|\zeta|)^{d-|\alpha|} \leq R^k$ applies eg for $|\alpha| = k$. This shows (4.18).

In case ψ is supported in a corona as described, $d - |\alpha| < 0$ and $\zeta \in \text{supp } \psi$ entail

$$(1 + |R\zeta|)^{d-|\alpha|} R^{|\alpha|} \leq (R\theta_0)^{d-|\alpha|} R^{|\alpha|} \leq \max(\theta_0^{d-k}, \theta_0^d) R^d. \quad (4.22)$$

This yields an improvement of (4.21) for terms with $|\alpha| > d$; thence (4.20). \square

As desired Corollary 4.7 shows that the a -factor $F_a(x)$ has its norm in L_∞ bounded by a symbol seminorm. This applies of course in $|a(x, D)u(x)| \leq F_a(x)u^*(x)$. Hereby one could take R equal to the spectral radius of u , or if possible R so large that the corona $\{\eta \mid \theta_1 R \leq |\eta| \leq \Theta_1 R\}$ is a neighbourhood of $\text{supp } a(x, \cdot) \hat{u}(\cdot)$ for all x ; cf (4.18) and (4.20).

A good choice of N is more delicate: eg $N \geq \text{order}(\mathcal{F}u)$ was seen to imply that $u^*(N, R; x)$ is finite everywhere; this was relaxed completely to $N > 0$ for $u \in L_p \cap \mathcal{F}^{-1}\mathcal{E}'$ in Lemma 2.2. For general $u \in L_p$, $1 \leq p \leq 2$, the order of $\mathcal{F}u$ is 0, so u^* is finite regardless of $N > 0$.

Especially for functions u in Sobolev spaces H^s the function u^* is always finite for $N > 0$. Therefore it is harmless that the estimates in Corollary 4.7 depend on N , for only seminorms $p_{\alpha, 0}(a)$ with $|\alpha| \leq 1 + [n/2 + N]$ enters there, and by taking $0 < N < 1/2$ in both odd and even dimensions estimates of $a(x, D)u(x)$ with $u \in \bigcup H^s$ only requires the well-known estimates of $D_\eta^\alpha a(x, \eta)$ for $|\alpha| \leq [n/2] + 1$.

However, going back to L_p -bounds of $u^*(x)$, one is often forced to take $N > n/p$ in the L_p -estimates of $a(x, D)u$; cf Theorem 2.6.

In addition to high frequencies removed by the spectral cut-off function χ in Theorem 4.1, the symbols dependence on x may be frequency modulated by means of a Fourier multiplier $\varphi(Q^{-1}D_x)$, which depends on a second spectral quantity $Q > 0$. For the modified symbol

$$a_Q(x, \eta) = \varphi(Q^{-1}D_x)a(x, \eta) \quad (4.23)$$

and the corresponding symbol factor one can as shown below find its asymptotics for $Q \rightarrow \infty$. In Littlewood–Paley theory, this is a frequently asked question for F_{a_Q} :

Corollary 4.9. *When $a \in S_{1,\delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$, $0 \leq 1 < \delta$ and $\varphi \in C_0^\infty(\mathbb{R}^n)$ with $\varphi = 0$ in a neighbourhood of $\xi = 0$, then there is a seminorm p on $S_{1,\delta}^d$ and constants c_M , depending only on M , n , N , ψ and φ , such that for $R \geq 1$, $M > 0$, $Q > 0$,*

$$0 \leq F_{a_Q}(N, R; x) \leq c_M p(a) Q^{-M} R^{\max(d+\delta M, [N+n/2]+1)}. \quad (4.24)$$

Here $d + \delta M$ can replace the maximum when the auxiliary function ψ in F_{a_Q} fulfils the corona condition in Corollary 4.7.

Proof. Because $a_Q(x, \eta) = \int Q^n \check{\varphi}(Qz) a(x - z, \eta) dz$, where $\check{\varphi}$ has vanishing moments of every order, Taylor's formula with remainder gives for any $M \in \mathbb{N}$

$$a_Q(x, \eta) = \sum_{|\beta|=M} \frac{M}{\beta!} \int (-z)^\beta Q^n \check{\varphi}(Qz) \int_0^1 (1-\tau)^{M-1} \partial_x^\beta a(x - \tau z, \eta) d\tau dz. \quad (4.25)$$

Letting z^β absorb Q^M before substitution of z by z/Q , one finds

$$\begin{aligned} Q^M F_{a_Q}(N, R; x) &\leq \sum_{|\beta|=M} \frac{M}{\beta!} \iiint (1-\tau)^{M-1} (1+|z|)^M |\check{\varphi}(z)| (1+|Ry|)^N \\ &\quad \times |\mathcal{F}_{\eta \rightarrow y}^{-1}(\partial_x^\beta a(x - \tau z/Q, \eta) \psi(\eta/R))| d\tau dz dy. \end{aligned} \quad (4.26)$$

Integrating first with respect to y it follows by applying Corollary 4.7 to $\partial_x^\beta a \in S_{1,\delta}^{d+\delta M}$ that, with $p(a) = \sum p_{\alpha,\beta}(a)$, where $|\alpha| \leq [N+n/2] + 1$ and $|\beta| = M$,

$$F_{a_Q}(N, R; x) \leq c_M p(a) R^{d+\delta M} Q^{-M}, \quad (4.27)$$

when ψ satisfies the corona condition. Otherwise the stated inequality results. \square

Remark 4.10. For general $\rho \in]0, 1]$ the asymptotics of $F_a(x)$ is here $\mathcal{O}(R^{\max(d+\delta M+(1-\rho)k, k)})$ and $\mathcal{O}(R^{d+\delta M+(1-\rho)k})$, respectively, for $k = [N+n/2] + 1$.

Remark 4.11. As an alternative to the techniques in this section, Marschall's inequality gives a pointwise estimate for symbols $b(x, \eta)$ in $L_{1,\text{loc}}(\mathbb{R}^{2n}) \cap \mathcal{S}'(\mathbb{R}^{2n})$ with support in $\mathbb{R}^n \times \bar{B}(0, 2^k)$ and $\text{supp } \mathcal{F}u \subset \bar{B}(0, 2^k)$, $k \in \mathbb{N}$:

$$|b(x, D)v(x)| \leq c \|b(x, 2^k \cdot)\|_{\dot{B}_{1,t}^{n/t}} M_t u(x), \quad 0 < t \leq 1. \quad (4.28)$$

This goes back to [Mar85, p.37] and was exploited in eg [Mar91, Mar95, Mar96]. In the above form it was proved in [Joh05] under the natural condition that the right-hand side is in $L_{1,\text{loc}}(\mathbb{R}^n)$.

While $M_t u$ is as in Remark 2.8, the norm of the homogeneous Besov space $\dot{B}_{1,t}^{n/t}$ on the symbol is defined analogously to that of $B_{p,q}^s$ in (5.14) below in terms of a partition of unity, though here with $1 = \sum_{j=-\infty}^\infty (\varphi(2^{-j}\eta) - \varphi(2^{1-j}\eta))$, $\eta \neq 0$ so that (5.14) should be read with ℓ_q over \mathbb{Z} . This yields the well-known dyadic scaling property that

$$\|b(x, 2^k \cdot)\|_{\dot{B}_{1,t}^{n/t}} = 2^{k(\frac{n}{t}-n)} \|b(x, \cdot)\|_{\dot{B}_{1,t}^{n/t}}. \quad (4.29)$$

While this can be useful, and indeed fits well into the framework of the next section, cf [Mar91, Mar95, Mar96], it is often simpler to invoke the factorisation inequality with F_b and u^* etc.

5. LITTLEWOOD–PALEY ANALYSIS

In order to obtain L_p -estimates, it is convenient to depart from the limit in (3.6). As usual the test function ψ there gives rise to a Littlewood–Paley decomposition $1 = \psi(\eta) + \sum_{j=1}^{\infty} \varphi(2^{-j}\eta)$ by setting $\varphi = \psi - \psi(2\cdot)$. Note here that if $\psi \equiv 1$ for $|\eta| \leq r$ while $\psi \equiv 0$ for $|\eta| \geq R$, one can fix an integer $h \geq 2$ so that $2R < r2^h$.

Inserting twice into (3.6) that $\psi(2^{-m}\eta) = \psi(\eta) + \varphi(2^{-1}\eta) + \dots + \varphi(2^{-m}\eta)$, the so-called paradifferential splitting from the 1980's is recovered: when $a(x, \eta)$ is in $S_{\rho, \delta}^d$, $0 \leq \delta < \rho \leq 1$, and $u \in \mathcal{S}'(\mathbb{R}^n)$,

$$a_\psi(x, D)u = a_\psi^{(1)}(x, D)u + a_\psi^{(2)}(x, D)u + a_\psi^{(3)}(x, D)u, \quad (5.1)$$

whereby the expressions are given by the three series below (they converge in \mathcal{S}'),

$$a_\psi^{(1)}(x, D)u = \sum_{k=h}^{\infty} \sum_{j \leq k-h} a_j(x, D)u_k = \sum_{k=h}^{\infty} a^{k-h}(x, D)u_k \quad (5.2)$$

$$a_\psi^{(2)}(x, D)u = \sum_{k=0}^{\infty} (a_{k-h+1}(x, D)u_k + \dots + a_{k-1}(x, D)u_k + a_k(x, D)u_k + a_k(x, D)u_{k-1} + \dots + a_k(x, D)u_{k-h+1}) \quad (5.3)$$

$$a_\psi^{(3)}(x, D)u = \sum_{j=h}^{\infty} \sum_{k \leq j-h} a_j(x, D)u_k = \sum_{j=h}^{\infty} a_j(x, D)u^{j-h}. \quad (5.4)$$

Here $u_k = \varphi(2^{-k}D)u$ while $a_k(x, \eta) = \varphi(2^{-k}D_x)a(x, \eta)$; by convention φ is replaced by ψ for $k = 0$ and $u_k \equiv 0 \equiv a_k$ for $k < 0$. In addition superscripts are used for the convenient shorthands u^{k-h} and $a^{k-h}(x, D)$; eg the latter is given by $\sum_{j \leq k-h} a_j(x, D) = \text{OP}(\psi(2^{h-k}D_x)a(x, \eta))$. Using this, there is a brief version of (5.3),

$$a_\psi^{(2)}(x, D)u = \sum_{k=0}^{\infty} ((a^k - a^{k-h})(x, D)u_k + a_k(x, D)(u^{k-1} - u^{k-h})). \quad (5.5)$$

The main property behind (5.1) is the following inclusions for the spectra of the summands in (5.2), (5.4) and (5.5), with $R_h = \frac{r}{2} - R2^{-h}$:

$$\text{supp } \mathcal{F}(a^{k-h}(x, D)u_k) \subset \{ \xi \mid R_h 2^k \leq |\xi| \leq \frac{5R}{4} 2^k \}, \quad (5.6)$$

$$\text{supp } \mathcal{F}(a_k(x, D)u^{k-h}) \subset \{ \xi \mid R_h 2^k \leq |\xi| \leq \frac{5R}{4} 2^k \}, \quad (5.7)$$

$$\text{supp } \mathcal{F}(a_k(x, D)(u^{k-1} - u^{k-h}) + (a^k - a^{k-h})(x, D)u_k) \subset \{ \xi \mid |\xi| \leq 2R2^k \}. \quad (5.8)$$

Details are omitted here since these spectral corona and ball properties have been known since the 1980's. Note that although they were verified then for symbols $a(x, \eta)$ that were elementary in the sense of Coifman and Meyer [CM78], this is now a redundant restriction because of the general spectral support rule for $u \in \mathcal{F}^{-1}\mathcal{E}'(\mathbb{R}^n)$,

$$\text{supp } \mathcal{F}(a(x, D)u) \subset \{ \xi + \eta \mid (\xi, \eta) \in \text{supp } \mathcal{F}_{x \rightarrow \xi} a, \eta \in \text{supp } \mathcal{F}u \}, \quad (5.9)$$

(A short proof of this can be found in [Joh10, App. B]. Cf [Joh05, Joh08, Joh10] for the full version.) (5.6)–(5.8) follow easily from (5.9), as shown in [Joh05, Joh10].

The novelty in relation to pointwise estimates is that the summands in the decomposition (5.2)–(5.4) can be controlled thus: for $a_\psi^{(1)}(x, D)u$ the fact that $k \geq h \geq 2$ allows the corona condition of Corollary 4.7 to be fulfilled for $\Theta_0 = r/2$ and $\Theta_1 = R$ (ie the auxiliary function there is 1 on $\text{supp } \hat{u}$), so (5.2) and the factorisation inequality simply give the first estimate:

$$|a^{k-h}(x, D)u_k(x)| \leq F_{a^{k-h}}(N, R2^k; x)u_k^*(N, R2^k; x) \leq cp(a)(R2^k)^d u_k^*(x). \quad (5.10)$$

Hereby $p(a^{k-h}) \leq \|\mathcal{F}^{-1}\psi\|_1 p(a)$ is a convolution estimate that only changes the constant.

In $a_\psi^{(2)}(x, D)u$ the terms may be treated similarly: in (5.5) it is clear that $(a^k - a^{k-h})(x, D)u_k$ for $k \geq 1$ only requires the constant to have $\|\mathcal{F}^{-1}(\psi - \psi(2^h \cdot))\|_1$ as a factor instead of $\|\mathcal{F}^{-1}\psi\|_1$, while for $k = 0$ it may just be increased by a fixed power of R using the full generality of Corollary 4.7. The remainders in (5.5) have $k > 0$ and can be written as in (5.3). Hence the second estimate,

$$\begin{aligned} |(a^k - a^{k-h})(x, D)u_k(x) + a_k(x, D)[u^{k-1} - u^{k-h}](x)| \\ \leq F_{a^k - a^{k-h}}(N, R2^k; x)u_k^*(N, R2^k; x) + \sum_{l=1}^{h-1} F_{a_k}(N, R2^{k-l}; x)u_{k-l}^*(N, R2^{k-l}; x) \\ \leq cp(a)(R2^k)^d \sum_{l=0}^{h-1} 2^{-ld} u_{k-l}^*(N, R2^{k-l}; x). \end{aligned} \quad (5.11)$$

Here the sum over l is harmless, because the number of terms is independent of k .

The improved asymptotics of Corollary 4.9 come into play as reinforcements for the series for $a_\psi^{(3)}(x, D)u$. Indeed, for $Q = 2^j$ the first part of (5.4) gives, for $M > 0$, the third estimate

$$\begin{aligned} |a_j(x, D)u^{j-h}(x)| &\leq \sum_{k=0}^{j-h} |a_j(x, D)u_k(x)| \leq \sum_{k=0}^j F_{a_j}(N, R2^k; x)u_k^*(N, R2^k; x) \\ &\leq c'_M p(a) 2^{-jM} \sum_{k=0}^j (R2^k)^{d+\delta M} u_k^*(N, R2^k; x). \end{aligned} \quad (5.12)$$

Though the number of terms on the right-hand side depends on j , this sum turns out to be manageable due to 2^{-jM} , which (as is known) serves as a summation factor.

Anyhow, the easy estimates (5.10), (5.11), (5.12) imply boundedness in several scales. This is perhaps most transparent for the Besov spaces $B_{p,q}^s(\mathbb{R}^n)$. These generalise both the Sobolev spaces $H^s(\mathbb{R}^n)$ and the Hölder spaces $C^s(\mathbb{R}^n)$ (with $0 < s < 1$, cf Proposition 2.1),

$$H^s = B_{2,2}^s, \quad C^s = B_{\infty,\infty}^s. \quad (5.13)$$

The spaces $B_{p,q}^s$ are for $s \in \mathbb{R}$, $p, q \in]0, \infty]$ defined by means of the Littlewood–Paley decomposition as the $u \in \mathcal{S}'$ for which the following (quasi-)norm is finite,

$$\|u\|_{B_{p,q}^s} = \left(\sum_{j=0}^{\infty} 2^{sjq} \|\varphi(2^{-j}D)u\|_p^q \right)^{1/q}; \quad (5.14)$$

hereby the norm in ℓ_q should be read as the supremum over j for $q = \infty$. (Often a specific choice of the function ψ is stipulated, but this is immaterial as they all lead to equivalent norms on the spaces). For $p, q \in [1, \infty]$ the space $B_{p,q}^s$ is a Banach space. Note that the first part of (5.13) is an easy consequence of (5.14); cf [Hör97] for the second.

Theorem 5.1. *When $a(x, \eta)$ belongs to $S_{1,\delta}^d(\mathbb{R}^n \times \mathbb{R}^n)$ and $0 \leq \delta < 1$, then*

$$a(x, D): H^{s+d}(\mathbb{R}^n) \rightarrow H^s(\mathbb{R}^n) \quad (5.15)$$

$$a(x, D): B_{p,q}^{s+d}(\mathbb{R}^n) \rightarrow B_{p,q}^s(\mathbb{R}^n) \quad (5.16)$$

is continuous for every $s \in \mathbb{R}$, $0 < p \leq \infty$, $0 < q \leq \infty$.

Proof. Taking norms of L_p and ℓ_q on both sides of (5.10), Theorem 2.6 gives for $N > n/p$,

$$\left(\sum_{k=0}^{\infty} 2^{skq} \|a^{k-h}(x, D)u_k\|_p^q \right)^{1/q} \leq cp(a) \left(\sum_{k=0}^{\infty} 2^{(s+d)kq} \|u_k^*\|_p^q \right)^{1/q} \quad (5.17)$$

$$\leq c'p(a) \left(\sum_{k=0}^{\infty} 2^{(s+d)kq} \|u_k\|_p^q \right)^{1/q} = c'p(a) \|u\|_{B_{p,q}^{s+d}}. \quad (5.18)$$

Because of the dyadic corona property (5.6), the above estimate implies that $a_{\psi}^{(1)}(x, D)u = \sum a^{k-h}(x, D)u_k$ converges to an element in $B_{p,q}^s$, the norm of which is estimated by the right-hand side (this is well known, cf [Yam86], [RS96, 2.3.2] or [Joh10]). So for $m = 1$,

$$\|a_{\psi}^{(m)}(x, D)u\|_{B_{p,q}^s} \leq c''p(a) \|u\|_{B_{p,q}^{s+d}}. \quad (5.19)$$

The contribution $a^{(3)}(x, D)$ in (5.1) is treated similarly, except for the sum over k . This is handled with a small lemma, namely $\sum_{j=0}^{\infty} 2^{sjq} (\sum_{k=0}^j |b_k|)^q \leq c \sum_{j=0}^{\infty} 2^{sjq} |b_j|^q$, valid for all $b_j \in \mathbb{C}$ and $0 < q \leq \infty$ provided $s < 0$; cf [Yam86]. So with $M > s$ in (5.12),

$$\begin{aligned} \sum_{j=0}^{\infty} 2^{sjq} \|a_j(x, D)u^{j-h}\|_p^q &\leq \sum_{j=0}^{\infty} 2^{(s-M)jq} \left(\sum_{k=0}^j c'_M p(a) (R2^k)^{d+\delta M} \|u_k^*(N, R2^k; \cdot)\|_p \right)^q \\ &\leq cp(a)^q \sum_{j=0}^{\infty} 2^{(s+d-(1-\delta)M)jq} \|u_j\|_p^q = cp(a)^q \|u\|_{B_{p,q}^{s+d-(1-\delta)M}}. \end{aligned} \quad (5.20)$$

Increasing to $M > 0$ if necessary, this implies (5.19) for $m = 3$.

For $a^{(2)}(x, D)u$ the estimate is a little simpler, for in (5.11) one only needs to apply the norms of L_p and ℓ_q with respect to x and k , respectively, and use the (quasi-)triangle inequality. As (5.8) is only a dyadic ball property, the resulting estimate gives (5.19) with $m = 2$ for $s > \max(0, \frac{n}{p} - n)$. Then this shows (5.16), hence (5.15), via (5.1).

However, one can reduce to such s by writing $a(x, D) = \Lambda^t (\Lambda^{-t} a(x, D))$ with $t = 2|s| + 1$ (or $t = 2|s| + 1 + \frac{n}{p} - n$ if $0 < p < 1$), for $\Lambda^t = \text{OP}((1 + |\eta|^2)^{t/2})$ is of order t in the $B_{p,q}^s$ -scale. \square

The theorem is well established, of course. Eg [Hör85, Thm. 18.1.13] or [SR91, Thm. 3.6] gives the H^s -part with a classical reduction to Schur's lemma. The above proof should be interesting because it combines the factorisation inequality etc with Littlewood–Paley theory.

The flexibility of the present proof is apparent from the fact that it extends *at once* to the $B_{p,q}^s$ with arbitrary parameters $p, q \in]0, \infty]$. This result first appeared in [Yam86], where the proof is cumbersome due to the use of elementary symbols and multiplier results.

6. THE CASE OF TYPE 1, 1-OPERATORS

The above results carry over to type 1, 1-operators with just a few necessary changes. The methods were in fact developed for such operators, which emphasizes the efficacy of pointwise estimates. Some brief remarks on the outcome are given here.

The reader may consult [Joh08, Joh10] for a review of *operators* of type 1, 1 and a systematic treatment. Here it suffices to recall that for $a \in S_{1,1}^d(\mathbb{R}^n \times \mathbb{R}^n)$ the identity (3.6) is used as the *definition*: when the limit exists in $\mathcal{D}'(\mathbb{R}^n)$ and is independent of ψ , then $u \in D(a(x, D))$ and the action of $a(x, D)$ on u is taken to equal the limit.

Secondly, for u in $\mathcal{S} + \mathcal{F}^{-1}\mathcal{E}'$ the limit in (3.6) exists and equals the right-hand side of (3.4). Since this does not depend on ψ , one has $\mathcal{S} + \mathcal{F}^{-1}\mathcal{E}' \subset D(a(x, D))$. Cf [Joh08, Cor. 4.7].

Therefore the proof of Theorem 4.1, that departs from (3.4), applies *verbatim* to type 1, 1-operators. Hence the factorisation inequality (4.6) is valid for them. Corollary 4.3 also holds in this case, but the proof needs to be changed to obtain the decisive addition that $\mathcal{O}_M \subset D(a(x, D))$, which was verified in [Joh10]. However, the proof of Corollary 4.4 gives without changes

Theorem 6.1. *For $a \in S_{1,1}^d$, $p \in]0, \infty]$, $R \geq 1$ and $N > n/p$ one has*

$$\|a(x, D)u\|_p \leq C(N, R)\|u\|_p \quad (6.1)$$

whenever $u \in L_p(\mathbb{R}^n) \cap \mathcal{S}'(\mathbb{R}^n)$, fulfils $\text{supp } \hat{u} \subset \bar{B}(0, R)$.

This result is a novelty in the type 1, 1-context. It is noteworthy because some operators in $\text{OP}(S_{1,1}^0)$ are unbounded on L_p , even for $p = 2$, by a construction of Ching [Chi72] — and therefore pointwise estimates seem indispensable for this theorem.

The basic estimates of $F_a(x)$ in Theorem 4.5 also carry over to $\delta = 1 = \rho$ with no change at all. The introduction of symbol seminorms in Corollaries 4.7 and 4.9 can also be used *verbatim*, so here too the corona condition gives the asymptotics

$$\mathcal{O}(R^d), \quad \mathcal{O}(Q^{-M}R^{d+M}). \quad (6.2)$$

Moreover, the paradifferential decomposition in Section 5 is unchanged, although for type 1, 1-operators it has to be made for arbitrary ψ because of their definition.

As a difference it holds in this context that the series for $a^{(2)}(x, D)u$ converges if and only if $u \in D(a(x, D))$. Those for $a^{(1)}(x, D)u$ and $a^{(3)}(x, D)u$ converge for all $u \in \mathcal{S}'(\mathbb{R}^n)$, which was proved in [Joh10] by duality as well as explicitly via the pointwise estimates above.

In addition the proof of boundedness in Theorem 5.1 carries over to type 1, 1-operators, but in general only for $s > \max(0, \frac{n}{p} - n)$. For by Ching's counter-example [Chi72], left-composition with the lift operator Λ^t does not leave the full operator class invariant.

However, under Hörmander's fundamental condition that $\mathcal{F}_{x \rightarrow \xi} a(x, \eta)$ be suitably small along the twisted diagonal $\xi = -\eta$, cf [Hör88, Hör89, Hör97], several results on boundedness for the

Besov scale $B_{p,q}^s$ and the Lizorkin–Triebel scale $F_{p,q}^s$, that for $q = 2$ and $1 < p < \infty$ restricts to H_p^s , were derived using pointwise estimates.

It would be outside the topic here to give the full statements, so the reader is referred to [Joh10] for more details on these results for operators of type 1, 1.

7. FINAL REMARKS

As mentioned the pointwise techniques were developed during the author’s work on type 1, 1-operators [Joh10], but used there with only brief explanations. A detailed presentation has been postponed to the present paper, because the techniques should be of interest in their own right. This is illustrated by the proofs of Corollary 4.3, Theorem 5.1 and Theorem 6.1, for example.

REFERENCES

- [Chi72] Chin Hung Ching, *Pseudo-differential operators with nonregular symbols*, J. Differential Equations **11** (1972), 436–447.
- [CM78] R. R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque, vol. 57, Société Mathématique de France, Paris, 1978.
- [FS72] C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), 137–193.
- [Hör85] L. Hörmander, *The analysis of linear partial differential operators*, Grundlehren der mathematischen Wissenschaften, Springer Verlag, Berlin, 1983, 1985.
- [Hör88] ———, *Pseudo-differential operators of type 1, 1*, Comm. Partial Differential Equations **13** (1988), no. 9, 1085–1111.
- [Hör89] ———, *Continuity of pseudo-differential operators of type 1, 1*, Comm. Partial Differential Equations **14** (1989), no. 2, 231–243.
- [Hör97] ———, *Lectures on nonlinear hyperbolic differential equations*, Mathématiques & Applications, vol. 26, Springer Verlag, Berlin, 1997.
- [Joh05] J. Johnsen, *Domains of pseudo-differential operators: a case for the Triebel–Lizorkin spaces*, J. Function Spaces Appl. **3** (2005), 263–286.
- [Joh08] ———, *Type 1, 1-operators defined by vanishing frequency modulation*, New Developments in Pseudo-Differential Operators (L. Rodino and M. W. Wong, eds.), Operator Theory: Advances and Applications, vol. 189, Birkhäuser, 2008, pp. 201–246.
- [Joh10] ———, *Type 1, 1-operators on spaces of temperate distributions*, Tech. Report R-2010-13, Aalborg University, 2010, (preprint).
- [JS07] J. Johnsen and W. Sickel, *A direct proof of Sobolev embeddings for quasi-homogeneous Lizorkin–Triebel spaces with mixed norms*, J. Function Spaces Appl. **5** (2007), 183–198.
- [Mar85] J. Marschall, *Pseudo-differential operators with non-regular symbols*, Ph.D. thesis, Free University of Berlin, 1985.
- [Mar91] ———, *Weighted parabolic Triebel spaces of product type. Fourier multipliers and pseudo-differential operators*, Forum Math. **3** (1991), no. 5, 479–511.
- [Mar95] ———, *On the boundedness and compactness of nonregular pseudo-differential operators*, Math. Nachr. **175** (1995), 231–262.
- [Mar96] ———, *Nonregular pseudo-differential operators*, Z. Anal. Anwendungen **15** (1996), no. 1, 109–148.
- [Pee75] J. Peetre, *On spaces of Triebel–Lizorkin type*, Ark. Mat. (1975), 123–130.
- [RS96] T. Runst and W. Sickel, *Sobolev spaces of fractional order, Nemytskiĭ operators and non-linear partial differential equations*, Nonlinear analysis and applications, vol. 3, de Gruyter, Berlin, 1996.
- [SR91] X. Saint Raymond, *Elementary introduction to the theory of pseudodifferential operators*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1991.
- [Tri83] H. Triebel, *Theory of function spaces*, Monographs in mathematics, vol. 78, Birkhäuser Verlag, Basel, 1983.
- [Tri92] ———, *Theory of function spaces II*, Monographs in mathematics, vol. 84, Birkhäuser Verlag, Basel, 1992.

- [Yam86] M. Yamazaki, *A quasi-homogeneous version of paradifferential operators, I. Boundedness on spaces of Besov type*, J. Fac. Sci. Univ. Tokyo Sect. IA, Math. **33** (1986), 131–174.

DEPT. OF MATHEMATICAL SCIENCES, FREDRIK BAJERS VEJ 7G, DK-9220 AALBORG ØST, DENMARK
E-mail address: `j.johnsen@math.aau.dk`