

Quantitative proteomic analysis of Ibuprofen-degrading *Patulibacter* sp. strain I11

Kjeldal, Henrik; Lolas, Ihab Bishara Yousef; Knudsen, Anders Dahl; Carvalho, Gilda; Nielsen, Kåre Lehmann; Barreto Crespo, Maria Teresa; Stensballe, Allan; Nielsen, Jeppe Lund

Publication date:
2012

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Kjeldal, H., Lolas, I. B. Y., Knudsen, A. D., Carvalho, G., Nielsen, K. L., Barreto Crespo, M. T., Stensballe, A., & Nielsen, J. L. (2012). *Quantitative proteomic analysis of Ibuprofen-degrading Patulibacter sp. strain I11*. Poster presented at 14th International Symposium on Microbial Ecology, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Quantitative proteomic analysis of Ibuprofen-degrading *Patulibacter* sp. strain I11

Kjeldal H^{1*}, Almeida B^{1,2,3**}, Lolas I¹, Knudsen A D¹, Carvalho G^{2,4}, Nielsen K L¹, Barreto Crespo M T^{2,3}, Stensballe A¹, and Nielsen J L¹

¹Department of Biotechnology, Chemistry and Environmental Engineering, Aalborg University, Aalborg, Denmark

²Instituto de Biologia Experimental e Tecnológica (IBET), Av. da República (EAN), Oeiras, Portugal

³Instituto de Tecnologia Química e Biológica (ITQB)-Universidade Nova de Lisboa (UNL), Av. da República, Estação Agronómica Nacional, Portugal

⁴REQUIMTE/CQFB, Chemistry Department, FCT, Universidade Nova de Lisboa, Caparica, Portugal

*kjeldal@bio.aau.dk

**Equal contributor



Introduction

The increase in diversity and quantity of Pharmaceutically Active Compounds (PhACs) detected in the effluents of wastewater treatment plants is an issue of great concern due to health and environmental associated risks of the PhACs ¹.

Ibuprofen, a non-steroidal anti-inflammatory drug, is considered one of the most frequently occurring PhACs in the influent wastewater, typically being found in the range of 10-400 µg/L. Typical Ibuprofen removal efficiencies range from 80-100%, depending on operational conditions and wastewater treatment plant configuration^{2,3}. The elimination of ibuprofen is being ascribed primarily to biodegradation. However, in order to investigate the conditions for better removal of compounds like ibuprofen, we need to know the identity of the organisms involved and how their ibuprofen degradation activity depend on the controlling parameters. For this purpose we wanted to identify the genes involved and develop quantitative molecular tools for determining the activity of these genes.

Objective

The main objective of this study was to investigate the biochemical pathway of ibuprofen degradation in the ibuprofen degrading strain *Patulibacter* sp. Strain I11 using quantitative tandem mass spectrometry.

Methods




Fig. 1 The differential changes in the proteome of *Patulibacter* sp. strain I11, grown in the presence and absence of ibuprofen, were characterised by the combination of stable isotope metabolic labelling and 1-D gelbased shotgun Proteomics. The genome of *Patulibacter* sp. Strain I11 was sequenced and annotated and used as the reference database for the subsequent MS-based protein identification (the sequencing and annotation part of the genome of *Patulibacter* sp. strain I11 have been omitted in the above flowchart). The setup was carried out in biological duplicates using a forward and reverse labelling strategy (only the forward labelled duplicate is depicted above). For the reverse labelled duplicate the metabolic labels were reversed, i.e. yielding (¹⁴N + Ibuprofen) and (¹⁵N + Ibuprofen). The forward and reverse labelling strategy served the purpose of evaluating potential bias of the ¹⁴N- and ¹⁵N- medium on protein expression levels.

Results




Fig. 2 A) Metabolic labelling was carried out in biological duplicate and proteins considered up-regulated (proteins with log₂ ratio ≥ 0.9) were B) pooled and Gene Ontology-annotated at three different levels: Biological Process, Cellular Component and Molecular Function. Each pie slice is labelled with the GO subcategory name, number of GO annotations within the category as well as the percentage fraction of annotations. Replicate 1 corresponds to the forward labelled replicate (14N + Ibuprofen, 15N + Ibuprofen) whereas Replicate 2 corresponds to the reverse labelled replicate (14N + Ibuprofen, 15N + Ibuprofen).

Accession ^A	Description ^B	Log ₂ Ratio ^C		Protein Score ^D		# ^E
		Replicate 1	Replicate 2	Replicate 1	Replicate 2	
D3F114	Enoyl-CoA hydratase/isomerase	0.9	-	145	-	2
D3F2P4	Nuclear export factor GLE1	1.0	-	335 (388)	-	17 (25)
D3F178	Putative uncharacterized protein	1.1	-	482	-	24
E6SEB9	Aminotransferase class-III	1.2	-	96	-	8
E1VK11	AMP-dependent synthetase and ligase	1.3	-	314	-	6
D3FA23	Putative uncharacterized protein	1.4	-	174 (382)	-	6 (10)
D3F6B1	Putative uncharacterized protein	1.5	-	311 (292)	-	7 (14)
Q47N65	Putative uncharacterized protein	1.6	-	235 (219)	-	4 (4)
D3F293	Putative uncharacterized protein	1.7	-	747 (723)	-	40 (46)
D3F3V3	Rieske (2Fe-2S) iron-sulphur domain protein	1.8	-	313 (366)	-	13 (13)
D3FFG0	Putative uncharacterized protein	1.9	-	2195 (1785)	-	45 (97)
B0CFJ3	Putative uncharacterized protein	2.0	-	113 (110)	-	4 (6)
F1YMM4	Glycosidase	2.1	-	749	-	16
D3FBK3	Collagen triple helix repeat protein	2.2	-	412 (316)	-	26 (21)
D3E272	Putative uncharacterized protein	2.3	-	(881)	-	(46)
D3FEE9	Putative uncharacterized protein	2.4	-	(252)	-	(8)
D3FEF4	PpiC-type peptidyl-prolyl cis-trans isomerase	2.5	-	958	-	64
D3FBI3	Daunorubicin resistance ABC transporter ATPase subunit	2.6	-	509 (473)	-	8 (10)
D3FBH9	Extracellular solute-binding protein family 1	2.7	-	369	-	8
D3FBI2	ABC-2 type transporter	2.8	-	161	-	4

Table 1 Differentially expressed proteins of the biological replicates of *Patulibacter* sp. I11 grown in presence/absence of ibuprofen. Only up-regulated proteins (Log₂ ratio ≥ 0.9) are shown in the table. No major influence of the ¹⁴N- and ¹⁵N- medium on the protein expression levels was observed.

^AUniProt accession number of the closest protein homologue, ^BDescription of the closest protein homologue, ^CLog₂ ratio obtained from the quantitative proteomics analysis, ^DProtein Score obtained from the quantitative proteomics analysis, ^EThe number of quantitated peptides upon which the quantitative value (Log₂ ratio) was determined.

Conclusion

- Several proteins related to uptake and degradation of aromatic acids as well as compound transport-related proteins were found among the proteins up-regulated in response to Ibuprofen.
- The high number of up-regulated putative uncharacterised proteins might suggest a novel pathway for the degradation of Ibuprofen in *Patulibacter* sp. Strain I11.

Acknowledgements

The authors acknowledge the support of FTP (Danish Research Council for Technology and Production) and Aalborg University. AB, CG and BCMT were supported by Fundação para a Ciência e Tecnologia through the grant Pest-OE/EQB/LA0004/2011, project grant PTDC/EBB-EBI/098862/2008, and fellowship grants SFRH/BD/47748/2008 and SFRH/BPD/30800/2006.

References

- Pal, A., K. Y. H. Gin, A. Y. C. Lin, and M. Reinhard. 2010. Impacts of emerging organic contaminants on freshwater resources: Review of recent occurrences, sources, fate and effects. *Science of the Total Environment* 408:6062-6069.
- Santos, L. H. M. L. M., A. N. Araujo, A. Fachini, A. Pena, C. Delerue-Matos, and M. C. B. S. M. Montenegro. 2010. Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. *Journal of Hazardous Materials* 175:45-95.
- Remes, A., M. Ihos, and F. Manea. 2010. Electrochemical Characterization of Some Electrode Materials for Pharmaceutically Active Compounds Degradation. *Chem. Bull. "POLITEHNICA" Univ. (Timisoara)* 69:152-155.