



**Aalborg Universitet**

**AALBORG UNIVERSITY**  
DENMARK

“Candidatus Competibacter”-lineage genomes retrieved from metagenomes reveal functional metabolic diversity

McIlroy, Simon Jon; Albertsen, Mads; Andresen, Eva Kammer; Saunders, Aaron Marc; Kristiansen, Rikke; Stokholm-Bjerregaard, Mikkel; Nielsen, Kåre Lehmann; Nielsen, Per Halkjær

*Publication date:*  
2013

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

McIlroy, S. J., Albertsen, M., Andresen, E. K., Saunders, A. M., Kristiansen, R., Stokholm-Bjerregaard, M., Nielsen, K. L., & Nielsen, P. H. (2013). “Candidatus Competibacter”-lineage genomes retrieved from metagenomes reveal functional metabolic diversity. Poster presented at Microbial Ecology and Water Engineering 2013, Ann Arbor, Michigan, United States. <http://www.mewe2013.org/>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

#### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# "*Candidatus* Competibacter"-lineage genomes retrieved from metagenomes reveal functional metabolic diversity

Simon Jon McIlroy<sup>1</sup>, Mads Albertsen<sup>1</sup>, Eva Kammer Andresen<sup>1</sup>, Aaron Marc Saunders<sup>1</sup>, Rikke Kristiansen<sup>1</sup>, Mikkel Stokholm-Bjerregaard<sup>1,2</sup>, Kåre Lehmann Nielsen<sup>1</sup> & Per Halkjær Nielsen<sup>1</sup>

<sup>1</sup> Center for Microbial Communities, Aalborg University, Aalborg, Denmark

<sup>2</sup> Kruger A/S, Veolia Water Solutions and Technologies, Aalborg, Denmark

AAU

## Introduction

The "*Candidatus* Competibacter"-lineage (henceforth Competibacter) exhibits a glycogen accumulating organism (GAO) phenotype, relying on aerobically stored glycogen to energize anaerobic carbon uptake and storage as polyhydroxyalkanoates (PHA).

This bi-phasic mode of life is important for Competibacter to survive in EBPR systems. However, as they do not contribute to phosphorus removal, but compete for organic resources with the PAOs, they theoretically reduce EBPR efficiency.





Fig. 1. FISH image of Competibacter (yellow)

## Aim

To enhance our understanding of the Competibacter-lineage through the construction of genome-based-metabolic models.

## Methods

### Metagenome



Metagenome datasets generated from 2 lab-scale enrichments, seeded from a full-scale EBPR waste water treatment plant (WWTP).

### Assembly




Fig 2. PCA plot for genome extraction.

*De novo* assembled genome fragments from 2 individual Competibacter strains extracted using tetranucleotide frequency, coverage, and GC content.

## Annotation

Both genomes annotated using the 'MicroScope' pipeline (GenoScope) and curated for pathways relevant to metabolic model construction (Fig. 5).



## Conclusions

This study is the first of its kind for organisms with the GAO phenotype and provides a solid foundation for future -omic based studies. The revealed potential for phenotypic diversity among these related *Competibacteraceae* species indicates potential niche partitioning and underlines the complex nature of interaction between the PAO and GAO populations in EBPR systems.

## Results

Table 1. Metabolic comparison of pathways in sequenced GAO and PAO

	GAO <sup>φ</sup>	PAO*
	C. denitrificans C. odensis Accumulibacter	T. elongata T. australiensis T. jenkinsii T. japonica
<b>Central metabolism</b>		
TCA cycle	+	+
Glyoxylate shunt	+	+
Pentose phosphate (oxidative)	-	+
Pentose phosphate (non-oxidative)	+	+
Glycolysis EMP pathway	+	+
Glycolysis ED pathway	+	+
Fermentation (glucose → lactate)	+	+
<b>Storage compound metabolism</b>		
Glycogen synthesis	+	+
Trehalose synthesis	-	+
PHA synthesis	+	+
TAG synthesis	-	+
Polyphosphate synthesis	+	+
- Pit transporter	-	+
<b>Nitrogen metabolism</b>		
Nitrate reduction	+	+
Nitrate reduction to nitrogen (denitrification)	+	-
Nitrite reduction (respiratory)	+	-
Nitrite reduction to ammonia (assimilatory)	-	+
Nitrogen fixation	-	+

<sup>φ</sup> This study; \* Garcia-Martin et al., (2006) *Nature Biotech.*; Kristiansen et al., (2013) *ISME J.*




Fig. 4 Diagrammatic representation of P metabolism in PAO. Comparison of polyphosphate metabolism genes in available PAO and GAO genomes reveals the absence of the *pit* gene in the latter. The Pit phosphate transporter is theoretically involved in anaerobic proton motive force generation energising carbon uptake in PAO.




Fig 5. Metabolic model for Competibacter-lineage genomes in EBPR WWTPs a. in the absence of electron acceptors (anaerobic) and b. in the presence of the electron acceptors oxygen (aerobic) or nitrate. E.T.C = electron transport chain; PEP = phosphoenolpyruvate; KDPG = 2-dehydro-3-deoxy-D-gluconate-6-phosphate; TAG = triacylglycerols; PHB = polyhydroxybutyrate; PHV = polyhydroxyvalerate; PH2MB = polyhydroxy-2-methylbutyrate; PH2MV = polyhydroxy-2-methylvalerate. Both genomes contain genes for glycogen and PHA cycling and VFA metabolism. Differences between the genomes relate to storage polymers, nitrogen metabolism and glucose fermentation.




Fig 3. Phylogenetic tree for Competibacter-lineage sequences. Genome derived sequences are presented in colour. Phylogenetic analyses reveal genomes are members of Competibacter sub-groups 1 and 5. Diversity among the 'genus' Competibacter is high and thus the group better represents a family - *Competibacteraceae*.