

## **Monitoring of pharmaceutical processes with image analysis**

*direct and indirect approaches*

Kucheryavskiy, Sergey; Wu, Jian-Xiong

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Kucheryavskiy, S., & Wu, J.-X. (2013). *Monitoring of pharmaceutical processes with image analysis: direct and indirect approaches*. Abstract from Scandinavian Symposium on Chemometrics, Stockholm, Sweden.

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Monitoring of pharmaceutical processes with image analysis: direct and indirect approaches

Sergey Kucheryavskiy<sup>1</sup>, Jian-Xiong Wu<sup>2,3</sup>

<sup>1</sup>Institute of Biotechnology, Chemistry and Environmental Engineering, Aalborg University, Denmark.

<sup>2</sup>FOSS Analytical A/S, Denmark.

<sup>3</sup>Department of Pharmacy, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.

[svk@bio.aau.dk](mailto:svk@bio.aau.dk)

Image analysis (IA) is an effective and non-invasive tool for extracting useful information about depicted objects and scenes. It is widely used in many industrial applications including on-line process monitoring. From a pharmaceutical research and development perspective, the interest for image analytical applications is increasingly being recognized. One of the success stories for IA is particle analysis, giving quantitative estimation of geometrical and morphological properties (e.g. diameter, perimeter, area, roundness, etc.) of objects of interest as well as counting the objects.

A usual analytical procedure for particle analysis includes preprocessing to enhance image quality, segmentation and morphological operations with segmented binary masks. This procedure can be considered as a “direct measurement” of the objects of interest, and it can be efficiently implemented if objects are separated among each other and from the rest of scene, e.g. using threshold operation or edge detection.

In many cases however, segmentation can be a challenging step, especially if images are noisy and/or objects are partially overlapped or merged. In this case, using more sophisticated segmentation methods, e.g. a combination of traditional histogram based segmentation with demerging algorithms (e.g. watershed transformation) or shape detectors, can solve the problem. But for certain complicated cases, these methods do not work satisfactory and indirect approach can be considered as an alternative.

The idea of indirect approach is to calculate integral properties of an image (statistics of intensity or color distributions, textural properties, etc.) and use them as predictors in multivariate models from which the particle parameters can be inferred. Although, this approach does not allow estimation of the particle properties with high precision, an average estimation over the whole image can be obtained.

In the present study, crystallization of a needle shaped active pharmaceutical ingredient (API) was monitored using polarized light microscopy with both direct and indirect IA approaches. The methods were validated on crystallization simulated images. Because the needle shaped particle has poor flowability, the presence of this crystalline morphology can compromise further downstream processability. Hough and Radon transformations were employed for direct measurements, and Angle Measure Technique (AMT) was used for extracting textural properties for indirect measurements. Both techniques were applied to several sets of simulated and real images (video frames) of nucleation and crystal growth process of needle shaped particles, and their performance were estimated and compared.