Aalborg Universitet



## BASES OF CREATIVE PARTICLES OF HIGGS THEORY (CPH THEORY)

Javadi, Hossein; Forouzbakhsh, Farshid

Published in: Hadronic Journal

Publication date: 2010

Document Version Early version, also known as pre-print

Link to publication from Aalborg University

Citation for published version (APA): Javadi, H., & Forouzbakhsh, F. (2010). BASES OF CREATIVE PARTICLES OF HIGGS THEORY (CPH THEORY). Hadronic Journal, 33(5), 589-611. https://getinfo.de/app/BASES-OF-CREATIVE-PARTICLES-OF-HIGGS-THEORY-CPH/id/BLSE%3ARN290199692

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain You may freely distribute the URL identifying the publication in the public portal -

#### Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

# **Bases of CPH Theory**

# **Creative Particles of Higgs Theory**

H. Javadi<sup>a</sup> and F. Forouzbakhsh<sup>b</sup>

a) Invited professor of the Faculty of Science at Azad Islamic University, Tehran campuses Tehran, Iran

Javadi\_hossein@hotmail.com

b) Academic Researcher, Office of the Vice Chancellor for Research and Technology, University of Tehran. Tehran, Iran

fforouz@ut.ac.ir

#### Abstract

Zero Point Energy (ZPE) describes the random electromagnetic oscillations that are left in the vacuum after all other energy has been removed. One way to explain this is by means of the uncertainty principle of quantum physics, which implies that it is impossible to have a zero energy condition.

In CPH Theory the ZPE explained by using a novel description of the gravitons. This is based on the behavior of photons in a gravitational field, leading to a new definition of the graviton. In effect, gravitons behave as if they have charge and magnetic effects.

These are referred to as negative color charge, positive color charge and magnetic color. From this, it can be shown that a photon is made of color charges and magnetic color. This definition of the structure of a photon then leads to an explanation of how the vacuum produces Zero Point Energy (ZPE).

According to the results of this looking on gravitons we can definitely say that the best way for unifying the interactions is generalizing color charge from the structure of photon to nuclear. These color charges and magnetic color form the electromagnetic energy. Electromagnetic energy converts to matter and anti-matter such as charged particles. Charged particles use gravitons and generate electromagnetic field. This way of looking at the problem show how two opposite charged particles repel each other in far distance and absorb each other at a very small distance. To conclude, this article shows how quarks produce vector bosons.

**Keywords:** Zero point energy, Dirac equation, Hawking radiation, graviton, pair production, color-charge, magnetism color, photon, virtual photon, interactions, vector bosons, strong interaction, electroweak, broken symmetric, electromagnetic, spin, negative and positive photon, Higgs.

#### **1** Introduction;

Scientists describe the universe in terms of two basic partial theories - the general relativity and quantum mechanics... The general theory of relativity describes the force of gravity and the large-scale structure of the universe. Quantum mechanics, on the other hands, deals with phenomena on extremely small scales. These two theories are known to be inconsistent with each other - they cannot both be correct. There are many ways to do combine these theories and many theories such as Loop Quantum Theory and String Theory had propounded.

But Theory of CPH (Creative Particles of Higgs) takes a new way. CPH Theory has reconsidered 4 theories (Classical Mechanics, Quantum Mechanics, Relativity and Higgs). In fact CPH Theory is a new looking and developing of **Quantum Chromodynamic**. So, CPH Theory is a **Sub Quantum Chromodynamic theory**. In fact we must do change our understanding of graviton.

We know there is a unit informer in universe that is photon, and all of our information of universe transfer by photon. Until we do not know everything about photon and its structure, our information's about universe is questionable. Notice that many physicists, such as Faraday and Planck noted the great similarities between electric fields and gravity. If a unified field theory can be found, someone must resolve whether or not it is based on particles and gravity fields or electromagnetic fields. But CPH theory is some in between these two concepts.

CPH Theory started of relationship between force and energy. Photon appears to have no further internal substructure. But many phenomenon as Compton's effect, pair production, red-shift and blue-shift... lead CPH Theory show photon has a structure.

Theory of CPH have proclaimed by a simply definition of CPH and a principle that calls CPH Principle. In fact CPH theory is an empiric and sensibility theory. And it does different CPH Theory with other theories. Shortly, CPH theory proclaims the following conceptions;

1- When we will be able to explain quantum level phenomenon, that we do thinking on sub quantum quantities.

2- To explaining relationship between fermions and bosons, we must do change our mind of gravity and graviton. In fact gravitons behave like charge or magnet force in sub quantum levels.

3- We never can do combine Quantum mechanics with General Relativity without attention to Higgs theory. In fact there is an especial relationship between force and energy like mass and energy in relativity. This shows we reconsider the second Newton's law. It shows a unified theory comes up of reconsideration the quantum mechanics, relativity, Higgs theory and classical mechanics.

In CPH Theory, the ZPE (Zero Point Energy) explained by using a novel description of the graviton. This is based on the behavior of photons in a gravitational field, leading to a new definition of the graviton. In effect, gravitons behave as if they have charge and magnetic effects.

These are referred to as negative color charge, positive color charge and magnetic color. From this, it can be shown that a photon is made of color charges and magnetic color. This definition of the structure of a photon leads to an explanation of how the vacuum produces Zero Point Energy (ZPE).

Also, these color charges and magnetic color form the electromagnetic energy. Electromagnetic energy converts to matter and anti-matter such as charged particles. Charged particles use gravitons and generate electromagnetic field. In fact, a charged particle is a generator for producing virtual photons, both negative and positive photons. This view shows how two oppositely charged particles repel each other in greater distance and absorb each other at a very small distance. To conclude, it shows how quarks produce vector bosons. In general, it appears that all known interactions between objects can be described through negative and positive color charges.

## 2 The Photon in a gravitational field

Looking at the behavior of a photon in a gravitational field can help resolve vacuum energy. The fields around a "ray of light" are electromagnetic waves, not static fields. The electromagnetic field generated by a photon is much stronger than the associated gravitational field. Suppose a photon falls in a gravitational field, its energy (mass) increases. According to;

$$W = \Delta mc^2$$

The force of gravity performs work on the photon, so the mass (energy) of the photon increases.

However, the energy of a photon depends on its electric and magnetic fields. Therefore, one part of the work done by gravity converts to electrical energy and the other part converts to magnetic energy. How can the Higgs boson show how particles acquire mass? Moreover, according to the Higgs boson, what happens in during the blue shift?

## 3 Color-charges and color-magnetism

The change of frequency of a photon in a gravitational field has been demonstrated by the Pound-Rebka experiment. When a photon falls in a gravitational field, it acquires energy equal to;

which separates into three parts; one part behaves like a positive electric field and another part behaves like a negative electric field. These neutralize each other in the structure of the photon (a photon itself is neutral) and the third part behaves like a magnetic field.

In quantum mechanical theory, every field is quantized. In addition, force is described as energy per distance shown by:

$$F = -\frac{dU}{dx}$$
(1)

If we consider this equation from the aspect of quantum mechanics, a graviton enters into the structure of a photon, carrying gravitational force. As a result, a graviton disappears and the energy of the photon increases.

Similarly, Red Shift has the opposite effect. As a photon escapes from a gravitational field, its frequency shifts to red and its energy converts to gravitons. How can we describe this interaction between photons and gravitons on a sub-quantum scale such as in the structure of a photon?

Another consideration is virtual particles, implied by the uncertainty principle in the form:

$$\Delta E \Delta t \ge \hbar, \hbar = \frac{h}{2\pi}$$
 (2)

In the vacuum, one or more particles with energy  $\Delta E$  above the vacuum may be created for a short time  $\Delta t$ . So, any analysis of the ZPE should be able to explain this particle production and even Hawking radiation in a strong gravitational field such as that of a black hole. To do this, the best way is to use the Dirac equation, which originally demonstrated the possibility of a particle pair, that is, a particle and antiparticle. The relationship between energy and momentum for a massless particle is given by;

$$\langle \mathbf{E} \rangle = \langle |\mathbf{P}| \rangle c \qquad (3)$$

Now it is possible to change the definition of the rest mass of a particle. As we know, some particles such as photons are never seen at rest in any reference frame. According to relativity however, they do have mass that derives from their energy. For example, a photon has a mass given by:

$$E=mc^{2}=h\nu \implies m=\frac{h\nu}{c^{2}} \qquad (4)$$

So, there are two kinds of particles in physics;

• Some particles like the photon move only with the speed of light c, in all inertial reference frames. Let's call this kind the NR particles or Never at Rest condition particles

• Other particles like the electron always move with speed v<c in all inertial reference frames; they have a rest mass, and could not call NR particles.

According the above definition, photons and gravitons are NR particles, while electrons and protons are not NR particles.

Suppose a photon with NR mass m and energy E=hy falls toward the earth relative to an inertial reference frame on the surface of earth. Its frequency increases from v tov ', and a large number of gravitons enter into the structure of the photon such that

 $\Delta v = v' - v$ . The problem is; how many gravitons enter into a photon to provide the least possible photon energy change (minimum $\Delta v$ )?

Now, according to this argument, in order to calculate the number of gravitons involved in  $\Delta v$  and to explain their properties, suppose a photon with frequency  $\Box$  is formed of  $n_1$  elements, and with a frequency  $v_1$  it contains  $n_2$  elements. These elements are not the same, because they exhibit different properties. Let's propose a 1×4 matrix

Now we need to calculate A, B, C and D so that they satisfy the properties of a photon. When gravity works on a photon, gravitons enter into the photon and the intensity of its electric field increases.

The photon has no electrical effect; therefore A and B must carry electric field around the photon with opposite effect. So, according to the relative intensity of electric and magnetic fields E=cB, we can write;

A=cH<sup>+</sup>, B=cH<sup>-</sup>, c is the speed of light

Here  $\mathbf{H}^+$  is positive color-charge and  $\mathbf{H}^-$  is negative color-charge. In addition, in the above relation c is a mathematical constant that relates E and B in electromagnetism. So, let's show  $c=\kappa$ . Then the above relation becomes:

 $A = \kappa H^+$ ,  $B = \kappa H^-$ ,  $\kappa$  is a mathematical constant

When a large count of  $H^+$  enters into a photon, the intensity of its positive electric field increases. According to the Maxwell equations, the intensity of its magnetic field increases as well.

Also, element C must carry a magnetic effect around the positive color-charges and the same applies to the D element for the negative color-charges. Therefore, C and D are the same but with opposite direction. So, according to the relationship between the intensity of electric and magnetic fields, we can write;

Then the matrix  $\Box A B C D \Box$  takes the following form here named the CPH matrix;

$$\begin{bmatrix} CPH \end{bmatrix} = \begin{bmatrix} \kappa H^+ & \kappa H^- & H^m & H^m \end{bmatrix}$$
(5)

According to the above expression, we are now able to define the least magnitude of a photon. A photon of minute energy contains some positive color-charges  $H^+$ , negative color-charges  $H^-$ , right rotation color-magnetism  $H^m$  and left rotation color-magnetism  $-H^m$  as shown in the CPH matrix (relation 5). This very small energy can be express as the following;

Minute electromagnetic energy 
$$=(2\kappa+2)E_{CPH}$$
 (6)

Note; the energy of a CPH  $(E_{CPH})$  is defined later in relation (8).

The argument is now in a position to offer some supportable propositions about the photon and a new definition of the graviton.

### 4 Gravitons

Many physicists believe the graviton does not exist, at least not in the simplistic manner in which it usually envisioned. Superficially speaking, quantum gravity using the gauge interaction of a spin-2 field (graviton) fails to work the way that the photon and other gauge bosons do.

Maxwell's equations always admit a spin-1, linear wave, but Einstein's equations rarely admit a spin-2, linear wave, and when they do it is not exact.

However, in the present article the photon is made of gravitons. To resolve this, we need to continue with the definition of CPH and the Principle of CPH and then return to properties of a graviton.

#### 5 Definition of a CPH

What is a CPH? It is the Creative Particle of Higgs, or, CPH is an existence unit of nature. In other words, everything is made of CPH. Therefore, a CPH is appropriately referred to as the unit of nature, although this not meant to be a "particle" as this concept has been traditionally referred to in physics.

A CPH is a NR particle of a kind nonethess, with a constant NR mass  $^{m}_{CPH}$ , which moves with a constant magnitude speed of  $V_{CPH} > c$  in any inertial reference frame, where c is the speed of light. According to the mass-energy relation, the NR mass of a CPH is defined relative a to photon's NR mass by;

$$m_{CPH} < m = \frac{h\nu}{c^2} \forall \nu \qquad (7)$$

And the relationship between energy and momentum for the NR mass CPH is given by;

$$\langle \mathbf{E}_{CPH} \rangle = \langle |\mathbf{P}_{CPH}| \rangle \mathbf{V}_{CPH} = \text{constant}$$
 (8)

Relation (8) shows that the energy of every CPH is constant in any interaction between two (or more) CPH or other particles. So, given  $\left|V_{CPH}\right|$  cannot alter, they must take on spin to conserve the total energy.

In other words, in any inertial reference frame and Cartesian components;

$$\left|\mathbf{V}_{\text{CPH}}(x)\right| + \left|\mathbf{V}_{\text{CPH}}(y)\right| + \left|\mathbf{V}_{\text{CPH}}(z)\right| = \left|\mathbf{V}_{\text{CPH}}\right|, \text{ CPH has no spin}$$
(9)

$$|\mathbf{V}_{\text{CPH}}(x)| + |\mathbf{V}_{\text{CPH}}(y)| + |\mathbf{V}_{\text{CPH}}(z)| < |\mathbf{V}_{\text{CPH}}|$$
, CPH has spin (10)

#### When a CPH has spin, it is called a graviton

Simply, a lone graviton without spin is a CPH. When;

$$|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| \ge c$$
 (11)

There is no difference between bosons and fermions. In this case, a CPH carries gravitational force and behaves like a fermion. Therefore, there are color-charges, only.

When;

$$|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| \le c$$
, the magnetic effect appears.

For particles like electrons or quarks;

$$|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| < c$$
 (12)

Other bosons also occur. For example, reconsider pair production. Before pair production, there is a photon only. After pair production there is an electron, positron and a virtual photon (boson) that carries electromagnetic force. So, we can write:

$$\left| \mathbf{V}_{\text{CPH}}(y) \right| + \left| \mathbf{V}_{\text{CPH}}(z) \right| < c \tag{13}$$

Spontaneous Symmetry Breaking has occurred.

Accordingly, a CPH with spin is called a graviton, so space is full of CPH. Increasing density of CPH in space causes their separation to decrease until they feel and absorb each other.

Suppose two CPH are moving in the x-axis direction and absorb each other, such that their paths change without decreasing the magnitude of  $V_{CPH}$ . According to relation (8), we are able to construct an operator  $R_{\star}(\Delta \varphi)$ , which rotates a CPH by an angle  $\Delta \varphi$  about the x-axis (toward z-axis or y-axis) in position space. Also, we can construct  $T_{z}(\Delta \varphi)$  which rotates the CPH by  $\Delta \varphi$  about the x-axis in spin space. We would expect such an operator to take the form;

$$T_z(\Delta \varphi) = \exp(-\frac{\mathrm{i} \mathrm{s}_z \Delta \varphi}{\hbar})$$
 (14)

Thus, according to this expression, two interacting CPH rotate each other, but they cannot have same direction of spin. They spin in opposite directions. If positive color-charge has up spin, then negative color-charge must take on down spin.

### 6 Principle of the CPH theory

CPH is a unit of minuscule energy with a constant NR mass  $(m_{CPH})$  that moves with a constant magnitude of speed such that  $|V_{CPH}| > |c|$ , in all inertial reference frames.

Any interaction between CPH and other existing particles represents a moment of inertia I where; the magnitude of  $V_{CPH}$  (or  $|V_{CPH}|$ ) is constant and never changes.

Therefore,

 $\nabla |V_{CPH}| = 0$  in all inertial reference frames and any space

Based on the principal of CPH, a CPH has two types of energy generated by its movement within its inertial frame. One is translational and the other is spin. In physics, we represent energy summation (both kinetic and potential) by a Hamiltonian equation and energy difference by a LaGrangian. Therefore, in the case of CPH, we use a Hamiltonian to describe the summation of energy generated by translation and spin as follows:

$$E_{CPH} = T + S \tag{15}$$

Where T is translational and S is spin energy of a CPH respectively. Since the speed and mass of CPH are constant, then  $E_{CPH} = constant$ . CPH produces energy and energy produces Matter and Anti-Matter. In fact, everything has been formed of CPH.

#### 7 CPH and the cyclic group

As explained above, gravitons in interaction with each other convert to color-charges and color-magnetism. In addition, when a CPH has spin, it is calling a graviton. Therefore, we can define a cyclic group for electric field that is generated by gravitons.

So, G<g> given by;

$$G(g) = \{nH^+, nH^- \mid n \in Z\}$$
 (16)

Suppose 2n color-charges  $(nH^* \text{ and } nH^*)$  combine and move in space. There are two electric fields with opposite sign in that space. About each such field a magnetic field forms, produced by gravitons. According to the signs of these fields, the direction of this magnetism is different, so their elements are same. Therefore, there is a cyclic group given by:

$$G\langle g \rangle = \{ k H^m \,|\, k \in Z \}$$
(17)

According to the above group and the CPH matrix (relation 5), we can explain zero point energy.

#### **8 Zero Point Energy**

Space is full of gravitons. Gravitons interact with each other and convert to colorcharges Interaction between gravitons depends on their density  $\rho(g)$  in a given volume. Energy production is given by:

$$E = \iiint_{V} \rho(g) dx dy dz \qquad (18)$$

According to the above expression, we are able to explain the mechanism of ZPE.Some gravitons with the same NR mass CPH m convert to color-charges, and two electric fields form. These fields neutralize each other. However, positive color-charges repel each other, and the same action applies to the negative color-charges.

Therefore, when the intensity of color-charges grows, about each field (negative and positive fields) a magnetic field forms. This magnetic field maintains the electric field. This mechanism is explainable by the Larmor radius (gyroradius or cyclotron radius) given by;

$$r_{g} = \frac{mv_{\perp}}{|q|B}$$
(19)

Where  $r_g$  is the gyroradius, m is the mass of the charged particle,  $v \square$  is the velocity component perpendicular to the direction of the magnetic field, q is the charge of the particle, and B is the constant magnetic field.

This defines the radius of circular motion of a charged particle in the presence of a uniform magnetic field. When; color-charges change in the structure of a photon, then magnetic-color changes too. Therefore the electric fields do not decay in the structure of a photon.

In general, a photon has been formed of two parts;

1- A large number of negative color charges and magnetic color. Magnetic color maintains color charges in a tube-like distribution, so negative magnetic color forms an appropriate negative electric field. In addition, the same happens for positive electric field in the opposite sense. So it is now possible to demonstrate the least possible negative color charges with their magnetic color by  $\triangleleft$ , so that;

$$\triangleleft = (\kappa H^{-}, -H^{m}) \tag{20}$$

2- Similarly to the above; positive color charges with their magnetic color can be shown by  $\triangleright$ , so that;

$$\triangleright = (\kappa H^+, +H^m) \tag{21}$$

The signs (+and -) of  $(+H^m)$ ,  $(-H^m)$  depend on the direction of movement around the color charges. In fact, there is a kind of magnetic color in the structure of a photon. Therefore, generally, a photon is given by;

$$\mathbf{n} | \triangleleft \rangle + \mathbf{n} | \triangleright \rangle = | \mathbf{E} \rangle \tag{22}$$

In the quantum mechanics of any general field, plane waves of specific spin can always be written in terms of photons with a simple spin state and a general spatial wave function. Thus the fundamental entity, the photon, can be considered quite generally to be a plane wave with a circularly polarized spin component (Any field can be built from these basic ingredients).

For simplicity, consider a photon traveling in the x direction, or consider the direction of the photon as choosing the coordinate axis so that x points along the photon's momentum vector. Every element in the photon (relation 22) moves with linear speed in

the same direction as the photon.

The Dirac equation results for total energy are given by;

$$E^{2} = \left(\alpha_{0}mc^{2} + \sum_{j=1}^{3}\alpha_{j}cp_{j}\right)^{2}$$
(23)

Comparing relationship (22) with equation (23), the energy equation can now be written as;

$$\mathbf{E}^{2} = \left( \mathbf{n} \left| \triangleleft \right\rangle + \mathbf{n} \left| \triangleright \right\rangle \right)^{2} \tag{24}$$

The roots of relations (23) and (24), are given by;

$$\mathbf{E}_{-} = -\sqrt{\alpha_0 \mathbf{m}_0 \mathbf{c}^2 + \sum_{j=1}^{3} \alpha_j c p_j} = \mathbf{n} \triangleleft = \mathbf{n}(\kappa \mathbf{H}^-, -\mathbf{H}^m)$$
(25)

$$E_{+} = + \sqrt{\alpha_0 m_0 c^2 + \sum_{j=1}^{3} \alpha_j c p_j} = n \triangleright = n(\kappa H^+, + H^m)$$
(26)

In pair production, relation (25) defines an electron and relation (26) defines a positron. We saw that a photon ( $\gamma$ -ray) produces two charged particles, an electron and positron with negative and positive charges. Before production, we have two electric and two magnetic fields. After production, there are two particles with electric field and two weak magnetic fields around them.

This phenomenon shows that an electric field has no charge effect when formed of two kinds of color-charge, namely negative color-charge and positive color-charge.

Moreover, in pair production, negative color-charges combine with each other to make a negatively charged particle, and positive color-charges combine with each other to make a positively charged particle. The magnetic colors with different direction move around the electron and positron. Consider relations (25) and (26), given by;

$$E_{-}=n \triangleleft = n(\kappa H^{-}, -H^{m})$$
$$E_{+}=n \triangleright = n(\kappa H^{+}, +H^{m})$$

The pair annihilates each other to form energy. In addition, there is no electric effect around the photons, so;

$$\mathbf{n} \triangleleft + \mathbf{n} \triangleright = \gamma + \gamma \tag{27}$$




Fig1: annihilation of a particle pair

In this process, each particle (electron and positron) decomposes to two parts. Each part of the electron combines with each part of the positron and converts to quantum energy (see figure1). This phenomenon shows that the electron is divisible.

Physicists use this phenomenon as a way to confirm the mass-energy equation  $E=mc^2$ , but in fact, there is another important concept inherent in pair annihilation since;

$$\mathbf{n} \triangleleft + \mathbf{n} \triangleright = \left(\frac{\mathbf{n}}{2} \triangleleft + \frac{\mathbf{n}}{2} \triangleright\right) + \left(\frac{\mathbf{n}}{2} \triangleleft + \frac{\mathbf{n}}{2} \triangleright\right) = \gamma + \gamma \tag{28}$$

These photons are neutral and they carry two electric and magnetic fields. This phenomenon is acceptable only where two opposite charged particles separate and recombine again.

#### **9 Hawking Radiation**

In a simplified version of the explanation, Hawking predicted that energy fluctuations from the vacuum cause the generation of particle-antiparticle pairs near the event horizon of a black hole. One of the particles falls into the black hole while the other escapes before they have an opportunity to annihilate each other. The net result is that to someone viewing the black hole, it would appear that a particle has been emitted.

How is Hawking radiation explainable by the equation for the ZPE? To resolve this problem, there are three aspects of a black hole to consider;

- The density of gravitons is extremely high around a black hole
- Gravitons convert to photons readily
- The Dirac equation shows how photons produce matter and anti-matter.

According to the above expression, the space around a black hole produces highenergy photons whose energy is enough for pair production.

In a black hole situation, n becomes large, so  $E^2$  (relation 24) is comparable to the total mass of a particle and anti-particle. This process does not need to take into account the time factor that the uncertainty principle dictates in relation (2). So, pair production is a common occurrence around a black hole.

#### 10 Dynamics of charge particles of CPH Theory view point

Consider the electron and positron that give by relations (20) and (21). The electron contains a set of negative color charges that keeps by magnetic colors. This rotational sphere-like (electron spinning) is in a sea of gravitons. Gravitons are negative and positive charges color. Around the negative color charges of an electron is a magnetic field.

The electron has two opposite effects on color charges around itself. Negative color charges of electron absorb positive color charges (of space) and repel negative color charges. Magnetic field contracts positive color charges and repels them (see Ampere law).

Now we can define an operator for producing positive electric force particle. Let us show this operator by  $(\triangleleft \$)$  per time that acts on the electron and produces positive electric force. So, it given by;

$$\frac{\mathrm{d}}{\mathrm{dt}} \triangleleft \mathrm{s=a} \triangleright = \mathrm{a}(\kappa \mathrm{H}^{+}, +\mathrm{H}^{\mathrm{m}}) \tag{29}$$

There, a is a natural number. Consider that  $\triangleleft s$  is a set of positive color charges, it makes a positive electric field around the electron. This electric field repels the negative charge particle, because every negative charge particle produces same electric field.

Same as above, positron produces a negative electric field around itself. So, it given by;

$$\frac{\mathrm{d}}{\mathrm{dt}} \triangleright s = a \triangleleft = a(\kappa \mathrm{H}^{\mathrm{c}}, -\mathrm{H}^{\mathrm{m}})$$
 (30)

When a negative electric force particle  $(a \triangleleft)$  reaches to positron, it combines with positive electric force particle  $a \triangleright$  and they convert to quantum energy, so that;

$$|a \triangleright\rangle + |a \triangleleft\rangle = E$$
 (31)

This quantum energy transfers to the positron. Then, the positron accelerates toward the electron. The same process happens for the electron, and they absorb each other. For understanding, this process reconsiders to the annihilation of pair. Opposite interaction happens between two same charge particles, and they repel each other.

In fact, the structure of a virtual photon is difference from a real photon. There are different kinds exchange particles that carry electromagnetic interaction, one is a positive photon and other one is a negative photon. This point of view is able to explain easily the interaction between charge particles. In other words;

$$\frac{d}{dt} \triangleleft s = \gamma^{+} \qquad (32)$$
$$\frac{d}{dt} \triangleright s = \gamma^{-} \qquad (33)$$

And;

$$\gamma^{+} + \gamma^{-} = \gamma \qquad (33)$$

This view shows why a virtual photon is invisible.

#### **11 Strong Interaction and CPH Theory**

This section shows how positive charge particles absorb each other in very small distance. Generally, two positive charged particles produce banding energy, in small distances.

According to quantum chromodynamics, a proton is made of two up quarks (u) with (+2/3) charge and a down quark with (-1/3) charge. How it is that two up quarks with positive charged do not repel each other?

Generally, suppose two positive charged particles A and B are at far distance from each other. As explained in section 10, any positive charged particle absorbs negative color charges and repels positive color charges. When the distance between A and B is greater than the atomic radius, they emit negative photon  $\gamma^-$ . There are three locations around each positive charged particle (figure2).




Fig2; Locations around each positive charged particle

In real space, every charged particle is plunging in a sea of gravitons. Location 3 (figure2) is full of gravitons that move with speed of  $v \ge c$ . When gravitons reach to location 2, electrical field (or magnetic field) of charged particle acts on them so that gravitons convert to positive and negative color charges. Positive charged particle repels positive color charges and absorbs negative color charges. Therefore, negative color charges enter into location 1 (figure2). In location 1, negative color charges convert to negative photon that given by (relation 33). In general, location 3 is full of gravitons, location 2 is full of negative and positive color charges, and positive charged particle generates negative photon in location 1. Now suppose two positive charged particles (A<sup>+</sup>, B<sup>+</sup>) are near each other that location 2 interferes with each other (Figure3).




Fig 3; interconnect two positive charged particles

Locations 2 of A and B interconnect, there is a set color charges generated by A, described as follows:

$$\left\{ (\mathrm{H}^{+},\mathrm{H}^{-}) \middle| \mathrm{H}^{+},\mathrm{H}^{-} \in \mathrm{field} \mathrm{A} \right\}$$

Charged particle A repels positive color charges  $(H^+)$ , they move toward B particle, and negative color charges  $(H^-)$  move toward A.

Also, charged particle B generates a set of positive and negative color charges  $(H^+, H^-)$  as follows;

$$\left\{ (\mathrm{H}^{+},\mathrm{H}^{-}) \middle| \mathrm{H}^{+},\mathrm{H}^{-} \in \mathrm{field} \; \mathrm{B} \right\}$$

Their direction movement is the opposite of A production. Therefore, in location 2, positive color charges  $H^+$  from A and negative color charges  $H^-$  from B, have the same direction movement that is toward the B particle. They combine and convert to electromagnetic energy and move toward the particle B. The same action happens for positive color charges  $H^+$  from B and negative color charges  $H^-$  from A; so, they form quantum energy that moves toward A. This shown as follows;

$$|a\rangle + |a\rangle = E$$

These energies form the banding energy between A and B. In a heavy nucleus, that contains a lot of protons, every quark interacts with each other and produces banding energy

Consider the centre of stars, two hydrogen ions (protons) move toward each other, when their distance decreases, then locations 2 of them interconnect and produce banding energy.

#### 12 Weak interaction

In the Standard Model of particle physics, it is due to the exchange of the heavy  $Z^0$ ,  $W^+$ ,  $W^-$ . Its most familiar effect is beta decay and the associated radioactivity. Consider a neutron (contents one up quark, two down quarks). Although the neutron is heavier than its sister nucleon, the proton (contents *uud*), it cannot decay into a proton without changing the flavor of one of its down quarks. Neither the strong interaction nor electromagnetic allow flavor changing, so this must proceed by weak decay. In this process, a down quark in the neutron changes into an up quark by

emitting a  $W^{-}$  boson, which then breaks up into a high-energy electron and an electron antineutrino. Since high-energy electrons are beta radiation, this is called beta decay.

$$^{137}_{55}$$
Cs  $\rightarrow ^{137}_{56}$ Ba+e<sup>-</sup> + $\overline{\nu_e}$ 

In the above formula, a neutron udd (two down quarks  $2 \times -\frac{1e}{3}$  and one up quark  $+\frac{2e}{3}$  electric charge) decays to a proton (uud  $2 \times +\frac{2e}{3}$  and  $-\frac{1e}{3}$  electric charges), with +e electric charge. Before reaction, there is no integer charge in neutron structure, but after reaction, there is an electron with -e electric charge. Suppose an electron formed of E\_=n <, therefore, a down quark given by  $\overline{3}$ , and an up quark shown by  $\overline{3}$ . It is attendable that a quark over that negative (or positive) color charges contains electromagnetic energy  $(|a > \rangle + |a < \rangle = E$ ), so the mass of up quark is not equal the  $2\pi \bowtie$ 

mass of  $\overline{3}$ .

For example; in  $\beta^+$  decay with +e electric charge that is equivalent of  $n \triangleright$ , consider the following relation;

energy + 
$$p^+ \rightarrow n^0 + e^+ + \nu_e$$

A n<sup>0</sup> contains two down quarks  $[2 \times (-\frac{1e}{3} \text{ or } \frac{n <}{3})]$  and one up quark  $(+\frac{2e}{3} \text{ or } \frac{2n >}{3})$ , but a proton contains  $[2 \times (\frac{2n >}{3} \text{ or } +\frac{2e}{3})]$  and one down quark  $(-\frac{1e}{3} \text{ or } \frac{n <}{3})$ .

So, in above formula, energy separates color charges and recombines them in a new form. There is a positron with integer  $(n \triangleright or + e)$  electric charge, in right side of above formula, but there is not so integer electric charge, in left side. Therefore, electric charge of particles is not stable, but the color charges conserved. It meant that the sum number of negative and positive color charges is equal in any reaction. Therefore, vector bosons  $z^0 \downarrow W^+$ ,  $W^-$  given by;

$$W^{+} \sim n \triangleright$$
$$W^{-} \sim n \triangleleft$$
$$Z^{0} \sim (k \triangleright + k \triangleleft)$$

#### Summary;

According to this article we have generalized color charge from the nuclear regime to the photon. This new view of color charge means that we can redefine the graviton and electromagnetic energy. Gravitons behave like charged particles and in the interaction between gravity and the photon, gravitons convert to negative and positive color charges and magnetic color. These color charges and magnetic color form the

electromagnetic energy. Electromagnetic energy converts to matter and anti-matter as charged particles. Space is full of gravitons. Gravitons interact with each other and convert to color-charges. These color charges and magnetic color form electromagnetic energy.

This view shows how particles appear and when Spontaneous Symmetry Breaking has occurred.

 $|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| = |V_{CPH}| > c$ , CPH has not spin

CPH is isolation that never seems.

 $|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| < |V_{CPH}|$ , CPH has spin

When CPH has spin, it calls graviton

Space (vacuum) is full of CPH (gravitons), and CPH produces energy.

$$|\mathbf{V}_{\text{CPH}}(x)| + |\mathbf{V}_{\text{CPH}}(y)| + |\mathbf{V}_{\text{CPH}}(z)| \ge c$$

Gravitons convert to negative and positive color charges and magnetic color.

 $|\mathbf{V}_{\text{CPH}}(x)| + |\mathbf{V}_{\text{CPH}}(y)| + |\mathbf{V}_{\text{CPH}}(z)| \le c$ 

Electric and magnetic fields appear.

$$|V_{CPH}(x)| + |V_{CPH}(y)| + |V_{CPH}(z)| < c$$

#### matter and anti-matter appear

Spontaneous Symmetry Breaking has occurred. Moreover, fundamental particles appear, then fermions and bosons interaction with each other. For example, reconsider pair production, before pair production, there is a photon only. There is an electron a positron and virtual photon (boson) that carries electric force, after pair production.

#### Acknowledgment:

The authors would like to express their sincere thanks to William M. MacArthur (wilmac@aapt.net.au) and Sean McCarthy (seanmac@cablenet.ie) for editing the current article.

#### **Refrences;**

1- Richard Cushman. and Wilberd van der Kallen. (2005). 'A new interpretation for the mass of a classical relativistic particle', Elsevier Journal, Physics Letters Volume 24, Issue 3, May 2006, Pages 230-234

2- Ravindran V. (2006) 'Higher-order threshold effects to inclusive processes in QCD', Elsevier Journal, Physics Letters B, Available online 11 July 2006

Perelstein M. (2006) 'Little Higgs models and their phenomenology', Elsevier Zero Point Energy and Dirac Equation

#### Zero Point Energy and Dirac Equation

#### 15/15

3- Journal, Progress in Particle and Nuclear Physics, Available online June 2006 4- Gupta G. P. and. Lal K. C. (1971) *'On the equivalence of gravitational red shift and temperature shift in the M\_ssbauer effect'*, Elsevier Journal, Physics Letters A Volume 36, Issue 5, September 1971, Pages 421-422

5 - Peter S. Riseborough, E-book, Advanced Quantum Mechanics March 14, 2007
6- Stefan Waner, E-book, Introduction to Differential Geometry & General Relativity
4th Printing January 2005

7- http://www.daviddarling.info/encyclopedia/C/Casimir.html

8- \* advanced string theory http://superstringtheory.com/basics/basic3a.html

9- F. Smarandache, V. Christianto, Fu Yuhua, R. Khrapko, J. Hutchison, E-book, Unfolding

the Labyrinth: Open Problems in Physics, (University of Microfilm International)

http://physics.ucr.edu http://physics.bu.edu/py106/Notes.html http://www.physics.csbsju.edu http://www.newadvent.org/cathen/06342b.htm http://archive.ncsa.uiuc.edu/ http://physics.syr.edu http://hyperphysics.phy-astr.gsu.edu/hbase/relativ/mmhist.html#c1 http://www.schoolscience.co.uk/content/4/physics/particles/particlesmodel6.html http://www.superstringtheory.com/experm/exper2.html http://www2.slac.stanford.edu/vvc/theory/model.html http://archive.ncsa.uiuc.edu/ http://www.toequest.com/ http://www-groups.dcs.st-and.ac.uk http://en.wikipedia.org http://csep10.phys.utk.edu/astr161/lect/index.html http://csep10.phys.utk.edu/