



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The Western Denmark Cardiac Computed Tomography Registry

a review and validation study

Nielsen, Lene Hüche; Nørgaard, Bjarne Linde; Tilsted, Hans-Henrik; Sand, Niels Peter; Jensen, Jesper Møller; Bøttcher, Morten; Diederichsen, Axel C.; Lambrechtsen, Jess; Kristensen, Lone Deibjerg; Mickley, Hans; Munkholm, Henrik; Gøtzsche, Ole; Knudsen, Lars Lyhne; Bøtker, Hans Erik; Pedersen, Lars; Schmidt, Morten

Published in:
Clinical Epidemiology

DOI (link to publication from Publisher):
[10.2147/CLEP.S73728](https://doi.org/10.2147/CLEP.S73728)

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Nielsen, L. H., Nørgaard, B. L., Tilsted, H.-H., Sand, N. P., Jensen, J. M., Bøttcher, M., Diederichsen, A. C., Lambrechtsen, J., Kristensen, L. D., Mickley, H., Munkholm, H., Gøtzsche, O., Knudsen, L. L., Bøtker, H. E., Pedersen, L., & Schmidt, M. (2015). The Western Denmark Cardiac Computed Tomography Registry: a review and validation study. *Clinical Epidemiology*, 7, 53-64. <https://doi.org/10.2147/CLEP.S73728>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

The Western Denmark Cardiac Computed Tomography Registry: a review and validation study

Lene Hüche Nielsen¹
 Bjarne Linde Nørgaard²
 Hans Henrik Tilsted³
 Niels Peter Sand⁴
 Jesper Møller Jensen²
 Morten Bøttcher⁵
 Axel C Diederichsen⁶
 Jess Lambrechtsen⁷
 Lone Deibjerg Kristensen⁸
 Hans Mickley⁶
 Henrik Munkholm¹
 Ole Gøtzsche²
 Lars Lyhne Knudsen⁵
 Hans Erik Bøtker²
 Lars Pedersen⁹
 Morten Schmidt^{2,9}

¹Department of Cardiology, Lillebaelt Hospital-Vejle, Vejle, Denmark;

²Department of Cardiology Aarhus University Hospital-Skejby, Aarhus, Denmark; ³Department of Cardiology, Aalborg University Hospital, Aalborg, Denmark;

⁴Department of Cardiology, Hospital of Southwestern Denmark-Esbjerg, Esbjerg, Denmark;

⁵Department of Cardiology, Regional Hospital Herning, Herning, Denmark;

⁶Department of Cardiology, Odense University Hospital, Odense, Denmark;

⁷Department of Cardiology, Odense University Hospital-Svendborg, Svendborg, Denmark;

⁸Department of Cardiology, Regional Hospital Silkeborg, Silkeborg, Denmark;

⁹Department of Clinical Epidemiology, Aarhus University Hospital, Aarhus, Denmark

Correspondence: Lene Hüche Nielsen
 Department of Cardiology,
 Lillebaelt Hospital-Vejle, Kappeltoft 25,
 DK-7100 Vejle, Denmark
 Email lenehuche@gmail.com

Background: As a subregistry to the Western Denmark Heart Registry (WDHR), the Western Denmark Cardiac Computed Tomography Registry (WDHR-CCTR) is a clinical database established in 2008 to monitor and improve the quality of cardiac computed tomography (CT) in Western Denmark.

Objective: We examined the content, data quality, and research potential of the WDHR-CCTR.

Methods: We retrieved 2008–2012 data to examine the 1) content; 2) completeness of procedure registration using the Danish National Patient Registry as reference; 3) completeness of variable registration comparing observed vs expected numbers; and 4) positive predictive values as well as negative predictive values of 19 main patient and procedure variables.

Results: By December 31, 2012, almost 22,000 cardiac CTs with up to 40 variables for each procedure have been registered. Of these, 87% were coronary CT angiography performed in patients with symptoms indicative of coronary artery disease. Compared with the Danish National Patient Registry, the overall procedure completeness was 72%. However, an additional medical record review of 282 patients registered in the Danish National Patient Registry, but not in the WDHR-CCTR, showed that coronary CT angiographies accounted for only 23% of all nonregistered cardiac CTs, indicating >90% completeness of coronary CT angiographies in the WDHR-CCTR. The completeness of individual variables varied substantially (range: 0%–100%), but was >85% for more than 70% of all variables. Using medical record review of 250 randomly selected patients as reference standard, the positive predictive value for the 19 variables ranged from 89% to 100% (overall 97%), whereas the negative predictive value ranged from 97% to 100% (overall 99%). Stratification by center status showed consistently high positive and negative predictive values for both university (96%/99%) and nonuniversity centers (97%/99%).

Conclusion: WDHR-CCTR provides ongoing prospective registration of all cardiac CTs performed in Western Denmark since 2008. Overall, the registry data have a high degree of completeness and validity, making it a valuable tool for clinical epidemiological research.

Keywords: coronary computed tomography angiography, database, epidemiology, registries

Introduction

Technological advances in computed tomography (CT) have facilitated the development of cardiac CT enabling rapid and accurate imaging of the cardiovascular system, including coronary arteries, coronary arterial wall, cardiac valves, myocardium, and associated structures.¹ The primary clinical indication for cardiac CT is the detection or exclusion of coronary artery disease (CAD) in patients suspected of angina (ie, coronary CT angiography [CTA]).^{2,3} As a subregistry of the Western Denmark

Heart Registry (WDHR),⁴ the Western Denmark Cardiac Computed Tomography Registry (WDHR-CCTR) was established in 2008.⁵ In addition to monitoring and improving the cardiac CT quality in Western Denmark, the aim of the WDHR-CCTR is to allow for real-world clinical and health-service research.⁵ The coronary interventional data of the WDHR have previously been reviewed,⁴ whereas the cardiac CT data have not. We therefore examined the content, data quality, and research potential of the WDHR-CCTR, focusing particularly on the coronary CTA data.

Methods

Setting

Western Denmark has a population of 3.3 million (55% of the total Danish population). The Danish National Health Service provides tax-supported health care for the entire Danish population including free access to hospitals, general practitioners and outpatient specialty clinics, and partial reimbursement of prescribed medications.⁶ The Danish Civil Registration System assigns a unique 10-digit Civil Personal Register (CPR) number to all residents at birth or upon immigration and records information on residents' vital and emigration status with daily update.⁷ The CPR number enables linkage between all registries in Denmark at the individual level.⁷

Data registration

The WDHR-CCTR is derived from an internet-based online system, running on an encrypted public net.⁴ A common interface form secures standardized data collection from the cardiac centers.⁴ To optimize the data quality, data are entered by the cardiac center at which the procedure is performed into a computer-based data management system using the patient's CPR number.⁴ Serial numbers for patients and procedures are automatically generated to protect patient confidentiality. Standardized forms are used for registration of data. Three registration forms (a referral form, a medical history form, and a procedural form) are completed when entering data into the WDHR-CCTR (Table S1).

Study population

As of 2014, all nine cardiac centers (three university and six nonuniversity centers) in Western Denmark report to the WDHR-CCTR (Figure 1). University cardiac centers are defined by performing coronary revascularization procedures. WDHR-CCTR previously included ten cardiac centers, but recently the two centers at Aarhus University Hospital (Skejby and Tage Hansens Gade) were merged into one center. The WDHR-CCTR includes information on adult




Figure 1 Cardiac centers reporting to Western Denmark Cardiac Computed Tomography Registry.

Notes: Color codings indicate the five regions of Denmark. Numbers indicate the cardiac centers reporting to the registry: (1) Vendsyssel Hospital; (2) Aalborg University Hospital; (3) Regional Hospital Herning; (4) Regional Hospital Silkeborg; (5) Aarhus University Hospital (previously comprising two separate cardiac centers); (6) Lillebaelt Hospital-Vejle; (7) Hospital of Southwestern Denmark-Esbjerg; (8) Odense University Hospital; and (9) Odense University Hospital-Svendborg Hospital.

patients (≥ 16 years) referred for cardiac CT in Western Denmark. We included all patients registered between January 1, 2008, and December 31, 2012. In addition, we examined separately the subgroup of patients undergoing coronary CTA on suspicion of angina.

Statistical analysis

Content

We reviewed the variables recorded in the WDHR-CCTR and presented categorical variables using frequencies and proportions, and continuous variables using medians and interquartile ranges.

Completeness

By completeness of data, we refer to the proportion of registrations completed. Data completeness was evaluated at two levels: 1) completeness of procedure registrations and 2) completeness of variable registrations.

To estimate completeness of procedure registration, we used the Danish National Patient Registry (DNPR) as reference standard. Since 1977, all admissions to Danish hospitals have been registered in the DNPR using one primary discharge diagnosis and when appropriate secondary discharge diagnoses according to the International Classification of Disease (eighth revision through 1993 and the

tenth revision thereafter).⁸ Outpatient specialty clinic visits have been included in the DNPR since 1995 and diagnostic procedures since 1999.⁸ Cardiac CT is coded in the DNPR according to the Nordic Medical Statistics Committees Classification of Surgical Procedures (code: UXCC00A).⁹ The DNPR forms the basis for hospitals financial reimbursement, and hence, the completeness of procedure registrations in the DNPR is likely high. We therefore considered the DNPR an appropriate reference standard.

The completeness of procedure registration in the WDHR-CCTR was calculated as the proportion of patients registered with a cardiac CT in the DNPR from hospitals in Western Denmark that was also registered in the WDHR-CCTR.¹⁰ Of note, coronary and noncoronary CTAs are not registered separately in the DNPR. Therefore, to evaluate the degree of underregistration of coronary and noncoronary CTAs in the WDHR-CCTR in WDHR-CCTR, we reviewed the medical records from all patients ($n=282$) who had cardiac CT registered in the DNPR, but not in the WDHR-CCTR during 2008–2012 at Aarhus University Hospital and Lillebaelt Hospital-Vejle.

The completeness of the individual variables was calculated as the number of patients with information on the variable divided by the total number of patients registered in the WDHR-CCTR. If patients underwent more than one cardiac CT within the study period, only data from the first procedure were included in the analyses.

Validity

For variable validation, we included only coronary CTAs because it is by far the most common indication for cardiac CT and the only one reported consecutively to the WDHR-CCTR since its establishment. We assessed the validity of the WDHR-CCTR data through review of hospital medical records (both paper and electronic files). A computer-generated random sample of 250 patients from five different centers (50 patients from each center) was drawn. We selected patients from both university (Aarhus, Aalborg, and Odense) and nonuniversity cardiac centers (Lillebaelt Hospital-Vejle and Hospital of Southwestern Denmark-Esbjerg). Within each cardiac center, patients were selected randomly regardless of whether data were missing on one or more variables. Correlating with the increasing number of procedures performed since 2008, the random sample included 4.8% of all patients from 2008, 13.6% from 2009, 26.0% from 2010, 26.4% from 2011, and 29.2% from 2012.

For each procedure, we selected 19 main patient and procedure variables (Table S1). One clinician (LHN) reviewed

all medical records. During validation, one of three outcomes was recorded: confirmed by medical record, not confirmed by medical record, or missing in medical record. To facilitate calculation of predictive values also for continuous variables and variables with more than two categories, the registry data were considered confirmed in the medical records only when there was an exact match between the variable values in the registry and medical records. We considered the procedure date correct if the differences between the dates in the WDHR-CCTR and the medical records were <1 week.

We constructed a 2×2 table with results for the true-positive, false-positive, false-negative, and true-negative scores, enabling calculation of positive predictive values (PPVs) and negative predictive values (NPVs) for each variable. PPV was defined as the number of true positives/(number of true positives + number of false positives). NPV was defined as the number of true negatives/(number of true negatives + number of false negatives). We used Wilson's score methods to calculate confidence intervals (CIs).¹¹

Patients with missing information in the medical files on a specific variable were omitted from the calculation of PPV and NPV. However, because missingness potentially could be associated with the accuracy of variable registration, we repeated the analyses by including all missing medical records as confirmed (best-case scenario) and nonconfirmed (worst-case scenario) cases in the calculations. Finally, we stratified the predictive values by cardiac center status and calendar year. All analyses were performed with STATA (version 13.0). The present study was approved by the Danish Data Protection Agency (2008-58-0035), and the National Board of Health (Department of Monitoring and Patient Safety) approved access to medical records (3-3013-377).

Results

Content

During 2008–2012, we identified 21,623 patients registered with a cardiac CT in the WDHR-CCTR. Of these, 1.5% underwent more than one procedure. Among all patients, 18,899 patients underwent a coronary CTA (87%). The annual number of patients examined by coronary CTA increased from 369 patients in 2008 to 6,454 patients in 2012. The characteristics of the cardiac CT and the coronary CTA cohorts are provided in Table 1.

Completeness

Throughout the study period, a consistently higher number of cardiac CTs were registered in the DNPR than in the WDHR-CCTR (Figure 2). The proportion of patients

Table 1 Patient characteristics and variable completeness in the Western Denmark Cardiac Computed Tomography Registry, 2008–2012

	Frequency of characteristics		Variable completeness, %	
	Cardiac CT cohort overall (n=21,623)	Coronary CTA cohort ^a (n=18,899)	Cardiac CT cohort overall	Coronary CTA cohort ^a
Male sex	10,169 (47.0)	8,506 (45.0)	100	100
Age, years ^b	58 (50–66)	58 (50–66)	100	100
County	–	–	97.1	97.6
Area code	–	–	97.1	97.6
Referral date	–	–	78.7	77.2
Family history of premature CAD	8,044 (37.2)	7,353 (38.9)	85.7	82.7
Current lipid-lowering therapy	6,589 (30.5)	5,883 (31.1)	85.6	85.5
Current medical treatment for hypertension	7,400 (34.2)	6,532 (34.6)	85.5	85.3
Smoking status (current or former)	10,648 (49.2)	9,504 (50.3)	85.9	85.8
Diabetes	1,208 (5.6)	1,035 (5.5)	85.9	85.4
Previous PCI	278 (1.3)	166 (0.9)	85.9	86.2
Previous cardiac surgery	354 (1.6)	156 (0.8)	85.9	87.0
Previous myocardial infarction	310 (1.4)	193 (1.0)	85.9	86.1
Height (cm) ^b	172 (165–178)	171 (165–178)	79.5	80.6
Weight (kg) ^b	78 (67–89)	78 (67–88)	79.8	80.9
Creatinine level, $\mu\text{mol/L}^b$	73 (64–84)	73 (63–83)	77.8	78.8
Systolic blood pressure, mmHg ^b	140 (126–151)	140 (126–151)	77.6	78.9
Diastolic blood pressure, mmHg ^b	81 (75–90)	81 (75–90)	77.5	78.8
Left ventricular ejection fraction ^b	60 (60–60)	60 (60–60)	58.2	59.7
Type of CT scanner	–	–	98.5	99.7
Cardiac CT indication	–	–	99.6	100
Angina/equivalent	18,899 (87.4)	–	–	–
Heart failure	292 (1.4)	–	–	–
Visualization of valves and/or aorta disease	239 (1.1)	–	–	–
Visualization of stents	71 (0.3)	–	–	–
Evaluation of cardiac structures	631 (2.9)	–	–	–
Others	1,410 (6.5)	–	–	–
Unknown	81 (0.4)	–	–	–
Angina type (if any)	–	–	61.4	70.3
Typical angina	–	1,764 (9.4)	–	–
Atypical angina	–	5,844 (30.9)	–	–
Unspecific chest pain	–	4,760 (25.2)	–	–
Dyspnea	–	916 (4.8)	–	–
Unknown	–	5,615 (29.7)	–	–
Priority	–	–	93.9	95.5
Acute	118 (0.5)	78 (0.4)	–	–
Subacute	860 (4.0)	755 (4.0)	–	–
Elective	19,317 (89.3)	17,209 (91.1)	–	–
Unknown	1,328 (6.2)	857 (4.5)	–	–
Date of procedure	–	–	100	100
Kilo-electron volt ^b	100 (100–120)	100 (100–120)	97.2	98.4
Radiation dose	– ^c	– ^c	93.8	94.8
Iodine concentration, mg/mL ^b	350 (350–370)	350 (350–370)	93.0	94.6
Contrast volume, mL ^b	80 (70–90)	80 (70–90)	92.9	94.5
Preprocedural beta-blocker	13,007 (60.2)	11,991 (63.4)	93.4	95.1
Heart rhythm	–	–	93.5	95.2
Sinus rhythm	19,855 (91.8)	17,794 (94.2)	–	–
Atrial fibrillation	266 (1.2)	108 (0.6)	–	–
Other	106 (0.5)	85 (0.4)	–	–
Unknown	1,396 (6.5)	912 (4.8)	–	–
Heart rate, bpm ^b	59 (54–65)	59 (54–64)	93.4	95.0
Cardiac CT quality	–	–	88.9	90.1
Good	16,120 (74.6)	14,304 (75.7)	–	–
Moderate	2,170 (10.0)	1,897 (10.0)	–	–

(Continued)

Table 1 (Continued)

	Frequency of characteristics		Variable completeness, %	
	Cardiac CT cohort overall (n=21,623)	Coronary CTA cohort ^a (n=18,899)	Cardiac CT cohort overall	Coronary CTA cohort ^a
Nonevaluable	923 (4.3)	828 (4.4)	–	–
Unknown	2,410 (11.1)	1,870 (9.9)	–	–
CACS (Agatston) result ^b	0 (0–95)	0 (0–90)	87.8	87.2
Coronary CTA result	–	–	95.4	96.2
Normal	13,901 (64.3)	12,550 (66.4)	–	–
Nonobstructive CAD	2,573 (11.9)	2,213 (11.7)	–	–
1-vessel obstructive CAD	2,176 (10.1)	2,012 (10.6)	–	–
2-vessel obstructive CAD	739 (3.4)	673 (3.6)	–	–
3-vessel/LM obstructive CAD	462 (2.1)	312 (1.7)	–	–
Only CACS performed	778 (3.6)	427 (2.3)	–	–
Unknown	994 (4.6)	712 (3.7)	–	–
Consequence of cardiac CT results	–	–	86.2	86.7
Nothing further	9,966 (46.1)	8,673 (45.9)	–	–
Medical treatment	3,820 (17.7)	3,313 (17.5)	–	–
Referral to invasive coronary angiography	3,520 (16.3)	3,216 (17.0)	–	–
Referral to myocardial perfusion imaging	679 (3.1)	632 (3.3)	–	–
CACS is too high – referral to further testing	199 (0.9)	157 (0.8)	–	–
Referral to other noninvasive diagnostic testing	461 (2.1)	399 (2.1)	–	–
Unknown	2,977 (13.8)	2,509 (13.4)	–	–

Notes: ^aPatients with angina suspicious symptoms; ^bvalues in median \pm interquartile range; ^cwas not estimated due to differences in registered dose units.

Abbreviations: CACS, coronary artery calcium scoring; CAD, coronary artery disease; CT, computed tomography; CTA, computed tomography angiography; LM, left main; PCI, percutaneous coronary intervention.

registered with a cardiac CT in the DNPR and who also were registered in the WDHR-CCTR was overall 72% (20,560/28,566) (Figure 3). Thus, 8,006 cardiac CTs were registered in the DNPR, but not in the WDHR-CCTR. The completeness of cardiac CT registration in the WDHR-CCTR

improved, however, from 17% (378/2,233) in 2008 to 82% (7,271/8,837) in 2012 (Figure 3).

The medical record review of the cardiac CTs registered in the DNPR but not in the WDHR-CCTR revealed that 79% (192/244) of the procedures at Aarhus University Hospital




Figure 2 Number of patients having cardiac computed tomography (CT) registered in the Western Denmark Cardiac Computed Tomography Registry (WDHR-CCTR) and the Danish National Patient Registry (DNPR), 2008–2012.




Figure 3 Proportion of patients having a cardiac computed tomography registered in both the Western Denmark Cardiac Computed Tomography Registry (WDHR-CCTR) and the Danish National Patient Registry (DNPR) during 2008–2012.

and 66% (25/38) at Lillebaelt Hospital-Vejle were performed on indications other than for coronary assessment in symptomatic patients (overall 77%). These indications included coronary CTA in asymptomatic patients (heart failure and preoperative evaluation) and evaluation of cardiac anatomy (congenital heart disease, planning of electrophysiological procedures, or trans-catheter aortic valve replacement). Among the later procedures, approximately two-thirds were performed as part of research projects. Assuming the proportion of missing coronary CTA procedures in our sample (23%) could be extrapolated to all 8,006 missing cardiac CTs in the WDHR-CCTR, the completeness of coronary CTA procedures reported to the WDHR-CCTR registry was approximately 91% ($18,899/(18,899 + 1,841)$). A secondary finding from the medical record review was that all the cardiac CTs registered in the DNPR had actually been performed (282/282), yielding a PPV for cardiac CT in the DNPR of 100% (95% CI: 98.7%–100%).

The completeness of individual variables is shown in Table 1. The completeness ranged from 60% for the registration of left ventricular ejection fraction to 100% for CPR number, sex, age, and date of procedure. The completeness was $\geq 85\%$ in more than two-thirds (71%) of all variables.

Validity

All 250 medical records requested from the five selected cardiac centers were available for review and thus were included in the validation. The variables and reference standards used for the validation are provided in Table S2. Patients in the random sample had a higher prevalence of diabetes mellitus when compared with the total coronary

CTA cohort. Otherwise, the patient and CT procedural characteristics were similar to the overall coronary CTA cohort, indicating that the random sample was representative (Table S3). Among the 250 patients, 61% (152/250) had complete data without missing values of any variables. A total of 7% (349/4,750) of variable values were missing in the WDHR-CCTR, of which 45% (157/349) also were missing in the medical records.

The PPV ranged from 89% (95% CI: 81%–94%) for family history of premature CAD to 100% for previous cardiac surgery (95% CI: 21%–100%) and coronary CTA indication (95% CI: 98.5%–100%) (Table 2).

The NPV (when calculation was possible) ranged from 97% (95% CI: 91%–100%) for smoking status to 100% for previous myocardial infarction (95% CI: 98%–100%) and previous percutaneous coronary intervention (95% CI: 98%–100%) (Table 2). Stratification by university status showed consistently high overall predictive values (PPV/NPV) for both university (96%/99%) and nonuniversity cardiac centers (97%/99%).

Less than 9% of the data in the random sample could not be verified from the medical records due to missing information. The variables that were least obtainable from the medical records were height (143/228=62.3%) and weight (142/228=62.3%) (Table 2). With the exception of family history of premature CAD (76%, 95% CI: 76%–84%), height (33.5%, 95% CI: 28%–40%), and weight (34%, 95% CI: 28%–41%), the PPVs remained consistently high also in worst-case scenarios where all missing records were included as nonconfirmed cases. The NPVs also remained high in the worst-case scenarios.

Table 2 Validity of 19 selected variables recorded in the Western Denmark Cardiac Computed Tomography Registry during 2008–2012 using medical record review as reference standard

	True positive		False negative		True negative		PPV, % (95% CI)		NPV, % (95% CI)	
	Available data	Worst-case scenario ^a	Available data	Worst-case scenario ^a	Available data	Worst-case scenario ^a	Available data	Worst-case scenario ^a	Available data	Worst-case scenario ^a
Medical history data	361	13	18	1,359	95.3 (92.6–97.0)	89.1 (85.7–91.8)	95.6 (93.1–97.2)	99.1 (98.4–99.5)	96.2 (95.1–97.1)	99.1 (98.4–99.5)
Family history of premature CAD	74	3	9	119	89.2 (80.7–94.2)	76.3 (66.9–83.7)	90.7 (83.3–95.0)	97.5 (93.0–99.2)	95.2 (89.9–97.8)	97.6 (93.2–97.8)
Current lipid-lowering therapy	74	1	2	143	97.4 (90.9–99.3)	93.7 (86.0–97.3)	97.5 (91.2–99.3)	99.3 (96.2–99.9)	96.6 (92.3–98.6)	99.0 (96.3–99.9)
Current medical treatment for hypertension	77	3	4	135	95.1 (88.0–98.1)	91.7 (83.8–95.9)	95.2 (88.4–98.1)	97.8 (93.8–99.3)	94.4 (89.4–97.1)	97.9 (94.0–99.3)
Smoking status (current/former)	122	3	2	89	98.4 (94.3–99.6)	93.9 (88.3–96.9)	98.5 (94.6–99.6)	96.7 (90.9–99.9)	90.8 (83.5–95.1)	96.9 (91.4–99.0)
Diabetes	13	1	1	206	92.9 (66.5–98.7)	92.9 (66.5–98.7)	92.9 (66.5–98.7)	99.5 (97.3–99.9)	96.7 (93.4–98.8)	99.5 (97.4–99.9)
Previous PCI	0	0	0	223	–	–	–	100 (98.3–100)	97.4 (94.4–98.8)	100 (98.4–100)
Previous cardiac surgery	1	2	0	221	100 (20.6–100)	100 (20.6–100)	100 (20.6–100)	99.1 (97.5–99.9)	96.5 (93.3–98.2)	99.1 (96.9–99.8)
Previous myocardial infarction	0	0	0	223	–	–	–	100 (98.3–100)	97.8 (95.0–99.1)	100 (98.3–100)
Procedure data^b	2,185	–	62	–	97.2 (96.5–97.8)	84.8 (83.4–86.2)	97.6 (96.9–98.1)	–	–	–
Coronary CTA indication (anginal/equivalent)	250	–	–	–	100 (98.5–100)	100 (98.5–100)	100 (98.5–100)	–	–	–
Date of procedure	243	–	7	–	93.6 (94.3–96.6)	93.6 (94.3–96.6)	93.6 (94.3–96.6)	–	–	–
Height	73	–	2	–	97.3 (90.8–99.3)	33.5 (27.6–40.0)	99.1 (96.7–99.8)	–	–	–
Weight	75	–	1	–	98.7 (92.9–99.8)	34.4 (28.4–40.9)	99.5 (97.5–99.9)	–	–	–
Type of CT scanner	243	–	6	–	97.6 (94.8–98.9)	95.5 (92.1–97.5)	95.5 (92.1–97.5)	–	–	–
Kilo-electron volt	232	–	11	–	95.5 (92.1–97.5)	94.7 (91.1–96.9)	95.5 (92.1–97.5)	–	–	–
Radiation dose	222	–	5	–	97.8 (95.0–99.1)	96.5 (93.3–98.2)	97.8 (95.0–99.1)	–	–	–
Contrast volume	210	–	6	–	97.2 (94.1–98.7)	86.4 (81.5–90.2)	97.8 (95.0–99.1)	–	–	–
CACS result	220	–	4	–	96.2 (95.5–99.3)	94.8 (91.2–97.0)	97.5 (94.7–98.9)	–	–	–
Coronary CTA result	227	–	9	–	95.0 (92.9–98.0)	94.6 (91.0–96.8)	96.3 (93.0–98.0)	–	–	–
Consequence of the coronary CTA results	190	–	11	–	94.5 (90.5–96.9)	94.5 (90.5–96.9)	94.5 (90.5–96.9)	–	–	–

Notes: The study sample consisted of 250 randomly selected symptomatic patients who underwent coronary CTA at three university and two nonuniversity cardiac centers in Western Denmark. Patients with missing information in the registry or medical records were not included in the analyses. ^aWorst- and best-case scenarios were calculated assuming that none or all of the missing values in the medical files confirmed the registry data; ^bthe procedure data variables comprise continuous variables and categorical variables with more than two levels. For these variables, PPV was calculated from the proportion of exact matches on the corresponding scale (yes/no).

Abbreviations: CACS, coronary artery calcium scoring; CAD, coronary artery disease; CI, confidence interval; CT, computed tomography; CTA, computed tomography angiography; NPV, negative predictive value; PCI, percutaneous coronary intervention; PPV, positive predictive value.

Stratification by calendar year (before 2010 vs 2010 and later) showed that the PPV improved for diabetes from 83% (95% CI: 35%–97%) to 100% (95% CI: 69%–100%), for date of procedure from 73% (95% CI: 64%–80%) to 96% (95% CI: 92%–98.5%), and for consequence of coronary CTA from 92% (95% CI: 84%–96%) to 97% (95% CI: 92%–99%). In contrast, the PPV decreased for current medical treatment for hypertension from 97% (95% CI: 86%–99.5%) to 93% (95% CI: 82%–98%) and for kilo-electron volt from 98% (95% CI: 94%–99%) to 93% (95% CI: 88%–96.5%). Over the same period, the NPV for family history of premature CAD increased from 94% (95% CI: 86%–98%) to 100% (95% CI: 94%–100%) and for current medical treatment for hypertension from 96.5% (95% CI: 88%–100%) to 98.8% (95% CI: 93%–100%). For the other variables, the predictive values remained fairly constant over time.

Discussion

In this first review and validation of the WDHR-CCTR, we demonstrated that the registry as of 2012 includes data on approximately 22,000 patients undergoing cardiac CT. Among these, nearly 19,000 were for coronary assessment in patients with symptoms suggestive of CAD. Importantly, the yearly number of procedures and the completeness of data registration have increased consistently since the establishment of the registry in 2008. Although the overall completeness of cardiac CT procedures in the WDHR-CCTR compared with the DNPR was 72%, additional subanalysis indicated that the completeness of coronary CTAs was more than 90%. Although the completeness of individual variables differed substantially, it was high for the vast majority of variables. Finally, we found that the PPV and NPV of most patient and CT procedural variables were high, even when including missing patient files in the analyses as falsely registered in the WDHR-CCTR.

Our finding of an increasing use of coronary CTA since 2008 is in accordance with other reports,¹² and we expect this increase to continue in years to come. The lower completeness of cardiac CT procedures in the WDHR-CCTR when compared with the DNPR may be explained by the fact that cardiac CT examinations performed as a part of research projects and for planning of noncoronary interventions were not registered until recently in the WDHR-CCTR. In addition, cardiac CT procedures performed in children <16 years are not reported to the WDHR-CCTR.

To our knowledge, the only other large population-based cardiac CT registry is the Coronary CT Angiography Evaluation for Clinical Outcomes: An International

Multicenter registry (CONFIRM).¹³ The CONFIRM Registry prospectively enrolls patients and procedural and outcome data from six countries in the world (USA, Germany, Canada, South Korea, Switzerland, and Italy).¹³ In 2012, the registry comprised more than 27,000 patients undergoing ≥ 64 -slice cardiac CT at 12 centers between 2003 and 2009.¹⁴ The CONFIRM registry is thus a valuable source to study the prognostic value of cardiac CT.^{14–16} However, complete follow-up of patients registered in CONFIRM is not possible,^{14,15} and outcome data are often collected retrospectively by, eg, telephone contact or email questionnaires.^{13–16} In comparison, the WDHR-CCTR is a semi-national, multicenter registry with detailed information on indication, patient characteristics, procedural data, CT outcome, and clinical consequence. It is based on data from both university and nonuniversity cardiac centers, making it representative for contemporary real-world cardiac CT practice. Individual-level data linkage (using the CPR number) to the Danish Civil Registration System enables complete patient follow-up with accurate accounting for censoring due to emigration or death.⁷ Moreover, linkage to other registries provides valuable information on comorbidity, socioeconomic data, and subsequent diagnostic and therapeutic resource utilization (including medication and interventions), clinical events, and mortality.^{4,6,8,17–19}

Although our study overall supports the use of the WDHR-CCTR for research, there are also limitations to be considered. During the study period, the database has continuously been optimized, ie, selected variables have changed from being optional to obligate for reporting to the WDHR-CCTR. These changes in reporting standards may in part explain some of the missing values over time, which reduced the overall completeness of some variables. We also found that although the completeness for radiation dose was greater than 90%, the use and recording of different dose units to the registry may hamper the use of this variable for research.

The predictive values for the main patient and procedure variables examined in this study were found to be high. The variables systolic and diastolic blood pressures could be of interest when studying coronary CTA outcomes. Nonetheless, these variables were not validated because blood pressure most often is measured in relation to the cardiac CT examination and registered directly into WDHR-CCTR without further documentation in the medical records.

Conclusion

WDHR-CCTR provides ongoing prospective registration of all cardiac CTs performed in Western Denmark since 2008. Overall, the registry data have a high degree of completeness

and validity, making it a valuable tool for clinical epidemiological research.

Acknowledgments

The study was supported by Aarhus University. The funding source had no role in the design, conduct, analysis, or reporting of the study.

Author contributions

MS conceived the study idea. LHN and MS designed the study and reviewed the literature. LP collected the data. MS directed the analyses, which was carried out by LHN. LHN performed medical record review, organized the writing, and wrote the initial draft. All authors participated in the discussion and interpretation of the results, critically revised the manuscript for intellectual content, and approved the final version before submission. LHN had full access to the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bonow RO, Mann DL, Zipes DP, Libby P. *Braunwald's Heart Disease: a Textbook of Cardiovascular Medicine*. 9th ed. Philadelphia, PA: Saunders; 2012.
- Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J*. 2013;34(38):2949–3003.
- Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2012;60(24):e44–e164.
- Schmidt M, Maeng M, Jakobsen CJ, et al. Existing data sources for clinical epidemiology: The Western Denmark Heart Registry. *Clin Epidemiol*. 2010;2:137–144.
- The board of Western Denmark Heart Registry. Western Denmark Heart Registry: Annual Report 2008 (Data from 2007 and 2008) 2009.
- Johannesdottir SA, Horvath-Puhó E, Ehrenstein V, Schmidt M, Pedersen L, Sørensen HT. Existing data sources for clinical epidemiology: The Danish National Database of Reimbursed Prescriptions. *Clin Epidemiol*. 2012;4:303–313.
- Schmidt M, Pedersen L, Sørensen HT. The Danish Civil Registration System as a tool in epidemiology. *Eur J Epidemiol*. 2014;29(8):541–549.
- Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scand J Public Health*. 2011;39(Suppl 7):30–33.
- The Nordic Medico-Statistical Committee (NOMESCO) Classification of Surgical Procedures. 2012. Available from: http://nowbase.org/~media/Projekt%20sites/Nowbase/Publikationer/NCSP/NCSP%201_16.ashx.
- Sørensen HT, Sabroe S, Olsen J. A framework for evaluation of secondary data sources for epidemiological research. *Int J Epidemiol*. 1996;25(2):1075–1082.
- Brown LD, Cai TT, DasGupta A. Interval estimation for a binomial proportion. *Stat Sci*. 2001;16(2):101–117.
- Shaw LJ, Marwick TH, Zoghbi WA, et al. Why all the focus on cardiac imaging? *JACC Cardiovasc Imaging*. 2010;3(7):789–794.
- Min JK, Dunning A, Lin FY, et al. Rationale and design of the CONFIRM (COronary CT Angiography Evaluation For Clinical Outcomes: An International Multicenter) Registry. *J Cardiovasc Comput Tomogr*. 2011;5(2):84–92.
- Cho I, Chang HJ, Sung JM, et al. Coronary computed tomographic angiography and risk of all-cause mortality and nonfatal myocardial infarction in subjects without chest pain syndrome from the CONFIRM Registry (coronary CT angiography evaluation for clinical outcomes: an international multicenter registry). *Circulation*. 2012;126(3):304–313.
- Min JK, Dunning A, Lin FY, et al. Age- and sex-related differences in all-cause mortality risk based on coronary computed tomography angiography findings results from the International Multicenter CONFIRM (Coronary CT Angiography Evaluation for Clinical Outcomes: An International Multicenter Registry) of 23,854 patients without known coronary artery disease. *J Am Coll Cardiol*. 2011;58:849–860.
- Chow BJ, Small G, Yam Y, et al. Incremental prognostic value of cardiac computed tomography in coronary artery disease using CONFIRM: COroNary computed tomography angiography evaluation for clinical outcomes: an International Multicenter registry. *Circ Cardiovasc Imaging*. 2011;4(5):463–472.
- Helweg-Larsen K. The Danish Register of Causes of Death. *Scand J Public Health*. 2011;39(Suppl 7):26–29.
- Statistics Denmark. The IDA database. (In Danish: Den Integrerede Database for Arbejdsmarkedsforskning). 2012. Available from: <http://www.dst.dk/da/Statistik/dokumentation/kvalitetsdeklarationer/ida-databasen.aspx>. Accessed September 1, 2014.
- Gjerstorff ML. The Danish Cancer Registry. *Scand J Public Health*. 2011;39(Suppl 7):42–45.

Supplementary materials

Table SI Variables included in the registration forms for the Western Denmark Cardiac Computed Tomography Registry

Registration forms		
Referral form	Medical history form	Procedure form
CPR number	Family history of premature CAD	Type of CT scanner
Sex	Current lipid-lowering therapy	Cardiac CT indication (angina/equivalent, heart failure, visualization of valves and/or aorta disease, visualization of stents, evaluation of cardiac structures, or others)
County	Current medical treatment for hypertension	Angina type (if any): typical angina, atypical angina, unspecific chest pain, or dyspnea
Area code	Smoking status	Priority (acute, subacute, or elective)
Referral date	Diabetes	Date of procedure
	Previous PCI	Kilo-electron volt
	Previous cardiac surgery	Radiation dose (DLP)
	Previous myocardial infarction	Iodine dose (mg)
	Height (cm)	Contrast volume (mL)
	Weight (kg)	Contrast manufacturer
	Creatinine level ($\mu\text{mol/L}$)	Beta-blocker (if any): name and administration
	Systolic and diastolic blood pressures (mmHg)	Heart rhythm (sinus rhythm, atrial fibrillation, supraventricular/ventricular premature beats/others)
	Left ventricular ejection fraction (%)	Heart rate (bpm)
	Cardiac devices	Ventriculography ejection fraction (%)
	Current medical treatments	Cardiac CT quality (good, moderate, or nonevaluable)
		Complications (descriptive)
		Vessel dominance (balanced, right, or left)
		CACS (Agatston score) result
		Consequence of CACS results (termination or proceeding to contrast-enhanced cardiac CT)
		Coronary CTA result (only CACS was performed, none CAD [0% luminal stenosis], nonobstructive CAD [1%–49% luminal stenosis], obstructive CAD [$\geq 50\%$ luminal stenosis] categorized as: 1-vessel obstructive CAD, 2-vessel obstructive CAD, 3-vessel/left main obstructive CAD)
		Consequence of coronary CTA results (none, medical treatment, referral to invasive coronary angiography, referral to myocardial perfusion imaging, referral to other noninvasive diagnostic testing, or other)
		Extracardiac findings (descriptive)

Abbreviations: CACS, coronary artery calcium scoring; CAD, coronary artery disease; CT, computed tomography; CTA, computed tomography angiography; CPR, Civil Personal Register; DLP, dose length product; PCI, percutaneous coronary intervention.

Table S2 Variable values and reference standards used to validate data from the Western Denmark Coronary Computed Tomography Angiography Registry

Variables	Values (reference standards for validation)
Family history of premature CAD	Yes/no/unknown ^a
Current lipid-lowering therapy	Yes/no/unknown ^a
Current medical treatment for hypertension	Yes/no/unknown ^a
Smoking status	Current/never/former/unknown ^a
Diabetes	None/recently elevated blood glucose/insulin (\pm oral antidiabetic treatment)/oral antidiabetic treatment (without insulin)/nonpharmacological treatment/unknown ^a
Previous PCI	Yes/no/unknown ^a
Previous cardiac surgery	Yes/no/unknown ^a
Previous myocardial infarction	Yes/no/unknown ^a
Height	Exact height in cm ^a
Weight	Exact weight in kg ^a
Type of CT scanner	Siemens 64-slice/Siemens 64-slice dual-source/Phillips 64-slice/Phillips 256-slice/GE 64-slice/Toshiba 64-slice/Toshiba 320-slice/Siemens Flash GE 750 HD/other ^b
Date of procedure	Day-month-year (\pm 1 week) ^b
Kilo-electron volt	Exact kilo-electron volt ^b
Radiation dose	Exact radiation dose in DLP/millisievert ^b
Contrast volume	Exact volume in mL ^b
CACS result	Exact score in Agatston ^b
Coronary CTA result	Only CACS was performed/none CAD (0% luminal stenosis)/nonobstructive CAD (1%–49% luminal stenosis)/obstructive CAD (\geq 50% luminal stenosis) categorized as 1-vessel obstructive CAD/2-vessel obstructive CAD/3-vessel or left main obstructive CAD ^a
Consequence of coronary CTA results	Nothing further/medical treatment/referral to invasive coronary angiography/referral to myocardial perfusion imaging/referral to other noninvasive diagnostic testing/other

Notes: ^aReference standard: medical records; ^breference standard: medical records including imaging reports from working stations.

Abbreviations: CACS, coronary artery calcium scoring; CAD, coronary artery disease; CT, computed tomography; CTA, computed tomography angiography; DLP, dose length product; PCI, percutaneous coronary intervention.

Table S3 Characteristics of symptomatic patients in the coronary CTA cohort and patients in the randomly selected sample from the Western Denmark Cardiac Computed Tomography Registry, 2008–2012

	Coronary CTA cohort overall (n=18,899)	Random sample ^a (n=250)
Male sex	10,169 (53.8)	107 (42.8)
Age, years ^b	58 (50–66)	56.5 (48–63)
Family history of premature CAD	7,353 (38.9)	97 (38.8)
Current lipid-lowering therapy	5,883 (31.1)	79 (31.6)
Current medical treatment for hypertension	6,532 (34.6)	84 (33.6)
Smoking status (current or former)	9,504 (50.3)	131 (52.4)
Diabetes	1,035 (5.5)	33 (13.2)
Previous PCI	166 (0.9)	0
Previous cardiac surgery	156 (0.8)	1 (0.4)
Previous myocardial infarction	193 (1.0)	0
Height (cm) ^b	171 (165–178)	171 (158–184)
Weight (kg) ^b	78 (67–88)	78 (56–100)
Creatinine level, $\mu\text{mol/L}^b$	73 (63–83)	72 (55–89)
Systolic blood pressure, mmHg ^b	140 (126–151)	138 (112–164)
Diastolic blood pressure, mmHg ^b	81 (75–90)	80 (65–95)
Left ventricular ejection fraction ^b	60 (60–60)	60 (60–60)
Angina type (if any)		
Typical angina	1,764 (9.4)	22 (8.8)
Atypical angina	5,844 (30.9)	78 (31.2)
Unspecific chest pain	4,760 (25.2)	58 (23.2)
Dyspnea	916 (4.8)	8 (3.2)
Unknown	5,615 (29.7)	84 (33.6)
Priority		
Acute	78 (0.4)	2 (0.8)
Subacute	755 (4.0)	2 (0.8)
Elective	17,209 (91.1)	230 (92.0)
Unknown	857 (4.5)	16 (6.4)
Kilo-electron volt ^b	100 (100–120)	120 (100–120)
Radiation dose	– ^c	– ^c
Iodine concentration, mg/mL ^b	350 (350–370)	370 (350–370)
Contrast volume, mL ^b	80 (70–90)	77.5 (70–90)
Preprocedural beta-blocker	11,991 (63.4)	164 (65.6)
Heart rhythm		
Sinus rhythm	17,794 (94.2)	229 (91.6)
Atrial fibrillation	108 (0.6)	2 (0.8)

(Continued)

Table S3 (Continued)

	Coronary CTA cohort overall (n=18,899)	Random sample ^a (n=250)
Other	85 (0.4)	2 (0.8)
Unknown	912 (4.8)	17 (6.8)
Heart rate (bpm) ^b	59 (54–64)	60 (55–65)
Cardiac CT quality		
Good	14,304 (75.7)	184 (73.6)
Moderate	1,897 (10.0)	20 (8.0)
Nonevaluable	828 (4.4)	18 (7.2)
Unknown	1,870 (9.9)	28 (11.2)
CACS (Agatston) result ^b	0 (0–90)	0 (0–75)
Coronary CTA result		
Normal	12,550 (66.4)	164 (65.6)
Nonobstructive CAD	2,213 (11.7)	39 (15.6)
1-vessel obstructive CAD	2,012 (10.6)	19 (7.6)
2-vessel obstructive CAD	673 (3.6)	7 (2.8)
3-vessel/LM obstructive CAD	312 (1.7)	6 (2.4)
Only CACS performed	427 (2.3)	6 (2.4)
Unknown	712 (3.7)	9 (3.6)
Consequence of coronary CTA results		
Nothing further	8,673 (45.9)	96 (38.4)
Medical treatment	3,313 (17.5)	39 (15.6)
Referral to invasive coronary angiography	3,216 (17.0)	42 (16.8)
Referral to myocardial perfusion imaging	632 (3.3)	10 (4.0)
CACS is too high – referral to further testing	157 (0.8)	1 (0.4)
Referral to other noninvasive diagnostic testing	399 (2.1)	9 (3.6)
Unknown	2,509 (13.4)	53 (21.2)

Notes: ^a250 randomly selected symptomatic patients who underwent coronary CTA at three university cardiac centers (Aarhus, Aalborg, and Odense) and two nonuniversity cardiac centers (Lillebaelt Hospital-Vejle and Hospital of Southwestern Denmark-Esbjerg); ^bvalues in median \pm interquartile range; ^cwas not estimated due to differences in registered dose units.

Abbreviations: CACS, coronary artery calcium scoring; CAD, coronary artery disease; CT, computed tomography; CTA, computed tomography angiography; LM, left main; PCI, percutaneous coronary intervention.

Clinical Epidemiology

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

Submit your manuscript here: <http://www.dovepress.com/clinical-epidemiology-journal>

systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Dovepress