



AALBORG UNIVERSITY
DENMARK

Aalborg Universitet

Exercise increases pressure pain tolerance but not pressure and heat pain thresholds in healthy young men

Vaegter, H. B.; Bement, M. Hoeger; Madsen, A. B.; Fridriksson, J.; Dasa, M.; Graven-Nielsen, Thomas

Published in:
European Journal of Pain

DOI (link to publication from Publisher):
[10.1002/ejp.901](https://doi.org/10.1002/ejp.901)

Publication date:
2017

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Vaegter, H. B., Bement, M. H., Madsen, A. B., Fridriksson, J., Dasa, M., & Graven-Nielsen, T. (2017). Exercise increases pressure pain tolerance but not pressure and heat pain thresholds in healthy young men. *European Journal of Pain*, 21(1), 73-81. <https://doi.org/10.1002/ejp.901>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

ABSTRACT

Background: Exercise causes an acute decrease in the pain sensitivity known as exercise-induced hypoalgesia (EIH) but the specificity to certain pain modalities is unknown. This study aimed to compare the effect of isometric exercise on the heat and pressure pain sensitivity.

Methods: On 3 different days, twenty healthy young men performed 2 submaximal isometric knee extensions (30% maximal voluntary contraction in 3 min) and a control condition (quiet rest). Before and immediately after exercise and rest, the sensitivity to heat pain and pressure pain were assessed in randomized and counterbalanced order. Cuff pressure pain threshold (cPPT) and pain tolerance (cPTT) were assessed on the ipsilateral lower leg by computer-controlled cuff algometry. Heat pain threshold (HPT) was recorded on the ipsilateral foot by a computer-controlled thermal stimulator.

Results: cPTT was significantly increased after exercise compared with baseline and rest ($P < 0.05$). Compared with rest, cPPT and HPT were not significantly increased by exercise. No significant correlation between exercise-induced changes in HPT and cPPT was found. Test-retest reliability before and after the rest condition was better for cPPT and CPTT (intraclass-correlation > 0.77) compared with HPT (intraclass-correlation = 0.54).

Conclusions: The results indicate that hypoalgesia after submaximal isometric exercise is primarily affecting tolerance of pressure pain compared with the pain threshold. These data contributes to the understanding of how isometric exercise influences pain perception, which is necessary to optimize the clinical utility of exercise in management of chronic pain.

Significance: The effect of exercise on pain tolerance may be relevant for patients in chronic musculoskeletal pain as a pain coping strategy.

Keywords: Exercise-induced hypoalgesia, cuff algometry, pressure pain sensitivity, pressure pain tolerance, heat pain thresholds

EXERCISE INCREASES PRESSURE PAIN TOLERANCE BUT NOT PRESSURE AND HEAT PAIN THRESHOLDS IN HEALTHY YOUNG MEN

H.B. Vaegter^{ab}, M. Hoeger Bement^c, A.B. Madsen^d, J. Fridriksson^d, M. Dasa^d, T. Graven-Nielsen^e

^a Pain Research Group, Pain Center South, Odense University Hospital, Odense, Denmark

^b Institute of Clinical Research, Faculty of Health Sciences, University of Southern Denmark

^c Department of Physical Therapy, College of Health Sciences, Marquette University, Milwaukee, Wisconsin, USA

^d School of Physiotherapy, University College Lillebaelt, Odense, Denmark

^e Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark

Original paper for European Journal of Pain

Running title: EIH, heat and pressure pain sensitivity

Corresponding author:

Henrik Bjarke Vægter Ph.D.

Pain Research Group, Pain Center South, University Hospital Odense

Department of Clinical Research, Faculty of Health Sciences, University of Southern Denmark

Heden 9, Indgang 201

DK – 5000 Odense C

Tel.: +45 65413869; fax: +45 65415064.

E-mail address: henrik.bjarke.vaegter@rsyd.dk

Funding: HBV was supported by grants from the philanthropic foundation TrygFonden (7-11-0990) and The Danish Rheumatism Association (R95-A1871). Funders were not involved in the design or conduct of this study. TGN is a part of Center for Neuroplasticity and Pain (CNAP) supported by the Danish National Research Foundation (DNRF121).

Conflict of interest: There are no actual or potential conflicts of interest for any of the authors. Nocitech is a company partly owned by Aalborg University.

What's already known about this topic?

- Exercise causes an acute decrease in the pain sensitivity but the specificity to certain pain modalities is unknown.

What does this study add?

- The results indicate that hypoalgesia after submaximal isometric exercise is primarily affecting tolerance of pressure pain compared with the heat and pressure pain threshold.
- These data contributes to the understanding of how isometric exercise influences pain perception, which is necessary to optimize the clinical utility of exercise in management of chronic pain.

ABSTRACT

Background: Exercise causes an acute decrease in the pain sensitivity known as exercise-induced hypoalgesia (EIH) but the specificity to certain pain modalities is unknown. This study aimed to compare the effect of isometric exercise on the heat and pressure pain sensitivity.

Methods: On 3 different days, twenty healthy young men performed 2 submaximal isometric knee extensions (30% maximal voluntary contraction in 3 min) and a control condition (quiet rest). Before and immediately after exercise and rest, the sensitivity to heat pain and pressure pain were assessed in randomized and counterbalanced order. Cuff pressure pain threshold (cPPT) and pain tolerance (cPTT) were assessed on the ipsilateral lower leg by computer-controlled cuff algometry. Heat pain threshold (HPT) was recorded on the ipsilateral foot by a computer-controlled thermal stimulator.

Results: cPTT was significantly increased after exercise compared with baseline and rest ($P < 0.05$). Compared with rest, cPPT and HPT were not significantly increased by exercise. No significant correlation between exercise-induced changes in HPT and cPPT was found. Test-retest reliability before and after the rest condition was better for cPPT and CPTT (intraclass-correlation > 0.77) compared with HPT (intraclass-correlation = 0.54).

Conclusions: The results indicate that hypoalgesia after submaximal isometric exercise is primarily affecting tolerance of pressure pain compared with the pain threshold. These data contributes to the understanding of how isometric exercise influences pain perception, which is necessary to optimize the clinical utility of exercise in management of chronic pain.

Significance: The effect of isometric exercise on pain tolerance may be relevant for patients in chronic musculoskeletal pain as a pain coping strategy.

Keywords: Exercise-induced hypoalgesia, cuff algometry, pressure pain sensitivity, pressure pain tolerance, heat pain thresholds

INTRODUCTION

Efficiency of the endogenous pain inhibitory pathways can be assessed by paradigms of exercise-induced hypoalgesia (EIH) (Lannersten and Kosek, 2010) with recordings of pain sensitivity before and after an exercise condition. Isometric muscle exercises have been linked to modulation of pain sensitivity in healthy subjects (Hoeger Bement et al., 2008, Hoeger Bement et al., 2014, Vaegter et al., 2014) and in patients with chronic pain (Vaegter et al., 2016, Hoeger Bement et al., 2011). Moreover, it has been hypothesized that impaired EIH may be indicative of a dysfunction of the pain inhibitory systems (Lannersten and Kosek, 2010). In healthy subjects, EIH after isometric exercises are often demonstrated as an increase in pressure pain thresholds (Koltyn et al., 2001, Kosek and Lundberg, 2003, Koltyn and Umeda, 2007, Hoeger Bement et al., 2008, Hoeger Bement et al., 2009, Umeda et al., 2010, Hoeger Bement et al., 2014, Lemley et al., 2014, Lemley et al., 2015, Vaegter et al., 2014) or a decrease in heat pain ratings (Naugle et al., 2014, Misra et al., 2014, Koltyn et al., 2014).

Few studies on EIH have assessed both heat pain and pressure pain sensitivity modalities (Kodesh and Weissman-Fogel, 2014, Cook et al., 2010, Naugle et al., 2014), and no studies have directly compared these modalities at the same time. Furthermore, heat pain thresholds and pain tolerance are rarely assessed, and no studies have compared the effect of isometric exercise on different aspects of pain sensitivity. Such a comparison will significantly contribute to the understanding of how physical activity influences pain perception, which is necessary to optimize the clinical utility of physical activity as a method of pain management. **The potential effect of exercise on pain tolerance could be relevant for patients in chronic pain as a pain coping strategy.** In addition, it has been recommended to include a range of stimulus intensities in the assessment of experimental pain sensitivity to reveal potential effects that are manifest with more painful stimuli (Greenspan et al., 2007). Moreover, different nociceptive pathways in skin and muscles are evoked by varying stimulation modalities, and responses to different experimental pain modalities should be assessed in combination to improve understanding of the pain experience (Neziri et al., 2011).

Previously, it has been demonstrated that input to dorsal horn neurons from muscle nociceptors is subject to stronger descending inhibition compared with input from cutaneous nociceptors (Yu and Mense, 1990), and it may be hypothesized that the magnitude of EIH would be greater for assessment in the deeper musculoskeletal structures compared with assessment on the skin.

The primary aim of this study was to compare heat pain threshold, pressure pain threshold and pressure pain tolerance before and after isometric exercise and quiet rest in healthy young men. It was hypothesized that 1) isometric exercise would increase pressure pain thresholds as well as pressure pain tolerance compared with quiet rest, 2) the hypoalgesic response to exercise would be greater in the deeper tissues compared with the skin, and 3) the exercise-induced changes in heat and pressure pain thresholds would not be correlated.

METHODS

Subjects

In this study 20 healthy young men (age: 24.4 ± 2.0 years; body-mass-index: 24.8 ± 2.1 kg/m²; 18 with right side dominance) were included. **Due to potential gender related differences in pain modulation capacity (Popescu et al., 2010) and EIH (Koltyn et al., 2001) only young men between 18 and 30 years of age were included in the study.** Subjects were recruited by advertisement at the local university and the local physiotherapy school. All subjects were naive to experimental pain testing. None of the included subjects suffered from neurological, psychological, cardiovascular diseases, had any pain or used any pain medication during the weeks prior to participation. All subjects were asked to refrain from physical exercises, coffee and nicotine on the days of participation. The study was conducted in accordance with the Declaration of Helsinki, approved by the local ethical committee (S-20140203) and all subjects provided written informed consent.

Procedure

Each subject was assessed at the same time of day on three different days separated by 1 week (Fig. 1). In the first session subjects were thoroughly introduced to the procedures for the pain sensitivity assessments by drawings as well as verbal instructions. All pain sensitivity assessments were performed with the subject seated on a plinth without foot support. In the beginning of each of the three sessions all subjects completed 1-2 practice trial with assessment of heat and pressure pain sensitivity on the leg not used for assessment of EIH to ensure that all participants understood the procedures. Each session lasted approximately 30 min. All assessments were performed by a male experimenter.

Session 1: Before and immediately after a 15 min control condition (quiet rest) the sensitivity to heat pain and pressure pain were assessed. The sequence between assessment of heat and pressure pain sensitivity were randomized and counterbalanced. Subjects were instructed to

relax comfortably in a supine position on a plinth for 15 min in a temperate and undisturbed room with the light subdued. Following the control condition, the maximal voluntary contraction (MVC) for an isometric knee extension with the dominant leg was determined. Subjects were seated on a table with full support of the whole thigh. The dominant leg was strapped above the ankle to the force transducer (Commander Muscle Tester, Powertrack II, JTECH Medical, USA). The MVC during isometric knee extension was determined in a position of ninety degrees of knee flexion. Three maximal contractions separated by one min between contractions were performed and the average MVC was used to determine the sub-maximal value.

Session 2 and 3: Before and immediately after a 15 min active condition (initiated with 12 min rest followed by a 3 min submaximal isometric knee extension at 30% of MVC with the dominant leg) the sensitivity to either heat pain or pressure pain was assessed on the exercised leg. The intensity and duration of contraction was chosen based on previous studies in healthy subjects, which have shown robust EIH at this intensity (Vaegter et al., 2014, Kosek and Ekholm, 1995). During the sustained sub-maximal isometric contractions, subjects were required to match the target force as displayed on the monitor of the force transducer. The subjects were verbally encouraged to sustain the force throughout the 3 min. The sequence between session 2 and 3 were randomized and counterbalanced. Rating of perceived exertion (RPE; Borg Scale: 6-20) and rating of perceived pain (0-10 numerical rating scale, NRS) during isometric knee extension were assessed just before completion of the knee extension.

Assessment of heat pain sensitivity

Heat pain threshold (HPT) was assessed by a computer controlled surface thermode (MSA Thermal Stimulator, SENSELab, Somedic Sales AB, Hörby, Sweden) covering a 25x50 mm skin area on the dorsum of the dominant foot. The method of limit was used where the temperature started at baseline of 32°C and increased by 1.0°C/s with a maximum of 50°C. As soon as the heat sensation was defined as first sensation of pain, the subjects were instructed to press a handheld switch. The peak temperature was stored and the thermode instantly decreased its temperature (3.0°C/s) to the baseline of 32°C. The thermal stimulus was repeated three times and the average heat pain thresholds were calculated.

Assessment of pressure pain sensitivity

Pressure pain thresholds (cPPT) and pressure pain tolerance (cPTT) were assessed by computer-controlled cuff algometry (Nocitech, Denmark and Aalborg University, Denmark). A 13-cm wide silicone tourniquet cuff (VBM, Sulz, Germany) with an equal-sized proximal and distal chamber was wrapped around the dominant lower leg. The cuff was mounted with a 5 cm distance between its upper rim and the tibial tuberosity. The cuff pressure was increased with a rate of 1 kPa/s in both chambers and the maximal pressure limit was 100 kPa. The maximal pressure limit was based on the maximum capacity of the system. Air was supplied from a 200 liters external air tank to avoid loud noises from the cuff system during assessment. The participants used an electronic visual analogue scale (VAS) to rate their pressure-induced pain intensity and a button to release the pressure. The electronic VAS was sampled at 10 Hz. Zero and 10 cm extremes on the VAS were defined as “no pain” and as “maximal pain”, respectively. The participants were instructed to rate the pain intensity continuously on the electronic VAS from when the pressure was defined as first sensation of pain and to press the pressure release button when the pain was intolerable. The pressure value, when the subject rated the sensation of pain as 1 cm on the VAS was defined as the pain threshold (cPPT) and when the subject terminated the pressure inflation was defined as the tolerance (cPTT). In case the maximum pressure stimulation was achieved before reaching the cPTT, 100 kPa was used for further analysis as a conservative estimate of the cPTT.

Statistics

The distribution of HPT, cPPT, cPTT, pain intensity scores (NRS), and the ratings of perceived exertion (RPE) during isometric contractions did not deviate significantly from normality (Shapiro-Wilks test: $P > 0.06$). The effect of sequence between assessment of heat pain and pressure pain sensitivity on HPT, cPPT, and cPTT prior to rest was analyzed with paired t-tests. The effects of exercise and rest on heat pain and pressure pain sensitivity were analyzed with mixed-model repeated-measures analysis of variances (RM-ANOVAs) with time (before and after) as repeated measure and condition (active and control) as group factor. Effect sizes between active and control conditions were determined using partial eta squared. Due to significant difference in HPT before rest and exercise conditions, the percentage change in heat and pressure pain sensitivity before and after isometric exercise and rest was calculated. The distribution of percentage change after isometric exercise deviated from normality (Shapiro-Wilks test: $P < 0.001$) and the percentage differences were compared with non-parametric Wilcoxon Signed Rank test. The Friedman test was used to analyze the percentage change in heat and pressure pain sensitivity after exercise with the

factor *modality* (heat pain threshold, pressure pain threshold, pressure pain tolerance). In case of significant factors or interactions in ANOVAs or Friedman test, Bonferroni corrected post-hoc tests were used for comparisons incorporating correction for the multiple comparisons. Paired t-tests were used to compare the pain intensity scores (NRS) and the ratings of perceived exertion (RPE) during isometric contractions in session 2 and 3. Furthermore, Pearson product-moment correlations were calculated to determine associations between exercise-induced percentage change in cPPT, cPTT, and HPT and between NRS and RPE scores during exercise and exercise-induced percentage change in cPPT, cPTT, and HPT. *P*-values less than 0.05 were considered significant. Finally, intraclass correlations (ICCs) based on a single rating, consistency, 2-way mixed effect model (ICC_{3,1}) and Bland-Altman methods were used for analysis of test-retest reliability of cPPT, cPTT and HPT before and after rest. An ICC above 0.75 was taken as excellent reliability, 0.40–0.75 was fair to good reliability, and less than 0.40 defined poor reliability (Fleiss, 1986). Data were analyzed using SPSS Statistics, version 21 (IBM, Armonk, NY, USA).

RESULTS

Heat pain and pressure pain sensitivity test-retest repeatability

Repeatability between tests of HPT was fair with ICC of 0.54 (Table 1). Results from Bland-Altman demonstrated reasonable agreement for HPT reflected in the 95 % CI of the mean difference, where zero lies within the interval. Repeatability between tests of cPPT and cPTT was excellent with ICCs of 0.86 and 0.77, respectively and results from Bland-Altman analysis demonstrated no systematic bias between assessments.

Isometric contractions

The average MVC was 455.0 ± 86.7 N. The pain intensity and rated perceived exertion reported during the submaximal isometric contractions in the session with assessment of pressure pain sensitivity (NRS: 6.1 ± 1.4 ; RPE: 15.2 ± 1.6) and heat pain sensitivity (NRS: 6.3 ± 1.5 ; RPE: 15.2 ± 1.5) were not significantly different (t-test: $P > 0.49$).

Heat pain sensitivity after quiet rest and isometric contraction

There was no significant effect of assessment sequence on HPT prior to the resting condition (t-test: $P > 0.4$). Baseline HPTs were significantly different during the quiet rest (46.6 ± 2.1 °C) and isometric contraction sessions (45.4 ± 2.9 °C; t-test: $P < 0.023$). The RM-ANOVA of HPT

demonstrated a significant main effect of time ($F(1,38) = 7.09$, $P < 0.011$, $\eta_p^2 = 0.16$). Post-hoc test showed that HPT increased during quiet rest (before: $46.6 \pm 2.1^\circ\text{C}$; after: $46.9 \pm 2.2^\circ\text{C}$) and during isometric contraction (before: $45.4 \pm 2.9^\circ\text{C}$; after: $46.8 \pm 2.0^\circ\text{C}$). The interaction between condition and time in the RM-ANOVA approached significance ($F(1,38) = 2.39$, $P < 0.13$, $\eta_p^2 = 0.06$). In addition, the difference in percentage change in HPT after rest ($0.9 \pm 4.5\%$) and isometric contraction ($3.4 \pm 5.9\%$) approached significance (Wilcoxon: $P = 0.08$).

Pressure pain threshold during quiet rest and isometric contraction

There was no significant effect of assessment sequence on cPPT prior to the resting condition (t-test: $P = 0.94$). Baseline cPPTs during the quiet rest (20.6 ± 8.5 kPa) were not significantly different compared with the isometric contraction sessions (24.4 ± 11.2 kPa; t-test: $P = 0.08$). **Pressure pain threshold (cPPT) increased during quiet rest (before: 20.6 ± 8.5 kPa; after: 21.7 ± 9.0 kPa) and during isometric contraction (before: 24.4 ± 11.2 kPa; after: 26.3 ± 11.7 kPa).** In the RM-ANOVA of cPPT a main effect of time approached significance (Fig. 2A; $F(1,38) = 3.56$, $P = 0.07$, $\eta_p^2 = 0.09$). There was no significant interaction between condition and time ($F(1,38) = 0.19$, $P = 0.67$, $\eta_p^2 = 0.005$). There was no significant difference in percentage change in cPPT after rest ($6.8 \pm 22.9\%$) and isometric contraction ($11.9 \pm 23.7\%$; Wilcoxon: $P > 0.3$).

Pressure pain tolerance during quiet rest and isometric contraction

There was no significant effect of assessment sequence on cPTT prior to the resting condition (t-test: $P = 0.81$). Baseline cPTTs were similar during the quiet rest (63.7 ± 18.4 kPa) and isometric contraction sessions (63.11 ± 18.3 kPa; t-test: $P = 0.84$). **Pressure pain tolerance (cPTT) increased during quiet rest (before: 63.7 ± 18.4 kPa; after: 64.2 ± 18.3 kPa) and during isometric contraction (before: 63.1 ± 18.3 kPa; after: 74.2 ± 18.3 kPa).** The RM-ANOVA of cPTT demonstrated a significant interaction between condition and time (Fig. 2B; $F(1,38) = 10.15$, $P < 0.003$, $\eta_p^2 = 0.21$). Post-hoc test showed that cPTT did not change during quiet rest but increased during isometric contraction. There was a significant difference in percentage change in cPTT after rest ($2.9 \pm 18.1\%$) and isometric contraction ($20.2 \pm 19.1\%$; Wilcoxon: $P < 0.01$).

Comparisons of EIH on HPT, cPPT, and cPTT

There was a statistically significant difference in EIH depending on the noxious stimulus used to assess EIH ($X^2(2) = 6.7$, $P = 0.035$; Fig 3). Post hoc test showed a significant percentage increase in cPTT compared with cPPT and HPT (Wilcoxon: $P < 0.05$).

Associations between exercise-induced changes in heat and pressure pain sensitivity

There was a significant positive correlation between the exercise-induced percentage change in cPPT and the change in cPTT ($r(18) = 0.50$, $P < 0.026$). There was no significant correlation between the exercise-induced percentage change in heat pain sensitivity and the percentage change in pressure pain sensitivity ($r(18) < 0.13$, $P > 0.59$). No significant correlations were found between ratings of pain intensity and perceived exertion during the submaximal isometric contractions and the exercise-induced changes in heat pain and pressure pain sensitivity ($r(18) < 0.42$, $P > 0.07$).

DISCUSSION

This is the first study to compare the effects of a submaximal isometric exercise condition on heat and pressure pain sensitivity in healthy young men. As hypothesized, an increase in pressure pain tolerance was found after exercise compared with baseline and the control condition. In contrast with the hypothesis, no significant effects were found for pressure pain and heat pain thresholds. The results indicate that hypoalgesia after submaximal isometric exercise is primarily affecting tolerance of pain compared with the pain threshold. Furthermore, no significant correlations between exercise-induced changes in heat pain and pressure pain sensitivity were found. Pressure pain sensitivity was not significantly affected by quiet rest and assessments of pressure pain sensitivity were more reliable than assessment of heat pain sensitivity. These findings have clinical implications as the deeper tissues play an important role in many musculoskeletal pain conditions (Graven-Nielsen and Arendt-Nielsen, 2010) where exercise often is an essential part of treatment and rehabilitation (Mannerkorpi and Henriksson, 2007). Furthermore, the effect of exercise on pain tolerance could be relevant for patients in chronic pain.

Exercise-induced hypoalgesia

The current findings are in agreement with a recent study demonstrating an increase in pressure pain tolerance after submaximal isometric exercise (Vaegter et al., 2015) indicating that the hypoalgesia after isometric exercise manifest with more intensely painful stimuli. However, the

results are in contrast to previous studies demonstrating increases in pressure pain thresholds (Kosek and Ekholm, 1995, Koltyn et al., 2001, Kosek and Lundberg, 2003, Koltyn and Umeda, 2007, Hoeger Bement et al., 2008, Hoeger Bement et al., 2009, Umeda et al., 2010, Naugle et al., 2013, Hoeger Bement et al., 2014, Lemley et al., 2014, Koltyn et al., 2014, Vaegter et al., 2014) after submaximal isometric exercise. **In the previous studies demonstrating increase in pressure pain thresholds after isometric exercise pressure pain thresholds are often assessed with manual algometry.** The contrast in findings with manual pressure and cuff algometry may suggest that the spatial integration is a major determinant for the hypoalgesic response after isometric exercise. In contrast to manual pressure algometry, computer-controlled cuff algometry stimulates a larger tissue volume (Polianskis et al., 2001). Moreover, cuff algometry is less likely to be influenced by local variations in pain sensitivity and is also an examiner-independent technique reducing the potential measurement bias. The current results are in agreement with a previous study demonstrating no hypoalgesic response after isometric hand exercises when compared with a rest condition (Umeda et al., 2009). In addition, Bartholemew et al. (1996) found that pressure pain tolerance but not pressure pain threshold increased after an exercise session with mixed types of exercises. **Although multisegmental increase in PPT after isometric exercise has been demonstrated (Vaegter et al., 2014, Kosek and Lundberg, 2003, Hoeger Bement et al., 2008) the increase in pressure pain thresholds is larger in the exercising body part compared with non-exercising body parts (Vaegter et al., 2014), indicating that local mechanisms play an important role in the EIH response after isometric exercise. Moreover, pronounced EIH responses at the contracting thigh muscle compared with the contralateral non-contracting thigh muscle has previously been demonstrated (Kosek and Lundberg, 2003). This could influence the results in this study as heat and pressure pain sensitivity was assessed on the foot and lower leg, respectively, and not on the thigh.**

Although heat pain threshold increased compared with baseline no significant difference was found compared with quiet rest, indicating that isometric exercise does not influence pain perception to pressure or heat stimulus near the threshold when compared with quiet rest. The effect of isometric exercise on heat pain threshold has not previously been investigated, but the results are in agreement with previous studies demonstrating no effect on heat pain threshold after aerobic exercise (Kodesh and Weissman-Fogel, 2014, Cook et al., 2010). However, previous studies have demonstrated reduced pain intensity to heat pain (Misra et al., 2014) and reduced temporal summation of heat pain (Koltyn et al., 2013) after isometric exercise indicating that isometric exercise can influence pain perception to heat stimulus above the pain threshold. Furthermore, the

study by Misra et al (2014) demonstrated a progressive increase in the hypoalgesic effect with an increase in exercise intensity and it is currently unknown whether higher intensity exercise (e.g. 60% MVC) would have influenced pain perception to pressure or heat stimulus near the pain threshold. Isometric exercise was the only exercise stimulus used in this study; thus, the results cannot be generalized to other modes of exercise.

Using other paradigms for assessment of endogenous pain modulation, such as conditioning pain modulation, similar differences in modality-specific findings have been demonstrated. In line with the current results, Kosek and Hansson (1997) and Tuveson et al. (2006) showed that pressure pain threshold but not heat pain threshold increased in healthy subjects after a tourniquet test used to assess the conditioning pain modulation. However, Leffler et al. (2002) demonstrated an increase in pressure pain and heat pain thresholds in healthy subjects during a cold pressor test and Oono et al. (2013) showed an increase in pressure pain thresholds and pain tolerance in healthy subjects when a compression device around the head was used to assess conditioning pain modulation. Although conflicting results, different mechanisms may underlie endogenous pain modulation for various types of noxious stimulation and further research in this area is warranted.

The non-significant correlation between heat pain thresholds and pressure pain thresholds indicate that heat stimulation and cuff algometry assess different mechanisms. Similar findings have been reported for pain thresholds assessed by electrical, thermal and mechanical modalities (Neziri et al., 2011).

Test-retest reliability

Cuff pressure pain threshold and tolerance demonstrated excellent ICCs and acceptable agreement between tests with no systematic mean difference before and after the resting condition in healthy young men. Previous studies on cuff pressure algometry have demonstrated high levels of reliability with ICC values above 0.7 for test-retest data in healthy subjects (Graven-Nielsen et al., 2015) and in patients with chronic musculoskeletal pain (Vaegter et al., 2016). Previous studies demonstrating good test-retest reliability have based the pressure algometry pain thresholds on the average of at least two trials (Nussbaum and Downes, 1998, Ohrbach and Gale, 1989). However, the present study showed high ICC and acceptable agreement based on just one repetition with computer-controlled cuff algometry.

Heat pain threshold demonstrated lower ICC compared with cuff algometry but acceptable agreement between tests with no systematic mean difference between the two sessions. A previous systematic review on the test-retest reliability of quantitative sensory testing including heat pain threshold demonstrated that the reliability of heat pain threshold ranged from fair to excellent. A possible explanation for the lower ICC for heat pain threshold compared to cuff algometry is that heat pain threshold is more easily affected by environmental factors, such as ambient temperature and noise; methodological factors, such as test protocol, test application, and test instructions; and the cooperation and attention of the individual being tested (Chong and Cros, 2004). This may also explain the differences in baseline HPT found in the two sessions. Furthermore, heat pain and pressure pain sensitivity was assessed at different sites (dorsum of foot vs. circumference around lower extremity). It is possible that the difference in test-retest reliability is related to whether the tests are applied at bony or more muscular body sites.

Limitations

Pain tolerance was only assessed with pressure stimulus and the effect of isometric exercise on heat pain tolerance is unclear in this sample. Heat pain tolerance was not assessed in this study due to ethical considerations. Previous research has shown that heat pain stimulation may influence subsequent responses to mechanical stimulation (Grone et al., 2012) causing a risk of carry over effect in the experimental design in session 1. However, such carry-over effect is unlike in the present study as no significant order effect was found on heat pain or pressure pain sensitivity. Finally, the results from the current study can only be generalized to healthy young men and it is unclear whether women, older subjects, and individuals with chronic pain would experience similar results. **Further research on gender differences in EIH after isometric exercises is warranted as previous studies have demonstrated mixed results (Koltyn et al., 2001; Kosek and Lundberg 2003).**

Conclusion

Isometric exercise significantly increased cuff pressure pain tolerance compared with baseline and the control condition. Although not known if related with the exercise dose the findings suggest that hypoalgesia after isometric exercise is primarily affecting tolerance of pain compared with the pain threshold. These findings indicate that mechanisms underlying exercise-induced hypoalgesia after isometric exercise are targeting pain perception above the threshold and contribute to the

understanding of how isometric exercise influences pain perception, which is necessary to optimize the clinical utility of exercise in management of chronic pain.

Author contributions

All authors contributed to the conception and design of the study as well as making intellectual contributions to its content. Henrik Bjarke Vaegter, Anders Bjarke Madsen, Jakob Fridriksson, and Marcus Dasa collected data, Henrik Bjarke Vaegter, Thomas Graven-Nielsen and Marie Hoeger Bement contributed to the analysis and interpretation. Henrik Bjarke Vaegter drafted the manuscript. All authors discussed the results, commented on and approved the final manuscript.

Figure Legends

Fig. 1: Illustration of the experimental procedure performed on the three testing days. Session 1: The sensitivity to heat pain and pressure pain were assessed before and immediately after a 15 min control condition (quiet rest). The sequence between assessment of heat and pressure pain sensitivity were randomized and counterbalanced. Following rest the maximal voluntary contraction (MVC) for isometric knee extension was determined. Session 2 and 3: Before and immediately after a 15 min active condition (a 3 min submaximal isometric knee extension at 30% of MVC with the dominant leg preceded by 12 min rest) the sensitivity to either heat pain or pressure pain was assessed. The sequence between session 2 and 3 were randomized and counterbalanced. ‘MVC’: Maximal voluntary contraction. ‘NRS’: Numerical rating scale. ‘RPE’: Rating of perceived exertion. ‘cPPT’: Cuff pressure pain threshold. ‘cPTT’: Cuff pressure pain tolerance. ‘HPT’: Heat pain threshold.

Fig. 2: Mean (+ SEM, n=20) cuff pressure pain threshold (**A**) and cuff pressure pain tolerance (**B**) assessed before and after a submaximal isometric knee extension (Active) and quiet rest (Control). The cuff pressure pain sensitivity was assessed at the dominant lower leg. Significantly different compared with baseline (*, $P < 0.05$) and significantly different compared with the control condition (†, $P < 0.05$).

Fig. 3: Mean (+ SEM, n=20) percentage increase in heat pain threshold (HPT), cuff pressure pain threshold (cPPT), and cuff pressure pain tolerance (cPTT) after submaximal isometric exercise. Significantly different compared with other assessment parameters (*, $P < 0.05$).

References

- Bartholomew, J. B., Lewis, B. P., Linder, D. E., Cook, D. B. (1996). Post-exercise analgesia: replication and extension. *J Sports Sci* **14**, 329-334.
- Chong, P. S., Cros, D. P. (2004). Technology literature review: quantitative sensory testing. *Muscle Nerve* **29**, 734-47.
- Cook, D. B., Stegner, A. J., Ellingson, L. D. (2010). Exercise alters pain sensitivity in Gulf War veterans with chronic musculoskeletal pain. *J Pain* **11**, 764-772.
- Fleiss, J. L. (1986). The design and analysis of clinical experiments, New York, Wiley.
- Graven-Nielsen, T., Arendt-Nielsen, L. (2010). Assessment of mechanisms in localized and widespread musculoskeletal pain. *Nat Rev Rheumatol* **6**, 599-606.
- Graven-Nielsen, T., Vaegter, H. B., Finocchietti, S., Handberg, G., Arendt-Nielsen, L. (2015). Assessment of musculoskeletal pain sensitivity and temporal summation by cuff pressure algometry: A reliability study. *Pain* In Press.
- Greenspan, J. D., Craft, R. M., Leresche, L., Arendt-Nielsen, L., Berkley, K. J., Fillingim, R. B., Gold, M. S., Holdcroft, A., Lautenbacher, S., Mayer, E. A., Mogil, J. S., Murphy, A. Z., Traub, R. J. (2007). Studying sex and gender differences in pain and analgesia: a consensus report. *Pain* **132**, S26-45.
- Grone, E., Crispin, A., Fleckenstein, J., Irnich, D., Treede, R. D., Lang, P. M. (2012). Test order of quantitative sensory testing facilitates mechanical hyperalgesia in healthy volunteers. *J Pain* **13**, 73-80.
- Hoeger Bement, M., Dicapio, J., Rasiarmos, R., Hunter, S. K. (2008). Dose response of isometric contractions on pain perception in healthy adults. *Med Sci Sports Exerc* **40**, 1880-1889.
- Hoeger Bement, M., Rasiarmos, R. L., Dicapio, J. M., Lewis, A., Keller, M. L., Harkins, A. L., Hunter, S. K. (2009). The role of the menstrual cycle phase in pain perception before and after an isometric fatiguing contraction. *Eur J Appl Physiol* **106**, 105-112.
- Hoeger Bement, M., Weyer, A., Hartley, S., Drewek, B., Harkins, A. L., Hunter, S. K. (2011). Pain perception after isometric exercise in women with fibromyalgia. *Arch Phys Med Rehabil* **92**, 89-95.
- Hoeger Bement, M., Drewek, B., Hunter, S. K. (2014). Men report greater pain relief following sustained static contractions than women when matched for baseline pain. *J Mot Behav* **46**, 107-13.

- Kodesh, E., Weissman-Fogel, I. (2014). Exercise-induced hypoalgesia - interval versus continuous mode. *Appl Physiol Nutr Metab* **39**, 829-34.
- Koltyn, K. F., Brellenthin, A. G., Cook, D. B., Sehgal, N., Hillard, C. (2014). Mechanisms of Exercise-Induced Hypoalgesia. *J Pain* **15**, 1294-1304.
- Koltyn, K. F., Knauf, M. T., Brellenthin, A. G. (2013). Temporal summation of heat pain modulated by isometric exercise. *Eur J Pain* **17**, 1005-11.
- Koltyn, K. F., Trine, M. R., Stegner, A. J., Tobar, D. A. 2001. Effect of isometric exercise on pain perception and blood pressure in men and women. *Med Sci Sports Exerc*, **33**, 282-290.
- Koltyn, K. F., Umeda, M. (2007). Contralateral attenuation of pain after short-duration submaximal isometric exercise. *J Pain* **8**, 887-892.
- Kosek, E., Ekholm, J. (1995). Modulation of pressure pain thresholds during and following isometric contraction. *Pain* **61**, 481-486.
- Kosek, E., Hansson, P. (1997). Modulatory influence on somatosensory perception from vibration and heterotopic noxious conditioning stimulation (HNCS) in fibromyalgia patients and healthy subjects. *Pain* **70**, 41-51.
- Kosek, E., Lundberg, L. (2003). Segmental and plurisegmental modulation of pressure pain thresholds during static muscle contractions in healthy individuals. *Eur J Pain* **7**, 251-258.
- Lannersten, L., Kosek, E. (2010). Dysfunction of endogenous pain inhibition during exercise with painful muscles in patients with shoulder myalgia and fibromyalgia. *Pain* **151**, 77-86.
- Leffler, A. S., Hansson, P., Kosek, E. (2002). Somatosensory perception in a remote pain-free area and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from long-term trapezius myalgia. *Eur J Pain* **6**, 149-159.
- Lemley, K. J., Drewek, B., Hunter, S. K., Hoeger Bement, M. (2014). Pain relief after isometric exercise is not task-dependent in older men and women. *Med Sci Sports Exerc* **46**, 185-91.
- Lemley, K. J., Hunter, S. K., Hoeger Bement, M. (2015). Conditioned Pain Modulation Predicts Exercise-Induced Hypoalgesia in Healthy Adults. *Med Sci Sports Exerc* **47**, 176-84.
- Mannerkorpi, K., Henriksson, C. (2007). Non-pharmacological treatment of chronic widespread musculoskeletal pain. *Best Pract Res Clin Rheumatol* **21**, 513-534.
- Misra, G., Paris, T. A., Archer, D. B., Coombes, S. A. (2014). Dose-response effect of isometric force production on the perception of pain. *PLoS One* **9**, e88105.

- Naugle, K. M., Naugle, K. E., Fillingim, R. B., Riley, J. L., (2014). Isometric Exercise as a Test of Pain Modulation: Effects of Experimental Pain Test, Psychological Variables, and Sex. *Pain Med* **15**, 692-701.
- Neziri, A. Y., Scaramozzino, P., Andersen, O. K., Dickenson, A. H., Arendt-Nielsen, L., Curatolo, M. (2011). Reference values of mechanical and thermal pain tests in a pain-free population. *Eur J Pain* **15**, 376-83.
- Nussbaum, E. L., Downes, L. (1998). Reliability of clinical pressure-pain algometric measurements obtained on consecutive days. *Phys Ther* **78**, 160-9.
- Ohrbach, R., Gale, E. N. (1989). Pressure pain thresholds in normal muscles: reliability, measurement effects, and topographic differences. *Pain* **37**, 257-63.
- Oono, Y., Baad-Hansen, L., Wang, K., Arendt-Nielsen, L., Svensson, P. (2013). Effect of conditioned pain modulation on trigeminal somatosensory function evaluated by quantitative sensory testing. *Pain* **154**, 2684-90.
- Polianskis, R., Graven-Nielsen, T., Arendt-Nielsen, L. (2001). Computer-controlled pneumatic pressure algometry--a new technique for quantitative sensory testing. *Eur J Pain* **5**, 267-77.
- Popescu, A., LeResche, L., Truelove, E. L., Drangsholt, M. T. (2010). Gender differences in pain modulation by diffuse noxious inhibitory controls: a systematic review. *Pain*, **150**, 309-318.
- Tuveson, B., Leffler, A. S., Hansson, P. (2006). Time dependent differences in pain sensitivity during unilateral ischemic pain provocation in healthy volunteers. *Eur J Pain* **10**, 225-32.
- Umeda, M., Newcomb, L. W., Ellingson, L. D., Koltyn, K. F. (2010). Examination of the dose-response relationship between pain perception and blood pressure elevations induced by isometric exercise in men and women. *Biol Psychol* **85**, 90-96.
- Umeda, M., Newcomb, L. W., Koltyn, K. F. (2009). Influence of blood pressure elevations by isometric exercise on pain perception in women. *Int J Psychophysiol* **74**, 45-52.
- Vaegter, H. B., Handberg, G., Graven-Nielsen, T. (2014). Similarities between exercise-induced hypoalgesia and conditioned pain modulation in humans. *Pain* **155**, 158-67.
- Vaegter, H. B., Handberg, G., Graven-Nielsen, T. (2015). Isometric exercises reduce temporal summation of pressure pain in humans. *Eur J Pain* **19**, 973-83.
- Vaegter, H. B., Handberg, G., Graven-Nielsen, T. (2016). Hypoalgesia After Exercise and Cold Pressor Test are Reduced in Chronic Musculoskeletal Pain Patients with High Pain Sensitivity. *Clin J Pain* **32**, 58-69.

Yu, X. M., Mense, S. (1990). Response properties and descending control of rat dorsal horn neurons with deep receptive fields. *Neuroscience* **39**, 823-31.

Table 1: Intraclass correlations (ICCs) and Bland/Altman analyses for assessment of pain sensitivity parameters before and after the resting condition in session 1. ‘HPT’: Heat pain threshold. ‘cPPT’: Cuff pressure pain threshold. ‘cPTT’: Cuff pressure pain tolerance.

Pain sensitivity parameter	ICC			Bland and Altman		
	Before resting Mean \pm SD	After resting Mean \pm SD	ICC _{3,1} (95 % CI)	Mean difference (95 % CI)	SD diff (kPa)	95 % limits of agreement
HPT (°C)	46.6 \pm 2.1	46.9 \pm 2.2	0.54 (0.14-0.79)	0.4 (-0.6 – 1.3)	2.1	-3.7 – 4.5
cPPT (kPa)	20.6 \pm 8.5	21.7 \pm 8.9	0.86 (0.67-0.94)	1.2 (-1.0 – 3.4)	4.7	-8.0 – 10.4
cPTT (kPa)	63.7 \pm 18.4	64.2 \pm 18.3	0.77 (0.50-0.90)	0.5 (-5.4 – 6.4)	12.6	-.24.2 – 25.2





