



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Coronary artery calcification in patients diagnosed with severe mental illness

Kugathasan, Pirathiv; Berg Johansen, Martin; Jensen, Mikkel Bak; Aagaard, Jørgen; Jensen, Svend Eggert

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Kugathasan, P., Berg Johansen, M., Jensen, M. B., Aagaard, J., & Jensen, S. E. (2018). *Coronary artery calcification in patients diagnosed with severe mental illness*. Poster presented at The 26th Congress of the European Psychiatric Association , Nice, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Coronary artery calcification in patients diagnosed with severe mental illness



Pirathiv Kugathasan^{1,2}, Martin B. Johansen^{3,4}, Mikkel B. Jensen¹, Jørgen Aagaard^{1,2}, Svend E. Jensen^{2,3}

- 1) Aalborg University Hospital, Psychiatry, Aalborg, Denmark
- 2) Department of Clinical Medicine, Aalborg University, Aalborg, Denmark
- 3) Department of Cardiology, Aalborg University Hospital, Aalborg, Denmark
- 4) Unit of Clinical Biostatistics, Aalborg University Hospital, Aalborg, Denmark

Background

- Patients diagnosed with severe mental illness (SMI) have an excess mortality, primarily caused by cardiovascular disease.
- The majority of deaths are related to an increased incidence of coronary artery disease, but efforts to predict and manage cardiovascular risk in patients with SMI have been ineffective.
- Coronary artery calcification (CAC) is a clinical predictor for coronary artery disease, which can be measured by cardiac CT.
- CAC is widely used in the general population to predict future cardiovascular events, but no others have compared any outcomes between patients with SMI and the general population to date.

Aim

- To investigate the effect of coronary artery calcification on mortality in patients diagnosed with SMI compared to the effect in the general population.

Methods

- Design
 - A Danish population based cohort study
- Study period
 - 1 January 2008 to 31 December 2016
- Study population
 - Patients diagnosed with SMI
 - Schizophrenia (ICD-10; F20)
 - Bipolar disorder (ICD-10; F30+F31)




Figure 1. Study flowchart for patient selection and eligibility.

Results

| | General population (n=48,193) | Severe mental illness (n=564) | P |
|-------------------------------|-------------------------------|-------------------------------|--------|
| Coronary calcium score, n (%) | | | <0.01 |
| 0-100 | 36,049 (74.80) | 452 (80.14) | |
| >100 | 12,144 (25.20) | 112 (19.86) | |
| Number of deaths, n (%) | 1009 (2.10) | 21 (3.72) | |
| Causes of death, n (%) | | | <0.001 |
| Cardiovascular death | 343 (33.99) | 6 (28.57) | |
| Other natural death | 628 (62.24) | 10 (47.62) | |
| Unnatural death | 38 (3.77) | 5 (23.81) | |

Table 1. Follow-up characteristics of the individuals included. The variables are further divided into low coronary calcium score (0-100) and high coronary calcium score (>100).

| Coronary calcium score | Crude HR | 95% CI | Adjusted HR* | 95% CI |
|------------------------------|----------|-----------|--------------|-----------|
| General population | | | | |
| 0-100 | 1.0 | Reference | 1.0 | Reference |
| >100 | 3.33 | 2.95-3.77 | 1.58 | 1.38-1.80 |
| Severe mental illness | | | | |
| 0-100 | 2.53 | 1.51-4.22 | 3.86 | 2.31-6.46 |
| >100 | 4.43 | 1.98-9.90 | 3.07 | 1.37-6.87 |

Table 2. Cox proportional hazards model of mortality rates in the general population and patients with SMI. *Adjusted for age, gender, and calendar period.




Figure 2. Kaplan-Meier estimates on survival following cardiac CT in low score- versus high score agatston groups for individuals in the general population and patients with SMI.

Conclusion

- Patients with SMI are not demonstrating signs of early coronary artery calcification.
- Mortality rates were still markedly higher in patients with SMI compared to the general population, suggesting a more rapid progression of ischemic heart disease in SMI patients.

Contact

Pirathiv Kugathasan, MSc, PhD student
Email: p.kugathasan@rn.dk