



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Intra-articular gold micro particles and hyaluronic acid improve osteoarthritic pain and social function

A pilot study

Rasmussen, Sten

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Rasmussen, S. (2019). *Intra-articular gold micro particles and hyaluronic acid improve osteoarthritic pain and social function: A pilot study*. Poster presented at Pain in Europe, Congress of the European Pain Federation, EFIC, Valencia, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Intra-articular gold micro particles and hyaluronic acid improve osteoarthritic pain and social function -A pilot study

Sten Rasmussen^{1, 2}

¹)Department of Clinical Medicine, School of Medicine, Aalborg University, Aalborg Denmark

²)Department of Orthopedic Surgery, Aalborg University Hospital, Aalborg, Denmark

Background and aims

Many patients suffering from osteoarthritis (OA) do not get adequate pain relieve. Evidence suggest an inflammatory component in OA pain. Animal studies prove the effect of gold implantation in arthritic joints and a stimulation of the immune system.

Animal studies indicate gold ions have a long-acting effect on OA pain. The immuno-modulatory effect of gold ions have for more than 50 years a known anti-inflammatory effect in the treatment of rheumatic arthritis. Gold ions alter the function of macrophages by inhibiting lysosomal enzymes and lowering production of pro-inflammatory cytokines (1-2).

Dissolucytotic metallic gold (DMG) ions have an immune-suppressive effect in laboratory testing (3-6) (Figure 1 - 4). Animal studies prove the effect of gold implantation in arthritic joints (7-10). Injection of DMG in animal models stimulate the immune system (11-12). The carrier for injecting the DMG micro particles is hyaluronic acid (13-14).

The present open pilot study aimed to investigate if gold ions released from intraarticular gold micro particles have a role in different joint OA.




Figure 1. Macrophages controls the dissolution membrane which liberate the gold ions by oxidation of the surface. Once the ions are liberated, most likely as Au(CN)₂⁻, they are free to diffuse through the immediate microenvironment. The gold-loaded molecules are taken up into the cells, primarily macrophages, mastcells and histocytes.




Figure 2. Close to the gold implant gold-loaded molecular clusters are located outside cells. The two loaded cells are believed to be macrophages loaded with gold ions. The gold ions accumulate primarily in the lysosomes (3).




Figure 3. Once in the intercellular fluid and the intracellular compartments, the gold ions act in the same ways that have been demonstrated for systemically administered gold ions. The effect is related to the ability of the gold ions to unfold the protein structures.




Figure 4. Gold ions suppress inflammation locally by affecting certain signalling molecules and binding enzymes essential for the inflammatory process. The DNA binding activity and transcription regulation of NF-κB is abolished when Au⁺ ions replace Zn²⁺ ions. (4)

Methods

A cohort of 39 patients, aged 58 (37-78) years, pain \geq 3 months, and Kellgren-Lawrence OA grade 3-4 were included. Metallic gold 20 mg, 72.000 pieces, 20-40 μ -meter (Berlock-Micro-Implants, HumanGoldInject) (13-14) were injected into the joints using 1-2 mL hyaluronic acid as the carrier. The primary outcome measure was change in pain at 8 weeks. The secondary outcome measures was change in social outcome assessed by a Danish 8 point version of the Glasgow Outcome Scale at 8 weeks.

Results

This study includes 31 knee, 6 hip, 4 shoulder, 4 ankle, 2 wrist and 10 small joints. Pain or social outcome was improved in 34/39 patients (95% CI 72.6 – 95.7 %). Pain was reduced in 44/57 joints (95 % CI 61 – 89 %). Social outcome was improved for 42/57 joints (95 % CI 57.2 – 86.5 %) (P < 0.001).

Conclusions

The significant improvements in pain and social outcome caused by the intraarticular gold micro particles and hyaluronic acid indicate an inhibition of inflammation.

| | Before | 8 weeks |
|---|--------|---------|
| <i>Dead</i> | | |
| <i>Vegetative</i> | | |
| Dependent in daily life activities | 9 | 1 |
| Independent and cannot resume normal activity | 7 | 2 |
| Previous activity at lower level | 19 | 10 |
| Normal activity with deficit | 20 | 18 |
| Normal activity with minor deficits or complaints | 2 | 19 |
| No signs or symptoms | | 7 |

Table 1. Short category description and results of the Danish eight-point version of the Glasgow Outcome Scale.

References

- Persellin RH. The effect of gold salt on lysosomal enzymes of the peritoneal macrophage. *Arthritis Rheum* 1966;9:57-65.
- Yanni G. Intramuscular gold decreases cytokine expression and macrophage numbers in the rheumatoid synovial membrane. *Ann Rheum Dis* 1994;53:315-22.
- Danscher G. In vivo liberation of gold ions from gold implants. *Histochem Cell Biol* 2002; 117:447-52.
- Yang JP, Merin JP, Nakano T, Kato T, Kitade Y, Okamoto T. Inhibition of the DNA-binding activity of NF-kappa B by gold compounds in vitro. *FEBS Lett.* 1995 Mar 13;361(1):89-96.
- Larsen A. Gold ions bio-released from metallic gold particles reduce inflammation and apoptosis and increase the regenerative responses in focal brain injury. *Histochem Cell Biol* 2008;130:681-92.
- Seifert O. Gene expression profiling of macrophages: implications for an immunosuppressive effect of dissolucytotic gold ions. *J Inflamm* 2012;9:43.
- Jaeger GT. Double-blind, placebo-controlled trial of the pain-relieving effects of the implantation of gold beads into dogs with hip dysplasia. *Vet Rec* 2006;158:722-6.
- Jaeger GT. Two years follow-up study of the pain-relieving effect of gold bead implantation in dogs with hip-joint arthritis. *Acta Vet Scand* 2007;49:9.
- Jaeger GT. Gold Bead Implantation in Acupoints for Coxofemoral Arthrosis in Dogs. *Animals* 2012;2:426-36.
- Lie KI. Inflammatory response to therapeutic goldbead implantation in canine hip joint osteoarthritis. *Vet Pathol* 2011;48:1118-24.
- Havarinasab S. Gold causes genetically determined autoimmune and immunostimulatory responses in mice. *Clin Exp Immunol* 2007;150:179-88.
- Hazer DB. Soft tissue response to the presence of polypropylene-G-poly(ethylene glycol) comb-type graft copolymers containing gold nanoparticles. *J Biomed Biotechnol* 2011;2011:956169.
- Danscher G. In: Patent US, editor. Medicament and method of treatment of patients with heavy metals 2010. United States Patent Number US 7,655,261 B2.
- Märki N. Safety of Intra-Articular Gold Microimplants in Horses. *J Equine Vet Res* 2018;60:59-66.



AALBORG UNIVERSITY HOSPITAL



AALBORG UNIVERSITY
DENMARK