



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## **Patellar fractures are associated with an increased risk of total knee arthroplasty**

*A matched cohort study of 6096 patellar fractures with a mean follow-up of 14.3 years*

Larsen, P; Rathleff, M S; Østgaard, S E; Johansen, M B; Elsøe, R

*Published in:*  
The Bone & Joint Journal

*DOI (link to publication from Publisher):*  
[10.1302/0301-620X.100B11.BJJ-2018-0312.R2](https://doi.org/10.1302/0301-620X.100B11.BJJ-2018-0312.R2)

*Publication date:*  
2018

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Larsen, P., Rathleff, M. S., Østgaard, S. E., Johansen, M. B., & Elsøe, R. (2018). Patellar fractures are associated with an increased risk of total knee arthroplasty: A matched cohort study of 6096 patellar fractures with a mean follow-up of 14.3 years. *The Bone & Joint Journal*, 100-B(11), 1477-1481. <https://doi.org/10.1302/0301-620X.100B11.BJJ-2018-0312.R2>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# The Bone & Joint Journal

70 years of orthopaedic excellence

**Patella fractures are associated with an increased risk of total knee arthroplasty—a matched cohort study of 6,096 patella fractures with an average of 14.3 years follow-up**

Journal:	<i>The Bone &amp; Joint Journal</i>
Manuscript ID	BJJ-2018-0312.R1
Manuscript Type:	Original Article
Keywords:	Total Knee Replacement, patella fracture, epidemiology

SCHOLARONE™  
Manuscripts

1  
2  
3 **Patella fractures are associated with an increased risk of total knee**  
4 **arthroplasty—a matched cohort study of 6,096 patella fractures with an average**  
5 **of 14.3 years follow-up**  
6  
7  
8

9 *ABSTRACT*  
10

11  
12 **Aims:** To investigate the incidence of total knee replacement (TKR) following patella  
13 fractures and compare this to an age- and gender-matched group without a prior  
14 patella fracture.  
15  
16

17  
18 **Patients and Methods:** A national matched cohort study based on the Danish  
19 National Patient Register including all citizens of Denmark (approximately 5.7  
20 million) was undertaken. 6096 patients who sustained a patella fracture in Denmark  
21 between January 1, 1996, and December 31, 2000, were included and followed until  
22 December 31, 2015, with regard to treatment with TKR and/or knee arthroscopy.  
23  
24  
25  
26  
27

28  
29 **Results:** Patients with a patella fracture had an increased risk of TKR (hazard ratio  
30 (HR): 1.83, 95%CI: 1.57-2.13) compared to citizens without a patella fracture, and the  
31 effect was strongest during the first five years (HR: 3.02, 95%CI: 2.26-4.03). Patients  
32 with a patella fracture also had a higher risk of knee arthroscopy (HR: 3.94, 95%CI:  
33 3.49-4.46), and the effect was highest during the first five years after the fracture (HR:  
34 7.40, 95%CI: 6.32-8.66).  
35  
36  
37  
38  
39  
40  
41

42  
43 **Conclusions:** Patella fractures are associated with a considerably increased risk of  
44 total knee arthroplasty and knee arthroscopy. The consequences of a patella fracture  
45 are more severe than previously documented, and patients must expect a lifelong  
46 increased risk of TKR.  
47  
48  
49  
50

- 51  
52  
53  
54
  - Patients suffering a patella fracture has an increased risk of TKR and knee  
55 arthroscopy throughout life.  
56  
57  
58  
59  
60

## ***INTRODUCTION***

The modern elderly population has an increased demand for a pain-free and active lifestyle with a high functional capacity.<sup>1</sup> They want to stay active without limitations due to joint pain. This may contribute to the steady increase in joint replacements around the world.<sup>1</sup> Identifying early factors associated with an increased risk of joint pain, osteoarthritis, and knee joint replacement has become increasingly important.

Patella fractures account for 0.7-1%<sup>2,3</sup> of all fractures, with an incidence of 13.1/100,000/year.<sup>4</sup> Despite the commonality, little information exists about the long-term consequences following a patella fracture.<sup>3,5-12</sup> Recent literature suggests that 50 percent of patients with patella fractures continue to experience symptoms affecting their function during activities of daily living and quality of life (QOL).<sup>5,9,13</sup>

Traumatic knee injuries that involve the tibia may increase the risk of posttraumatic knee osteoarthritis (OA).<sup>10,14,15</sup> The injury mechanisms underpinning the development of OA may be related to the forces applied to the chondral surfaces at the time of injury, fracture comminution, and mode of injury.<sup>15</sup> Post-injury factors such as the quality of articular surface reduction and the choice of treatment method could also contribute to the risk of posttraumatic knee OA following a patella fracture.<sup>5,6</sup>

Total knee replacement (TKR) is commonly required in the treatment of end-stage knee osteoarthritis. A large-scale retrospective study of 19,641 TKR procedures showed that 0.5% of all patients treated with a TKR reported a history of a patella fracture.<sup>14</sup> This suggests that a patella fracture may increase the risk of TKR later in life.

1  
2  
3 The primary aim of this study was to investigate the incidence of TKR following  
4 patella fractures compared to an age- and gender-matched group without a prior  
5 patella fracture.  
6  
7  
8  
9

10  
11 The secondary aims were to investigate the incidence of knee arthroscopy following a  
12 patella fracture and compare this to an age- and gender-matched group without a prior  
13 patella fracture and, furthermore, to compare the time to TKR and arthroscopy of the  
14 knee following a patella fracture compared to that of an age- and gender-matched  
15 non-exposed group.  
16  
17  
18  
19  
20  
21  
22  
23

## 24 ***PATIENTS AND METHODS***

### 25 ***Study design***

26  
27 The study was designed as a matched cohort study using prospectively obtained  
28 registry data from the Danish National Patient Register on all citizens of Denmark.  
29  
30  
31  
32  
33  
34

35 All patients who sustained a patella fracture in Denmark between January 1, 1996,  
36 and December 31, 2000, were included and followed until December 31, 2015,  
37 regarding treatment with TKR (including patellofemoral replacement) and/or knee  
38 arthroscopy.  
39  
40  
41  
42  
43  
44  
45

46 Danish law requires that all patient contacts with hospital and outpatient clinics in  
47 Denmark are registered in the Danish National Patient Register.<sup>16</sup> Hospital  
48 identification, date and time of activity, and patient's municipality (among other  
49 characteristics) are registered. A Civil Registration Number (CPR) is given to all  
50 residents of Denmark and registered in the Civil Registration System, and information  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 on emigration and death is recorded in this registry.<sup>17</sup> This enables researchers to have  
4  
5 a complete and valid registration of all health-related issues on an individual level in a  
6  
7 complete population.<sup>18</sup>  
8  
9

10  
11 The Local Ethics Committee was asked to approve the study and answered that the  
12  
13 study design does not need notification. The Danish Data Protection Agency  
14  
15 approved the study (J. nr. 2008-58-0028, Id: 2016-176). The reporting of the study  
16  
17 complies with the Strengthening the Reporting of Observational Studies in  
18  
19 Epidemiology (STROBE) Statement.<sup>19</sup>  
20  
21

22  
23  
24 The full study protocol and study analysis plan was published online before the start  
25  
26 of the study.<sup>XX</sup> The reporting of the present study lacks information considering  
27  
28 ipsilateral and contralateral information as predefined because side-specific  
29  
30 information was not a required part of data in the Danish National Patient Register.  
31  
32  
33

### 34 35 ***Study population and data retrieval*** 36

37 The exposed group of patients with a patella fracture was identified through a  
38  
39 retrospective review of all Danish patients diagnosed with a patella fracture between  
40  
41 January 1, 1996, and December 31, 2000. All patients who were registered regardless  
42  
43 of contact type with the ICD-10 diagnosis code for “patella fracture” (S82.0) were  
44  
45 identified in the Danish National Patient Register. Basic characteristics, gender, and  
46  
47 age at the time of diagnosis were obtained. Patients were included at the time of first  
48  
49 contact with a patella fracture reported to the Danish National Patient Register.  
50  
51 Patients with prior patella fractures and patients with prior TKR were excluded from  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 the study. All patients were followed with regard to surgery with TKR and/or  
4 arthroscopic surgery of the knees identified in the Danish National Patient Register.  
5  
6  
7

8  
9 The non-exposed group consisted of individuals identified from the Civil Registration  
10 System matched to the exposed case based on age and gender. For each exposed case,  
11 we included 10 non-exposed citizens who had not experienced a patella fracture or a  
12 TKR before the inclusion date. Information regarding arthroscopy of the knees and  
13 TKR were obtained from the Danish National Patient Register.  
14  
15  
16  
17  
18  
19

20  
21 Both exposed and non-exposed individuals were censored in case of emigration from  
22 the country or at the end of follow-up. Death was considered a competing event and  
23 so was receiving a TKR when considering knee arthroscopy as the outcome.  
24  
25  
26  
27

### 28 ***Primary and secondary outcome***

29  
30 The primary outcome was treatment with a TKR. The secondary outcome was  
31 treatment with knee arthroscopy.  
32  
33  
34  
35

### 36 ***Statistical methods***

37  
38 The risk of experiencing a TKR over time was assessed using cumulative incidence  
39 proportions which were calculated using the Aalen-Johansen estimator.<sup>20</sup>  
40  
41  
42  
43  
44

45  
46 The analysis evaluating the effect of patella fracture on the incidence of TKR was  
47 performed using a Cox proportional hazards regression model comparing the exposed  
48 and non-exposed groups. The effect estimate was reported as a hazard ratio with a  
49 corresponding 95% confidence interval to estimate the incidence rate ratio. Results  
50 from the crude analysis without adjustments are reported. Furthermore, we repeated  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 the analysis stratified by age groups (0-50, 51+) and gender. To investigate the  
4  
5 assumption of proportional hazards, the follow-up time is divided into five-year  
6  
7 periods.  
8  
9

10  
11 Additional analyses were performed to investigate the effect of patella fracture on the  
12  
13 secondary outcome (knee arthroscopy) following the same methods as for the main  
14  
15 analysis except that TKR, in addition to death, was considered as a competing event  
16  
17 for arthroscopy.  
18  
19

20  
21  
22 All analyses were performed using Stata statistical software (StataCorp LP), and the  
23  
24 significance level for analyses ( $\alpha$ ) was set to 0.05.  
25  
26  
27

## 28 **RESULTS**

29  
30  
31 Between January 1, 1996, and December 31, 2000, we identified 6,096 patients who  
32  
33 sustained a patella fracture in Denmark. The median age of patients was 50.6 (IQR:  
34  
35 28.5-68.9) years, and 49.1% were women. The average observational period was 14.3  
36  
37 years.  
38  
39

### 40 **Primary outcome**

41  
42  
43 The analysis showed that 3.3% of patients with a prior patella fracture were treated  
44  
45 with a TKR (N=200) and that 2% of patients without a prior patella fracture were  
46  
47 treated with a TKR (N=1,239) during the observational period. Patients with a prior  
48  
49 patella fracture had a 1.83 (95%CI: 1.57-2.13) times higher hazard ratio (HR)  
50  
51 compared to patients without a prior patella fracture. The effect was highest during  
52  
53 the first five years after the fracture (HR: 3.02 (95%CI: 2.26-4.03)) (Table 1).  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5 The cumulative incidence of TKR during the entire observational period expressed for  
6  
7 the two groups is shown in Figure 1. The figure shows a significantly increased risk  
8  
9 of TKR in patients with a prior patella fracture compared to patients without a prior  
10  
11 patella fracture throughout the observational period.  
12  
13

#### 14 15 *Analyses of age and gender difference on TKR*

16  
17 Both genders show a significantly increased incidence of TKR in patients with a prior  
18  
19 patella fracture compared with patients without a prior patella fracture. Compared to  
20  
21 women, men presented with an increased incidence of TKR throughout the  
22  
23 observational period. The age group above 50 years presented with a substantially  
24  
25 increased risk compared to the age group below 50 years. However, the age group  
26  
27 below 50 years presented with a proportionately greater risk of TKR compared to the  
28  
29 reference population when compared to the age group above 50 years. (Supplemental  
30  
31 figure 1-4).  
32  
33  
34  
35  
36

#### 37 *Secondary outcomes—knee arthroscopy*

38  
39 The analysis showed that 6.8% of patients with a prior patella fracture were treated  
40  
41 with a knee arthroscopy (N=416) and 2.1% of patients without a prior patella fracture  
42  
43 were treated with a knee arthroscopy (N=1,249). Patients with a prior patella fracture  
44  
45 had a 3.94 (95%CI: 3.49-4.46) times higher hazard compared to patients without a  
46  
47 prior patella fracture. The effect was highest during the first five years after the  
48  
49 fracture (HR: 7.40 (95%CI: 6.32-8.66)) (Table 2).  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The cumulative incidence of knee arthroscopy during the entire observational period  
4 is shown in Figure 2. The figure shows a significantly increased risk of knee  
5 arthroscopy in patients with a prior patella fracture compared to patients who had not  
6 experienced a prior patella fracture throughout the observational period.

7  
8  
9  
10  
11 The analyses of age and gender differences in the incidence of knee arthroscopy show  
12 an almost equal risk between genders. The analysis stratified on age showed that the  
13 age group below 50 years of age had a considerably increased risk compared to the  
14 age group above 50. The vast majority of increased risk of arthroscopy is observed  
15 within the first three years following fracture. (Supplemental Figure 5-8).

## 22 23 24 **DISCUSSION**

25  
26 Patients with a prior patella fracture had an 83% higher incidence of TKR in the years  
27 following their injury compared to a similar group of individuals without a patella  
28 fracture. The incidence was highest during the first five years following a patella  
29 fracture. Collectively, these results indicate that patients with a patella fracture may  
30 be at risk of developing posttraumatic osteoarthritis, long-term pain, and may  
31 experience early treatment with TKR.  
32  
33  
34  
35  
36  
37  
38  
39  
40

41  
42 The long-term consequences of patella fractures may be more serious than previously  
43 anticipated, especially in younger patients. Julin et al.<sup>21</sup> reported that young age  
44 impairs the prognosis of TKR and is associated with increased revision rates. The  
45 outcome of revision surgery following TKR has been reported with impaired  
46 function, increased pain, and decreased QOL.<sup>22,23</sup> Moreover, Houdeck et al.<sup>14</sup> reported  
47 that a patella fracture before TKR was associated with a worse outcome but  
48 equivalent implant survival. Consequently, surgeons may be reluctant to offer  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 younger patients with severe posttraumatic osteoarthritis following a patella fracture  
4  
5 treatment with TKR. Further research needs to address this and focus on other  
6  
7 treatment modalities which might postpone or reduce the need for TKR following a  
8  
9 patella fracture.  
10

11  
12  
13 National preference regarding the use of patella resurfacing in primary TKR might  
14 influence the threshold when considering the possibility to treat patients experiencing  
15 pain following a patella fracture with a primary TKR. Compared to other  
16 Scandinavian countries, Denmark has a high resurfacing rate of 76%, which might  
17 influence the number of patients treated with a TKR following a patella fracture.<sup>24</sup>  
18  
19  
20  
21  
22  
23  
24  
25

26 Patients with a previous patella fracture had a four times higher incidence of a knee  
27 arthroscopy compared to the control group. This is a new finding, and the literature  
28 lacks information about this association. These findings may be a strong indicator that  
29 many patients experience pain and decreased knee function, especially in the first  
30 years following a patella fracture. However, this study did not include clinical  
31 information regarding the underlying causes leading to surgery and the subsequent  
32 outcomes following the procedures.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 A significant strength is the design of this study. Using high-quality data registries in  
46 Denmark, we included the entire population of patients with patella fractures and  
47 compared this to a non-exposed age- and gender-matched control group. Coding in  
48 the Danish National Patient Register is required by national law since 1978, and  
49 allocation of payment to health care providers is partly based on this coding. In  
50 general, the Danish National Patient Register is widely accepted as one of the world's  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 most valid health registries with several decades of follow-up.<sup>18</sup> However, a small  
4 private activity, especially regarding arthroscopy, might have eluded the registry until  
5 mandatory registration by private hospitals was introduced in 2003. Although this  
6 might have had some effect on the crude incidence of surgery, this effect would have  
7 been present in both groups. A further limitation is the observation period of 14.3  
8 years on average which may be too short to capture the full long-term consequences,  
9 and hence, our results may be conservative estimates. Moreover, ipsilateral and  
10 contralateral information of TKR and arthroscopy is missing as side-specific  
11 information was not a required part of data in the Danish National Patient Register.  
12 This is a limitation and as result, the ipsilateral risk of TKR and arthroscopy is likely  
13 higher than the risk estimates reported in the present study. Due to the nature of the  
14 study, information regarding fracture classification, treatment methods and outcome is  
15 not available. As a consequence, no subgroup analysis with regards to type of fracture  
16 and treatment modalities on the risk of TKR and arthroscopy is possible.

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36 In conclusion, patella fractures are associated with a considerably increased risk of  
37 total knee replacement and a highly increased risk of knee arthroscopy compared with  
38 an age- and gender-matched control group with no prior patella fracture.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**REFERENCES**

1. **Cram P, Lu X, Kates SL, Singh JA, Li Y, Wolf BR.** Total knee arthroplasty volume, utilization, and outcomes among Medicare beneficiaries, 1991-2010. *JAMA* 2012;308(12):1227–36.
2. **Court-Brown CM.** *Rockwood and Green's Fractures in adults*. USA: Walters Kluwer, 2015.
3. **Melvin JS, Mehta S.** Patellar fractures in adults. *J Am Acad Orthop Surg* 2011;19(4):198–207.
4. **Larsen P, Court-Brown CM, Vedel JO, Vistrup S, Elsoe R.** Incidence and Epidemiology of Patellar Fractures. *Orthopedics* 2016;39(6):e1154–e1158.
5. **Petrie J, Sassoan A, Langford J.** Complications of patellar fracture repair: treatment and results. *J Knee Surg* 2013;26(5):309–12.
6. **Mehdi M, Husson JL, Polard JL, Ouahmed A, Poncer R, Lombard J.** [Treatment results of fractures of the patella using pre-patellar tension wiring. Analysis of a series of 203 cases]. *Acta Orthop Belg* 1999;65(2):188–96.
7. **SORENSEN KH.** THE LATE PROGNOSIS AFTER FRACTURE OF THE PATELLA. *Acta Orthop Scand* 1964;34:198–212.
8. **Scolaro J, Bernstein J, Ahn J.** In brief: Patellar fractures. *Clin Orthop Relat Res* 2011;469(4):1213–1215.
9. **Larsen P, Vedel JO, Vistrup S, Elsoe R.** Long-Lasting Hyperalgesia Is Common in Patients Following Patella Fractures. *Pain Med* 2017;
10. **LeBrun CT, Langford JR, Sagi HC.** Functional outcomes after operatively treated patella fractures. *J Orthop Trauma* 2012;26(7):422–6.
11. **Anand S, Hahnel JCR, Giannoudis P V.** Open patellar fractures: high energy injuries with a poor outcome? *Injury* 2008;39(4):480–4.
12. **Vedel, JO; Vistrup, S; Larsen, P; Elsoe R.** Altered long-term health-related quality of life in patients following patella fractures: a long-term follow-up study of 49 patients. *Eur J Trauma Emerg Surg* 2017;
13. **Vedel JO, Vistrup S, Larsen P, Elsoe R.** Altered long-term health-related quality of life in patients following patella fractures: a long-term follow-up study of 49 patients. *Eur J Trauma Emerg Surg* 2017;
14. **Houdek MT, Shannon SF, Watts CD, Wagner ER, Sems SA, Sierra RJ.** Patella Fractures Prior to Total Knee Arthroplasty: Worse Outcomes but

- 1  
2  
3 Equivalent Survivorship. *J Arthroplasty* 2015;30(12):2167–9.
- 4  
5 15. **Honkonen SE.** Degenerative arthritis after tibial plateau fractures. *J Orthop*  
6 *Trauma* Tampere University Hospital, Finland., 1995;9(4):273–277.
- 7  
8 16. **Lynge E, Sandegaard JL, Rebolj M.** The Danish National Patient Register.  
9 *Scand J Public Health* Institute of Public Health, University of Copenhagen,  
10 Copenhagen, Denmark. elsebeth@pubhealth.ku.dk, 2011;39(7 Suppl):30–33.
- 11  
12 17. **Pedersen CB.** The Danish Civil Registration System. *Scand J Public Health*  
13 2011;39(7 Suppl):22–5.
- 14  
15 18. **Andersen JS, Olivarius NDF, Krasnik A.** The Danish National Health  
16 Service Register. *Scand J Public Health* 2011;39(7 Suppl):34–37.
- 17  
18 19. **Elm E von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC,**  
19 **Vandenbroucke JP, et al.** The Strengthening the Reporting of Observational  
20 Studies in Epidemiology (STROBE) Statement: guidelines for reporting  
21 observational studies. *Int J Surg* 2014;12(12):1495–9.
- 22  
23 20. **Andersen PK, Geskus RB, Witte T de, Putter H.** Competing risks in  
24 epidemiology: possibilities and pitfalls. *Int J Epidemiol* 2012;41(3):861–870.
- 25  
26 21. **Julin J, Jämsen E, Puolakka T, Konttinen YT, Moilanen T.** Younger age  
27 increases the risk of early prosthesis failure following primary total knee  
28 replacement for osteoarthritis. A follow-up study of 32,019 total knee  
29 replacements in the Finnish Arthroplasty Register. *Acta Orthop*  
30 2010;81(4):413–9.
- 31  
32 22. **Hartley RC, Barton-Hanson NG, Finley R, Parkinson RW.** Early patient  
33 outcomes after primary and revision total knee arthroplasty. A prospective  
34 study. *J Bone Joint Surg Br* 2002;84(7):994–9.
- 35  
36 23. **Baker P, Cowling P, Kurtz S, Jameson S, Gregg P, Deehan D.** Reason for  
37 revision influences early patient outcomes after aseptic knee revision. *Clin*  
38 *Orthop Relat Res* 2012;470(8):2244–52.
- 39  
40 24. **Delaunay C.** Registries in orthopaedics. *Orthop Traumatol Surg Res*  
41 2015;101(1 Suppl):S69-75.
- 42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *Legends:*  
4  
5

6 Table 1—Cox proportional hazards regression model of TKR comparing the exposed  
7 group with the non-exposed group.  
8  
9

10  
11 Table 2—Cox proportional hazards regression model of knee arthroscopy comparing  
12 the exposed group with the non-exposed group.  
13  
14


15  
16 Figure 1—Cumulative incidence of TKR

17 Legends: -- exposed group, -- non-exposed group  
18  
19

20  
21 Figure 2— Cumulative incidence of arthroscopy

22 Legends: -- exposed group, -- non-exposed group  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


Figure 1—Cumulative incidence of TKR



Legends: -- exposed group, -- non-exposed group



Figure 2– Cumulative incidence of arthroscopy



Legends: -- exposed group, -- non-exposed group

Table 1:

Cox proportional hazards regression of TKR comparing the exposed group with the non-exposed group

Follow-up time	N at beginning of interval	# events in interval	HR (95% CI)
0-5 years	66958	277	3.02 [2.26 - 4.03]
5-10 years	58917	454	1.56 [1.17 - 2.07]
10-15 years	52107	490	1.50 [1.13 - 2.00]
15-20 years	46083	216	1.67 [1.08 - 2.57]
0-20 years	66958	1437	1.83 [1.57 - 2.13]

For Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

up

For Review Only

Table 2

Cox proportional hazards regression of knee arthroscopy comparing the exposed group with the non-

Follow-up time	N at beginning of interval	# events in interval	HR (95% CI)
0-5 years	66680	661	7.40 [6.32 - 8.66]
5-10 years	58027	388	1.51 [1.11 - 2.05]
10-15 years	50913	233	1.57 [1.04 - 2.36]
15-20 years	44762	98	2.06 [1.15 - 3.69]
0-20 years	66680	1380	3.94 [3.49 - 4.46]


For Review Only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


-exposed group.

For Review Only

## SUPPLEMENTAL FIGURES 1-8


Supplemental Figure 1: *Cumulative incidence of TKR age 0-50*

Legends: -- exposed group, -- non-exposed group

Supplemental Figure 2: *Cumulative incidence of TKR age 50+*


Legends: -- exposed group, -- non-exposed group

Supplemental Figure 3: *Cumulative incidence of TKR in men*




Legends: -- exposed group, -- non-exposed group

Supplemental Figure 4: *Cumulative incidence of TKR in women*




Legends: -- exposed group, -- non-exposed group

Supplemental Figure 5: *Cumulative incidence of knee arthroscopy age 0-50*



Legends: -- exposed group, -- non-exposed group


Supplemental Figure 6: *Cumulative incidence of knee arthroscopy age 50+*



Legends: -- exposed group, -- non-exposed group




Supplemental Figure 7: *Cumulative incidence of knee arthroscopy in men*



Legends: -- exposed group, -- non-exposed group

Supplemental Figure 8: *Cumulative incidence of knee arthroscopy in women*



Legends: -- exposed group, -- non-exposed group

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Review Only