



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Clinical value of serum hyaluronan and propeptide of type III collagen in patients with pancreatic cancer

Chen, Inna Markovna; Willumsen, Nicholas; Dehlendorff, Christian; Johansen, Astrid Zedlitz; Jensen, Benny Vittrup; Hansen, Carsten Palnaes; Hasselby, Jane Preuss; Bojesen, Stig Egil; Pfeiffer, Per; Nielsen, Svend Erik; Holländer, Niels Henrik; Yilmaz, Mette Karen; Karsdal, Morten; Johansen, Julia Sidenius

*Published in:*  
International Journal of Cancer

*DOI (link to publication from Publisher):*  
[10.1002/ijc.32751](https://doi.org/10.1002/ijc.32751)

*Publication date:*  
2020

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Chen, I. M., Willumsen, N., Dehlendorff, C., Johansen, A. Z., Jensen, B. V., Hansen, C. P., Hasselby, J. P., Bojesen, S. E., Pfeiffer, P., Nielsen, S. E., Holländer, N. H., Yilmaz, M. K., Karsdal, M., & Johansen, J. S. (2020). Clinical value of serum hyaluronan and propeptide of type III collagen in patients with pancreatic cancer. *International Journal of Cancer*, 146(10), 2913-2922. Advance online publication. <https://doi.org/10.1002/ijc.32751>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

To be submitted to International Journal of Cancer as an original manuscript

**Title: Clinical value of serum hyaluronan and propeptide of type III collagen in patients with pancreatic cancer**

**Running title:** Hyaluronan and propeptide of type III collagen in patients with pancreatic cancer

**Authors:**

Inna Markovna Chen<sup>1\*</sup>, Nicholas Willumsen<sup>2</sup>, Christian Dehlendorff<sup>3</sup>, Astrid Zedlitz Johansen<sup>1</sup>, Benny Vittrup Jensen<sup>1</sup>, Carsten Palnæs Hansen<sup>4</sup>, Jane Preuss Hasselby<sup>5</sup>, Stig Egil Bojesen<sup>6,7</sup>, Per Pfeiffer<sup>8</sup>, Svend Erik Nielsen<sup>1,9</sup>, Niels Henrik Holländer<sup>10</sup>, Mette Karen Yilmaz<sup>11</sup>, Morten Karsdal<sup>2</sup>, Julia Sidenius Johansen<sup>1,7,12</sup>

**Affiliations:**

<sup>1</sup>Department of Oncology, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark.

<sup>2</sup>Nordic Bioscience A/S, Herlev, Denmark.

<sup>3</sup>Unit of Statistics and Pharmacoepidemiology, Danish Cancer Society Research Center, Copenhagen, Denmark.

<sup>4</sup>Department of Surgery, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark.

<sup>5</sup>Department of Pathology, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark.

<sup>6</sup>Department of Clinical Biochemistry, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ijc.32751

<sup>7</sup>Institute of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark.

<sup>8</sup>Department of Oncology, Odense University Hospital, Odense, Denmark.

<sup>9</sup>Department of Oncology, North Zealand University Hospital, Hillerød, Denmark.

<sup>10</sup>Department of Oncology, Zealand University Hospital, Næstved, Denmark.

<sup>11</sup>Department of Oncology, Aalborg University Hospital, Aalborg, Denmark.

<sup>12</sup>Department of Medicine, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev, Denmark.

**Correspondence addressed to:**

Inna Markovna Chen: Department of Oncology, Herlev Hospital, Borgmester Ib Juuls Vej 1, DK-2730 Herlev, Denmark. Telephone: +45 38682898. E-mail: inna.chen@regionh.dk

**Keywords:** Biomarkers, collagen type III, diagnostic biomarker, hyaluronan, pancreatic cancer, PRO-C3, prognostic biomarker.

**Abbreviations:**

AC	Ampullary carcinoma
AUC	Area under the ROC
BIOPAC	BIOMarkers in patients with pancreatic cancer
BMI	Body mass index
CI	Confidence interval
CP	Chronic pancreatitis
CRP	C-reactive protein
CV	Coefficient of variations
DBTC	Distal biliary tract cancer
ELISA	Enzyme-linked immunosorbent assay

FOLFIRINOX	5-FU, leucovorin, irinotecan, and oxaliplatin
ECM	Extracellular matrix
HA	Hyaluronic acid or hyaluronan
DA	Duodenal adenoma
HR	Hazard ratio
IFP	Interstitial fluid pressure
IL-6	Interleukin-6
IPMN	Intraductal papillary mucinous neoplasm
OS	Overall survival
PC	Pancreatic cancer
PS	Performance status
PRO-C3	Propeptide of type III collagen
ROC	Receiver operating characteristics curve
TME	Tumor microenvironment
YKL-40	Chitinase 3-like 1 protein (CHI3L1)

**“Novelty and Impact” What’s new? (max. 75 words)**

Hyaluronan (HA) and collagen are highly expressed in pancreatic cancer (PC) stroma. HA and collagen accumulation increase tumor interstitial fluid pressure, compromising blood flow and drug penetration. High levels of serum HA and propeptide of type III collagen (PRO-C3) were associated with short overall survival. Serum HA and PRO-C3 were higher in PC patients than patients with benign conditions. The findings warrant further exploration to help clarify links between HA, collagen type III, and PC.

## Abstract

Hyaluronan (HA) and collagen are highly expressed in pancreatic cancer (PC) stroma. HA and collagen accumulation increase tumor interstitial fluid pressure, compromising blood flow and drug penetration. The aim of this biomarker study was to determine the clinical utility of serum HA and the propeptide of type III collagen (PRO-C3) in patients with PC. A cohort from the Danish BIOPAC study (NCT03311776) including patients with histologically confirmed pancreatic ductal adenocarcinoma (n=809), ampullary carcinoma (n=44), distal biliary tract cancer (n=31), chronic pancreatitis (n=15), intraductal papillary mucinous neoplasm (n=41), duodenal adenoma (n=7), and no cancer (n=25). Healthy controls were available for serum HA (n=141) and PRO-C3 (n=8). The main outcome was overall survival (OS) of patients with PC in relation to pretreatment serum HA and PRO-C3 levels. Patients with PC had higher baseline serum HA and PRO-C3 than healthy subjects and patients with benign conditions. Pretreatment serum baseline HA and PRO-C3 in patients with PC were associated with poorer survival and PRO-C3 remained prognostic also after adjusting for age, performance status, stage, presence of liver and peritoneum metastasis, and CA19-9. Detection of HA and PRO-C3 may be useful in differentiating between malignant and benign pancreatic conditions. Serum HA and PRO-C3 were prognostic for OS in patients with PC.

## Introduction

Pancreatic cancer (PC) is one of the most lethal of all major cancers<sup>11, 36, 40</sup>. Pancreatic ductal adenocarcinomas account for 90% of all PC tumors and are highly resistant to all standard treatments including chemotherapy, surgery, radiation, and targeted agents, partly due to the multifaceted and immunosuppressive tumor microenvironment (TME)<sup>4, 6, 39, 49</sup>. The prognosis has only improved slightly during the last decades, and the overall 5-year survival remains low at 8%<sup>36</sup>. PC is characterized by pronounced inflammation and disperse tumor cells in a dense desmoplastic stroma consisting of pancreatic stellate cells (PSCs), carcinoma-associated fibroblasts, extracellular matrix (ECM), endothelial cells, and immune cells. Interaction between stroma and inflammation is reported<sup>3, 13, 33, 43, 46</sup>. Stroma contributes to the hypoxic and avascular microenvironment in both primary tumors and metastases<sup>10, 49</sup>. However, stromal biology and its prognostic impact are still under discussion<sup>44</sup>. Multiple stroma-targeted therapies are under active investigation<sup>21, 47</sup>.

Hyaluronic acid (HA), also called hyaluronan, and collagen are major components of the ECM of the tumor stroma in PC, and high levels of each are found in both primary tumors and metastatic lesions<sup>17, 20, 49</sup>. HA is a glycosaminoglycan that is synthesized at high levels by cancer cells and fibroblasts in the TME and promotes tumor growth, angiogenesis, inflammation, immunosuppression as well as preserves growth factors and cytokines<sup>20, 23, 38, 42</sup>.

The collagen types I, III, and IV are concomitantly overexpressed together with HA in PC<sup>49</sup>. Collagen type IV is the major component of the basement membrane, a specialized ECM that underlies the epithelial and endothelial cell layer and plays a major role in tumor cell invasion and angiogenesis. Collagen type I and type III are the major components of the stromal ECM and the interstitial matrix. They are produced by PSCs within the desmoplastic stroma and promote PC cell adhesion, proliferation, migration, and chemo-resistance<sup>2, 14, 24, 50, 54</sup>. The ECM in the tumor stroma undergoes

Accepted Article

excessive remodeling. The increased activity of PSC/fibroblasts leads to excessive formation and accumulation of interstitial collagens<sup>27</sup>. HA and collagen accumulation increase tumor solid stress and interstitial fluid pressure (IFP), compromising blood vessels, circulation, perfusion, and drug penetration<sup>8, 25, 29, 37, 42</sup>. The prognostic value of HA and collagen deposition in PC tissue has been described in a few studies<sup>5, 7, 12, 35, 49</sup>. HA was hypothesized to be a potential therapeutic target to decrease tumor IFP and enhance drug delivery and immune cells' access into the tumor<sup>31, 34, 37</sup>. Reprogramming of PSC and downregulation of collagen-related signals by the vitamin D receptor ligand, calcipotriol, have been shown to improve gemcitabine penetration into the tumor, resulting in prolonged survival in PC mouse models<sup>35</sup>.

Promising results of targeting of HA in combination with chemotherapy were reported in a randomized phase II study, which showed that patients with high HA levels in the tumor microenvironment had a largest improvement in progression free survival after addition of pegvorhialuronidase alfa (PEGPH20) to a standard combination of nab-paclitaxel/gemcitabine<sup>15</sup>. Preliminary data based on retrospective analysis of this trial suggested that type III collagen turnover measured in serum was able to identify patients that would benefit from the addition of PEGPH20<sup>48</sup>.

A phase III study targeting HA is ongoing (NCT02715804). Although HA and collagen expression in the TME are considered prognostic biomarkers, there are no/few studies examining serum concentrations of HA and type III collagen in patients with PC. Previous studies conducted in other malignancies have shown a negative correlation between serum HA and outcome<sup>1, 19, 28, 45, 52, 53</sup> and serum collagen fragments and outcome<sup>7, 18, 22, 51</sup>. The aim of this biomarker study was to investigate the clinical utility of serum HA and propeptide of type III collagen (PRO-C3) in patients with PC.

## **Materials and Methods**

### **Patients**

We analyzed pretreatment serum samples from a cohort of patients included in the Danish BIOPAC study “Biomarkers in Patients with Pancreatic Cancer” (NCT03311776) from six hospitals in Denmark admitted from December 2008 until September 2017. A total of 972 participants, including consecutive BIOPAC patients with histologically confirmed PC, ampullary carcinoma (AC), distal biliary tract cancer (DBTC), chronic pancreatitis (CP), intraductal papillary mucinous neoplasm (IPMN), patients subjected to surgery in whom PC was suspected but not histologically confirmed (no pathologic finding), and patients with duodenal adenomas (DA), were studied.

Serum samples and clinical data from patients were collected prospectively. Retrospective HA and PRO-C3 analyses and the clinical data review of electronic medical records were performed separately (or independent of each other). Cancer patients were treated with various types of chemotherapy according to national guidelines [www.gicancer.dk](http://www.gicancer.dk).

### **Biological samples**

Blood samples were obtained at time of diagnosis and/or before operation if available. Sequential samples collected during and after treatment (before 2nd cycle and at the time of first CT scan) were available for the group of patients receiving palliative chemotherapy. Samples were processed according to the nationally approved standard operation procedures for blood ([www.herlevhospital.dk/biopac.dk](http://www.herlevhospital.dk/biopac.dk)).

### **Serum HA analysis**



Pre-treatment and longitudinal serum concentrations of HA were determined by a human HA high-sensitive solid-phase immunoassay (Quantikine® Immunoassay, R&D Systems, Abingdon, OX, UK) in accordance with the manufacturer's instructions. The lower level of detection (LLOD) for HA was 0.068 ng/mL. The inter-assay coefficient of variation (CV) was  $\leq 7.2\%$  and the intra-assay CV was  $\leq 4.9\%$ . The molecular weight range of HA detected by the Quantikine kit was 0.2 – 40.0 ng/mL. The median serum HA in 141 healthy blood donors was 41 ng/ml.

### **Serum PRO-C3 analysis**

Serum PRO-C3 was determined by enzyme-linked immunosorbent assay (ELISA) using a monoclonal antibody to detect the N-protease mediated cleavage of the N-terminal propeptide of type III collagen (Nordic Bioscience, PRO-C3 assay)<sup>26</sup>. The LLOD was 4.0 ng/mL. The inter- and intra-CVs were 11.0% and 4.1%. The molecular weight range of PRO-C3 detected by the Nordic Bioscience ELISA kit was 1.1 – 8.5 ng/mL. The median serum PRO-C3 level in eight healthy subjects was 7 ng/mL.

### **ETHICAL ISSUES**

The BIOPAC study was approved by the Regional Ethics Committee (VEK ref. KA-20060113) and the Danish Data Protection Agency (j.nr. 2006-41-6848). Informed consent for research use was obtained from all patients at the enrolling hospitals before prospective serum banking. Serum HA and PRO-C3 assessments were not predefined in the protocol but approval was obtained before analysis. The responsible investigators ensured that the clinical studies were conducted in agreement with the Declaration of Helsinki, version 8.

## Statistical analysis

Results are reported in accordance with the REMARK (reporting recommendations for tumor marker prognostic study) guidelines<sup>32</sup>. The objective was to examine the clinical value of serum HA and PRO-C3 and the association between baseline and longitudinal serum HA levels and overall survival (OS) in the PC patients. Patients were followed from the date of inclusion in the BIOPAC study to the end of follow-up (October 24, 2018) or until death from any cause, whichever came first.

Descriptive statistics were performed to describe patients' demographics and clinical baseline characteristics. Receiver operating characteristic (ROC) analysis was conducted to assess the diagnostic values of serum HA. Changes of serum HA and PRO-C3 over time during and after systemic treatment and their prognostic value were investigated. Hazard ratios (HRs) adjusted for baseline characteristics (age, gender, performance status (PS), stage, history of diabetes, smoking and alcohol abuse, serum CA19-9, as well as chemotherapy regimen (gemcitabine, mFOLFIRINOX, gemcitabine with nab-paclitaxel, or other), and body mass index (BMI) were estimated with Cox proportional-hazard regression. Kaplan–Meier curves were used to illustrate the survival curves in patients with PC in relation to measured values of serum HA and PRO-C3 levels, comparing patients with the 25% highest values with those with the 75% lowest for each biomarker separately. The association between changes in serum HA and PRO-C3 during palliative chemotherapy and OS was analyzed in patients in whom sequential (baseline, before 2nd cycle, and at the time of first CT scan) data were available. Differences in serum concentration over time from baseline were measured; negative values correspond to a decrease in serum HA and PRO-C3. Patients were grouped according to change from baseline. Statistical analyses were conducted using the latest version of R (currently 3.2.4).

## **Data availability**

The data that support the findings of this study are available upon reasonable request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

## **Results**

### **Characteristics of study population**

The study included 972 participants with the following diagnoses: histologically confirmed PC (n=809), AC (n=44), DBTC (n=31), CP (n=15), IPMN (n=41), no pathologic finding (n=25), DA (n=7 including six patients with high-grade dysplasia and one patient with ungraded dysplasia). Clinicopathological data of PC patients are shown in Table 1. The median age was 67 years (range 37–89 years), and most of the PC patients had PS 0 or 1. The majority of PC patients had advanced stage III or IV disease, had normal weight, and were registered as present or earlier smokers. Fifty-one patients across all stages were never treated with chemotherapy. Of 598 patients who received palliative chemotherapy, 404 (67.6%) were treated with gemcitabine, 120 (20.1%) with FOLFIRINOX (5-FU, leucovorin, irinotecan, and oxaliplatin), 55 (9.2%) with combination of gemcitabine and nab-paclitaxel, and 19 (3.2%) with other combinations.

### **Pretreatment serum HA and PRO-C3 and clinical characteristics and overall survival**

First, we compared serum HA levels between the study groups (Table 2). The median pretreatment HA level was 84 ng/mL (IQR: 48-131). Pretreatment serum HA in patients with cancer including PC, AC, DBTC was increased compared to patients with benign conditions or healthy controls. The median serum HA level in PC patients was about two-fold higher than in healthy controls ( $p<0.001$ )

and was increased compared to patients with benign conditions including CP, IPMN, DA and those subjects without pathological findings ( $p<0.001$ ). No statistically significant difference in HA was observed comparing PC with other malignant diseases. We further stratified the PC patients according to stage, PS, and presence of liver metastases. Higher serum HA levels were associated with metastatic disease, presence of liver metastases, and worse PS (Table 2). Thirty-four percent of patients with liver metastases had the highest quartile of serum HA compared to 19% patients without liver metastases ( $p<0.001$ ). Correlation analyses displayed a pairwise positive relation between baseline serum HA and PRO-C3, CA19-9 and inflammation markers in patients with PC (Table S1). ROC analysis showed that serum HA level had a better potential to discriminate PC patients from healthy controls (AUC=0.75, 95% CI 0.72–0.79, Figure 1A) than from patients with benign conditions (AUC=0.63, 95% CI 0.57–0.69, Figure 1B).

We subsequently investigated the clinical value of serum PRO-C3 in the same cohort. The median pretreatment PRO-C3 level was 18 ng/mL (IQR: 11-33). Serum PRO-C3 was higher in patients with malignant diseases than in those with benign conditions (Table 2). PC patients had higher median serum PRO-C3 levels compared to patients with benign conditions ( $p<0.001$ ). PRO-C3 was positively correlated with higher stage and presence of liver metastases ( $p<0.001$ ). Double as many patients with liver metastases had serum PRO-C3 in the highest quartile compared to patients without liver metastases (37% versus 18%,  $p<0.001$ ).  $PS\geq 2$  was associated with higher serum PRO-C3 values. Detection of PRO-C3 performed slightly better as a diagnostic marker than serum HA. ROC analyses comparing PC patients with healthy controls and patients with benign conditions yielded an AUC=0.87, 95% CI 0.71–1.00 (Figure 1C) and AUC=0.71, 95% CI 0.65–0.76 (Figure 1D).

Overall survival according to levels of HA and PRO-3C is shown in Figure 2. Univariate analysis showed that high baseline serum HA (1.18, 95% CI: 1.13–1.25) and PRO-C3 (1.28, 95%: CI 1.11–

1.49) were associated with worse prognosis. After allocating patients into stages, a significance association was still seen for patients with stage IV (Figure 2). High baseline HA and PRO-C3 levels according to types of treatments in 579 patients undergoing palliative chemotherapy were associated with short survival (Figures S1). After adjusting factors associated with OS, the HRs for serum HA and PRO-C3 were 1.04 (95% CI: 0.97–1.13) and 1.42 (95% CI: 1.14–1.76), respectively (Table 3). Further adjustment for inflammatory markers (CRP, IL-6, and YKL-40) changed the HRs to 1.02 (95% CI: 0.93–1.11) and 1.35 (95% CI: 1.05–1.73) for serum HA and PRO-C3, respectively (Table S2 and Supplementary Materials & Methods). Patients with combined 25% highest levels of both serum HA and PRO-C3 had the poorest survival (Figure S2).

#### **Prognostic value of changes in serum HA and PRO-C3 over time**

Serum HA levels decreased after the first treatment cycle, with a median difference of  $-11.50$  [IQR,  $-41.98, 12.23$ ] ng/mL, and at the time of first CT evaluation 2 or 3 months after chemotherapy start of  $-34.10$  [IQR,  $-72.95, -5.25$ ] ng/mL compared to the HA level at baseline. In contrast, no decrease was observed for longitudinal serum PRO-C3 concentrations:  $-0.90$  [IQR,  $-3.80, 3.05$ ] and  $-1.90$  [IQR,  $-9.30, 4.30$ ] after the first treatment cycle and at the time of first CT evaluation, respectively. Changes in HA and PRO-3C were not associated with overall survival (Figure 3).

## Discussion

PC remains one of the most aggressive tumors and is known to be highly enriched with ECM in the desmoplastic stroma, and HA and collagen are essential components of the ECM. Biomarker availability, reliability, and function of the serum levels of HA and N-terminal propeptide of type III collagen (PRO-C3) in patients with PC have not been extensively investigated so far. In this study, we demonstrated that higher values of serum PRO-C3 and HA at the time of diagnosis are associated with an increased mortality. However, serum HA was not associated with OS after adjusting for potential confounding factors. This suggests that serum PRO-C3 may serve as prognostic marker in patients with PC. In support for this hypothesis, collagen turnover rate needs to be maintained in normal tissue homeostasis, and changes in the deposition of collagens result in the tissue fibrosis that drives tumorigenesis<sup>41</sup>. Hence type III collagen synthesis may be a major component of the dense stroma (desmoplastic reaction) in the tumor and linked to tumor growth, metastasis, and drug resistance.

Our findings are in line with reports on the facilitating role of HA and PRO-C3 in the malignant behavior of tumor cells and the IFP increase within a tumor, limiting perfusion and compromising drug delivery<sup>8, 25, 29, 37, 42</sup>. Analyses of HA and collagen levels in tumors from patients with PC revealed that the survival was worse among patients with high HA and collagen accumulation compared to subjects with low HA and collagen expression, suggesting that HA and collagen levels in tumors of patients with PC may be predictive of survival<sup>5, 49</sup>. In contrast, prognostic significant relevance was not observed for HA accumulation, found in the majority of the PDAC-resected tumors<sup>12</sup>. Surgical examination of tissues from 114 PC patients identified a particular rearrangement of collagen fibers surrounding the tumor as a "biomarker" of early death<sup>7</sup>. By staining 53 patients' primaries and 57 patients' metastases, Whatcott et al. found highly fibrotic stroma with high levels of HA and collagen type I in both primary tumors and metastases<sup>49</sup>. PDAC patients with a high

Accepted Article

expression of HA were more susceptible to therapy targeting HA<sup>15</sup>. Addition of HA blockade to FOLFIRINOX did not show any benefit in patients unselected for tumor HA status<sup>30</sup>. Interestingly, in our study the more pronounced association with survival of baseline HA according to types of treatments in patients undergoing palliative chemotherapy was observed in participants treated with gemcitabine and nab-paclitaxel combination (Figures S1). However, due to the small number of patients undergoing combination palliative chemotherapy the survival data should be viewed with caution. A phase III trial investigating the efficacy of HA blockade in combination with gemcitabine and nab-paclitaxel in PDAC is pending (<https://clinicaltrials.gov/ct2/show/NCT02715804>). Collagen type III in PC was found to be linked to fibrosis within ECM and associated with poor prognosis and tumorigenesis<sup>16, 26</sup>. Targeting of collagen compartments in combination with various drugs is under extensive investigation.

We demonstrated that serum HA was increased in PC patients from the BIOPAC cohort compared with patients with benign conditions and healthy subjects. Likewise, a significantly higher concentration of serum PRO-C3 was observed in PC patients compared to patient with non-malignant conditions. Specific protease-generated fragments of collagen were significantly elevated in serum from patients with PC compared to controls, and a combination of serum biomarkers reflecting matrix metalloproteinase-mediated collagen turnover resulted in complete separation between healthy controls and patients with PC<sup>50</sup>.

We also demonstrated that high serum HA and PRO-C3 levels at diagnosis were not restricted to patients with metastatic PC but were present in all PC patients regardless of stage. However, our study suggests an association between higher concentrations of serum HA and PRO-C3 levels and advanced stage of disease, many metastatic sites, as well as presence of liver metastases. This is in agreement with the suggestion that the increase in matrix components is due to the ECM degradation that occurs

during metastasizing<sup>9</sup>. Furthermore, our findings were compared to patients with other upper gastrointestinal cancer diagnoses, including AC and DBTC. No significant difference was observed for AC and DBTC compared to patients with PC. Finally, serum HA and PRO-C3 also correlated with serum CRP, IL-6, and YKL-40, suggesting mutual interplay between inflammation and stroma. While treatment of patients with PC is based on a balance between tolerability and benefit, the optimization of non-invasive cost-effective monitoring would be of useful clinical value. Prognosis of PC patients often displays a dynamic pattern during systemic treatment. We observed a reduction in serum HA but not PRO-C3 levels in longitudinal samples; however, neither a decrease in HA nor in PRO-C3 was associated with survival. Pretreatment serum HA and PRO-C3 in patients undergoing palliative chemotherapy are more robustly associated with survival than serum HA and PRO-C3 after first cycle chemotherapy and at the time of the first evaluation CT scanning. This suggests that pretreatment serum HA and PRO-C3 would be more useful in identifying patients at risk of poor survival due to primary aggressive biology. However, due to the small number of patients with available longitudinal serum samples, natural selection, and the limited number of patients who received gemcitabine with nab-paclitaxel or FOLFIRINOX, the survival data according to serum HA and PRO-C3 changes should be interpreted with caution. To our knowledge, this is the first large-scale study to describe serum HA and PRO-C3 in samples derived from population-based consecutive patients with PC that can be applied to future research addressing stroma and ECM.

Several limitations must be noted. First, causal relationships cannot be determined in an observational study, and residual confounding is always a concern. However, prospective well-characterized data sets with clinico-pathological characteristics and treatment data on consecutive patients across the different centers including patients with PC in the BIOPAC study in a well-defined period of time reduce residual confounding. Second, staging is based on established routine practice, and it is well



known that clinical staging systems are not as clear-cut. However, we demonstrated consistent associations between serum HA and PRO-C3 levels with survival across all stages. Third, selection bias may affect the generalizability of the results since only around 20% of all registered Danish patients with PC are included in the BIOPAC study. Thus, many patients are not referred to systemic therapy due to rapid deterioration and die within a short time after diagnosis. Enrolled patients were more likely to have better PS, and more likely to receive chemotherapy. Thus, it is possible that the relationship between serum HA and PRO-C3 and survival in this subset is not generalizable to all patients with PC. Finally, whether serum HA and PRO-C3 values are correlated with HA and PRO-C3 expression in tumors has not yet been determined.

In conclusion, we have shown that high serum HA and PRO-C3 levels are associated with increased mortality in patients with PC. Moreover, compared to healthy subjects and a group of patients with benign conditions, high values of these parameters are measured in patients with malignant lesions, including PC, AC, and DBTC. Serum HA and PRO-C3 were associated with PC stage. Our findings suggest that serum HA and PRO-C3 levels may have a clinical value and motivate further research on serum HA and PRO-C3 as novel, minimally invasive and cost-effective prognostic and diagnostic markers of PC. Our study also supports evidence that encourages ongoing investigation of stroma targeting strategies.

## **Declarations**

### **Consent for publication**

Not applicable.

### **Competing interests**

NWI, MK are employees at Nordic Bioscience involved in biomarker discovery and development.

MK holds stocks of Nordic Bioscience. All the other authors do not have any conflict to declare.

### **Funding**

No external funding was used in the establishment of the BIOPAC study (Biobank and Database) and the continued collection of biological samples and clinical data. Biomarker analysis was funded by the Department of Oncology, Herlev and Gentofte Hospital, Denmark.

### **Authors' contributions**

All authors contributed to inclusion of patients in the BIOPAC study. AZJ, IMC, and JSJ entered the clinical data in the database. IMC drafted and wrote the manuscript and revised the manuscript after feedback from all authors. All authors contributed to review of the present manuscript and approved the final version of the manuscript.

### **Acknowledgements**

Many thanks to the biomedical laboratory scientists Charlotte Falk, Vibeke H. Holm, Dorte B. Nielsen, Helle B. Sørensen, and Mie B. Krüger for their assistance with the blood sampling and analysis. The nurses Hanne M. Michaelsen, Lars M. Larsen, Lene T. Voxen, Finn Hejlesen, Signe

Gamborg Nielsen, Jette L. Hansen, the physician Dan Calatayud, the and medical students Camilla Palmquist and Karin Bagni are acknowledged for their contribution to inclusions of patients, data registration, or sample collection. We thank Edwin Stanton Spencer for English language editing.

**Provenance and peer review**

Not commissioned; external peer review.

Accepted Article

## References

- 1 Anagnostopoulou E, Papanastasopoulou C, Papastamataki M, Kotsiou A, Topouzoglou Z, Anagnostopoulos N *et al.* Serum Hyaluronic Acid Levels Are Altered in Acute Leukemia Patients: Potential Prognostic Implications. *Acta Haematol* 2017; 138: 44-51.
- 2 Apte MV, Park S, Phillips PA, Santucci N, Goldstein D, Kumar RK *et al.* Desmoplastic reaction in pancreatic cancer: role of pancreatic stellate cells. *Pancreas* 2004; 29: 179-187.
- 3 Babic A, Schnure N, Neupane NP, Zaman MM, Rifai N, Welch MW *et al.* Plasma inflammatory cytokines and survival of pancreatic cancer patients. *Clin Transl Gastroenterol* 2018; 9: 145.
- 4 Bardeesy N, DePinho RA. Pancreatic cancer biology and genetics. *Nature reviews Cancer* 2002; 2: 897-909.
- 5 Cheng XB, Sato N, Kohi S, Yamaguchi K. Prognostic impact of hyaluronan and its regulators in pancreatic ductal adenocarcinoma. *PLoS One* 2013; 8: e80765.
- 6 Clark CE, Hingorani SR, Mick R, Combs C, Tuveson DA, Vonderheide RH. Dynamics of the immune reaction to pancreatic cancer from inception to invasion. *Cancer research* 2007; 67: 9518-9527.
- 7 Drifka CR, Loeffler AG, Mathewson K, Keikhosravi A, Eickhoff JC, Liu Y *et al.* Highly aligned stromal collagen is a negative prognostic factor following pancreatic ductal adenocarcinoma resection. *Oncotarget* 2016; 7: 76197-76213.
- 8 DuFort CC, DelGiorno KE, Hingorani SR. Mounting Pressure in the Microenvironment: Fluids, Solids, and Cells in Pancreatic Ductal Adenocarcinoma. *Gastroenterology* 2016; 150: 1545-1557 e1542.
- 9 El-Mezayen HA, Toson el SA, Darwish H, Metwally FM. Development of a novel metastatic breast cancer score based on hyaluronic acid metabolism. *Medical oncology (Northwood, London, England)* 2013; 30: 404.
- 10 Feig C, Gopinathan A, Neesse A, Chan DS, Cook N, Tuveson DA. The pancreas cancer microenvironment. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2012; 18: 4266-4276.
- 11 Ferlay J, Partensky C, Bray F. More deaths from pancreatic cancer than breast cancer in the EU by 2017. *Acta oncologica (Stockholm, Sweden)* 2016; 55: 1158-1160.
- 12 Gebauer F, Kemper M, Sauter G, Prehm P, Schumacher U. Is hyaluronan deposition in the stroma of pancreatic ductal adenocarcinoma of prognostic significance? *PLoS One* 2017; 12: e0178703.

- 13 Greer JB, Whitcomb DC. Inflammation and pancreatic cancer: an evidence-based review. *Curr Opin Pharmacol* 2009; 9: 411-418.
- 14 Hamada S, Masamune A. Elucidating the link between collagen and pancreatic cancer: what's next? *Expert Rev Gastroenterol Hepatol* 2018; 12: 315-317.
- 15 Hingorani SR, Zheng L, Bullock AJ, Seery TE, Harris WP, Sigal DS *et al.* HALO 202: Randomized Phase II Study of PEGPH20 Plus Nab-Paclitaxel/Gemcitabine Versus Nab-Paclitaxel/Gemcitabine in Patients With Untreated, Metastatic Pancreatic Ductal Adenocarcinoma. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2018; 36: 359-366.
- 16 Imamura T, Iguchi H, Manabe T, Ohshio G, Yoshimura T, Wang ZH *et al.* Quantitative analysis of collagen and collagen subtypes I, III, and V in human pancreatic cancer, tumor-associated chronic pancreatitis, and alcoholic chronic pancreatitis. *Pancreas* 1995; 11: 357-364.
- 17 Jacobetz MA, Chan DS, Neesse A, Bapiro TE, Cook N, Frese KK *et al.* Hyaluronan impairs vascular function and drug delivery in a mouse model of pancreatic cancer. *Gut* 2013; 62: 112-120.
- 18 Jensen C, Madsen DH, Hansen M, Schmidt H, Svane IM, Karsdal MA *et al.* Non-invasive biomarkers derived from the extracellular matrix associate with response to immune checkpoint blockade (anti-CTLA-4) in metastatic melanoma patients. *J Immunother Cancer* 2018; 6: 152.
- 19 Kehlet SN, Sanz-Pamplona R, Brix S, Leeming DJ, Karsdal MA, Moreno V. Excessive collagen turnover products are released during colorectal cancer progression and elevated in serum from metastatic colorectal cancer patients. *Sci Rep* 2016; 6: 30599.
- 20 Knudson CB, Knudson W. Hyaluronan-binding proteins in development, tissue homeostasis, and disease. *FASEB J* 1993; 7: 1233-1241.
- 21 Kota J, Hancock J, Kwon J, Korc M. Pancreatic cancer: Stroma and its current and emerging targeted therapies. *Cancer letters* 2017; 391: 38-49.
- 22 Lipton A, Leitzel K, Ali SM, Polimera HV, Nagabhairu V, Marks E *et al.* High turnover of extracellular matrix reflected by specific protein fragments measured in serum is associated with poor outcomes in two metastatic breast cancer cohorts. *International journal of cancer* 2018; 143: 3027-3034.
- 23 Mahlbacher V, Sewing A, Elsasser HP, Kern HF. Hyaluronan is a secretory product of human pancreatic adenocarcinoma cells. *Eur J Cell Biol* 1992; 58: 28-34.

- 24 Menke A, Philippi C, Vogelmann R, Seidel B, Lutz MP, Adler G *et al.* Down-regulation of E-cadherin gene expression by collagen type I and type III in pancreatic cancer cell lines. *Cancer research* 2001; 61: 3508-3517.
- 25 Minchinton AI, Tannock IF. Drug penetration in solid tumours. *Nature reviews Cancer* 2006; 6: 583-592.
- 26 Nielsen MJ, Nedergaard AF, Sun S, Veidal SS, Larsen L, Zheng Q *et al.* The neo-epitope specific PRO-C3 ELISA measures true formation of type III collagen associated with liver and muscle parameters. *Am J Transl Res* 2013; 5: 303-315.
- 27 Nissen NI, Karsdal M, Willumsen N. Collagens and Cancer associated fibroblasts in the reactive stroma and its relation to Cancer biology. *J Exp Clin Cancer Res* 2019; 38: 115.
- 28 Peng C, Wallwiener M, Rudolph A, Cuk K, Eilber U, Celik M *et al.* Plasma hyaluronic acid level as a prognostic and monitoring marker of metastatic breast cancer. *International journal of cancer* 2016; 138: 2499-2509.
- 29 Provenzano PP, Cuevas C, Chang AE, Goel VK, Von Hoff DD, Hingorani SR. Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma. *Cancer cell* 2012; 21: 418-429.
- 30 Ramanathan RK, McDonough SL, Philip PA, Hingorani SR, Lacy J, Kortmansky JS *et al.* Phase IB/II Randomized Study of FOLFIRINOX Plus Pegylated Recombinant Human Hyaluronidase Versus FOLFIRINOX Alone in Patients With Metastatic Pancreatic Adenocarcinoma: SWOG S1313. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2019; 37: 1062-1069.
- 31 Rosengren S, Clift R, Zimmerman SJ, Souratha J, Thompson BJ, Blouw B *et al.* Abstract 4886: PEGylated recombinant hyaluronidase PH20 (PEGPH20) enhances checkpoint inhibitor efficacy in syngeneic mouse models of cancer 2016; 76: 4886-4886.
- 32 Sauerbrei W, Taube SE, McShane LM, Cavenagh MM, Altman DG. Reporting Recommendations for Tumor Marker Prognostic Studies (REMARK): An Abridged Explanation and Elaboration. *Journal of the National Cancer Institute* 2018; 110: 803-811.
- 33 Schultz NA, Christensen IJ, Werner J, Giese N, Jensen BV, Larsen O *et al.* Diagnostic and Prognostic Impact of Circulating YKL-40, IL-6, and CA 19.9 in Patients with Pancreatic Cancer. *PLoS One* 2013; 8: e67059.
- 34 Shepard HM. Breaching the Castle Walls: Hyaluronan Depletion as a Therapeutic Approach to Cancer Therapy. *Front Oncol* 2015; 5: 192.

- 35 Sherman MH, Yu RT, Engle DD, Ding N, Atkins AR, Tiriach H *et al.* Vitamin D receptor-mediated stromal reprogramming suppresses pancreatitis and enhances pancreatic cancer therapy. *Cell* 2014; 159: 80-93.
- 36 Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019; 69: 7-34.
- 37 Singha NC, Nekoroski T, Zhao C, Symons R, Jiang P, Frost GI *et al.* Tumor-associated hyaluronan limits efficacy of monoclonal antibody therapy. *Mol Cancer Ther* 2015; 14: 523-532.
- 38 Spinelli FM, Vitale DL, Demarchi G, Cristina C, Alaniz L. The immunological effect of hyaluronan in tumor angiogenesis. *Clin Transl Immunology* 2015; 4: e52.
- 39 Stromnes IM, DelGiorno KE, Greenberg PD, Hingorani SR. Stromal reengineering to treat pancreas cancer. *Carcinogenesis* 2014; 35: 1451-1460.
- 40 The Lancet O. Pancreatic cancer: cause for optimism? *The Lancet Oncology* 2016; 17: 845.
- 41 Thomas D, Radhakrishnan P. Tumor-stromal crosstalk in pancreatic cancer and tissue fibrosis. *Mol Cancer* 2019; 18: 14.
- 42 Thompson CB, Shepard HM, O'Connor PM, Kadhim S, Jiang P, Osgood RJ *et al.* Enzymatic depletion of tumor hyaluronan induces antitumor responses in preclinical animal models. *Mol Cancer Ther* 2010; 9: 3052-3064.
- 43 Tjomsland V, Niklasson L, Sandstrom P, Borch K, Druid H, Bratthall C *et al.* The desmoplastic stroma plays an essential role in the accumulation and modulation of infiltrated immune cells in pancreatic adenocarcinoma. *Clin Dev Immunol* 2011; 2011: 212810.
- 44 Torphy RJ, Wang Z, True-Yasaki A, Volmar KE, Rashid N, Yeh B *et al.* Stromal Content Is Correlated With Tissue Site, Contrast Retention, and Survival in Pancreatic Adenocarcinoma. *JCO Precis Oncol* 2018; 2018.
- 45 Ueda J, Yoshida H, Mamada Y, Tani ai N, Yoshioka M, Hirakata A *et al.* Evaluation of the Impact of Preoperative Values of Hyaluronic Acid and Type IV Collagen on the Outcome of Patients with Hepatocellular Carcinoma After Hepatectomy. *Journal of Nippon Medical School = Nippon Ika Daigaku zasshi* 2018; 85: 221-227.
- 46 Vainer N, Dehlendorff C, Johansen JS. Systematic literature review of IL-6 as a biomarker or treatment target in patients with gastric, bile duct, pancreatic and colorectal cancer. *Oncotarget* 2018; 9: 29820-29841.
- 47 Vennin C, Murphy KJ, Morton JP, Cox TR, Pajic M, Timpson P. Reshaping the Tumor Stroma for Treatment of Pancreatic Cancer. *Gastroenterology* 2018; 154: 820-838.

- 48 Wang S, Willumsen N, Bager C, Karsdal MA, Chondros D, Taverna D. Extracellular matrix (ECM) circulating peptide biomarkers as potential predictors of survival in patients (pts) with untreated metastatic pancreatic ductal adenocarcinoma (mPDA) receiving pegvorhialuronidase alfa (PEGPH20), nab-paclitaxel (A), and gemcitabine (G) 2018; 36: 12030-12030.
- 49 Whatcott CJ, Diep CH, Jiang P, Watanabe A, LoBello J, Sima C *et al.* Desmoplasia in Primary Tumors and Metastatic Lesions of Pancreatic Cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2015; 21: 3561-3568.
- 50 Willumsen N, Bager CL, Leeming DJ, Smith V, Karsdal MA, Dornan D *et al.* Extracellular matrix specific protein fingerprints measured in serum can separate pancreatic cancer patients from healthy controls. *BMC cancer* 2013; 13: 554.
- 51 Willumsen N, Bager CL, Kehlet SN, Dragsbaek K, Neergaard JS, Hansen HB *et al.* Excessive matrix metalloprotease-mediated degradation of interstitial tissue (type I collagen) independently predicts short-term survival in an observational study of postmenopausal women diagnosed with cancer. *Oncotarget* 2017; 8: 52501-52510.
- 52 Wu RL, Huang L, Zhao HC, Geng XP. Hyaluronic acid in digestive cancers. *Journal of cancer research and clinical oncology* 2017; 143: 1-16.
- 53 Xing RD, Chang SM, Li JH, Li H, Han ZX. Serum hyaluronan levels in oral cancer patients. *Chin Med J (Engl)* 2008; 121: 327-330.
- 54 Zheng X, Carstens JL, Kim J, Scheible M, Kaye J, Sugimoto H *et al.* Epithelial-to-mesenchymal transition is dispensable for metastasis but induces chemoresistance in pancreatic cancer. *Nature* 2015; 527: 525-530.

Table 1. **Clinical characteristics of 809 patients with pancreatic cancer.**

<b>Characteristic</b>	<b>Number (%)</b>
<b>Age, median (range) years</b>	67 (37-89)
>75 years	135 (16.7)
<b>Sex</b>	
<b>Male</b>	440 (54.4)
<b>Female</b>	369 (45.6)
<b>ECOG PS</b>	
<b>0</b>	283 (35.0)
<b>1</b>	346 (42.8)
<b>2+</b>	103 (12.7)
<b>unknown</b>	77 (9.5)
<b>Stage</b>	
<b>I+II</b>	197 (24.4)
<b>III+IV</b>	595 (73.5)



<b>unknown</b>	17 (2.1)
<b>Diabetes</b>	
<b>yes</b>	210 (26.0)
<b>no</b>	599 (74.0)
<b>Smoking</b>	
<b>yes</b>	483 (59.7)
<b>no</b>	251 (31.0)
<b>unknown</b>	74 (9.1)
<b>Alcohol</b>	
<b>yes</b>	179 (22.1)
<b>no</b>	554 (68.5)
<b>unknown</b>	76 (9.4)
<b>BMI</b>	
<b>underweight</b>	63 (7.8)
<b>normal weight</b>	434 (53.6)
<b>overweight</b>	252 (31.1)
<b>unknown</b>	60 (7.4)
<b>Metastatic sites</b>	
<b>Liver</b>	337 (41.7)
<b>Peritoneum</b>	83 (10.3)
<b>Chemotherapy type</b>	
<b>Never received chemotherapy</b>	51 (6.3)
<b>Adjuvant gemcitabine</b>	160 (19.8)
<b>1st line palliative</b>	
<b>Gemcitabine</b>	404 (49.9)
<b>FOLFIRINOX</b>	120 (14.8)
<b>Gemcitabine and nab-paclitaxel</b>	55 (6.8)
<b>Other</b>	19 (2.3)
<b>CA19-9 median [IQR] U/ml</b>	519 [76, 3230]

Abbreviations: BMI, body mass index; ECOG PS, Eastern Cooperative Oncology Group performance status; IQR, interquartile range.

**Table 2.** Baseline serum levels of HA and PRO-C3 in patients with PC stratified by stage, presence of liver metastases, and ECOG PS compared to patients with other upper gastrointestinal cancer, benign diseases, and healthy controls.

<b>Characteristic</b>	<b>HA median [IQR] (ng/mL)</b>	<b>PRO-C3 median [IQR] (ng/mL)</b>
-----------------------	------------------------------------	--

<b>Pancreatic cancer (n=809)</b>	84 [48, 131] <sup>a,b</sup>	18 [11, 33] <sup>a,b</sup>
<b>Stage</b>		
I+II	75 [46, 126]	15 [10, 29]
III	69 [39, 101]	14 [10, 22]
IV	94 [51, 140] <sup>c</sup>	20 [13, 41] <sup>c</sup>
<b>Liver metastases</b>		
No	74 [43, 114]	15 [10, 25]
Yes	100 [57, 148] <sup>d</sup>	22 [14, 49] <sup>d</sup>
<b>ECOG PS</b>		
0	76 [44, 121]	17 [11, 35]
1	87 [48, 132]	17 [11, 31]
2+	98 [57, 140] <sup>e</sup>	19 [15, 36] <sup>f</sup>
<b>Ampullary carcinoma (n=44)</b>	102 [70, 154]	17 [12, 38]
<b>Distal biliary tract cancer (n=31)</b>	95 [66, 147]	30 [18, 52]
<b>Chronic pancreatitis (n=15)</b>	72 [36, 105]	11 [10, 17]
<b>Intraductal papillary mucinous neoplasm (n=41)</b>	66 [32, 101]	11 [8, 15]
<b>Duodenal adenoma (n=7)</b>	47 [32, 53]	14 [11, 17]
<b>No pathologic finding (N=25)</b>	53 [37, 100]	11 [9, 14]
<b>Healthy controls (n=141 and n=7)</b>	41 [27, 60]	7 [5, 9]

<sup>a</sup>P<0.001 compared with healthy controls.

<sup>b</sup>P<0.001 compared with benign conditions.

<sup>c</sup>P<0.001 across different stages.

<sup>d</sup>P<0.001 compared with no liver metastases.

<sup>e</sup>P=0.02 across different PS.

<sup>f</sup>P<0.05 across different PS.

Abbreviations: HA, hyaluronan; IQR, interquartile range; PRO-C3, propeptide of type III collagen; ECOG PS, Eastern Cooperative Oncology Group performance status.

**Table 3.** Univariate and multivariate Cox analyses for overall survival according to clinical characteristics and pretreatment serum HA and PRO-C3 (HRs are reported per 100 unit increases). The significant terms are included in the multivariate analyses.

Variable	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P-value	HR (95% CI)	P-value
Age ≥ 75 years (vs. <75 years)	1.42 (1.18-1.72)	<0.01	1.33 (1.06-1.66)	<0.01
Sex (female vs. male)	1.01 (0.88-1.17)	0.88	-	-
PS 1 (vs. PS 0)	1.60 (1.35-1.89)	<0.01	1.34 (1.12-1.61)	<0.01
PS 2 (vs. PS 0)	2.71 (2.15-3.43)	<0.01	2.58 (1.99-3.35)	<0.01
BMI (underweight vs. normal weight)	1.27 (0.97-1.66)	0.09	-	-
BMI (overweight vs. normal weight)	1.02 (0.87-1.20)	0.82	-	-
Diabetes (yes vs. no diabetes)	1.14 (0.97-1.34)	0.11	-	-
Smoking ((ever vs. never)	1.08 (0.92-1.26)	0.37	-	-

Alcohol abuse (ever vs. never)	1.02 (0.86-1.21)	0.84	-	-
Stages 3 and 4 (vs. stage 1 and 2)	3.67 (3.03-4.45)	<0.01	3.71 (2.27-6.09)	<0.01
Liver metastasis (yes vs. no met)	2.42 (2.08-2.81)	<0.01	1.69 (1.40-2.04)	<0.01
Peritoneum metastasis (yes vs. no met)	1.58 (1.25-1.99)	<0.01	1.28 (0.99-1.66)	0.06
Never received chemotherapy (vs. Res)	3.39 (2.41-4.78)	<0.01	4.21 (2.09-8.50)	<0.01
Gemcitabine (P) (vs. Res)	4.24 (3.43-5.26)	<0.01	1.05 (0.64-1.72)	0.85
FOLFIRINOX (P) (vs. Res)	2.48 (1.91-3.22)	<0.01	0.64 (0.38-1.09)	0.10
Gemcitabine+nab-paclitaxel (P) (vs. Res)	2.86 (2.02-4.04)	<0.01	0.63 (0.35-1.13)	0.12
Other combinations (P) (vs. Res)	3.31 (2.03-5.41)	<0.01	0.94 (0.49-1.82)	0.86
CA19-9 >median	1.93 (1.65-2.26)	<0.01	1.39 (1.17-1.65)	<0.01
HA (per 100 ng/mL)	1.18 (1.13-1.25)	<0.01	1.04 (0.97-1.13)	0.27
PRO-C3 (per 100 ng/mL)	1.28 (1.11-1.49)	<0.01	1.42 (1.14-1.76)	<0.01





Abbreviations: BMI, body mass index; PS, performance status; HA 100, additional risk of death for every 100-unit increase in hyaluronan; HR, hazard ratio; CI, confidence interval; met, metastasis; P, palliative; PRO-C3 100, additional risk of death for every 100-unit increase in pro-peptide of type III collagen; Res, resection.


## Figure Legends





**Figure 1.** Receiver operating characteristic (ROC) curve of baseline serum HA (A, B) and PRO-C3 (C, D) detection in discriminating PC patients from healthy controls and from patients with benign conditions. Abbreviations: AUC, area under the ROC; CI, confidence interval; HA, hyaluronan; PRO-C3, propeptide of type III collagen.

**Figure 2.** Kaplan–Meier survival curves showing the association between pre-treatment serum HA (2A) and PRO-C3 (2B) in patients with PC across all stages and for stage 4 (Figures 2C and 2D, HA and PRO-C3, respectively). Cut-off is the 75 percentile.

**Figure 3.** Kaplan–Meier survival curves in patients with PC receiving palliative chemotherapy, grouped according to decreased versus increased serum levels of HA and PRO-C3 after the first treatment cycle (Figures 3A and 3B) and at the time of first CT evaluation (Figures 3C and 3D).







6:43 PM DEC 01, 2022

THE MOMENT YOUR GUT INSTINCT  
WAS BACKED BY SCIENTIFIC  
RESULTS\_



## THE DIFFERENCE OF BREAKTHROUGH MOMENTS

**WITH COMPLETE SOLUTIONS FOR GROUNDBREAKING DISCOVERIES FROM A TRUSTED PARTNER.**

Your next breakthrough could be closer than you imagine, especially with the right resources to help you advance your research. At BD, we are dedicated to helping you get the data you need, when, where and how you need it. Our integrated solutions in instrumentation, software and reagents are optimized to work together to bring you closer to your next breakthrough moment. And you can depend on us for world-class training, service and support to help you get the most from the results your research depends on. Discover a range of optimized solutions that open endless possibilities for your future research. **Discover the new BD.**

Learn how you can advance your research >

