



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Development of an approach for characterisation of striae in bubble containing glasses

Jensen, Martin; Keding, Ralf; Yue, Yuanzheng

Publication date:
2010

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Jensen, M., Keding, R., & Yue, Y. (2010). *Development of an approach for characterisation of striae in bubble containing glasses*. Abstract from 10th ESG Glass Conference, Magdeburg, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Development of an approach for characterisation of striae in bubble containing glasses

Martin Jensen, Ralf Keding, Yuanzheng Yue
Section of Chemistry, Aalborg University, DK-9000 Aalborg, Denmark
E-mail address: maj@bio.aau.dk (M. Jensen)

Inhomogeneities such as striae and bubbles in glass-forming melts affect the physical, chemical, mechanical, and optical properties of the produced glass. Chemical striae are a major component of the inhomogeneity present in a wide variety of natural and manmade glasses and are often the source of the poor quality of glass products. To detect and quantify the extent of striae in glasses, a picture processing method has been established by the present authors. However, it is found that the presence of the bubbles (gas inclusions) results in considerable uncertainties of the quantification of striae. The present work demonstrates how the occurrence of bubbles dramatically affects the outcome of the striae quantification. To eliminate the disturbing effect of bubbles, an improved picture processing method for precise assessment of the striae in bubble containing glasses is proposed. Thus, accurate characterisation, and hence, control of the inhomogeneity of glasses on a routine basis is made feasible.