



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Atrial fibrillation is more frequently associated with nonobstructive coronary lesions

The Bialystok Coronary Project

Tomaszuk-Kazberuk, Anna; Koziański, Marek; Kuzma, Łukasz; Bujno, Elżbieta; Lopatowska, Paulina; Rogalska, Ewelina; Dobrzycki, Sławomir; Sobkowicz, Bożena; Lip, Gregory Y H

Published in:

Polskie Archiwum Medycyny Wewnętrznej

DOI (link to publication from Publisher):

[10.20452/pamw.15635](https://doi.org/10.20452/pamw.15635)

Creative Commons License
CC BY-NC-SA 4.0

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Tomaszuk-Kazberuk, A., Koziański, M., Kuzma, Ł., Bujno, E., Lopatowska, P., Rogalska, E., Dobrzycki, S., Sobkowicz, B., & Lip, G. Y. H. (2020). Atrial fibrillation is more frequently associated with nonobstructive coronary lesions: The Bialystok Coronary Project. *Polskie Archiwum Medycyny Wewnętrznej*, 130(12), 1029-1036. Advance online publication. <https://doi.org/10.20452/pamw.15635>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Atrial fibrillation is more frequently associated with nonobstructive coronary lesions: the Białystok Coronary Project

Anna Tomaszuk-Kazberuk¹, Marek Koziński², Łukasz Kuźma³,
Elżbieta Bujno⁴, Paulina Łopatowska⁴, Ewelina Rogalska⁴,
Sławomir Dobrzycki³, Bożena Sobkowicz¹, Gregory Y. H. Lip^{5,6}

1 Department of Cardiology, Medical University of Białystok, Białystok, Poland

2 Department of Cardiology and Internal Medicine, Medical University of Gdańsk, Gdynia, Poland

3 Department of Invasive Cardiology, Medical University of Białystok, Białystok, Poland

4 Department of Cardiology, University Hospital in Białystok, Białystok, Poland

5 Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Liverpool, United Kingdom

6 Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

KEY WORDS

atrial fibrillation,
coronary angiography,
coronary artery
disease

EDITORIAL

by Koziński, see p. 1024

ABSTRACT

INTRODUCTION Atrial fibrillation (AF) and chronic coronary syndrome (CCS) share common risk factors and frequently coexist. Additionally, AF symptoms may mimic CCS.

OBJECTIVES The aim of the study was to investigate the hypothesis indicating absence of significant coronary lesions in patients with AF as compared with those with sinus rhythm.

PATIENTS AND METHODS We conducted a single-center retrospective study including consecutive patients referred for elective coronary angiography between 2007 and 2016.

RESULTS The study population included 8288 patients out of whom 1674 had AF. There were substantial differences between groups with and without AF. Patients with AF were significantly older, more often were men and had diabetes, and more frequently were diagnosed with both chronic kidney disease and heart failure. On the other hand, they had history of hyperlipidemia less often. CCS was less frequently detected in patients with AF as compared with those with sinus rhythm (37.5% vs 41.1%; $P < 0.001$). Additionally, the latter group more often underwent subsequent coronary angioplasty (19.2% vs 22.3%; $P = 0.004$). Multivariable analysis identified AF as an independent factor associated with absence of significant coronary lesions (odds ratio, 1.57; 95% CI, 1.32–1.87; $P < 0.001$). Moreover, a comparison between patients with and without angiographically significant CCS revealed a higher prevalence of AF in the latter group (18.7% vs 21.2%; $P = 0.006$).

CONCLUSIONS In our study, AF was associated with the absence of significant coronary lesions on angiography, reflecting difficulties with qualifying patients with AF for invasive CCS diagnostic workup. Our findings suggest the need for more efficacious noninvasive diagnostic approach for patients with AF and suspected CCS.

INTRODUCTION Atrial fibrillation (AF) is the most common cardiac arrhythmia associated with excessive risk of ischemic stroke and heart failure as well as reduced life expectancy.^{1,2} On the other hand, chronic coronary syndrome (CCS) remains the main cause of morbidity and mortality in an aging population. Both disease entities share common risk factors such as hypertension, diabetes, and obesity.³⁻⁵

The prevalence of CCS in patients with AF varies from 17% to 47%,^{3,6,7} whereas AF in patients with CCS is much less common, ranging from 0.2% to 5%.⁸⁻¹² Patients with AF more often have concomitant CCS as compared with sinus rhythm controls.^{13,14} Additionally, AF is more frequently present in patients with either peripheral or cerebrovascular artery disease than in nonatherosclerotic individuals.¹⁵

Correspondence to:

Prof. Anna Tomaszuk-Kazberuk, MD,
PhD, Department of Cardiology,
Medical University of Białystok,
ul. M. Skłodowskiej-Curie 24A,
15-276 Białystok, Poland,
phone: +48 85 746 86 56,
email: a.tomaszuk@poczta.fm

Received: August 25, 2020.

Revision accepted:

September 28, 2020.

Published online: October 5, 2020.

Pol Arch Intern Med. 2020;

130 (12): 1029-1036

doi:10.20452/pamw.15635

Copyright by the Author(s), 2020

WHAT'S NEW?

In this study, the number of nonsignificant findings on coronary angiography was 1.6-fold higher in patients with atrial fibrillation (AF) than in patients without AF. Atrial fibrillation was associated with absence of significant coronary lesions and patients with AF less frequently underwent coronary angioplasty as compared with those with sinus rhythm. This raises the possibility that it may be difficult to select patients with AF for invasive procedures, given the increased findings of nonsignificant chronic coronary syndromes (CCS) on coronary angiography. The reason for these findings might be multifactorial: AF symptoms mimic CCS symptoms, ST-segment depression during AF poorly predicts obstructive CCS, stress tests are rarely proceeded in patients with AF, and rapid rhythm makes computed tomography scan difficult to interpret. A more efficacious noninvasive diagnostic approach is needed in patients with AF and suspected CCS.

Patients with AF in the course of arrhythmia may present with chest pain, which can be accompanied by transient ischemic-type ST-segment changes, sometimes with marginally elevated cardiac necrosis markers, thus mimicking symptoms of CCS.¹⁶ There are contradictory data on angiographic findings among patients with AF.¹⁴ It is important to look at the prevalence of significant CCS lesions in patients with AF compared with those with sinus rhythm to prevent unnecessary coronary angiography in patients with AF.

On the other hand, there are several studies including the ARIC (the Atherosclerosis Risk in Communities) study, the MESA (Multi-Ethnic Study of Atherosclerosis), and the Rotterdam Study which found an association between subclinical atherosclerosis and incident AF.¹⁷

The aim of the present study was to investigate the hypothesis indicating the absence of significant coronary lesions in patients with AF as compared with those with sinus rhythm referred for elective coronary angiography. This indicates that some patients with AF may have undergone coronary angiography needlessly.

PATIENTS AND METHODS We reviewed medical records of 26 985 patients hospitalized in the Department of Invasive Cardiology of the Medical University of Białystok, Poland, who had coronary angiography performed because of exacerbated angina (recurrent chest pain, classical stable angina, long history of chest pain / angina, or other symptoms such as dyspnea) between 2007 and 2016. The Białystok Coronary Project is an observational research project that includes hospitalized patients with AF and CCS. It is focused on diagnostics and therapy of this population. We excluded patients with acute coronary syndromes (ACS), Takotsubo cardiomyopathy, and history of ischemic heart disease, as well as those referred for coronary angiography before heart valve surgery. Prior cardiosurgical valve replacement was also the exclusion criterion. The set of extracted variables included demographic data,

medical history, physical examination, resting electrocardiogram, routine transthoracic echocardiogram, coronary angiogram, and percutaneous coronary intervention (PCI). Eventually, our final study cohort included 8288 patients. Concomitant diseases were defined via the *International Statistical Classification of Diseases, Tenth Revision (ICD-10)* codes, not prescription claims.

We analyzed whether history of AF was associated with the absence of significant coronary lesions on coronary angiography. We also investigated factors connected with the absence of significant coronary lesions on angiography. The study protocol conformed to the ethical guidelines of the 1975 Declaration of Helsinki, and was approved by the local ethics committee.

Coronary angiography All patients underwent coronary angiography and the number of diseased vessels were assessed. Coronary angiography was performed according to the Judkins technique.¹⁸ Diagnosis of CCS and indication for PCI was performed according to the current European guidelines.¹⁹ Significant stenosis of the coronary vessel was defined as more than 50% in the main stem of the left coronary artery and 70% in the rest of the epicardial arteries. The degree of CCS was classified as single-, double-, or triple-vessel disease.

Echocardiographic analysis Left ventricular ejection fraction (LVEF) was assessed in transthoracic echocardiography using the modified biplane Simpson method (Philips Ultrasound System, Sonos 5500, Aalborg, Denmark, equipped for harmonic imaging with a 3.6 MHz transducer) and was derived in accordance with the recommendations of the European Society of Echocardiography.²⁰

Definition of atrial fibrillation Atrial fibrillation was defined as the presence of AF on electrocardiogram during the index hospitalization and/or as indicated by a diagnosis found in medical records, the hospital inpatient database, or outpatient databases. Electrocardiographic AF was defined as an irregular rhythm with fibrillatory waves and no defined P waves. Diagnoses and AF classification were based on physician-assigned diagnoses in the medical records and/or the presence of corresponding codes from the *International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM)* for AF (427.31) in the hospital discharge or outpatient databases. Atrial fibrillation was further divided into the following categories: paroxysmal AF or chronic AF (persistent and permanent).¹

Statistical analysis Continuous data were expressed as means (SD) or medians and interquartile ranges (IQRs) when appropriate. Relative frequencies were used to present categorical variables. Normality of the distribution was assessed with the Kolmogorov–Smirnov test, while the 2-tailed *t* test was applied for

FIGURE 1 Flow chart of the study population selection
Abbreviations: AF, atrial fibrillation




TABLE 1 Characteristics of the screened population

Patients	Value
All	26 985 (100)
Included in the final analysis	8288 (30.7)
Admitted for coronary angiography before heart valve surgery	3499 (13)
Hospitalized due to NSTEMI	3126 (11.6)
Hospitalized due to STEMI	3102 (11.5)
With history of ischemic heart disease	2940 (10.9)
Hospitalized due to UA	2503 (9.3)
Admitted for scheduled percutaneous coronary intervention	2211 (8.2)
Other	1316 (4.9)

Data are presented as number (percentage) of patients.

Abbreviations: NSTEMI, non-ST-segment elevation myocardial infarction; STEMI, ST-segment elevation myocardial infarction; UA, unstable angina

the comparative analysis of variables. Non-normally distributed data were compared with the Mann–Whitney test. Categorical variables were compared using the χ^2 test. The multivariable logistic regression backward stepwise Wald method was used to determine the odds ratio (OR) for the absence of significant coronary lesions. A 2-tailed *P* value of less than 0.05 was considered statistically significant. The IBM SPSS version 25.0 software package (SPSS, Chicago, Illinois, United States) was used.

RESULTS Out of 26 985 patients admitted for elective coronary angiography, a total of 8288 patients were included in the final analysis (FIGURE 1, TABLE 1). Over 80% of study participants had hypertension, 66.9% had hyperlipidemia, and 25.7% had diabetes. Chronic kidney disease (CKD) and chronic heart failure (CHF) were diagnosed in 32.5% and 20.6% of the study population, respectively. Detailed characteristics of the study population is provided in TABLE 2.

A total of 1674 (20.2%) patients had AF. Paroxysmal AF was the most frequent type of the analyzed arrhythmia. However, we found only a marginally lower prevalence of permanent AF (FIGURE 1).

Atrial fibrillation population There were substantial differences in terms of clinical characteristics between groups with and without AF. Patients with AF were older, more often men, and more often had diabetes, and had higher body mass index and more frequently were diagnosed with both CKD and CHF. On the other hand, they were less likely to have history of hyperlipidemia. Mean (SD) HAS-BLED score was 2.14 (0.87), $\text{CHA}_2\text{DS}_2\text{-VASc}$ score was 3.99 (1.6).

The highest mean (SD) HAS-BLED and $\text{CHA}_2\text{DS}_2\text{-VASc}$ scores were noted in patients with permanent AF: 2.2 (0.91) and 4.3 (1.6), respectively. In patients with paroxysmal AF, the mean (SD) HAS-BLED was 2.15 (0.84) and in persistent AF, it was 1.9 (0.81).

TABLE 2 Comparison between patients with and without atrial fibrillation (continued on the next page)

Variable	All study participants (n = 8288)	Patients with AF (n = 1674)	Patients without AF (n = 6614)	P value
Male sex, n (%)	4500 (54)	1018 (61)	3482 (53)	<0.001
Age, y, mean (SD)	65.2 (10.2)	68.6 (9.6)	64.3 (10.2)	<0.001
BMI, kg/m ² , mean (SD)	29.1 (4.9)	30 (5.2)	28.9 (4.7)	<0.001
NYHA class, median (IQR)	2 (2–3)	3 (2–3)	2 (2–3)	0.22
CCS class, median (IQR)	2 (2–2)	2 (2–2)	2 (2–2)	0.26
CAD with significant stenosis on coronary angiography	3345(40.4)	627 (37.5)	2718 (41.1)	<0.001
Single-vessel CAD	1507 (18.2)	342 (20.4)	1165 (17.6)	<0.001
Double-vessel CAD	807 (9.7)	126 (7.5)	681 (10.3)	<0.001
Triple-vessel and/or significant LM stenosis	1031 (12.4)	159 (9.5)	872 (13.2)	<0.001
Hyperlipidemia	5548 (66.9)	961 (57.4)	4587 (69.4)	<0.001
Hypertension	6886 (83.1)	1370 (81.8)	5516 (83.4)	0.14
Diabetes mellitus	2132 (25.7)	477 (28.5)	1655 (25)	0.005
Chronic heart failure	2690 (32.5)	1038 (62)	1652 (24.9)	<0.001
HF _r EF	1381 (51.3)	621 (59.8)	760 (46)	<0.001
HF _m rEF	599 (22.3)	220 (21.2)	379 (22.9)	
HF _p EF	710 (26.4)	197 (18.9)	513 (31.1)	
Left ventricular ejection fraction, %, median (IQR)	55 (42–60)	45 (30–55)	56 (50–60)	<0.001
Chronic kidney disease	1706 (20.6)	556 (33.2)	1150 (17.4)	<0.001
Creatinine, mg/dl, median (IQR)	0.89 (0.78–1.05)	0.97 (0.82–1.17)	0.87 (0.78–1.02)	<0.001
eGFR, ml/min/1.73 m ² , mean (SD)	76.4 (19.1)	70 (20)	77.9 (18.5)	<0.001
eGFR >90 ml/min/1.73 m ²	2095 (26)	266 (16.4)	1829 (28.6)	<0.001
eGFR 60–90 ml/min/1.73 m ²	4455 (55.2)	855 (52.8)	3600 (55.9)	
eGFR 45–60 ml/min/1.73 m ²	967 (12)	302 (18.6)	665 (10.3)	
eGFR 30–45 ml/min/1.73 m ²	406 (5)	161 (9.9)	245 (3.8)	
eGFR 15–30 ml/min/1.73 m ²	79 (0.9)	27 (1.7)	52 (0.1)	
eGFR <15 ml/min/1.73 m ²	63 (0.8)	10 (0.6)	53 (0.9)	
RBC, 10 ⁹ /mm ³ , median (IQR)	4.58 (4.28–4.88)	4.58 (4.23–4.89)	4.58 (4.29–4.8)	<0.001
Total cholesterol, mg/dl, median (IQR)	171 (146–203)	167 (141–198)	172 (148–204)	0.006
High-density lipoprotein cholesterol, mg/dl, median (IQR)	46 (39–55)	45 (38–53)	47 (39–56)	<0.001
Low-density lipoprotein cholesterol, mg/dl, median (IQR)	99 (77–127)	97 (75–124)	99 (78–128)	0.037
Triglycerides, mg/dl, median (IQR)	117 (85–165)	113 (82–159)	118 (85–167)	0.4
Presence of any significant stenosis on coronary angiography	3345 (40.4)	627 (37.5)	2718 (41.1)	<0.001
LM stenosis	245 (3)	42 (2.5)	203 (3.1)	0.2
LAD stenosis	2013 (24.3)	336 (20.1)	1677 (25.3)	<0.001
DIAG stenosis	735 (8.9)	132 (7.9)	603 (9.1)	0.09
Cx stenosis	1096 (13.2)	174 (10.4)	922 (14)	<0.001
OM stenosis	664 (8)	104 (6.2)	560 (8.5)	<0.001
RCA stenosis	1481 (17.9)	224 (13.4)	1257 (19)	<0.001
Patients qualified for conservative management	5330 (64.3)	1126 (67.3)	4204 (63.6)	0.004
Patients qualified for the Heart Team discussion	1161 (14)	227 (13.6)	934 (14.1)	0.55
Patients treated with PCI	1796 (21.7)	321 (19.2)	1475 (22.3)	0.004
PCI LM	34 (0.4)	7 (0.4)	27 (0.4)	0.95
PCI LAD	1115 (13.4)	206 (12.3)	909 (13.7)	0.11
PCI DIAG	182 (2.2)	34 (2)	148 (2.2)	0.6
PCI Cx	451 (5.4)	81 (4.9)	370 (5.6)	0.2
PCI OM	61 (0.7)	13 (0.8)	48 (0.7)	0.83
PCI RCA	404 (4.9)	64 (3.8)	340 (5.1)	0.015

TABLE 2 Comparison between patients with and without atrial fibrillation (continued from the previous page)

Data are presented as number (percentage) of patients unless otherwise indicated. The percentage is calculated within the subgroup.

Abbreviations: BMI, body mass index; CAD, coronary artery disease; CCS, Canadian Cardiovascular Society Angina Grading Scale; Cx, circumflex artery; DIAG, diagonal artery; eGFR, estimated glomerular filtration rate; HFmrEF, heart failure with moderate ejection fraction; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; IQR, interquartile range; LAD, left anterior descending; LM, left main artery; NYHA, New York Heart Association Functional Classification; OM, left marginal artery; PCI, percutaneous coronary intervention; RBC, red blood cells; RCA, right coronary artery; others, see [FIGURE 1](#)

The lowest mean (SD) CHA₂DS₂-VASc score was noted in the group of patients with paroxysmal AF (3.74 [1.5]). In the group of patients with persistent AF, the mean (SD) CHA₂DS₂-VASc score was 3.87 (1.5). A detailed comparison between the 2 groups is shown in [TABLE 2](#).

Coronary angiography Out of 8288 patients who underwent coronary angiography, significant lesions were found in 3345 study participants (40.4%). Importantly, patients without significant CCS were more frequently women and more commonly presented with AF ([TABLE 3](#)). Additionally, CCS was markedly less frequently detected on coronary angiography in patients with AF than in those with sinus rhythm. Furthermore, the latter group significantly more often underwent subsequent coronary angioplasty ([TABLE 2](#)).

Predictors of nonsignificant coronary findings in the multivariable analysis In the logistic regression model, the odds ratio of nonsignificant findings on coronary angiography were 1.6-fold higher in patients with AF than in those without AF (OR, 1.57; 95% CI, 1.32–1.87; $P < 0.001$). The indexed OR for a 5-kg/m² increase in body mass index was 1.24 (95% CI, 1.15–1.35; $P < 0.001$), and for a 5-mg/ml increase in high-density lipoprotein cholesterol concentration, 1.16 (95% CI, 1.1–1.24; $P < 0.001$). The OR for CHF was 0.68 (95% CI, 0.56–0.83; $P < 0.001$), other cardiovascular risk factors such as hypercholesterolemia, 0.64 (95% CI, 0.55–0.75; $P < 0.001$), diabetes, 0.69 (95% CI, 0.59–0.81; $P < 0.001$), hypertension, 0.68 (95% CI, 0.56–0.82; $P < 0.001$), CKD, 0.79 (95% CI, 0.66–0.94; $P < 0.001$) and women as compared with men, 0.39 (95% CI, 0.34–0.46; $P < 0.001$). Additionally, for a 10-year increase in age, the OR was 0.71 (95% CI, 0.65–0.76; $P < 0.001$; [FIGURE 2](#))

Complications of coronary angiography There were 48 cases of significant vascular complications related to the puncture of the site (0.5%). In 75% of complications, the femoral artery was affected, and in 25% of cases, it was the radial artery. Out of those cases, 16 (1%) were patients with AF and 32 (0.4%) had sinus rhythm ($P = 0.004$).

In 9 patients (0.11%), neurological disorders associated with the procedure were recorded in 7 patients (0.1%) with sinus rhythm and in 2 patients (0.12%) with AF. All computed tomography (CT) examinations performed in these patients failed to show any signs of intracranial

bleeding. In 19 patients, cardiac arrest occurred during hospitalization: 7 patients (0.4%) with AF and 12 patients with sinus rhythm (0.2%). Overall composite complications were more common in patients with AF.

DISCUSSION In this study, nonsignificant findings on coronary angiography were 1.6-fold higher in patients with AF than in patients without AF. Second, AF was associated with significantly less atherosclerotic narrowing on coronary angiography (627 [37.5%] vs 2718 [41.1%] in patients without AF; $P < 0.001$) and patients with AF less frequently underwent coronary angioplasty as compared with the population with sinus rhythm. This raises the possibility that AF may result in difficulties in selecting patients with AF for invasive procedures, given the increased findings of nonsignificant CCS on coronary angiography.

The reason for these findings might be complex and multifactorial. Patients with AF often present with typical or atypical chest pain, transient ST-segment depression, and elevated cardiac markers. These findings may mimic symptoms of CCS.²¹ Moreover, rapid tachycardia is often observed in patients with AF, and ST-segment changes at these rates have been linked with myocardial ischemia. For example, Tsigkas et al²¹ reported ST-segment depression in 38% of patients with AF and rapid rhythm and only about 50% of these patients had significant CCS on coronary angiography. ST-segment depression may frequently occur during significant tachycardia, even without CCS, and it is not necessarily typical for cardiac ischemia if the ST segment depression is less than 2 mm.^{21,22}

Moreover, ST-segment depression during rapid AF has less significant value than ST-segment depression during exercise tests in patients with sinus rhythm.²³ Only 4% of patients without ST-segment depression during rapid AF had positive noninvasive tests for myocardial ischemia and CCS on angiography.²³ Troponin release was shown in 15% of patients with AF and symptoms of ischemia, although CCS at angiography was absent.²⁵ In a database of patients referred for coronary angiography, history of AF correlated with no obstructive CCS.^{24,26}

On the other hand, Kravlev et al²⁷ showed that left main stem and right coronary artery (RCA) disease was present more frequently in patients with AF compared with controls. They also suggested that significant atherosclerotic narrowing

TABLE 3 Comparison between patients with and without angiographically significant coronary artery disease

Variable	Patients with CAD (n = 3345)	Patients without CAD (n = 5483)	P value
Male sex	2238 (67)	2262 (46)	<0.001
Age, y, mean (SD)	66.9 (9.7)	64 (10.3)	<0.001
BMI, kg/m ² , mean (SD)	28.8 (4.6)	29.33 (5)	<0.001
NYHA class, median (IQR)	3 (4–3)	2 (2–3)	0.04
CCS class, median (IQR)	2 (2–2)	2 (2–2)	0.03
Left ventricular ejection fraction, %, median (IQR)	55 (44–60)	55 (42–60)	0.5
Chronic heart failure	1247 (37.3)	1443 (29.2)	<0.001
HFrEF	601 (48.2)	780 (54.1)	<0.001
HFmrEF	301 (24.1)	298 (20.6)	
HFpEF	345 (27.7)	365 (25.3)	
Hyperlipidemia	2348 (70.2)	3200 (64.7)	0.001
Hypertension	2891 (86.4)	3995 (80.8)	<0.001
Diabetes mellitus	1014 (30.3)	1118 (22.6)	<0.001
AF			
Any	627 (18.7)	1047 (21.2)	0.006
Paroxysmal	267 (42.6) ^a	490 (46.8) ^a	0.09
Persistent	69 (11) ^a	125 (11.9) ^a	0.56
Permanent	291 (46.4) ^a	432 (41.3) ^a	0.04
Chronic kidney disease	824 (24.6)	882 (17.9)	<0.001
Creatinine, mg/dl, median (IQR)	0.93 (0.81–1.1)	0.86 (0.77–1.01)	<0.001
eGFR, ml/min/1.73 m ² , mean (SD)	70 (20)	77.8 (18.6)	<0.001
eGFR >90 ml/min/1.73 m ²	754 (23.2)	1341 (27.9)	<0.001
eGFR 60–90 ml/min/1.73 m ²	1775 (54.5)	2680 (55.7)	
eGFR 45–60 ml/min/1.73 m ²	441 (13.6)	526 (10.9)	
eGFR 30–45 ml/min/1.73 m ²	214 (6.6)	192 (4)	
eGFR 15–30 ml/min/1.73 m ²	37 (1.1)	42 (0.1)	
eGFR <15 ml/min/1.73 m ²	32 (1)	31 (0.6)	
RBC, 10 ⁶ /mm ³ , median (IQR)	4.58 (4.26–4.89)	4.58 (4.28–4.87)	0.55
Total cholesterol, mg/dl, median (IQR)	169 (144–200)	173 (147–204)	0.008
High-density lipoprotein cholesterol, mg/dl, median (IQR)	44 (38–52)	48 (40–57)	0.075
Low-density lipoprotein cholesterol, mg/dl, median (IQR)	99 (77–125)	99 (77–128)	0.006
Triglycerides, mg/dl, median (IQR)	121 (87–170)	114 (83–162)	0.34

Data are presented as number (percentage) of patients unless otherwise indicated.

a The percentage is calculated within the subgroup.

Abbreviations: see **FIGURE 1** and **TABLE 2**


in the proximal RCA and the circumflex artery (Cx) prior to the takeoff of the atrial arteries raises the risk of AF. In our study, significant RCA and Cx disease were less common than in patients with sinus rhythm. Another analysis of 3220 patients referred for coronary angiography revealed that only 43% of patients with CCS and AF had a diseased RCA or Cx.²¹

Does AF contribute to CCS or vice versa? There is a debate on this topic and opinions are often contradictory. Motloch et al²⁸ found that the anatomical distribution of coronary artery stenoses did not contribute to AF in CCS patients; however, AF was linked to worse CCS severity, which might predispose individuals to AF by driving ischemic heart disease and changes in left ventricular function. In our study, patients with AF had less prevalent multivessel CCS. It is possible

that in patients with AF, ischemia at the level of microcirculation of the myocardial muscle (rather than epicardial vessels) is responsible for the presence of typical angina. Unfortunately, our study did not explore the impact of cardiac ischemia induced by coronary small-vessel disease.

Our findings reflect difficulties with selecting AF patients for invasive coronary investigations. In the general population, the sensitivity of exercise testing ranges from 60% to 70% and specificity is approximately 85%.²⁹ As mentioned before, ST-segment depression is a common finding in AF patients,³⁰ and ST-depression during rapid AF has relatively higher sensitivity and lower specificity, when compared with exercise testing. Thus, the absence of ST-depression during rapid AF has satisfactory sensitivity for excluding significant CCS.²¹ Low positive predictive values

FIGURE 2 Multivariable logistic regression analysis of nonsignificant lesions in the coronary artery. Data are presented as odds ratios and their 95% CI. Abbreviations: CHF, chronic heart failure; CKD, chronic kidney disease; DM, diabetes mellitus; HA, hypertension; HDL, high-density lipoprotein; HL, hyperlipidemia; others, see [FIGURE 1](#) and [TABLE 2](#)



for significant CCS of ST-segment depression have been reported at approximately 32%.^{23,24} Patients with AF are not optimal candidates for coronary CT angiography during a rapid cardiac rhythm. However, it is a good diagnostic modality for patients in sinus rhythm with low CCS probability.

Not only are patients with AF improperly referred for invasive coronary investigations, but they are at an increased risk of complications. In a study of 13 498 patients, it was shown that patients with a history of AF were more likely to have post procedural bleeding, more often needed blood transfusion, and more often had exacerbation of heart failure, cardiogenic shock, and in-hospital mortality, but there were no differences in the incidence of postprocedural stroke or vascular complications in matched cohorts.^{30,31} Similarly, several studies reported that AF patients were more likely to experience postprocedural heart failure and cardiogenic shock.³²⁻³⁴ In the present study, there were 48 cases of significant vascular complications concerning femoral and radial artery related to the puncture of the site, and were more frequent in patients with AF than patients with sinus rhythm. Also, neurological disorders associated with the procedure were recorded more frequently in patients with AF than those with sinus rhythm. All CT examinations performed thereafter did not show signs of intracranial bleeding. Cardiac arrest during hospitalization was numerically higher in patients with AF. Apart from this, it was already demonstrated that AF patients with NSTEMI undergoing PCI can be predisposed to contrast induced nephropathy.³⁵

Limitations Our study has several limitations. First, our findings were obtained in a retrospective single-center study and should be confirmed in a multicenter prospective study. However, our data represent real life, as they were obtained

during daily clinical practice. Second, we excluded from our analysis patients with ACS. Third, fractional flow reserve measurements were not performed on regular basis. This could cause an inaccurate assessment of the significance of coronary stenosis. Therefore, only cases of stenosis with a reduction of 70% were defined as significant.³² Fourth, we were able neither to reliably assess stress tests before invasive diagnostics nor to measure concentrations of natriuretic peptides. Fifth, our study did not explore the impact of myocardial ischemia induced by coronary microvascular disease. Sixth, patients with AF were less likely to have lipid disorders, which may affect the results. Seventh, due to a very large group of patients included in the study (nearly 10 000) and nearly 10 years of analysis (2007–2016), we were unable to obtain reliable data on smoking in the analyzed group. Over the years, the approach to smoking addiction and the way it is coded has changed. Our diagnoses were based on the diagnoses established by the physicians in charge and were not verified again. Additionally, it would be interesting to perform more detailed echocardiographic examinations of all study participants. Furthermore, retrospective chart databases provide easy and cheap access to large numbers of patients although limitations such as potential selection bias should be taken into consideration. The last thing, one of the subtypes of AF is newly diagnosed AF. As there is no specific ICD code for it, we were not able to distinguish this type of AF.

Conclusions In our study, AF was associated with the absence of significant coronary lesions on angiography and less frequent need for revascularization, reflecting difficulties with qualifying AF patients for invasive CCS diagnostic workup. Multiple factors might be the reason for these findings, namely, AF symptoms may mimic CCS symptoms, ST-segment depression during AF poorly

predicts obstructive CCS, stress tests are rarely proceeded in patients with AF, and rapid rhythm makes CT scan difficult to interpret. Our findings suggest the need for a more efficacious noninvasive diagnostic approach for patients with AF and suspected CCS.

ARTICLE INFORMATION

CONTRIBUTION STATEMENT ATK, MK, EB, and PL conceived the idea for the study and contributed to the design of the research. ATK, EB, LK, ER, SD, and PL were involved in data collection. ATK, MK, LK, and GYHL analyzed the data. ATK, MK, MG, AS, BS, and GYHL edited the manuscript. All authors approved the final version of the manuscript.

CONFLICT OF INTEREST None declared.

OPEN ACCESS This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (CC BY-NC-SA 4.0), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited, distributed under the same license, and used for noncommercial purposes only. For commercial use, please contact the journal office at pamw@mp.pl.

HOW TO CITE Tomaszuk-Kazberuk A, Koziański M, Kuźma Ł, et al. Atrial fibrillation is more frequently associated with nonobstructive coronary lesions: the Białystok Coronary Project. *Pol Arch Intern Med.* 2020; 130: 1029-1036. doi:10.20452/pamw.15635

REFERENCES

- 1 Kirchoff P, Benussi S, Kotecha D, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Europace.* 2016; 18: 1609-1678. [↗](#)
- 2 Potpara TS, Mujovic N, Lip GYH. Meeting the unmet needs to improve management and outcomes of patients with atrial fibrillation: fitting global solutions to local settings. *Pol Arch Intern Med.* 2019; 129: 574-576. [↗](#)
- 3 Michniewicz E, Młodawska E, Lopatowska P, et al. Patients with atrial fibrillation and coronary artery disease – double trouble. *Adv Med Sci.* 2018; 63: 30-35. [↗](#)
- 4 Bednarski J, Balsam P, Tymińska A, et al. District versus academic hospitals: differences in the clinical characteristics of patients with atrial fibrillation without valvular heart disease treated with oral anticoagulants. *Pol Arch Intern Med.* 2018; 128: 274-279. [↗](#)
- 5 Knutti J, Wijns W, Saraste A, et al. 2019 ESC Guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J.* 2020; 41: 407-477.
- 6 AFFIRM Investigators. Atrial Fibrillation Follow-up Investigation of Rhythm Management. Baseline characteristics of patients with atrial fibrillation: the AFFIRM study. *Am Heart J.* 2002; 143: 991-1001. [↗](#)
- 7 Lip GYH, Beevers DG. History, epidemiology and importance of atrial fibrillation. *BMJ.* 1995; 311: 1361-1363. [↗](#)
- 8 Krahn AD, Manfreda J, Tate RB, et al. The natural history of atrial fibrillation: incidence, risk factors and prognosis in the Manitoba follow-up study. *Am J Med.* 1995; 98: 476-484. [↗](#)
- 9 Otterstad JE, Kirwan BA, Lubsen J, et al; Action Investigators. Incidence and outcome of atrial fibrillation in stable symptomatic coronary disease. *Scand Cardiovasc J.* 2006; 40: 152-159. [↗](#)
- 10 Cameron A, Schwartz MJ, Kronmal RA, et al. Prevalence and significance of atrial fibrillation in coronary artery disease (CASS Registry). *Am J Cardiol.* 1988; 6: 714-717. [↗](#)
- 11 Haddad AH, Prchokov VK, Dean DC. Chronic atrial fibrillation and coronary artery disease. *J Electrocardiol.* 1978; 11: 67-69. [↗](#)
- 12 Cheng TO. Coronary artery disease as an uncommon cause of chronic atrial fibrillation. *Clin Res.* 1974; 22: 268A.
- 13 Weijts B, Pisters R, Haest RJ, et al. Patients originally diagnosed with idiopathic atrial fibrillation more often suffer from insidious coronary artery disease compared to healthy sinus rhythm controls. *Heart Rhythm.* 2012; 9: 1923-1929. [↗](#)
- 14 Kakkar AK, Mueller I, Bassand JP, et al; GARFIELD Registry Investigators. Risk profiles and antithrombotic treatment of patients newly diagnosed with atrial fibrillation at risk of stroke: perspectives from the international, observational, prospective GARFIELD registry. *PLoS ONE.* 2013; 8: e63479. [↗](#)
- 15 Goto S, Bhatt DL, Röther J, et al. Prevalence, clinical profile, and cardiovascular outcomes of atrial fibrillation patients with atherothrombosis. *Am Heart J.* 2008; 156: 855-863. [↗](#)
- 16 Kanjwal K, Imran N, Grubb B, et al. Troponin elevation in patients with various tachycardias and normal epicardial coronaries. *Indian Pacing Electrophysiol J.* 2008; 8: 172-174.
- 17 Chen LY, Leening MJ, Norby FL, et al. Carotid intima-media thickness and arterial stiffness and the risk of atrial fibrillation: the Atherosclerosis Risk

in Communities (ARIC) study, Multi-Ethnic Study of Atherosclerosis (MESA), and the Rotterdam Study. *J Am Heart Assoc.* 2016; 5: e002907. [↗](#)

- 18 Judkins MP. Percutaneous transfemoral selective coronary arteriography. *Radiol Clin North Am.* 1968; 6: 467-492.
- 19 Montalescot G, Sechtem U, Achenbach S, et al. 2013 ESC guidelines on the management of stable coronary artery disease: the Task Force on the management of stable coronary artery disease of the European Society of Cardiology. *Eur Heart J.* 2013; 34: 2949-3003. [↗](#)
- 20 Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging.* 2015; 16: 233-270. [↗](#)
- 21 Tsigkas G, Kopsida G, Xanthopoulou I, et al. Diagnostic accuracy of electrocardiographic ST-segment depression in patients with rapid atrial fibrillation for the prediction of coronary artery disease. *Can J Cardiol.* 2014; 30: 920-924. [↗](#)
- 22 Androulakis A, Azaouridis KA, Aggeli CJ, et al. Transient ST-segment depression during paroxysms of atrial fibrillation in otherwise normal individuals: relation with underlying coronary artery disease. *J Am Coll Cardiol.* 2007; 6: 1909-1911. [↗](#)
- 23 Mendes LA, Connelly GP, McKenney PA, et al. Right coronary artery stenosis: an independent predictor of atrial fibrillation after coronary artery bypass surgery. *J Am Coll Cardiol.* 1995; 25: 198-202. [↗](#)
- 24 Pradhan R, Chaudhary A, Donato AA. Predictive accuracy of ST depression during rapid atrial fibrillation on the presence of obstructive coronary artery disease. *Am J Emerg Med.* 2012; 30: 1042-1047. [↗](#)
- 25 Wierzbowska-Drabik K, Cygulska K, Ciešlik-Guerra U, et al. Circumferential strain of carotid arteries does not differ between patients with advanced coronary artery disease and group without coronary stenoses. *Adv Med Sci.* 2016; 61: 203-206. [↗](#)
- 26 Abidov A, Hachamovitch R, Rozanski A, et al. Prognostic implications of atrial fibrillation in patients undergoing myocardial perfusion single-photon emission computed tomography. *J Am Coll Cardiol.* 2004; 44: 1062-1070. [↗](#)
- 27 Krale V, Schneider K, Lang S, et al. Incidence and severity of coronary artery disease in patients with atrial fibrillation undergoing first-time coronary angiography. *PLoS One.* 2011; 6: e24964. [↗](#)
- 28 Motloch LJ, Reda S, Larbig R, et al. Characteristics of coronary artery disease among patients with atrial fibrillation compared to patients with sinus rhythm. *Hellenic J Cardiol.* 2017; 58: 204-212. [↗](#)
- 29 Morise AP, Diamond GA. Comparison of the sensitivity and specificity of exercise electrocardiography in biased and unbiased populations of men and women. *Am Heart J.* 1995; 130: 741-747. [↗](#)
- 30 Lopatowska P, Tomaszuk-Kazberuk A, Młodawska E, et al. Management of patients with valvular and non-valvular atrial fibrillation in Poland: Results from Reference Cardiology University Center. *Cardiol J.* 2015; 22: 296-305. [↗](#)
- 31 Zimetbaum PJ, Josephson ME, McDonald MJ, et al. Incidence and predictors of myocardial infarction among patients with atrial fibrillation. *J Am Coll Cardiol.* 2000; 36: 1223-1227. [↗](#)
- 32 Sutton NR, Seth M, Ruwende C, et al. Outcomes of patients with atrial fibrillation undergoing percutaneous coronary intervention. *J Am Coll Cardiol.* 2016; 68: 895-904. [↗](#)
- 33 Lopes RD, Elliott LE, White HD, et al. Antithrombotic therapy and outcomes of patients with atrial fibrillation following primary percutaneous coronary intervention: results from the APEX-AMI trial. *Eur Heart J.* 2009; 30: 2019-2028. [↗](#)
- 34 Tonino PA, Fearon WF, De Bruyne B, et al. Angiographic versus functional severity of coronary artery stenoses in the FAME study fractional flow reserve versus angiography in multivessel evaluation. *J Am Coll Cardiol.* 2010; 55: 2816-2821. [↗](#)
- 35 Düzel B, Emren SV, Berlinger R. Effect of atrial fibrillation on contrast-induced nephropathy development in patients with non-ST-segment elevation myocardial infarction. *Angiology.* 2017; 68: 871-876. [↗](#)