



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Prolonged Corticomotor Homeostatic Plasticity - Effects Of Different Protocols And Their Reliability

Wittkopf, Priscilla G; Larsen, Dennis B; Gregoret, Luisina; Graven-Nielsen, Thomas

Published in:
Brain Stimulation

DOI (link to publication from Publisher):
[10.1016/j.brs.2021.01.017](https://doi.org/10.1016/j.brs.2021.01.017)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Wittkopf, P. G., Larsen, D. B., Gregoret, L., & Graven-Nielsen, T. (2021). Prolonged Corticomotor Homeostatic Plasticity - Effects Of Different Protocols And Their Reliability. *Brain Stimulation*, 14(2), 327-329. Advance online publication. <https://doi.org/10.1016/j.brs.2021.01.017>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Prolonged corticomotor homeostatic plasticity – Effects of different protocols and their reliability



Keywords:

Homeostatic plasticity
Plasticity
Transcranial magnetic stimulation
Transcranial direct current stimulation

Homeostatic plasticity complements synaptic plasticity by stabilizing neural activity within a physiological range. Human homeostatic plasticity has been investigated using two blocks of non-invasive brain stimulation (NIBS) with an interval of no stimulation in-between blocks, and the homeostatic response captured by changes in M1 excitability by quantifying the amplitude of motor-evoked potentials (MEPs) [1]. A homeostatic response of reduced MEPs amplitude when a block of excitatory stimulation is primed by another block of excitatory stimulation has been demonstrated in humans [2–4]. Conversely, a homeostatic response of increased MEPs amplitude has been observed when a block of inhibitory stimulation is primed by another inhibitory block [4,5]. An impaired homeostatic response has been detected in subjects with migraine, and chronic low back pain [2,6]. It is possible that homeostatic plasticity mechanisms are affected over time, throughout the development of the disease [2,3]. To date, studies investigating homeostatic plasticity have been conducted using cross-sectional designs [1,7]. In order to investigate long-term homeostatic plasticity changes, it is important to verify the reliability of homeostatic plasticity induction protocols. No studies have investigated the reliability of protocols using two blocks of inhibitory transcranial direct current stimulation (tDCS), and little is known about the reliability of the duration of corticomotor excitability changes post-homeostatic plasticity induction. Here we present the duration of homeostatic response and the test-retest reliability of corticomotor excitability changes, provoked by two homeostatic plasticity induction protocols (cathodal tDCS and anodal tDCS).

A sample size calculation was conducted ($\alpha = 0.05$, $\beta = 0.80$, effect size = 0.48 based on MEPs comparisons reported in previous studies [2,3]) resulting in a needed sample size of 13 participants. To account for differences in study designs and possible dropouts, recruitment of 15 participants for each experiment was targeted. Participants were recruited continuously for both experiments and data was collected from fifteen participants for cathodal tDCS experiment (6 males, 25.3 ± 3.8 years) and fifteen participants for anodal tDCS experiment (4 males, 24.6 ± 4.24 years). Each

participant took part in two experimental sessions on the same time on two consecutive days. The corticomotor excitability in response to the homeostatic plasticity protocol (cathodal-tDCS or anodal-tDCS) was measured before and immediately post homeostatic plasticity induction (time point 0-min), and then every 10 minutes for 70 minutes (ethical approval VN-20190069).

Homeostatic plasticity was induced in the left M1 using tDCS applied for 7 minutes followed by 3 minutes of no stimulation and another block of 5 minutes of tDCS [2–4]. A constant current of 1 mA was transmitted through the tDCS system (Starstim 32, Neuroelectronics, Barcelona, Spain), using two 3.14 cm² Ag/AgCl gelled electrodes placed into a neoprene cap corresponding to the international 10/10 EEG system (NE056 Headcap R, Neuroelectronics, Barcelona, Spain). For cathodal-tDCS, the cathode was placed at C3 and the anode placed at Fp2. In the anodal-tDCS experiment, the anode was placed at C3 and the cathode was placed at Fp2.

A magnetic stimulator (Magstim 200², Magstim Company, Whitland, UK) was used to deliver monophasic pulses, using a focal figure-of-8 coil. A stimulation intensity of 120% of the resting motor threshold was used for MEP assessment of the right first dorsal interosseous muscle. Peak-to-peak amplitude was extracted for each MEP and averaged across 15 sequential MEPs recorded at each time point (5–7 s inter-stimulus intervals). The averaged MEPs were used for analysis. All TMS methods were conducted according to the TMS guidelines [8].

MEPs at all-time points were log-transformed and used for subsequent analyses. Two-way repeated measures analysis of variance (ANOVA) was conducted on MEPs with factors *Day* (Day1 and Day2) and *Time* (pre-tDCS, 0-min, 10-min, 20-min, 30-min, 40-min, 50-min, 60-min, and 70-min post-tDCS). Bonferroni corrections were used and level of statistical significance considered $p \leq 0.05$. Intra-class correlation coefficient (ICC; 3,k) and Bland–Altman methods were used for the analysis of test-retest and absolute reliability (Appendix).

Increased MEPs were observed from 10 to 30 minutes post-cathodal tDCS compared with pre-cathodal tDCS (main effect of Time $F_{3,8, 54.5} = 6.51$; $p < 0.001$, Fig. 1A). Main effect of Day and Day \times Time interaction were not significant ($F_{1,14} = 1.22$; $p = 0.287$ and $F_{8,112} = 0.343$; $p = 0.947$, respectively). Reduced MEPs were observed from 10 to 40 minutes post-anodal tDCS compared with pre-anodal tDCS (main effect of Time $F_{3,6,51.2} = 8.9$; $p < 0.001$, Fig. 1B). Main effect of Day and Day \times Time interaction were not significant ($F_{1,14} = 0.67$; $p = 0.428$ and $F_{8,112} = 1.83$; $p = 0.078$, respectively).

Excellent test-retest reliability was observed for resting motor threshold and MEP amplitudes measured pre-tDCS. Moderate-to-excellent reliability was observed for post-tDCS MEPs at each




Fig. 1. Mean (+/- standard error of the mean, N = 15) of motor evoked potential (MEP) ratios for each time point at Day1 and Day2 following cathodal-tDCS (A) and anodal-tDCS (B). Significantly different MEPs compared with pre-tDCS (*, $p \leq 0.05$).

time point normalised by pre-tDCS (Table A1, Appendix). Bland-Altman analysis indicates no systematic bias when assessing RMT, pre-tDCS MEPs, and post-tDCS MEPs normalised to pre-tDCS MEP at any time point (Table A1; Fig A2 and A3, Appendix).

Results indicate that reliable homeostatic response that lasts up to 40 minutes, can be induced using two blocks of cathodal or anodal M1 tDCS with an interval of no stimulation in between blocks. Similar homeostatic responses have been identified in human M1 with a variety of homeostatic plasticity induction protocols

[1,3,5]. Nonetheless, non-homeostatic responses have been observed when using similar protocols in healthy participants [9,10]. It is possible that differences between studies are due to homeostatic and non-homeostatic mechanisms that may interact when using NIBS [1]. Furthermore, differences in findings may be related to variability in responses to protocols of NIBS. Further studies are needed to investigate the relationship between homeostatic and non-homeostatic mechanisms when using NIBS to investigate homeostatic mechanism in humans.

Lasting effects of homeostatic plasticity up to 40 minutes post-tDCS can be reliably induced and assessed using tDCS over an interval of 24 hours without carry over effects. This finding is in agreement with a study investigating reliability of a homeostatic plasticity induction protocol at intervals of 2, 7, and 14 days, when using the same anodal tDCS protocol as in the present study [3]. Taken together, these findings indicate that the present protocols may be used in studies investigating homeostatic plasticity over time.

Research funding

Center for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121).

Declaration of competing interest

None.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.brs.2021.01.017>.

References

- [1] Karabanov A, et al. Consensus paper: probing homeostatic plasticity of human cortex with non-invasive transcranial brain stimulation (in eng) *Brain Stimul* 2015;8(5):993–1006. <https://doi.org/10.1016/j.brs.2015.06.017>. 2015 Sep-Oct.
- [2] Thapa T, Graven-Nielsen T, Chipchase LS, Schabrun SM. Disruption of cortical synaptic homeostasis in individuals with chronic low back pain. *Clin Neurophysiol* 2018;129(5):1090–6. <https://doi.org/10.1016/j.clinph.2018.01.060>. 2018/05/01/.
- [3] Thapa T, Schabrun SM. Test-retest reliability of homeostatic plasticity in the human primary motor cortex. *Neural Plast* 2018;2018.
- [4] Fricke K, Seeber AA, Thirugnanasambandam N, Paulus W, Nitsche MA, Rothwell JC. Time course of the induction of homeostatic plasticity generated by repeated transcranial direct current stimulation of the human motor cortex (in eng) *J Neurophysiol* Mar 2011;105(3):1141–9. <https://doi.org/10.1152/jn.00608.2009>.
- [5] Siebner HR, et al. Preconditioning of low-frequency repetitive transcranial magnetic stimulation with transcranial direct current stimulation: evidence for homeostatic plasticity in the human motor cortex (in eng) *J Neurosci* Mar 31 2004;24(13):3379–85. <https://doi.org/10.1523/jneurosci.5316-03.2004>.
- [6] Antal A, Lang N, Boros K, Nitsche M, Siebner HR, Paulus W. Homeostatic metaplasticity of the motor cortex is altered during headache-free intervals in migraine with aura (in eng) *Cerebr Cortex* Nov 2008;18(11):2701–5. <https://doi.org/10.1093/cercor/bhn032>.
- [7] Hassanzahraee M, Zoghi M, Jaberzadeh S. How different priming stimulations affect the corticospinal excitability induced by noninvasive brain stimulation techniques: a systematic review and meta-analysis (in eng) *Rev Neurosci* Nov 27 2018;29(8):883–99. <https://doi.org/10.1515/revneuro-2017-0111>.
- [8] Chipchase L, et al. A checklist for assessing the methodological quality of studies using transcranial magnetic stimulation to study the motor system: an international consensus study (in eng) *Clin Neurophysiol Sep* 2012;123(9):1698–704. <https://doi.org/10.1016/j.clinph.2012.05.003>.
- [9] Monte-Silva K, Kuo M-F, Liebetanz D, Paulus W, Nitsche MA. Shaping the optimal repetition interval for cathodal transcranial direct current stimulation (tDCS). *J Neurophysiol* 2010;103(4):1735–40.
- [10] Monte-Silva K, et al. Induction of late LTP-like plasticity in the human motor cortex by repeated non-invasive brain stimulation (in eng) *Brain Stimul* May 2013;6(3):424–32. <https://doi.org/10.1016/j.brs.2012.04.011>.

Priscilla G. Wittkopf, Dennis B. Larsen, Luisina Gregoret,
Thomas Graven-Nielsen*
Center for Neuroplasticity and Pain (CNAP), Department of Health
Science and Technology, Aalborg University, Aalborg, Denmark

* Corresponding author. Center for Neuroplasticity and Pain
(CNAP), Department of Health Science and Technology Faculty of
Medicine, Aalborg University, Fredrik Bajers Vej 7 D3, DK-9220,
Aalborg, Denmark.
E-mail address: tgn@hst.aau.dk (T. Graven-Nielsen).

10 December 2020
Available online 24 January 2021