



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Physical activity and perceived stress at work in university workers

a cross-sectional study

López-Bueno, Rubén; Andersen, Lars L; Smith, Lee; López-Sánchez, Guillermo F; Mompel, Javier; Casedas, Luis; Casajús, José A

Published in:
The Journal of Sports Medicine and Physical Fitness

DOI (link to publication from Publisher):
[10.23736/S0022-4707.19.10259-9](https://doi.org/10.23736/S0022-4707.19.10259-9)

Publication date:
2020

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):
López-Bueno, R., Andersen, L. L., Smith, L., López-Sánchez, G. F., Mompel, J., Casedas, L., & Casajús, J. A. (2020). Physical activity and perceived stress at work in university workers: a cross-sectional study. *The Journal of Sports Medicine and Physical Fitness*, 60(2), 314-319. Advance online publication. <https://doi.org/10.23736/S0022-4707.19.10259-9>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Physical activity and perceived stress at work in university workers: a cross-sectional study

Rubén LÓPEZ-BUENO ^{1,2} *, Lars L. ANDERSEN ^{2,3}, Lee SMITH ⁴, Guillermo F. LÓPEZ-SÁNCHEZ ⁵, Javier MOMPEL ⁶, Luis CASEDAS ⁶, José A. CASAJÚS ^{7,8,9}

¹Department of Physical Medicine and Nursing, University of Zaragoza, Zaragoza, Spain;

²National Research Centre for the Working Environment, Copenhagen, Denmark;

³Department of Health Science and Technology, Aalborg University, Copenhagen, Denmark; ⁴The Cambridge Centre for Sports & Exercise Sciences, Anglia Ruskin University, Cambridge, UK;

⁵Faculty of Sports Sciences, University of Murcia, Murcia, Spain; ⁶Occupational Risk Division, University of Zaragoza, Zaragoza, Spain. ⁷GENUD

(Growth, Exercise, Nutrition and Development) Research Group, University of Zaragoza, Zaragoza, Spain; ⁸Biomedical Research Networking Centre about Nutrition and Obesity

Physiopathology (CIBER-OBN), Madrid, Spain; ⁹AgriFood Institute of Aragon (IA2), Zaragoza, Spain.

*Corresponding author: Rubén López-Bueno, Department of Physical Medicine and Nursing, University of Zaragoza, San Francisco Campus, 50009, Zaragoza, Spain. E-mail: rlopezbu@unizar.es

ABSTRACT

BACKGROUND: Previous research has suggested high levels of physical activity (PA), either in occupational or leisure-time, to be associated with low levels of perceived stress at work (PSW). However, because studies have been set in particular conditions, there is no possibility to generalise results on other populations of workers. This study investigated the association between PA and PSW in university workers.

METHODS: University employees (N=757) aged from 26 to 65 years (47% female) at a large public Spanish university. Data were collected between January 2017 and December 2017. Physical Activity Vital Sign (PAVS) questionnaire and a single-item scale were used to assess PA levels and PSW. Associations were examined through an adjusted logistic regression.

RESULTS: Results showed the strongest association between high PSW and low PA levels after adjusting for age, gender and profession (Odds Ratio [OR]=2.60, 95%CI 1.44-3.68). Around half of the employees (51.9%) performed at least 150 minutes of PA per week, which is higher than in most other Spanish and European worker populations.

CONCLUSIONS: Adequately high levels of PA may be beneficial for stress management in university workers as previously seen in other types of workers. Promoting PA strategies at the workplace could improve the working environment and the health of the workers.

Key words: Physical activity - Mental health - Job strain – Workplace - Occupational hazards - Perceived stress.

TEXT

Introduction

Physical activity (PA) aids in the prevention of non-communicable disease and risk factors. The World Health Organization (WHO) ranks lack of PA as risk factor number four concerning global mortality ¹. Dose-response associations exist, i.e., people who are more active live longer and are less prone to developing several types of cancer, type II diabetes, cardiovascular diseases, or suffering from poor mental health ²⁻⁵. Moreover, PA promotes healthy ageing, and importantly, older people who exercise tend to be less frail ⁶. Saint-Maurice et al. ⁷ found that PA levels tracked from adolescence to later adulthood and those with high PA levels had a 29-36% lower risk for all-cause mortality. PA during adulthood after being inactive also decreases the risk of mortality by 32-35%.

Figures from the latest European Commission report on PA estimates that 60% of the European adults never or rarely exercise ⁸. Even more concerning, a downhill trend of 7.7% less cardiorespiratory fitness in the last five decades has been detected within high- and upper-middle-income countries ⁹. The International Sport and Culture Association Health (ISCA) and the Centre for Economics and Business Research (CEBR) report an annual cost of 80 billion euros derived from medical care, medical treatments, functional limitations, disability, loss of dependence and a loss of working hours and productivity for European countries ¹⁰. Lindegård et al. ¹¹ reported stress-related mental disorders and musculoskeletal disorders as the two most important factors behind long-term sick leave in Sweden. Therefore, implementing measures to diminish the impact of the absenteeism regarding these two issues would be desirable.

Because jobs tend to be less physically active in western countries ¹⁰ and more workers are diagnosed with mental disorders such as too much work-related stress, anxiety, depression or psychosocial disorders ¹², institutions are trying to tackle those issues by implementing different measures in the workplace. Sit-stand desks, workplace policy changes, provision of counselling or multi-component interventions show inconsistent results when improving PA and avoiding sitting ¹³, whereas implemented measures at an organisational level resulted in no significant improvement when lowering perceived stress at work (PSW) ¹⁴. However, PA-focused interventions seem to be one of the most promising actions to reduce work-related stress and psychosocial hazards, as current research report PA to improve figures on mental health ^{3,15,16} and absenteeism ^{14,17-19}. Easy to implement, encouraging employees to be active (walking through the office) and to climb stairs (instead of using the elevator) are two of the principal strategies regarding PA promotion in the workplace among white-collar workers ²⁰. Nevertheless, PA promotion has been reported to have a low priority for small and large companies likely because they are unaware of the relationship between PA with absenteeism and productivity ²¹. Thus, as occupational PA is usually less than the weekly recommended 150 minutes of moderate PA or 75 minutes of vigorous PA, to assess overall PA (leisure, occupational and commuting time) would be desirable to know whether workers are reaching recommended levels of PA ²². Johnson et al. ²³ reported that expensive tools were a barrier for companies as the balance between investment and cost are usually unknown. Therefore, the use of more accessible assessment and standardised instruments such as Physical Activity Vital Sign (PAVS) ²⁴ or its electronic version (EVS) ²⁵ could help to gain reliability and generability on the matter.

Moderate evidence has suggested high stress at work to be associated with lack of PA either in occupational or leisure-time ^{14,26-30}. Moreover, job stress has been associated with the

development of cardiovascular diseases such as atrial fibrillation, stroke, and coronary heart disease^{27,31-34}. The demand-control model, i.e., an imbalance between low control over working conditions, and the effort-reward divergence model are the two most used models to assess work-related stress³⁴. Individual factors and working conditions could contribute to the personal perception of job stress, but also both organisational culture and structure¹². Little is known about PA levels among university workers because research is usually focused on university students³⁵. Also, only a few low-quality studies have investigated PA determinants in the workplace within southern Europe countries such as Spain, the need for more research on the issue has been recently suggested by governmental institutions²². The present study aimed to investigate possible associations between PSW and PA levels among Spanish university employees.

Materials and methods

The current research was based on a cross-sectional design. The study got the approval of the Aragonese Ethics Research Committee (CEICA; Identification Code PI18/027). The sample consisted of university workers from a Northeastern Spanish university, including both academic and university staff. All data were de-identified and analysed anonymously. University workers who got the voluntary university health check and fulfilled the medical questionnaire from January 1, 2017, to December 31, 2017, were included in the sample. Seven hundred fifty-seven employees, including lecturers and service staff from all campuses, agreed to wholly or partly complete the survey. All questionnaires were anonymous, allocated in medical centres, and supervised by medical personnel during medical check-ups.

The tool used to estimate PA levels among workers was the Physical Activity Vital Sign (PAVS) short version ²⁴. This questionnaire consisted of two questions related to the number of days and minutes employees usually take part in PA in a regular week. Employees could choose among 0,1,2,3,4,5,6 or 7 days of weekly PA and 10,20,30,40,50,60,90 and 150 or more for minutes they typically do PA on average each of those selected days. Weekly minutes of PA were calculated by multiplying days with minutes. PAVS has previously been validated using accelerometry, showing a moderate agreement among the United States clinic administrative staff (Cohen's $\kappa=0.46$, $P<0.001$) ³⁶. For a better understanding of the results, obtained PA levels were categorised into three groups regarding WHO recommendations about moderate and vigorous PA. As PAVS estimates PA in bouts of 10 minutes, and WHO recommendations on PA are set in a range of 75 to 150 weekly minutes when combining both moderate to vigorous PA, 80 and 150 minutes were considered as cut-off points.

Levels of PSW were estimated through the following question: "How much labour stress are you experiencing?" Possible answers comprised a 10-point scale varying from 1 "a little" to 10 "very much" perceived work-related stress. According to previous research, single-item scales have shown content, criterion, and construct validity for group-level analysis when monitoring stress at work ³⁷. Several advantages such as reduced costs, increased face validity for the participant, and problems concerning the designing of sum scales support the use of single-item scales ³⁸. The question used for this study offered high reliability and validity (Cronbach's $\alpha = .83$) when tested in 24 university employees voluntarily chosen from the sample, and showed moderate correlation (Pearson's $r= 0.64$, $p<0.001$) when compared to the validated Spanish version of the effort-reward imbalance model questionnaire (intrinsic section), which demonstrated high reliability and validity among

Spanish hospital personnel (Cronbach's $\alpha = .81$)³⁹. According to the expertise of the Occupational Risk Division on medical checks from previous years, PSW was categorized into three equal groups using 4 and 7 score as two cut-off points. Both PAVS and PSW questions were included in the institutional medical written questionnaire designed by the University Occupational Risk Division. On the questionnaire, respondents chose from three groups related to age, two types of occupations (i.e. academic staff and service staff), and gender. Due to confidentiality reasons, the study did not register age by date of birth and occupation by a more complete wide range of options. For the same reason, other possible covariates such as lifestyle or demographic characteristics could not be implemented.

Statistical analyses were conducted through the Statistical Package for Social Science (SPSS) version 22.0. To check for differences between different groups regarding PA, age, gender, profession, and PSW, a one-way analysis of variance was computed. To investigate the association between the dependent variable, PSW, and the independent variable, PA, a multiple logistic regression adjusted by factors (gender and profession) and covariate (age) was implemented. Significance was set at $p < 0.05$.

Questionnaires with missing data ($n=289$) were excluded. Therefore, four hundred sixty-eight university employees fully answered to the questions related to variables and covariates. The final cohort did not vary substantially from those who were discarded in terms of mean age (45.6; $SD=11.2$ in the final sample vs 47.1; $SD=10.2$ in the excluded employees), the proportion of men (55.9% vs 55.1%) and academic staff (54.4% vs 52.3%).

Results

The initial sample was composed of 757 university employees. According to data provided by the institution, participants in this study represent 12.8% of all employees contractually linked to the university during 2017. Table 1 shows the features of the sample data with missing values. The study population was, on average, 46.6 years old (SD=10.8), and 53.4% were academic staff. The average level of PA was 177.2 (SD=145.6) minutes per week.

Table 2 shows the differences between the three categories of PA. No significant differences in PA were detected in age, gender, or occupation. However, a significant difference was found among PSW groups ($p=0.002$).

A further analysis to estimate the association between PA and PSW adjusted for gender, age and profession showed that those employees reaching 70 minutes of weekly PA or less were more likely to have high PSW than low PSW (OR 2.60, 95% CI 1.44-3.68). A weaker but still relevant association was also found between high PSW and medium PA (OR 2.12, 95% CI. 1.22-2.93) and medium PSW and low PA (OR 2.08, 95% CI. 1.46-2.57) (Table 3).

Discussion

The present study examined levels of physical activity and perceived stress at work of Spanish university workers, and this is the first known study to investigate these areas in Spain. The present study found 70 or less weekly PA minutes to be associated with 7 to 10 PSW points in a ten graded scale when taking low PSW group as reference. University staff showed an average PA of 177 minutes per week and around half of the participants (51.90%) met the goal of 150 weekly PA minutes. According to the present study, no significant differences in gender, age, or profession on PA levels were detected among employees.

Although PA has been usually reported using different measurement tools, a previous study used electronic PAVS version (Exercise Vital Sign) on a broader adult sample finding 30.4% participants meeting the 150 minutes criteria ²⁵. PA has also been estimated with the British adult population through the GPAQ (Global Physical Activity Questionnaire). Estimated PA for the British population aiming for WHO recommendations were 35.40% ⁴⁰. Also, a multicenter study using accelerometry-based measurements of PA found an average of 51 weekly minutes (SD 29.5) in a sample composed of 329 Spanish adults ⁴¹. A more recent study using AerobePAR questionnaire estimated that 33.20% of Spanish adults met the recommended PA ⁴². These figures are considerably lower than those observed in the present study.

Regarding university staff, a study by Cooper & Barton ³⁵ revealed that around half of the adult university employees reported sufficient PA concerning the recommended activity guidelines. Although that study measured PA through IPAQ (International Physical Activity Questionnaire), those figures are comparable to those found in the present population of university workers. The use of various measurement tools and a wide range of sociodemographic differences among cohorts might explain PA differences among studies. Several studies remark the importance of sociodemographic features such as education, and occupation when explaining differences among PA population levels ^{42,43}. It could clarify why figures are very close when comparing between the two university staff cohorts and quite different when comparing to a less homogeneous population.

Previous research shows that employees with low PA and high PSW are more prone to develop cardiovascular diseases such as ischaemic heart disease or atrial fibrillation ^{31,33}. Other studies focus more on workability, finding a significant difference in favour of those

with higher exercise levels ⁴⁴. In any case, high PSW is generally related to a less PA at and outside of the workplace ^{16,26,28,29,45,46}. A meta-analysis found that physical inactivity was 26% and 21% more likely among those with high-strain jobs and defined passive jobs, respectively ⁴⁷. Jonsdottir et al. ⁴⁸ observed either light PA or moderate to vigorous PA as the only significant factor correlated with Sweden health care and social insurance workers perceived stress after adjusting by age, gender, body mass index, and educational level. Moreover, Kouvounen et al. ⁴⁹ demonstrated public sectors employee with high strain, passive jobs or low control jobs to be less active than their counterparts with differences ranging between 2.6 to 5.2 MET (Metabolic Equivalent Task index) hours/week, even when adjusting for occupational and health factors.

The fact that questionnaires were fulfilled in a medical setting, which may contribute to gain certainty as subjects are less likely to overstate their level of PA during a health check ²⁵. Another strength of our study is related to the high number of assessed workers, which is quite challenging to reach with objective tools. On the other side, since a questionnaire was used to estimate levels of PA and PSW, the results of this research should be considered carefully because of the inherent risk of reporting bias. The PSW question has not been validated in previous research, whereas tools such as questionnaires and surveys usually lead to overestimating PA ^{50,51}. Consequently, the true levels of PA may be lower than those observed in this study. Because there are different methods for assessing stress, e.g., the job strain and ERI (effort-reward imbalance) model ³⁴, comparisons between studies are not easy to accomplish. Overall, the typical observational risk of bias involving issues such as reverse causation, residual confounding, and selection may also affect the present results. Because PAVS do not discriminate between moderate and vigorous exercise, some of the participants in the PA range between 80 and 140 minutes of weekly PA could potentially have reached

recommended levels of PA. Besides, the number of missing PA values is quite high, probably as a result of several employees not having enough time to complete the whole questionnaire during the medical check. Despite that, small differences were detected between missing and valid values concerning age, gender, and profession.

Conclusions

This research observed a strong association when matching high PSW and low PA within a university workers population. There were also relationships between high PSW and medium PA, and the same occurred between medium PSW and low PA. A possible dose-response association has been detected regarding high and medium PSW and the three levels of PA. University employees PA levels might be higher than those assessed in other adult populations as universities usually develop PA programs aimed at workers.

REFERENCES

1. WHO. Global Recommendations on Physical Activity for Health. Geneva; 2014. <https://www.who.int/dietphysicalactivity/global-PA-recs-2010.pdf>.
2. Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med.* 2016;176(6):816-825.
3. Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: An analysis of burden of disease and life expectancy. *Lancet.* 2012;380(9838):219-229.
4. Ekelund U, Brown WJ, Steene-Johannessen J, et al. Do the associations of sedentary behaviour with cardiovascular disease mortality and cancer mortality differ by physical activity level? A systematic review and harmonised meta-analysis of data from 850 060 participants. *Br J Sports Med.* 2018:1-9.
5. Wilmot EG, Edwardson CL, Achana FA, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: Systematic review and meta-analysis. *Diabetologia.* 2012;55(11):2895-2905.
6. Izquierdo M, Cadore EL, Rech A, et al. Benefits of resistance training in physically frail elderly: a systematic review. *Aging Clin Exp Res.* 2017;30(8):889-899.
7. Saint-Maurice PF, Coughlan D, Kelly SP, et al. Association of Leisure-Time Physical Activity Across the Adult Life Course With All-Cause and Cause-Specific Mortality. *JAMA Netw Open.* 2019;2(3):e190355.
8. European Commission. Special Eurobarometer 472 Report Sport and Physical Activity Fieldwork. Brussels; 2018.
9. Lamoureux NR, Fitzgerald JS, Norton KI, Sabato T, Tremblay MS, Tomkinson GR. Temporal Trends in the Cardiorespiratory Fitness of 2,525,827 Adults Between 1967 and 2016: A Systematic Review. *Sport Med.* 2018;49(1):41-45.
10. Centre for Economics and Business Research. The Economic cost of physical inactivity in Europe. London; 2015. [http://inactivity-time-bomb.nowwemove.com/download-report/The Economic Costs of Physical Inactivity in Europe \(June 2015\).pdf](http://inactivity-time-bomb.nowwemove.com/download-report/The Economic Costs of Physical Inactivity in Europe (June 2015).pdf).

11. Lindegård A, Larsman P, Hadzibajramovic E, Ahlborg G. The influence of perceived stress and musculoskeletal pain on work performance and work ability in Swedish health care workers. *Int Arch Occup Environ Health*. 2014;87(4):373-379.
12. European Agency for Safety and Health at Work (EU-OSHA). Calculating the Cost of Work-Related Stress and Psychosocial Risks. Luxembourg; 2014. https://osha.europa.eu/en/tools-and-publications/publications/literature_reviews/calculating-the-cost-of-work-related-stress-and-psychosocial-risks
13. Shrestha N, Kukkonen-Harjula, KT, Verbeek JH, Ijaz S, Hermans V, Bhaumik S. Workplace interventions for reducing sitting at work (Review). *Cochrane Database Syst Rev*. 2016;(3).
14. Bhui KS, Dinos S, Stansfeld SA, White PD. A Synthesis of the Evidence for Managing Stress at Work: A Review of the Reviews Reporting on Anxiety, Depression, and Absenteeism. *J Environ Public Health*. 2012;2012:1-21.
15. Chekroud SR, Gueorguieva R, Zheutlin AB, et al. Association between physical exercise and mental health in 1·2 million individuals in the USA between 2011 and 2015: a cross-sectional study. *The Lancet Psychiatry*. 2018;5(9):739-746.
16. Chou LP, Tsai CC, Li CY, Hu SC. Prevalence of cardiovascular health and its relationship with job strain: A cross sectional study in Taiwanese medical employees. *BMJ Open*. 2016;6(4):1-9.
17. Losina E, Yang HY, Deshpande BR, Katz JN, Collins JE. Physical activity and unplanned illness-related work absenteeism: Data from an employee wellness program. *PLoS One*. 2017;12(5):1-13.
18. Von Thiele Schwarz U, Hasson H. Employee self-rated productivity and objective organizational production levels: Effects of worksite health interventions involving reduced work hours and physical exercise. *J Occup Environ Med*. 2011;53(8):838-844.
19. Tolonen A, Rahkonen O, Lahti J. Leisure-time physical activity and direct cost of short-term sickness absence among Finnish municipal employees. *Arch Environ Occup Heal*. 2017;72(2):93-98.
20. Smith L, Sawyer A, Gardner B, et al. Occupational physical activity habits of UK office workers: Cross-sectional data from the active buildings study. *Int J Environ Res Public Health*. 2018;15(6):1214.

21. Bailey MM, Collier RK, Pollack Porter KM. A qualitative study of facilitators and barriers to implementing worksite policies that support physical activity. *BMC Public Health*. 2018;18(1):1145.
22. Instituto Nacional de Seguridad Salud y Bienestar en el Trabajo (INSSBT). Beneficios del Fomento de la actividad física y la práctica deportiva en términos de mejora de la salud, el bienestar y la productividad empresarial. Madrid; 2017. [http://www.insht.es/InshtWeb/Contenidos/Documentacion/NUEVO/Beneficios del fomento de la actividad fisica.pdf](http://www.insht.es/InshtWeb/Contenidos/Documentacion/NUEVO/Beneficios%20del%20fomento%20de%20la%20actividad%20fisica.pdf).
23. Johnson S, Regnaud JP, Marck A, Berthelot G, Ungureanu J, Toussaint JF. Understanding how outcomes are measured in workplace physical activity interventions: A scoping review. *BMC Public Health*. 2018;18(1):1-12.
24. Greenwood JLJ, Joy EA, Stanford JB. The Physical Activity Vital Sign: A Primary Care Tool to Guide Counseling for Obesity. *J Phys Act Heal*. 2010;7(5):571-576.
25. Coleman KJ, Ngor E, Reynolds K, et al. Initial validation of an exercise “vital sign” in electronic medical records. *Med Sci Sports Exerc*. 2012;44(11):2071-2076.
26. Choi B, Schnall PL, Yang H, et al. Psychosocial working conditions and active leisure-time physical activity in middle-aged us workers. *Int J Occup Med Environ Health*. 2010;23(3):239-253.
27. Kivimäki M, Nyberg ST, Batty GD, et al. Job strain as a risk factor for coronary heart disease: A collaborative meta-analysis of individual participant data. *Lancet*. 2012;380(9852):1491-1497.
28. Kouvonen A, Vahtera J, Oksanen T, et al. Chronic workplace stress and insufficient physical activity: A cohort study. *Occup Environ Med*. 2013;70(1):3-8.
29. Oshio T, Tsutsumi A, Inoue A. The association between job stress and leisure-time physical inactivity adjusted for individual attributes: evidence from a Japanese occupational cohort survey. *Scand J Work Environ Health*. 2016;42(3):228-236.
30. Kouvonen A, Kivimäki M, Elovainio M, et al. Effort/reward imbalance and sedentary lifestyle: An observational study in a large occupational cohort. *Occup Environ Med*. 2006;63(6):422-427.
31. Fransson EI, Nordin M, Magnusson Hanson LL, Westerlund H. Job strain and atrial fibrillation – Results from the Swedish Longitudinal Occupational Survey of Health and meta-analysis of three studies. *Eur J Prev Cardiol*. 2018;25(11):1142-1149.

32. Fransson EI, Nyberg ST, Heikkilä K, et al. Job Strain and the risk of stroke: An individual-participant data meta-analysis. *Stroke*. 2015;46(2):557-559.
33. Theorell T, Jood K, Järholm LS, et al. A systematic review of studies in the contributions of the work environment to ischaemic heart disease development. *Eur J Public Health*. 2016;26(3):470-477.
34. Backé EM, Seidler A, Latza U, Rossnagel K, Schumann B. The role of psychosocial stress at work for the development of cardiovascular diseases: A systematic review. *Int Arch Occup Environ Health*. 2012;85(1):67-79.
35. Cooper K, Barton GC. An exploration of physical activity and wellbeing in university employees. *Perspect Public Health*. 2016;136(3):152-160.
36. Ball TJ, Joy EA, Goh TL, Hannon JC, Gren LH, Shaw JM. Validity of two brief primary care physical activity questionnaires with accelerometry in clinic staff. *Prim Health Care Res Dev*. 2015;16(1):100-108.
37. Elo AL, Leppänen A, Jahkola A. Validity of a single-item measure of stress symptoms. *Scand J Work Environ Health*. 2003;29(6):444-451.
38. Wanous JP, Reichers AE, Hudy M. Overall job satisfaction: how good are single-item measures. *J Appl Psychol* 1997;82(2):247-52.
39. Rancaño I, Siegrist J, Hernández-Mejía R, Cueto-Espinar A, Macías Robles MD, Fernández-López JA. Evaluación del estrés laboral en trabajadores de un hospital público español. Estudio de las propiedades psicométricas de la versión española del modelo «Desequilibrio Esfuerzo-Recompensa». *Med Clin (Barc)*. 2013;120(17):652-657.
40. Hunter RF, Tully MA, Donnelly P, Stevenson M, Kee F. Knowledge of UK physical activity guidelines: Implications for better targeted health promotion. *Prev Med (Baltim)*. 2014;65:33-39.
41. Cerin E, Cain KL, Conway TL, et al. Neighborhood Environments and Objectively Measured Physical Activity in 11 Countries. *Med Sci Sports Exerc*. 2015;46(12):2253-2264.
42. Fernandez-Navarro P, Aragonés MT, Ley V. Leisure-time physical activity and prevalence of non-communicable pathologies and prescription medication in Spain. *PLoS One*. 2018;13(1):1-13.
43. Sabia S, Van Hees VT, Shipley MJ, et al. Association between questionnaire-and accelerometer-assessed physical activity: The role of sociodemographic factors. *Am J Epidemiol*. 2014;179(6):781-790.

44. Calatayud J, Jakobsen MD, Sundstrup E, Casana J, Andersen LL. Dose-response association between leisure time physical activity and work ability: Cross-sectional study among 3000 workers. *Scand J Public Health*. 2015;43(8):819-824.
45. Gimeno D, Elovainio M, Jokela M, De Vogli R, Marmot MG, Kivimäki M. Association between passive jobs and low levels of leisure-time physical activity: The Whitehall II cohort study. *Occup Environ Med*. 2009;66(11):772-776.
46. Lindberg CM, Srinivasan K, Gilligan B, et al. Effects of office workstation type on physical activity and stress. *Occup Environ Med*. 2018;75(10):689-695.
47. Fransson EI, Heikkilä K, Nyberg ST, et al. Job strain as a risk factor for leisure-time physical inactivity: An individual-participant meta-analysis of up to 170,000 men and women. *Am J Epidemiol*. 2012;176(12):1078-1089.
48. Jonsdottir IH, Rödger L, Hadzibajramovic E, Börjesson M, Ahlborg G. A prospective study of leisure-time physical activity and mental health in Swedish health care workers and social insurance officers. *Prev Med (Baltim)*. 2010;51(5):373-377.
49. Kouvonen A, Kivimäki M, Elovainio M, Virtanen M, Linna A, Vahtera J. Job strain and leisure-time physical activity in female and male public sector employees. *Prev Med (Baltim)*. 2005;41(2):532-539.
50. Dyrstad SM, Hansen BH, Holme IM, Anderssen SA. Comparison of self-reported versus accelerometer-measured physical activity. *Med Sci Sports Exerc*. 2014;46(1):99-106.
51. Tucker JM, Welk GJ, Beyler NK. Physical activity in U.S. adults: Compliance with the physical activity guidelines for Americans. *Am J Prev Med*. 2011;40(4):454-461.

NOTES

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organisation regarding the material discussed in the manuscript.

Authors' contributions.—Rubén López-Bueno: Study concepts/study design, data acquisition, data interpretation, literature research. Lars L. Andersen: Study concepts/study design, manuscript revision for important intellectual content. Lee Smith: Data acquisition, data analysis, data interpretation. Guillermo F. López-Sánchez: Data acquisition, data analysis, draft the work. Javier Mompel: Data acquisition, data interpretation, study design. Luis Casedas: Data acquisition, data interpretation, study design. José A. Casajús: Study concepts/study design, manuscript revision for important intellectual content, guarantor of integrity of entire study.

Table I. Descriptive statistics for the Spanish university workers sample (N=757).

Variable	n (%)
Age, M (SD)	46.6 (10.8)
Missing	0 (0)
Gender	
Male	396 (52.3)
Female	361 (47.7)
Missing	0 (0)
Job role	
Service Staff	319 (42.1)
Academic Staff	404 (53.4)
Missing	34 (4.5)
PA, M (SD)	177.2 (145.6)
Missing	289 (38.2)
PSW, M (SD)	5.29 (2.1)
Missing	12 (1.6)

Note: PA=Physical Activity; SD=Standard Deviation; PSW=Perceived Stress at Work; M=Mean

Table II. Cohort personal features and association with categorised levels of PA in Spanish university workers.

Variable	Low PA (≤ 70 min/week)	Medium PA (80- 150min/week)	High PA (≥ 160 min/week)	P Value
Gender (n, %)				
Male	58 (24.2)	69 (28.7)	113 (47.1)	0.895
Female	54 (23.7)	70 (30.7)	104 (45.6)	
Age, in years (n, %)				
≤ 41	45 (26.2)	42 (24.4)	85 (49.4)	0.985
42-53	33 (22.9)	46 (31.9)	65 (45.1)	
> 53	34 (22.4)	51 (33.6)	67 (44.1)	
Profession (n, %)				
Service Staff	47 (24.7)	53 (27.9)	90 (47.4)	0.915
Academic Staff	60 (23.1)	79 (30.4)	121 (46.5)	
PSW, in scale points (n, %)				
Low ≤ 4	26 (18.6)	38 (27.1)	76 (54.3)	0.002*
Medium 5-6	34 (20.4)	52 (31.1)	81 (48.5)	
High ≥ 7	51 (33.3)	45 (29.4)	57 (37.3)	

* $p < 0.005$.

Note: PA=Physical Activity; PSW=Perceived Stress at Work

Table III. Associations between different categorised levels of PA and PSW in Spanish university workers adjusted for gender, age, and profession. (n=468).

	Low PA (≤ 70 min/week)		Medium PA (80-150min/week)		High PA (≥ 160 min/week)	
	OR	95% CI	OR	95% CI	OR	95% CI
Low PSW	1		1		1	
Medium PSW	2.08	[1.46, 2.57]	1.45	[0.93, 2.26]	1.39	[0.73, 2.03]
High PSW	2.60	[1.44, 3.68]	2.12	[1.22, 2.93]	1.04	[0.71, 1.35]

Note: PA=Physical Activity; PSW=Perceived Stress at Work; OR=Odds Ratio; CI=Confidence Interval