



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The genomics of heart failure

design and rationale of the HERMES consortium

Lumbers, R. Thomas; Shah, Sonia; Lin, Honghuang; Czuba, Tomasz; Henry, Albert; Swerdlow, Daniel I.; Mälarstig, Anders; Andersson, Charlotte; Verweij, Niek; Holmes, Michael V.; Ärnlöv, Johan; Svensson, Per; Hemingway, Harry; Sallah, Neneh; Almgren, Peter; Aragam, Krishna G.; Asselin, Geraldine; Backman, Joshua D.; Biggs, Mary L.; Bloom, Heather L.; Boersma, Eric; Brandimarto, Jeffrey; Brown, Michael R.; Brunner-La Rocca, Hans Peter; Carey, David J.; Chaffin, Mark D.; Chasman, Daniel I.; Chazara, Olympe; Chen, Xing; Chen, Xu; Chung, Jonathan H.; Chutkow, William; Cleland, John G.F.; Cook, James P.; de Denus, Simon; Dehghan, Abbas; Delgado, Graciela E.; Denaxas, Spiros; Doney, Alexander S.; Dörr, Marcus; Dudley, Samuel C.; Engström, Gunnar; Esko, Tõnu; Fatemifar, Ghazaleh; Felix, Stephan B.; Finan, Chris; Ford, Ian; Fougerousse, Françoise; Fouodjio, René; Ghanbari, Mohsen; Ghasemi, Sahar; Giedraitis, Vilmantas; Giulianini, Franco; Gottdiener, John S.; Gross, Stefan; Guðbjartsson, Daníel F.; Gui, Hongsheng; Gutmann, Rebecca; Haggerty, Christopher M.; van der Harst, Pim; Hedman, Åsa K.; Helgadottir, Anna; Hillege, Hans; Hyde, Craig L.; Jacob, Jaison; Jukema, J. Wouter; Kamanu, Frederick; Kardys, Isabella; Kavousi, Maryam; Khaw, Kay Tee; Kleber, Marcus E.; Køber, Lars; Koekemoer, Andrea; Kraus, Bill; Kuchenbaecker, Karoline; Langenberg, Claudia; Lind, Lars; Lindgren, Cecilia M.; London, Barry; Lotta, Luca A.; Lovering, Ruth C.; Luan, Jian'an; Magnusson, Patrik; Mahajan, Anubha; Mann, Douglas; Margulies, Kenneth B.; Marston, Nicholas A.; März, Winfried; McMurray, John J.V.; Melander, Olle; Melloni, Giorgio; Mordi, Ify R.; Morley, Michael P.; Morris, Andrew D.; Morris, Andrew P.; Morrison, Alanna C.; Nagle, Michael W.; Nelson, Christopher P.; Newton-Cheh, Christopher; Niessner, Alexander; Niiranen, Teemu; Nowak, Christoph; O'Donoghue, Michelle L.; Owens, Anjali T.; Palmer, Colin N.A.; Paré, Guillaume; Perola, Markus; Perreault, Louis Philippe Lemieux; Portilla-Fernandez, Eliana; Psaty, Bruce M.; Rice, Kenneth M.; Ridker, Paul M.; Romaine, Simon P.R.; Roselli, Carolina; Rotter, Jerome I.; Ruff, Christian T.; Sabatine, Marc S.; Salo, Perttu; Salomaa, Veikko; van Setten, Jessica; Shalaby, Alaa A.; Smelser, Diane T.; Smith, Nicholas L.; Stefansson, Kari; Stender, Steen; Stott, David J.; Sveinbjörnsson, Garðar; Tammesoo, Mari Liis; Tardif, Jean Claude; Taylor, Kent D.; Teder-Laving, Maris; Teumer, Alexander; Thorgeirsson, Guðmundur; Thorsteinsdottir, Unnur; Torp-Pedersen, Christian; Trompet, Stella; Tuckwell, Danny; Tyl, Benoit; Uitterlinden, Andre G.; Vaura, Felix; Veluchamy, Abirami; Visscher, Peter M.; Völker, Uwe; Voors, Adriaan A.; Wang, Xiaosong; Wareham, Nicholas J.; Weeke, Peter E.; Weiss, Raul; White, Harvey D.; Wiggins, Kerri L.; Xing, Heming; Yang, Jian; Yang, Yifan; Yerges-Armstrong, Laura M.; Yu, Bing; Zannad, Faiez; Zhao, Faye; Regeneron Genetics Center; Wilk, Jemma B.; Holm, Hilma; Sattar, Naveed; Lubitz, Steven A.; Lanfear, David E.; Shah, Svati; Dunn, Michael E.; Wells, Quinn S.; Asselbergs, Folkert W.; Hingorani, Aron D.; Dubé, Marie Pierre; Samani, Nilesh J.; Lang, Chim C.; Cappola, Thomas P.; Ellinor, Patrick T.; Vasani, Ramachandran S.; Smith, J. Gustav

Published in:
ESC Heart Failure

DOI (link to publication from Publisher):
[10.1002/ehf2.13517](https://doi.org/10.1002/ehf2.13517)

Creative Commons License
CC BY 4.0

The genomics of heart failure: design and rationale of the HERMES consortium

R. Thomas Lumbers^{1,2,3*}, Sonia Shah^{4,5}, Honghuang Lin^{6,7}, Tomasz Czuba⁸, Albert Henry^{1,5}, Daniel I. Swerdlow^{5,9}, Anders Mälarstig^{10,11}, Charlotte Andersson^{7,12}, Niek Verweij¹³, Michael V. Holmes^{14,15,16}, Johan Ärnlöv^{17,18}, Per Svensson^{19,20}, Harry Hemingway^{1,2,21}, Neneh Sallah^{1,2,22}, Peter Almgren²³, Krishna G. Aragam^{24,25,26}, Geraldine Asselin²⁷, Joshua D. Backman²⁸, Mary L. Biggs^{29,30}, Heather L. Bloom³¹, Eric Boersma³², Jeffrey Brandimarto³³, Michael R. Brown³⁴, Hans-Peter Brunner-La Rocca³⁵, David J. Carey³⁶, Mark D. Chaffin³⁷, Daniel I. Chasman^{38,39}, Olympe Chazara⁴⁰, Xing Chen¹⁰, Xu Chen⁴¹, Jonathan H. Chung²⁸, William Chutkow⁴², John G.F. Cleland^{43,44}, James P. Cook⁴⁵, Simon de Denus^{27,46}, Abbas Dehghan^{47,48}, Graciela E. Delgado⁴⁹, Spiros Denaxas^{1,2,21,50}, Alexander S. Doney⁵¹, Marcus Dörr^{52,53}, Samuel C. Dudley⁵⁴, Gunnar Engström²³, Tõnu Esko^{24,55}, Ghazaleh Fatemifar^{1,2}, Stephan B. Felix^{52,53}, Chris Finan⁵, Ian Ford⁴³, Francoise Fougerousse⁵⁶, René Fouodjio²⁷, Mohsen Ghanbari⁵⁷, Sahar Ghasemi^{53,58}, Vilmantas Giedraitis⁵⁹, Franco Giuliani³⁸, John S. Gottdiener⁶⁰, Stefan Gross^{52,53}, Daniel F. Gudbjartsson^{61,62}, Hongsheng Gui⁶³, Rebecca Gutmann⁶⁴, Christopher M. Haggerty⁶⁵, Pim van der Harst^{13,66,67}, Åsa K. Hedman¹¹, Anna Helgadottir⁶¹, Hans Hillege¹³, Craig L. Hyde¹⁰, Jaison Jacob⁴², J. Wouter Jukema^{68,69}, Frederick Kamanu^{37,70}, Isabella Kardys³², Maryam Kavousi⁵⁷, Kay-Tee Khaw⁷¹, Marcus E. Kleber⁴⁹, Lars Køber⁷², Andrea Koekemoer⁷³, Bill Kraus⁷⁴, Karoline Kuchenbaecker^{22,75}, Claudia Langenberg⁷⁶, Lars Lind⁷⁷, Cecilia M. Lindgren^{24,78,79}, Barry London⁸⁰, Luca A. Lotta⁷⁶, Ruth C. Lovering⁵, Jian'an Luan⁷⁶, Patrik Magnusson⁴¹, Anubha Mahajan⁸¹, Douglas Mann⁸², Kenneth B. Margulies³³, Nicholas A. Marston⁷⁰, Winfried März^{49,83,84}, John J.V. McMurray⁸⁵, Olle Melander⁸⁶, Giorgio Melloni^{37,70}, Ify R. Mordi⁵¹, Michael P. Morley³³, Andrew D. Morris⁸⁷, Andrew P. Morris^{45,79}, Alanna C. Morrison³⁴, Michael W. Nagle¹⁰, Christopher P. Nelson⁷³, Christopher Newton-Cheh^{26,88}, Alexander Niessner⁸⁹, Teemu Niiranen^{90,91}, Christoph Nowak¹⁷, Michelle L. O'Donoghue⁷⁰, Anjali T. Owens³³, Colin N.A. Palmer⁵¹, Guillaume Paré⁹², Markus Perola⁹³, Louis-Philippe Lemieux Perreault²⁷, Eliana Portilla-Fernandez^{57,94}, Bruce M. Psaty^{30,95}, Kenneth M. Rice²⁹, Paul M. Ridker^{38,39}, Simon P.R. Romaine⁷³, Carolina Roselli^{13,37}, Jerome I. Rotter⁹⁶, Christian T. Ruff⁷⁰, Marc S. Sabatine⁷⁰, Perttu Salo⁹⁰, Veikko Salomaa⁹⁰, Jessica van Setten⁹⁷, Alaa A. Shalaby⁹⁸, Diane T. Smelser³⁶, Nicholas L. Smith^{95,99,100}, Kari Stefansson^{61,101}, Steen Stender¹⁰², David J. Stott¹⁰³, Garðar Sveinbjörnsson⁶¹, Mari-Liis Tammesoo⁵⁵, Jean-Claude Tardif^{27,104}, Kent D. Taylor⁹⁶, Maris Teder-Laving⁵⁵, Alexander Teumer^{53,58}, Guðmundur Thorgeirsson^{61,101}, Unnur Thorsteinsdottir^{61,101}, Christian Torp-Pedersen^{105,106,107}, Stella Trompet^{68,108}, Danny Tuckwell⁴², Benoit Tyl⁵⁶, Andre G. Uitterlinden^{57,109}, Felix Vaura^{90,110}, Abirami Veluchamy⁵¹, Peter M. Visscher⁴, Uwe Völker^{53,111}, Adriaan A. Voors¹³, Xiaosong Wang⁴², Nicholas J. Wareham⁷⁶, Peter E. Weeke⁷², Raul Weiss¹¹², Harvey D. White¹¹³, Kerri L. Wiggins¹¹⁴, Heming Xing⁴², Jian Yang⁴, Yifan Yang³³, Laura M. Yerges-Armstrong¹¹⁵, Bing Yu³⁴, Faiez Zannad¹¹⁶, Faye Zhao⁴², Regeneron Genetics Center²⁸, Jemma B. Wilk¹⁰, Hilma Holm⁶¹, Naveed Sattar⁸⁵, Steven A. Lubitz^{24,117}, David E. Lanfear^{63,118}, Svati Shah^{74,119,120}, Michael E. Dunn¹²¹, Quinn S. Wells¹²², Folkert W. Asselbergs^{2,3,5,97}, Aroon D. Hingorani^{3,5}, Marie-Pierre Dubé^{27,104}, Nilesh J. Samani⁷³, Chim C. Lang⁵¹, Thomas P. Cappola³³, Patrick T. Ellinor^{37,117}, Ramachandran S. Vasan^{7,123} and J. Gustav Smith^{8,24,124,125*}

¹Institute of Health Informatics, University College London, Gower St, London, WC1E 7HB, UK; ²Health Data Research UK London, University College London, London, UK; ³BHF Research Accelerator, University College London, London, UK; ⁴Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland, Australia; ⁵Institute of Cardiovascular Science, University College London, London, UK; ⁶Section of Computational Biomedicine, Department of Medicine, Boston University School of Medicine, Boston, MA, USA; ⁷National Heart, Lung, and Blood Institute's and Boston University's Framingham Heart Study, Framingham, MA, USA; ⁸Department of Cardiology, Clinical Sciences, Lund University and Skåne University Hospital, Lund, Sweden; ⁹Department of Medicine, Imperial College London, London, UK; ¹⁰Pfizer Worldwide Research & Development, Cambridge, MA, USA; ¹¹Cardiovascular Medicine Unit, Department of Medicine Solna, Karolinska Institute, Stockholm, Sweden; ¹²Department of Cardiology, Herlev Gentofte Hospital, Herlev, Denmark; ¹³Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ¹⁴Medical Research Council Population Health Research Unit at the University of Oxford, Oxford, UK; ¹⁵Clinical Trial Service Unit and Epidemiological Studies Unit, Nuffield Department of Population Health, Big Data Institute, University of Oxford, Oxford, UK; ¹⁶National Institute for Health Research Oxford Biomedical Research Centre, Oxford University Hospital, Oxford, UK; ¹⁷Department of Neurobiology, Care Sciences and Society/Section of Family Medicine and Primary Care, Karolinska Institutet, Stockholm, Sweden; ¹⁸School of Health and Social Sciences, Dalarna University, Falun, Sweden; ¹⁹Department of Clinical Science and Education, Karolinska Institutet, Södersjukhuset, Stockholm, Sweden; ²⁰Department of Cardiology, Södersjukhuset, Stockholm, Sweden; ²¹The National Institute for Health Research, University College London Hospitals Biomedical Research Centre, University College London, London, UK; ²²UCL Genetics Institute, University College London, London, UK; ²³Department of Clinical Sciences, Lund University, Malmö, Sweden; ²⁴Program in Medical and Population Genetics, The Broad Institute of MIT and Harvard, Cambridge, MA, USA; ²⁵Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA; ²⁶Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA; ²⁷Montreal Heart Institute, Montreal, Quebec, Canada; ²⁸Regeneron Genetics Center, Tarrytown, NY, USA; ²⁹Department of Biostatistics, University of Washington, Seattle, WA, USA; ³⁰Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology and Health Services, University of Washington, Seattle, WA, USA; ³¹Division of Cardiology, Department of Medicine, Emory University Medical Center, Atlanta, GA, USA; ³²Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands; ³³Penn Cardiovascular Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA; ³⁴Human Genetics Center, Department of Epidemiology, Human Genetics, and Environmental Sciences, School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX, USA; ³⁵Maastricht University Medical Center, Maastricht, The Netherlands; ³⁶Department of Molecular and Functional Genomics, Geisinger, Danville, PA, USA; ³⁷Cardiovascular Disease Initiative, The Broad Institute of MIT and Harvard, Cambridge, MA, USA; ³⁸Division of Preventive Medicine, Brigham and Women's Hospital, Boston, MA, USA; ³⁹Harvard Medical School, Boston, MA, USA; ⁴⁰Centre for Genomics Research, Discovery Sciences, BioPharmaceuticals R&D, AstraZeneca, Cambridge, UK; ⁴¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden; ⁴²Novartis Institutes for Biomedical Research, Cambridge, MA, USA; ⁴³Robertson Centre for Biostatistics & Glasgow Clinical Trials Unit, Institute of Health and Wellbeing, University of Glasgow, Glasgow Royal Infirmary, Glasgow, UK; ⁴⁴National Heart and Lung Institute, Imperial College, London, UK; ⁴⁵Department of Biostatistics, University of Liverpool, Liverpool, UK; ⁴⁶Faculty of Pharmacy, Université de Montréal, Montreal, Quebec, Canada; ⁴⁷Department of Epidemiology and Biostatistics, Imperial College London, St Mary's Campus, London, UK; ⁴⁸MRC-PHE Centre for Environment and Health, Department of Epidemiology and Biostatistics, Imperial College London, St Mary's Campus, London, UK; ⁴⁹Vth Department of Medicine (Nephrology, Hypertensiology, Endocrinology, Diabetology, Rheumatology), Medical Faculty of Mannheim, University of Heidelberg, Heidelberg, Germany; ⁵⁰The Alan Turing Institute, British Library, London, UK; ⁵¹Division of Molecular and Clinical Medicine, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK; ⁵²Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany; ⁵³DZHK (German Center for Cardiovascular Research), partner site Greifswald, Greifswald, Germany; ⁵⁴Cardiovascular Division, Department of Medicine, University of Minnesota, Minneapolis, MN, USA; ⁵⁵Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Estonia; ⁵⁶Translational and Clinical Research, Servier Cardiovascular Center for Therapeutic Innovation, Suresnes, France; ⁵⁷Department of Epidemiology, Erasmus University Medical Center, Rotterdam, The Netherlands; ⁵⁸Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany; ⁵⁹Department of Public Health and Caring Sciences, Geriatrics, Uppsala University, Uppsala, Sweden; ⁶⁰Department of Medicine, Division of Cardiology, University of Maryland School of Medicine, Baltimore, MD, USA; ⁶¹deCODE genetics/Amgen Inc., Reykjavik, Iceland; ⁶²School of Engineering and Natural Sciences, University of Iceland, Reykjavik, Iceland; ⁶³Center for Individualized and Genomic Medicine Research, Department of Internal Medicine, Henry Ford Hospital, Detroit, MI, USA; ⁶⁴Division of Cardiovascular Medicine, University of Iowa Carver College of Medicine, Iowa City, IA, USA; ⁶⁵Department of Translational Data Science and Informatics, Geisinger, Danville, PA, USA; ⁶⁶Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands; ⁶⁷Durrer Center for Cardiogenetic Research, ICIN-Netherlands Heart Institute, Utrecht, The Netherlands; ⁶⁸Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands; ⁶⁹Netherlands Heart Institute, Utrecht, The Netherlands; ⁷⁰TIMI Study Group, Division of Cardiovascular Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; ⁷¹Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK; ⁷²Department of Cardiology, Copenhagen University Hospital Rigshospitalet, Copenhagen, Denmark; ⁷³Department of Cardiovascular Sciences, University of Leicester and NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester, UK; ⁷⁴Duke Molecular Physiology Institute, Durham, NC, USA; ⁷⁵Division of Psychiatry, University College of London, London, UK; ⁷⁶MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge School of Clinical Medicine, Cambridge, UK; ⁷⁷Department of Medical Sciences, Uppsala University, Uppsala, Sweden; ⁷⁸Big Data Institute at the Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK; ⁷⁹Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK; ⁸⁰Division of Cardiovascular Medicine and Abboud Cardiovascular Research Center, University of Iowa, Iowa City, IA, USA; ⁸¹Genentech Inc., San Francisco, CA, USA; ⁸²Center for Cardiovascular Research, Division of Cardiology, Department of Medicine, Washington University School of Medicine, St. Louis, MO, USA; ⁸³Synlab Academy, Synlab Holding Deutschland GmbH, Mannheim, Germany; ⁸⁴Clinical Institute of Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Graz, Austria; ⁸⁵BHF Cardiovascular Research Centre, University of Glasgow, Glasgow, UK; ⁸⁶Department of Internal Medicine, Clinical Sciences, Lund University and Skåne University Hospital, Malmö, Sweden; ⁸⁷Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK; ⁸⁸Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA, USA; ⁸⁹Department of Internal Medicine II, Division of Cardiology, Medical University of Vienna, Vienna, Austria; ⁹⁰Finnish Institute for Health and Welfare, Helsinki, Finland; ⁹¹Department of Medicine, Turku University Hospital and University of Turku, Turku, Finland; ⁹²Department of Pathology and Molecular Medicine, McMaster University, Hamilton, Ontario, Canada; ⁹³National Institute for Health and Welfare, Helsinki, Finland; ⁹⁴Division of Vascular Medicine and Pharmacology, Department of Internal Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands; ⁹⁵Kaiser Permanente Washington Health Research Institute, Kaiser Permanente Washington, Seattle, WA, USA; ⁹⁶The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA, USA; ⁹⁷Department of Cardiology, Division Heart and Lungs, University Medical Center Utrecht, University of Utrecht, Utrecht, The Netherlands; ⁹⁸Division of Cardiology, Department of Medicine, University of Pittsburgh Medical Center and VA Pittsburgh HCS, Pittsburgh, PA, USA; ⁹⁹Department of Epidemiology, University of Washington, Seattle, WA, USA; ¹⁰⁰Department of Veterans Affairs Office of Research and Development, Seattle Epidemiologic Research and Information Center, Seattle, WA, USA; ¹⁰¹Faculty of Medicine, Department of Medicine, University of Iceland, Reykjavik, Iceland; ¹⁰²Department of Clinical Biochemistry, Copenhagen University Hospital, Herlev and Gentofte, Denmark; ¹⁰³Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK; ¹⁰⁴Faculty of Medicine, Université de Montréal, Montreal, Quebec, Canada; ¹⁰⁵Department of Epidemiology and Biostatistics, Aalborg University Hospital, Aalborg, Denmark; ¹⁰⁶Department of Health, Science and Technology, Aalborg University Hospital, Aalborg, Denmark; ¹⁰⁷Department of Cardiology, Aalborg University Hospital, Aalborg, Denmark; ¹⁰⁸Section of Gerontology and Geriatrics, Department of Internal Medicine, Leiden University Medical Center, Leiden, The Netherlands; ¹⁰⁹Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands; ¹¹⁰Department of Clinical Medicine, University of Turku, Turku, Finland; ¹¹¹Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, Greifswald, Germany; ¹¹²Division of Cardiovascular Medicine, Department of Internal Medicine, The Ohio State University Medical Center, Columbus, OH, USA; ¹¹³Green Lane Cardiovascular Service, Auckland City Hospital, Auckland, New Zealand; ¹¹⁴Department of Medicine, University of Washington, Seattle, WA, USA; ¹¹⁵Human Genetics, GlaxoSmithKline, Collegetown, PA, USA; ¹¹⁶CHU de Nancy, Inserm and IRI-CRCT (F-CRIN), Institut Lorrain du Coeur et des Vaisseaux, Université de Lorraine, Nancy, France; ¹¹⁷Cardiac Arrhythmia Service and Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA; ¹¹⁸Heart and Vascular Institute, Henry Ford Hospital, Detroit, MI, USA; ¹¹⁹Division of Cardiology, Department of Medicine, Duke University Medical Center, Durham, NC, USA; ¹²⁰Duke Clinical Research Institute, Durham, NC, USA; ¹²¹Regeneron Pharmaceuticals, Cardiovascular Research, Tarrytown, NY, USA; ¹²²Division of Cardiovascular Medicine and the Vanderbilt Translational and Clinical Cardiovascular Research Center, Vanderbilt University, Nashville, TN, USA; ¹²³Sections of Cardiology, Preventive Medicine and Epidemiology, Department of Medicine, Boston University Schools of Medicine and Public Health, Boston, MA, USA; ¹²⁴Wallenberg Center for Molecular Medicine and Lund University Diabetes Center, Lund University, Lund, Sweden; and ¹²⁵The Wallenberg Laboratory/Department of Molecular and Clinical Medicine, Institute of Medicine, Gothenburg University and the Department of Cardiology, Sahlgrenska University Hospital, Gothenburg, Sweden

Abstract

Aims The HERMES (HEart failure Molecular Epidemiology for Therapeutic targetS) consortium aims to identify the genomic and molecular basis of heart failure.

Methods and results The consortium currently includes 51 studies from 11 countries, including 68 157 heart failure cases and 949 888 controls, with data on heart failure events and prognosis. All studies collected biological samples and performed genome-wide genotyping of common genetic variants. The enrolment of subjects into participating studies ranged from 1948 to the present day, and the median follow-up following heart failure diagnosis ranged from 2 to 116 months. Forty-nine of 51 individual studies enrolled participants of both sexes; in these studies, participants with heart failure were predominantly male (34–90%). The mean age at diagnosis or ascertainment across all studies ranged from 54 to 84 years. Based on the aggregate sample, we estimated 80% power to genetic variant associations with risk of heart failure with an odds ratio of ≥ 1.10 for common variants (allele frequency ≥ 0.05) and ≥ 1.20 for low-frequency variants (allele frequency 0.01–0.05) at $P < 5 \times 10^{-8}$ under an additive genetic model.

Conclusions HERMES is a global collaboration aiming to (i) identify the genetic determinants of heart failure; (ii) generate insights into the causal pathways leading to heart failure and enable genetic approaches to target prioritization; and (iii) develop genomic tools for disease stratification and risk prediction.

Keywords Heart failure; Cardiomyopathy; Genetics; Biomarkers; Association studies

Received: 27 January 2021; Revised: 9 June 2021; Accepted: 5 July 2021

*Correspondence to: R. Thomas Lumbers, MD PhD, Institute of Health Informatics, University College London, Gower St, WC1E 7HB London, UK.

Email: t.lumbers@ucl.ac.uk J. Gustav Smith, MD PhD FESC, The Wallenberg Laboratory/Department of Molecular and Clinical Medicine, Institute of Medicine, Gothenburg University and the Department of Cardiology, Sahlgrenska University Hospital, Gothenburg, Sweden. Email: gustav.smith@wlab.gu.se

Introduction


Heart failure (HF) is a complex clinical syndrome that imposes a substantial burden on public health; an estimated 30 million people worldwide are living with HF, and the prevalence is expected to rise with the aging of the global population.¹ HF is associated with substantial morbidity and mortality, underscoring the importance of mitigating the disease burden. Despite the advent of disease-modifying treatments for HF with reduced ejection fraction, considerable unmet need remains.² For HF with preserved ejection fraction, an increasingly prevalent subtype, no treatments are available to improve patient outcomes.³ Decades of research, based on preclinical models of HF, have uncovered numerous potential therapeutic targets; however, few have been successfully validated in phase III outcomes trials, reflecting, in part, the challenge of modelling complex age-associated multi-morbid disease processes.⁴ Human genetics provides a means to study causal biology in the patient: informing target selection and the formulation of a mechanism-based taxonomy of disease subtypes to help identify new therapeutic targets.⁵

Heart failure generally occurs when changes in cardiac structure or function result in impairment of ventricular filling and/or contraction and in impaired cardiac output and/or increased cardiac filling pressures.² Coronary artery disease and diseases causing abnormal cardiac loading (such as hypertension, valvular heart disease, and congenital heart disease) are established and common causes of HF. Many other factors can increase the risk of HF through direct effects on

myocardial structure and function (cardiomyopathy), including, for a small proportion of cases, monogenic cardiomyopathy syndromes.⁶ Familial aggregation and adoption studies suggest a heritable component to HF risk and disease progression with estimates for heritability up to 26%.^{7–9} Linkage studies of familial cardiomyopathies and genome-wide association studies (GWASs) have identified a number of rare and common variants associated with increased HF risk (Figure 1); however, the genetic architecture remains largely unknown.^{10–13}

It is a feature of many complex traits and diseases that common genetic variants account for a proportion of the population genetic variance.¹⁴ The genetic background of individual patients with respect to HF risk may modify the effects of HF risk factors, including influencing the penetrance and expression of Mendelian gene disorders, as has been observed for other common complex diseases.¹⁵ Furthermore, the identification of common disease-associated variants implicates regions of the genome that harbour causal genes and enables the appraisal of the causal role of risk factors and pharmacological targets by Mendelian randomization (MR) analysis.¹⁶ GWASs offer a robust and reproducible approach for the discovery of common disease-associated variants. Large samples, typically achieved by combining multiple studies through meta-analysis, are required to achieve sufficient statistical power to discern genotype–disease associations with modest effects.¹⁷ These approaches help inform a mechanism-based taxonomy of HF to support the development of effective targeted therapeutics.¹⁸

Figure 1 Genetic architecture of heart failure (HF) onset and progression. Examples of genes in which common (allele frequency 5% and greater), low-frequency (1–5%), or rare variants (<1%) have been shown to influence risk for HF onset or progression. Effect sizes are expressed in odds ratios for HF risk and hazard ratios for HF progression. Common variants can be identified in genome-wide association studies as exemplified by *BAG3*, *CLCNKA*, and *TSLP* loci, whereas variations with low population allele frequencies such as familial variants in the *MYH7*, *LMNA*, and *LAMP2* genes associated with cardiomyopathy will typically require sequencing-based approaches (based on a recent review article).²⁷ Familial variants in *LMNA* and *LAMP2* have been associated poor prognosis and particular cardiac phenotypes, manifesting with cardiomyopathy and Danon disease, respectively. *Although individually rare, protein-truncating variants in the large gene encoding Titin (*TTN*) collectively have a reported prevalence of 1% in the population, confer increased risk of HF, and have evidence of interaction with environmental factors such as alcohol, chemotherapy, and pregnancy. **A 25-basepair deletion of the gene encoding cardiac myosin-binding protein C (*MYBPC3*) conferring risk for HF has been reported to have an allele frequency of 4% in Southern Asian populations, highlighting how low-frequency variants of large effect may be population specific.



Here, we describe the HERMES (HEart failure Molecular Epidemiology for Therapeutic targetS) consortium: a global scientific collaboration of genetic studies linked to HF and related phenotypes. The consortium will develop tools and methods to enable the definition of HF subtypes and related traits across multi-modal datasets, including derivation of validated phenotypes from genomic biobanks linked to electronic health records.¹⁹ HERMES aims to unlock the potential for human genetics to inform the identification and validation of novel therapeutic approaches in HF by creating an open collaborative resource for the scientific community.^{20,21}

Study design

Aims


The core objective of the HERMES consortium is to conduct large-scale genetic association studies of HF and related phenotypes in order to identify common and low-frequency

genetic variants associated with HF risk and prognosis (Figure 2). In subsequent stages, we will extend these analyses to include rare variant association studies, based on sequence data available in a subset of studies. GWASs will be complemented by a range of follow-up analyses, including MR and rare variant burden tests, in order to identify novel disease mechanisms and to test existing therapeutic hypotheses.

Addressing syndromic heterogeneity

A stepwise approach to the genetic study of HF phenotypes and sub-phenotypes will be employed. The first completed analysis addressed the undifferentiated HF syndrome, without subtyping according to conventional classifiers of aetiology or phenotypes of left ventricular ejection fraction (LVEF).²² This study maximizes statistical power for the discovery of genetic factors influencing common pathophysiological mechanisms, such as left ventricular fibrotic remodeling, increased filling pressures, neurohormonal activation, and extracellular fluid retention (systemic and pulmonary

Figure 2 Component phenotypes of heart failure (HF). Schematic representation of HF phenotypes across the life course that will be studied in HERMES. HF diagnosis is typically preceded by cardiometabolic risk factors and genetic susceptibility factors for endophenotypes of structural and functional cardiac dysfunction. Circles on the left represent common structural endophenotypes, from top to bottom: normal ventricle, ventricle with symmetric hypertrophy, ventricle with asymmetric (septal) hypertrophy, and dilated ventricle. The natural history of HF extends from the initial time point of diagnosis (Dx) through a gradual decline with increasing episodes of worsening typically necessitating in-hospital care (decompensations) towards terminal pump failure. Sudden death from arrhythmia may occur at any point. Heritable contributions have been described for both risk factors, endophenotypes, HF onset, and HF progression.



vascular congestion) that may modify risk associated with upstream HF risk factors. Subsequent studies will address HF subtypes, including established and novel aetiological and cardiac morpho-functional phenotypes.

HERMES collaborating studies

At present, HERMES is a collection of 51 studies that have derived genome-wide genotyping data from community-based participants or hospitalized patients with clinical HF, including longitudinal population-based cohort studies, hospital-based electronic health record cohorts, case-control studies, and clinical trials. Detailed case ascertainment for HF and related cardiovascular phenotypes has been done for most studies; in others, phenotyping is based on routinely collected data from clinical care, national quality registers, or public data repositories. In addition to studies based in academic institutions, the collaboration includes many clinical trial datasets, providing a unique opportunity to study the genetic determinants of disease progression in HF. Due to the provenance of data currently available in contributing cohorts, currently ongoing initial analyses are limited to individuals of European ancestry; however, a central objective will be to include subjects of non-European ancestry as data from ancestrally diverse populations become available. Each contributing study in HERMES has appropriate ethical approval from the respective institutional review boards, and all participants provided informed consent for the use of their genetic data for research.


Organization

The collaborative framework of HERMES is similar to that of other collaborative consortia for genetic investigations, as shown in *Figure 3*.²³ All studies participate on an equal basis and operate under mutually agreed policies concerning project management, results sharing, and publication, which are articulated in a Memorandum of Understanding (see Supporting Information).

Data sharing and governance

To obviate the need for sharing of individual participant-level data and attendant data governance considerations, the consortium has adopted a distributed analysis model based on pre-planned meta-analysis of summary-level data contributed by each participating cohort study (*Figure 3*). Common analysis plans, methods and analytical scripts for quality control, phenotype and sub-phenotype derivation, and genetic association analyses are implemented in each study by local analysts. The resulting within-study summary data are then returned to the coordinating centre for quality control and meta-analysis. Meta-analysis is conducted at two independent centres to enable validation of results. Following the publication of results, summary data from HERMES meta-analyses are published in full on the Cardiovascular Disease Knowledge Portal (<http://www.broadcvdi.org/>).

Figure 3 International participation in HERMES and distributed analysis workflow. The HERMES consortium includes investigators from 12 countries from North America and Europe. Activities are overseen by a scientific committee with representatives from each contributing cohort and an executive committee. Common analysis plans are developed by the analysis group and deployed by participating studies. Meta-analysis is conducted by the analysis group and results shared with project working groups. Upon publication, the full genome-wide association summary estimates from meta-analysis are made available publicly through the Cardiovascular Disease Knowledge Portal (<http://www.broadcvdi.org/>).



Heart failure phenotype definition

While formal, international definitions of HF are in use,² case definitions vary across participating studies, as do methods for ascertainment, reflecting differences in study design and data availability (Supporting Information, *Table S1*). The performance of several HF ascertainment criteria in widespread use has, however, been shown to be similar.²⁴ For the initial GWAS meta-analysis, a broad definition was used based on physician adjudication, electronic health records-based phenotype algorithms, and corroborated self-report. Subsequent studies will follow a stepwise strategy for phenotype definition to address HF subtypes based on aetiology, LVEF, and disease progression (*Figure 2*). Mobilizing HF subtype data from electronic health records, leveraging large genomic biobanks, will be necessary to ensure sufficient statistical power for subtype analysis, and this will be achieved through the deployment of validated multi-modal rule-based phenotyping algorithms.²⁵

Given the mortality associated with HF, inclusion of incident and prevalent cases in analyses may lead to attenuation of effect estimates, due to survivorship or collider bias and increased heterogeneity^{26,27}; however, this bias is partially mitigated by the increased power associated with a larger sample size that can be achieved when prevalent cases are included.

Genotyping and imputation

Participants have been genotyped with a range of genome-wide single nucleotide polymorphism (SNP) arrays (Supporting Information, *Table S1*). All collaborating studies conducted imputation from directly measured genotype using public reference panels (1000 Genomes Project,

Haplotype Reference Consortium) or from local whole-genome sequence-based reference panels; for each meta-analysis project, genotype imputation was performed against a common pre-specified reference panel. Phasing and imputation were conducted using Eagle, MaCH, SHAPEIT, minimac2, or IMPUTE2 software at the discretion of participating cohorts.

Approach to genetic analyses

For GWASs, the analysis plan specifies quality filters to be applied to the data and the regression models for association testing. Once study-specific GWAS results have been uploaded to the central analytic team, these datasets undergo a second round of QC in order to identify and rectify any study-specific issues, align effect alleles across studies, and apply minor allele frequency and imputation quality filters, prior to meta-analysis. Analyses are conducted in parallel at two independent sites and are subsequently reconciled.

In study-specific GWAS analyses, logistic regression or Cox proportional hazards regression analyses are used, assuming additive genetic effects. Models are adjusted for age, sex, and principal components and family structure as appropriate for individual cohorts. Analytical softwares are left to the discretion of individual cohorts and include genetest, ProbABEL, mach2dat, QuickTest, PLINK2, SNPTEST, or R.

Quality controls of study-specific results are conducted according to accepted guidance, as previously reported.²⁸ In brief, variant identifiers and alleles are harmonized using the EasyQC tool and allele frequencies compared with the European reference panel of the 1000 Genomes Project. Distributions of reported *P*-values are plotted against *P*-values derived from Z-scores and reviewed, as well as distributions of beta estimates and standard errors, and Manhattan plots.

Variants with low imputation quality (<0.5) and with extreme betas and standard errors (>10) are excluded. Genomic control is applied at the study level where genomic inflation is identified ($\lambda_{GC} > 1.1$). Single-variant tests are limited to common and low-frequency variants (minor allele frequency $\geq 1\%$).

Meta-analyses are conducted using inverse-variance weighting using METAL software (https://genome.sph.umich.edu/wiki/METAL_Documentation). Heterogeneity of effect estimates across studies is evaluated from Cochrane's Q and I^2 statistics. The contribution of cryptic population structure to test statistics is estimated based on the linkage disequilibrium score (LDSC) regression intercept (<https://github.com/bulik/ldsc>). Statistical significance thresholds are based on the Bonferroni adjustment for the number of tests performed.

Power for statistical analyses

Power calculations for HF onset were based on R implementation of the widely used algorithms from the CaTS power calculator for one-stage association studies, with power calculations from the standard normal distribution.²⁹ Power to detect genome-wide significant associations ($P < 5 \times 10^{-8}$), based on the current HERMES sample size for cases with corresponding control subjects, was calculated as a function of effect allele frequency under different effect sizes (odds ratios of 1.05, 1.1, 1.2, 1.3, 1.4, and 1.5 in additive models). Similar power will apply to the reciprocal of the odds ratios < 1.0 for protective alleles. Additive-model odds ratios of identified common variants have typically been in the range of 1.1–1.2, with larger studies further identifying even smaller effects. Power calculations for HF mortality were based on the *survSNP* package in R,³⁰ included all cases, and plotted similarly to HF onset. Power calculations were conducted using the computing environment R Version 3.5.1 (R Core Team, Vienna, Austria), and results were plotted using STATA Version 15 (StataCorp, College Station, Texas, USA).

Study description

Participating studies

The HERMES consortium currently includes investigators from 12 countries (*Figure 3*) including 7 industry partners, representing 16 population-based cohorts, 1 hospital-based electronic health record cohort, 9 case cohorts of which 6 with control samples, and 25 clinical trials of which 9 with non-HF control samples (Supporting Information, *Table S1*). Ten of the clinical trials of HF were conducted within the NHLBI HF clinical research network. Detailed cohort

descriptions are provided in the Supporting Information. For a continuously updated list of included cohorts, please refer to the consortium webpage (www.hermesconsortium.org).

In aggregate, the 51 HERMES cohorts comprise 68 157 HF cases and 949 888 controls of European ancestry with array-based genotyping (Supporting Information, *Table S1*). Most of the 16 population-based cohorts identified cases based on ICD codes in hospital registers (10 cohorts), while a few had adjudicated events from patient records (4 cohorts) or included re-exams (2 cohorts). Of the nine case collections, seven were primarily focused on HF while two identified HF cases from an at-risk population (COGEN and LURIC). Of the 25 clinical trials, 17 had HF as inclusion criterion, whereas 8 included broader groups of patients with cardiometabolic diseases and identified HF from adjudicated outcomes (three trials) or case report forms (five trials).

Characteristics of participating studies

Baseline characteristics of the contributing studies are presented in Supporting Information, *Table S2*. As expected, clinical trials typically included younger cases (median age < 70 years in most trials) and had a lower burden of co-morbid disease compared with population-based cohorts. Risk factor distributions were largely as expected, with a particularly high burden of hypertension and coronary artery disease in all studies. Information on LVEF was available in a subset of cohorts: 16 151 had LVEF $< 40\%$, 4113 had LVEF 40–50%, and 9676 had LVEF $> 50\%$, corresponding to HF with reduced ejection fraction, HF with mid-range ejection fraction, and HF with preserved ejection fraction.²

Follow-up times and mortality of HF cases are presented in Supporting Information, *Table S1*. Overall, mortality among HF cases was 27%; however, the duration of follow-up was highly variable across studies, with median study follow-up ranging from 1 to 116 months.

Genotypic information

Genotyping was conducted on different high-density SNP platforms (Supporting Information, *Table S1*) and imputed based on European ancestry imputation panels for up to 8 246 881 common or low-frequency variants (minor allele frequency $> 1\%$) in the combined dataset. Detailed sequence data were available in at least 30 000 subjects from eight cohorts with exome-wide coverage and 140 000 subjects from six cohorts with whole-genome coverage (Supporting Information, *Table S1*) and were planned or ongoing in several additional cohorts.

Statistical power

Power calculations were conducted based on all 68 157 cases described earlier for HF progression, with an average mortality of 27%, and all cases with corresponding controls for HF onset (949 888 controls, 44 016 cases). For HF risk, HERMES is powered (>0.8) to detect effects down to odds ratios of 1.10 for common variants (minor allele frequency > 0.05) and 1.20 for low-frequency variants (0.01–0.05) (Figure 4A). For HF mortality, HERMES is powered to detect effects down to hazard ratios of 1.20 for variants with minor allele frequency > 0.08 and 1.40 for low-frequency variants (Figure 4B).

Discussion

With the recent exception of combination angiotensin receptor blockade and neprilysin inhibition and sodium-glucose transport protein 2 inhibitors, successful drug development in HF has, for many years, been limited. Almost all current therapies are repurposed from other indications (e.g. angiotensin-converting enzyme inhibitors, beta-blockers, and mineralocorticoid receptor antagonists for systemic hypertension and sodium-glucose transport protein 2 inhibitors for T2DM) and may not directly target processes leading to adverse cardiac remodelling. Human genetic and genomic studies provide unique opportunities to explore the causal biology in patients; the HERMES consortium provides a collaborative platform that enables these approaches.


Heart failure is a broadly defined syndromic disorder with diverse causes leading to a range of phenotypes. While this complexity is mirrored in other common cardiovascular diseases, such as coronary artery disease, heterogeneity is

particularly marked for HF. Beyond the scope of conventional GWAS consortia, HERMES has a strong focus on the development and clinical validation of multi-modal definitions for HF in an effort to harmonize across different study designs and healthcare contexts. It is recognized that existing clinical classifiers may not optimally enrich for common disease mechanisms,¹⁸ and HERMES seeks new opportunities to dissect out disease heterogeneity using genomic and data science approaches.¹⁹ We describe a stepwise strategy for phenotype definition, starting with the clinical syndrome of HF and moving towards disease subtypes defined with precision. The approach allows for the definition of HF subtypes based on our emerging understanding, without prior assumptions about disease stratification.¹⁸

A substantial number of individuals with Mendelian disorders causing HF, such as dilated or hypertrophic cardiomyopathies, are included. We aim to develop polygenic scores for HF and component traits that may be useful in anticipating the likely penetrance and expression of rare variants associated with Mendelian cardiomyopathies. Inclusion of large longitudinal studies, including clinical trials and electronic health records-linked datasets, offers an opportunity to explore longitudinal phenotypes of HF onset and progression, which are likely to be essential for clarifying the key underlying causal mechanisms.

In future work, we aim to build on the HERMES collaborative platform through more detailed harmonization of covariates and imaging data across studies, enabling analysis at the individual participant level or under a distributed analysis model. Such a framework will enable the platform to support analysis of emerging data-driven definitions of HF subtypes with complex specifications, including those relating to trajectories of disease. We plan to extend our collaborative efforts to include other genome-scale molecular measurements, including serum proteomics and metabolomics, and to include populations with diverse ancestry.

Figure 4 Power estimates across the allele frequency spectrum for genome-wide association studies of heart failure risk and prognosis in HERMES. Figure illustrating empirical power for detecting different genetic variant effect sizes by varying minor allele frequencies (MAF), for (A) heart failure risk (odds ratio) and (B) heart failure prognosis (hazard ratio). Based on current HERMES sample size, with 949 888 controls compared with 44 016 cases for risk and 68 157 cases for prognosis.



The emergence of large genetic studies linked to information on HF and related traits presents an exciting opportunity to explore the causal biology of this increasingly prevalent disorder. HERMES provides a framework for scientific collaboration in support of this aim, bringing together relevant data resources and leading domain experts to address this challenging phenotype. The collaboration is open; we invite interested patients, providers, and researchers to participate and join in our efforts to inform new approaches to the prevention and treatment of HF.

Acknowledgements

The HERMES investigators express their gratitude to participants across all cohorts for making this work possible.

Conflict of interest

Daniel I. Swerdlow is an employee of Silence Therapeutics plc. Joshua D. Backman and Jonathan H. Chung are employees of Regeneron Genetics Center. Simon de Denus was supported through grants from Pfizer, AstraZeneca, Roche Molecular Science, DalCor, and Novartis. Bruce M. Psaty serves on the Steering Committee of the Yale Open Data Access Project funded by Johnson & Johnson. Carolina Roselli is supported by a grant from Bayer AG to the Broad Institute focused on the development of therapeutics for cardiovascular disease. Jean-Claude Tardif has received research support from Amarin, AstraZeneca, DalCor, Ionis, Pfizer, RegenexBio, Sanofi, and Servier and honoraria from AstraZeneca, DalCor, Pfizer, Sanofi, and Servier; holds minor equity interest in DalCor; and is an author of a patent on pharmacogenomics-guided CETP inhibition. Benoit Tyl receives full-time salary from Servier. Harvey D. White reports grants and personal fees from Eli Lilly and Company, Omthera Pharmaceuticals, Pfizer USA, Eisai Inc., DalCor Pharma UK Inc, CSL Behring LLC, American Regent, Sanofi-Aventis Australia Pty Ltd, and Esperion Therapeutics Inc. and personal fees from Genentech, Inc., outside the submitted work. Steven A. Lubitz receives sponsored research support from Bristol Myers Squibb/Pfizer, Bayer AG, Boehringer Ingelheim, and Fitbit and has consulted for Bristol Myers Squibb/Pfizer and Bayer AG. Michael E. Dunn is an employee of Regeneron Pharmaceuticals. Marie-Pierre Dubé has received honoraria from Dalcor, holds minor equity interest in DalCor, is an author of a patent on pharmacogenomics-guided CETP inhibition, and has received research support (access to samples and data) from AstraZeneca, Pfizer, Servier, Sanofi, and GlaxoSmithKline. Authors affiliated with deCODE genetics are employed by deCODE genetics/Amgen Inc.

Funding

R. Thomas Lumbers is supported by a UKRI Rutherford Fellowship hosted by Health Data Research UK (MR/S003754/1), the NIHR UCLH Biomedical Research Centre, and the EU/EFPIA Innovative Medicines Initiative 2 Joint Undertaking BigData@Heart grant no. 116074. The Heart Failure Clinical Research Network and the research reported in this article were supported by the National Heart, Lung, and Blood Institute of the National Institutes of Health (NIH) under award U10 HL084904 (for the Coordinating Center) and awards U10 HL110297, U10 HL110342, U10 HL110309, U10 HL110262, U10 HL110338, U10 HL110312, U10 HL110302, U10 HL110336, and U10 HL110337 (for Regional Clinical Centers). Albert Henry is supported by the British Heart Foundation Cardiovascular Biomedicine PhD studentship. Simon de Denus holds the Université de Montréal Chair in Pharmacogenomics. John J.V. McMurray is supported by a British Heart Foundation Centre of Research Excellence Grant RE/18/6/34217. Jerome I. Rotter was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. Jean-Claude Tardif holds the Canada Research Chair in Personalized Medicine and the Université de Montréal Pfizer-endowed research chair in atherosclerosis. Kent D. Taylor is supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. Harvey D. White reports grants from National Heart, Lung, and Blood Institute. Steven A. Lubitz is supported by NIH grant 1R01HL139731 and American Heart Association 18SFRN34250007. Marie-Pierre Dubé holds the Canada Research Chair in Precision medicine data analysis. Ramachandran S. Vasan acknowledges the support of contracts for the Framingham Heart Study (FHS) NO1-HC-25195, HHSN268201500001, and 75N92019D00031 from the National Heart, Lung, and Blood Institute. He is also supported in part by the Evans Medical Foundation and the Jay and Louis Coffman Endowment from the Department of Medicine, Boston University School of Medicine. J. Gustav Smith was supported by grants from the Swedish Heart-Lung Foundation (2016-0134, 2016-0315, and 2019-0526), the Swedish Research Council (2017-02554), the European-Research Council (ERC-STG-2015-679242), the Crafoord Foundation, Skåne University Hospital, the Scania county, governmental funding of clinical research within the Swedish National Health Service, a generous donation from the Knut and Alice Wallenberg Foundation to the Wallenberg Center for Molecular Medicine in Lund, and funding from the Swedish Research Council (Linnaeus grant Dnr 349-2006-

237, Strategic Research Area Exodiab Dnr 2009-1039) and Swedish Foundation for Strategic Research (Dnr IRC15-0067) to the Lund University Diabetes Center.

Table S1. Summary of participating studies: design, case ascertainment, genotyping and follow-up.

Table S2. Characteristics of participating studies.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

References

- Ziaiean B, Fonarow GC. Epidemiology and aetiology of heart failure. *Nat Rev Cardiol* 2016; **13**: 368–378.
- Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JG, Coats AJ, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, Jessup M, Linde C, Nihoyannopoulos P, Parissis JT, Pieske B, Riley JP, Rosano GM, Ruilope LM, Ruschitzka F, Rutten FH, van der Meer P, Authors/Task Force Members. Document Reviewers 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail* 2016; **18**: 891–975.
- Martin N, Manoharan K, Thomas J, Davies C, Lumbers RT. Beta-blockers and inhibitors of the renin-angiotensin aldosterone system for chronic heart failure with preserved ejection fraction. *Cochrane Database Syst Rev (Online)* 2018; **6**: CD012721.
- Chandrasekera PC, Pippin JJ. The human subject: an integrative animal model for 21(st) century heart failure research. *Am J Transl Res* 2015; **7**: 1636–1647.
- Smith JG, Newton-Cheh C. Genome-wide association studies of late-onset cardiovascular disease. *J Mol Cell Cardiol* 2015; **83**: 131–141.
- Elliott P, Andersson B, Arbustini E, Bilinska Z, Cecchi F, Charron P, Dubourg O, Kühl U, Maisch B, McKenna W, Monserrat L, Pankuweit S, Rapezzi C, Seferovic P, Tavazzi L, Keren A. Classification of the cardiomyopathies: a position statement from the European Society Of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2008; **29**: 270–276.
- Lee DS, Pencina MJ, Benjamin EJ, Wang TJ, Levy D, O'Donnell CJ, Nam BH, Larson MG, D'Agostino RB, Vasani RS. Association of parental heart failure with risk of heart failure in offspring. *N Engl J Med* 2006; **355**: 138–147.
- Lindgren MP, PirouziFard M, Smith JG, Sundquist J, Sundquist K, Zoller B. A Swedish Nationwide Adoption Study of the heritability of heart failure. *JAMA Cardiol* 2018; **3**: 703–710.
- Lindgren MP, Smith JG, Li X, Sundquist J, Sundquist K, Zoller B. Familial mortality risks in patients with heart failure—a Swedish sibling study. *J Am Heart Assoc* 2018; **7**: e010181.
- Smith NL, Felix JF, Morrison AC, Demissie S, Glazer NL, Loehr LR, Cupples LA, Dehghan A, Lumley T, Rosamond WD, Lieb W, Rivadeneira F, Bis JC, Folsom AR, Benjamin E, Aulchenko YS, Haritunians T, Couper D, Murabito J, Wang YA, Stricker BH, Gottdiener JS, Chang PP, Wang TJ, Rice KM, Hofman A, Heckbert SR, Fox ER, O'Donnell CJ, Uitterlinden AG, Rotter JI, Willerson JT, Levy D, van Duijn CM, Psaty BM, Witteman JCM, Boerwinkle E, Vasani RS. Association of genome-wide variation with the risk of incident heart failure in adults of European and African ancestry: a prospective meta-analysis of the cohorts for heart and aging research in genomic epidemiology (CHARGE) consortium. *Circ Cardiovasc Genet* 2010; **3**: 256–266.
- Morrison AC, Felix JF, Cupples LA, Glazer NL, Loehr LR, Dehghan A, Demissie S, Bis JC, Rosamond WD, Aulchenko YS, Wang YA, Haritunians T, Folsom AR, Rivadeneira F, Benjamin EJ, Lumley T, Couper D, Stricker BH, O'Donnell CJ, Rice KM, Chang PP, Hofman A, Levy D, Rotter JI, Fox ER, Uitterlinden AG, Wang TJ, Psaty BM, Willerson JT, van Duijn CM, Boerwinkle E, Witteman JCM, Vasani RS, Smith NL. Genomic variation associated with mortality among adults of European and African ancestry with heart failure: the cohorts for heart and aging research in genomic epidemiology consortium. *Circ Cardiovasc Genet* 2010; **3**: 248–255.
- Smith JG, Felix JF, Morrison AC, Kalogeropoulos A, Trompet S, Wilk JB, Gidlöf O, Wang X, Morley M, Mendelson M, Joeannes R, Ligthart S, Shan X, Bis JC, Wang YA, Sjögren M, Ngwa J, Brandimarto J, Stott DJ, Aguilar D, Rice KM, Sesso HD, Demissie S, Buckley BM, Taylor KD, Ford I, Yao C, Liu C, CHARGE-SCD consortium, EchoGen consortium, QT-IGC consortium, CHARGE-QRS consortium, Sotoodehnia N, van der Harst P, Stricker BHC, Kritchevsky SB, Liu Y, Gaziano JM, Hofman A, Moravec CS, Uitterlinden AG, Kellis M, van Meurs JB, Margulies KB, Dehghan A, Levy D, Olde B, Psaty BM, Cupples LA, Jukema JW, Djousse L, Franco OH, Boerwinkle E, Boyer LA, Newton-Cheh C, Butler J, Vasani RS, Cappola TP, Smith NL. Discovery of genetic variation on chromosome 5q22 associated with mortality in heart failure. *PLoS Genet* 2016; **12**: e1006034.
- Aragam KG, Chaffin M, Levinson RT, McDermott G, Choi SH, Shoemaker MB, Haas ME, Weng LC, Lindsay ME, Smith JG, Newton-Cheh C, Roden DM, London B, For the GRADE Investigators, Wells QS, Ellinor PT, Kathiresan S, Lubitz SA, Bloom HL, Dudley SC, Shalaby AA, Weiss R, Gutmann R, Saba S. Phenotypic refinement of heart failure in a national biobank facilitates genetic discovery. *Circulation* 2019; **139**: 489–501.
- Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath AC, Martin NG, Montgomery GW, Goddard ME, Visscher PM. Common SNPs explain a large proportion of the heritability for human height. *Nat Genet* 2010; **42**: 565–569.
- Kuchenbaecker KB, McGuffog L, Barrowdale D, Lee A, Soucy P, Dennis J, Lush M, Robson M, Spurdle AB, Ramus SJ. Evaluation of polygenic risk scores for breast and ovarian cancer risk prediction in BRCA1 and BRCA2 mutation carriers. *J Natl Cancer Inst* 2017; **109**: djw302.
- Smith GD, Hemani G. Mendelian randomization: genetic anchors for causal inference in epidemiological studies. *Hum Mol Genet* 2014; **23**: R89–R98.

17. de Bakker PI, Ferreira MA, Jia X, Neale BM, Raychaudhuri S, Voight BF. Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum Mol Genet* 2008; **17**: R122–R128.
18. Triposkiadis F, Butler J, Abboud FM, Armstrong PW, Adamopoulos S, Atherton JJ, Backs J, Bauersachs J, Burkhoff D, Bonow RO, Chopra VK, de Boer RA, de Windt L, Hamdani N, Hasenfuss G, Heymans S, Hulot JS, Konstam M, Lee RT, Linke WA, Lunde IG, Lyon AR, Maack C, Mann DL, Mebazaa A, Mentz RJ, Nihoyannopoulos P, Papp Z, Parissis J, Pedrazzini T, Rosano G, Rouleau J, Seferovic PM, Shah AM, Starling RC, Tocchetti CG, Trochu JN, Thum T, Zannad F, Brutsaert DL, Segers VF, de Keulenaer GW. The continuous heart failure spectrum: moving beyond an ejection fraction classification. *Eur Heart J* 2019; **40**: 2155–2163.
19. Hemingway H, Asselbergs FW, Danesh J, Dobson R, Maniadarakis N, Maggioni A, van Thiel GJM, Cronin M, Brobert G, Vardas P, Anker SD, Grobbee DE, Denaxas S, Innovative Medicines Initiative 2nd programme, Big Data for Better Outcomes, BigData@Heart Consortium of 20 academic and industry partners including ESC. Big data from electronic health records for early and late translational cardiovascular research: challenges and potential. *Eur Heart J* 2018; **39**: 1481–1495.
20. Ledford H. AstraZeneca launches project to sequence 2 million genomes. *Nature* 2016; **532**: 427.
21. Nelson MR, Johnson T, Warren L, Hughes AR, Chisoe SL, Xu CF, Waterworth DM. The genetics of drug efficacy: opportunities and challenges. *Nature Rev Genet* 2016; **17**: 197–206.
22. Regeneron Genetics Center, Shah S, Henry A, Roselli C, Lin H, Sveinbjörnsson G, Fatemifar G, Hedman ÅK, Wilk JB, Morley MP, Chaffin MD, Helgadottir A, Verweij N, Dehghan A, Almgren P, Andersson C, Aragam KG, Årnlöv J, Backman JD, Biggs ML, Bloom HL, Brandimarto J, Brown MR, Buckbinder L, Carey DJ, Chasman DI, Chen X, Chen X, Chung J, Chutkow W, Cook JP, Delgado GE, Denaxas S, Doney AS, Dörr M, Dudley SC, Dunn ME, Engström G, Esko T, Felix SB, Finan C, Ford I, Ghanbari M, Ghasemi S, Giedraitis V, Giulianini F, Gottdiener JS, Gross S, Guðbjartsson DF, Gutmann R, Haggerty CM, van der Harst P, Hyde CL, Ingelsson E, Jukema JW, Kavousi M, Khaw KT, Kleber ME, Köber L, Koekemoer A, Langenberg C, Lind L, Lindgren CM, London B, Lotta LA, Loring RC, Luan J, Magnusson P, Mahajan A, Margulies KB, März W, Melander O, Mordi IR, Morgan T, Morris AD, Morris AP, Morrison AC, Nagle MW, Nelson CP, Niessner A, Niiranen T, O'Donoghue ML, Owens AT, Palmer CNA, Parry HM, Perola M, Portilla-Fernandez E, Psaty BM, Rice KM, Ridker PM, Romaine SPR, Rotter JI, Salo P, Salomaa V, van Setten J, Shalaby AA, Smelser DT, Smith NL, Stender S, Stott DJ, Svensson P, Tammesoo ML, Taylor KD, Teder-Laving M, Teumer A, Thorgeirsson G, Thorsteinsdottir U, Torp-Pedersen C, Trompet S, Tyl B, Uitterlinden AG, Veluchamy A, Völker U, Voors AA, Wang X, Wareham NJ, Waterworth D, Weeke PE, Weiss R, Wiggins KL, Xing H, Yerges-Armstrong LM, Yu B, Zannad F, Zhao JH, Hemingway H, Samani NJ, McMurray JJV, Yang J, Visscher PM, Newton-Cheh C, Malarstig A, Holm H, Lubitz SA, Sattar N, Holmes MV, Cappola TP, Asselbergs FW, Hingorani AD, Kuchenbaecker K, Ellinor PT, Lang CC, Stefansson K, Smith JG, Vasana RS, Swerdlow DI, Lumbers RT. Genome-wide association and Mendelian randomisation analysis provide insights into the pathogenesis of heart failure. *Nat Commun* 2020; **11**: 163.
23. Preuss M, König IR, Thompson JR, Erdmann J, Absher D, Assimes TL, Blankenberg S, Boerwinkle E, Chen L, Cupples LA, Hall AS, Halperin E, Hengstenberg C, Holm H, Laaksonen R, Li M, März W, McPherson R, Musunuru K, Nelson CP, Burnett MS, Epstein SE, O'Donnell CJ, Quertermous T, Rader DJ, Roberts R, Schillert A, Stefansson K, Stewart AF, Thorleifsson G, Voight BF, Wells GA, Ziegler A, Kathiresan S, Reilly MP, Samani NJ, Schunkert H, CARDIoGRAM Consortium. Design of the Coronary ARtery Disease Genome-Wide Replication And Meta-Analysis (CARDIoGRAM) study: a genome-wide association meta-analysis involving more than 22 000 cases and 60 000 controls. *Circ Cardiovasc Genet* 2010; **3**: 475–483.
24. Mosaterd A, Deckers JW, Hoes AW, Nederpel A, Smeets A, Linker DT, Grobbee DE. Classification of heart failure in population based research: an assessment of six heart failure scores. *Eur J Epidemiol* 1997; **13**: 491–502.
25. Denaxas S, Gonzalez-Izquierdo A, Direk K, Fitzpatrick NK, Fatemifar G, Banerjee A, Dobson RJB, Howe LJ, Kuan V, Lumbers RT, Pasa L, Patel RS, Shah AD, Hingorani AD, Sudlow C, Hemingway H. UK phenomics platform for developing and validating EHR phenotypes: CALIBER. *J Am Med Inform Assoc* 2019; **26**: 1545–1559.
26. Dehghan A, Bis JC, White CC, Smith AV, Morrison A, Cupples LA, Trompet S, Chasman DI, Lumley T, Völker U, Buckley BM, Ding J, Jensen MK, Folsom AR, Kritchevsky SB, Girman CJ, Ford I, Dörr M, Salomaa V, Uitterlinden AG, Eiriksdottir G, Vasana RS, Franceschini N, Carty CL, Virtamo J, Demissie S, Amouyel P, Arveiler D, Heckbert SR, Ferrières J, Ducimetière P, Smith NL, Wang YA, Siscovick DS, Rice KM, Wiklund PG, Taylor KD, Evans A, Kee F, Rotter JI, Karvanen J, Kuulasmaa K, Heiss G, Kraft P, Launer LJ, Hofman A, Markus MRP, Rose LM, Silander K, Wagner P, Benjamin EJ, Lohman K, Stott DJ, Rivadeneira F, Harris TB, Levy D, Liu Y, Rimm EB, Jukema JW, Völzke H, Ridker PM, Blankenberg S, Franco OH, Gudnason V, Psaty BM, Boerwinkle E, O'Donnell CJ. Genome-wide association study for incident myocardial infarction and coronary heart disease in prospective cohort studies: the CHARGE consortium. *PLoS One* 2016; **11**: e0144997.
27. Smith JG. Molecular epidemiology of heart failure: translational challenges and opportunities. *JACC Basic Transl Sci* 2017; **2**: 757–769.
28. The Genetic Investigation of Anthropometric Traits (GIANT) Consortium, Winkler TW, Day FR, Croteau-Chonka DC, Wood AR, Locke AE, Mägi R, Ferreira T, Fall T, Graff M, Justice AE, Luan J, Gustafsson S, Randall JC, Vedantam S, Workalemahu T, Kilpeläinen TO, Scherag A, Esko T, Kutalik Z, Heid IM, Luos RJJ. Quality control and conduct of genome-wide association meta-analyses. *Nat Protoc* 2014; **9**: 1192–1212.
29. Skol AD, Scott LJ, Abecasis GR, Boehnke M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet* 2006; **38**: 209–213.
30. Owzar K, Li Z, Cox N, Jung SH. Power and sample size calculations for SNP association studies with censored time-to-event outcomes. *Genet Epidemiol* 2012; **36**: 538–548.