



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Supplementing prediction by EuroSCORE with social and patient-reported measures among patients undergoing cardiac surgery

Cromhout, Pernille F.; Thygesen, Lau C.; Moons, Philip; Nashef, Samer; Damgaard, Sune; Christensen, Anne V.; Rasmussen, Trine B.; Borregaard, Britt; Thrysoee, Lars; Thorup, Charlotte B.; Mols, Rikke E.; Juel, Knud; Berg, Selina K.

Published in:
Journal of Cardiac Surgery

DOI (link to publication from Publisher):
[10.1111/jocs.15227](https://doi.org/10.1111/jocs.15227)

Publication date:
2021

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Cromhout, P. F., Thygesen, L. C., Moons, P., Nashef, S., Damgaard, S., Christensen, A. V., Rasmussen, T. B., Borregaard, B., Thrysoee, L., Thorup, C. B., Mols, R. E., Juel, K., & Berg, S. K. (2021). Supplementing prediction by EuroSCORE with social and patient-reported measures among patients undergoing cardiac surgery. *Journal of Cardiac Surgery*, 36(2), 509-521. Advance online publication. <https://doi.org/10.1111/jocs.15227>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

This is the peer reviewed version of the following article: Cromhout, PF, Thygesen, LC, Moons, P, et al. Supplementing prediction by EuroSCORE with social and patient-reported measures among patients undergoing cardiac surgery. *J Card Surg.* 2021; 36: 509– 521., which has been published in final form at <https://doi.org/10.1111/jocs.15227>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

Supplementing prediction by EuroSCORE with social and patient reported measures among patients undergoing cardiac surgery

Pernille Fevejle Cromhout¹, Lau Caspar Thygesen², Philip Moons^{3,4}, Samer Nashef⁵, Sune Damgaard⁶, Anne V. Christensen⁷, Trine B. Rasmussen⁸, Britt Borregaard⁹, Lars Thrysoee¹⁰, Charlotte B. Thorup¹¹, Rikke Elmose Mols¹², Knud Juel², Selina Kikkenborg Berg^{2,7}

Original research

Affiliations

¹ Department of Cardiothoracic Anaesthesiology, Copenhagen University Hospital, Copenhagen, Denmark

² The National Institute of Public Health, University of Southern Denmark, Copenhagen., Denmark

³ KU Leuven Department of Public Health and Primary Care, KU Leuven - University of Leuven, Leuven, Belgium

⁴ Institute of Health and Care Sciences, University of Gothenburg, Gothenburg, Sweden

⁵ Department of Cardiothoracic Surgery, Papworth Hospital, Cambridge, United Kingdom

⁶ Department of Cardiothoracic Surgery, Copenhagen University Hospital, Copenhagen, Denmark

⁷ Department of Cardiology, Copenhagen University Hospital, Copenhagen, Denmark

⁸ Department of Cardiology, Herlev and Gentofte University Hospital, Hellerup, Denmark

⁹ Department of Cardiothoracic and Vascular Surgery, Odense University Hospital, Odense, Denmark

¹⁰ Department of Cardiology, Odense University Hospital, Odense, Denmark

¹¹ Department of Cardiology, Cardiac surgery & Clinical Nursing Research Unit, Aalborg University Hospital, Aalborg, Denmark

¹² Department of Cardiology, Aarhus University Hospital, Aarhus N, Denmark

Corresponding author

Pernille F. Cromhout. Department of Thoracic Anaesthesiology, Copenhagen University Hospital, Blegdamsvej 9, 2100 Copenhagen, Denmark. Pernille.fevejle.cromhout@regionh.dk

Word count

Abstract, 249

Text, 3787 (including title page, abstract, text, figures and legends)

Abstract

Objectives

The risk of poor outcomes is traditionally attributed to biological and physiological processes in cardiac surgery. However, evidence exists that other factors, such as emotional, behavioral, social and functional, are predictive of poor outcomes. Objectives were to evaluate the predictive value of several emotional, social, functional and behavioral factors on four outcomes; death within 90 days, prolonged stay in intensive care, prolonged hospital admission and readmission within 90 days following cardiac surgery.

Methods

This prospective study included adults undergoing cardiac surgery 2013-2014, including information on register-based socio-economic factors and self-reported health in a nested subsample. Logistic regression analyses to determine the association and incremental value of each candidate predictor variable were conducted. Multiple regression analyses were used to determine the incremental value of each candidate predictor variable, as well as discrimination and calibration based on AUC and Brier score.

Results

Of 3217 patients, 3% died, 9% had prolonged intensive care stay, 51% had prolonged hospital admission and 39% were readmitted to hospital. Patients living alone (OR, 1.19; 95% CI, 1.02-1.38), with lower educational levels (1.27; 1.04-1.54) and low health-related quality of life (1.43; 1.02-2.01) had prolonged hospital admission. Analyses revealed living alone as predictive of prolonged ICU stay (Brier, 0.08; AUC, 0.68), death (0.03; 0.71) and prolonged hospital admission (0.24; 0.62).

Conclusion

Living alone was found to supplement EuroSCORE in predicting death, prolonged hospital admission and prolonged ICU stay following cardiac surgery. Low educational level and impaired health-related quality of life were, furthermore, predictive of prolonged hospital admission.

Key words

Risk assessment; Cardiac surgery; Risk factors; Prognostic research; Patient-reported outcomes

Introduction

In cardiac surgery, surgical advances and an increasingly older population have resulted in more complex and high-risk patients being offered surgical procedures. Risk assessment in cardiac surgery aims to reduce poor outcomes by including individual differences in patient profiles and surgical complexity, as well as in surgeons' technical performance ¹.

There are several prognostic screening tools currently available for patients undergoing cardiac surgery. Established clinical risk factors included are history of previous heart surgery, the severity of coronary artery disease, and the degree of comorbidity. In most of Europe, EuroSCORE ^{2,3} is used in clinical practice. The total risk is calculated by adding scores from several risk factors (e.g. chronic pulmonary disease, serum creatinine and left ventricular dysfunction) resulting in a predicted percentage of surgical mortality. The score is defined to distinguish low (<3), moderate (3-5) and high-risk (6+) groups ^{2,3}. However, according to several validation studies ⁴⁻⁶ EuroSCORE is inaccurate in predicting mortality rates. Increasing evidence indicates that non-physiological factors, such as patients' emotional, behavioral, social and functional status are predictive of poor outcomes following cardiac surgery ⁷. Thus, disparities exist in traditional risk assessment in cardiac surgery designed to advise patients of their operative risk of death.

The complex nature of risk assessment and the lack of parameters representing the emotional, social, functional and behavioral lives of patients point toward the need for new definitions of risk and new approaches to risk management in cardiac surgery. The current study is a comprehensive research project developing a risk stratification model as a supplement to EuroSCORE. Emotional, social, functional and behavioral factors will be included to investigate the predictive value on mortality, prolonged stay in the intensive care unit (ICU), prolonged hospitalization and readmissions within 90 days after cardiac surgery.

The aim of this study is therefore to describe the predictive value of emotional, social, functional and behavioral factors and outcomes of death, prolonged length of stay in the ICU, prolonged length of

hospital admission and readmissions following cardiac surgery. The potential predictive factors are tested as a supplement to EuroSCORE.

Methods

This is a prospective study with follow-up until 90 days after cardiac surgery. The protocol for this study has been published.⁸

Study population and data sources

Patients having undergone cardiac surgery from April 2013 to 2014, aged 18 or above were eligible for inclusion. Information were obtained through national Danish registers for the total cohort: the Danish National Patient Register (DNPR) (cardiac surgery procedures, length of hospitalization)⁹, the Danish Civil Registration System (date of birth, sex, cohabitation status, migration, vital status)¹⁰, Danish Education Registers (educational level)¹¹ and the Danish Register on Personal Income (disposable income)¹². Information on socio-economic factors included educational level, income and cohabitation status. Educational level was categorized as basic school (≤ 10 years), upper secondary or vocational education, and higher education. Income was categorized in three groups according to median; $\leq 50\%$, $>50\%-150\%$, and $>150\%$. Cohabitation was defined as being married or living with a partner. Living alone included singles, divorced and widowed.

Furthermore, analyses were done for a nested subsample of patients that participated in the national cross-sectional survey DenHeart¹³. All patients discharged from a Danish Heart Centre were asked to fill out a questionnaire at hospital discharge to evaluate patient reported outcomes, reporting information on health-related quality of life (HRQoL), emotional and cognitive functioning, as well as questions about health behavior.¹³

Information on EuroSCORE and length of ICU stay was received from two clinical databases¹⁴.

Patient-reported outcomes (DenHeart)

HRQoL was measured using the 12-Item Short-Form Health Survey (SF12) and the HeartQoL questionnaires.

The (SF-12) is a generic measure of self-rated health constituting a measure of mental (MCS) and physical (PCS) health. Higher scores (0-100) indicate better perceived health.¹⁵ As recommended the cut-off was set as the mean minus one standard deviation, using the Danish normed score¹⁶.

The HeartQol is a disease-specific tool, scored from 0 (poor) to 3 (best)¹⁷. Scores are summarized in a global, a physical, and an emotional subscale score. For this study HeartQol quantities were converted to binary quantities based on the median score. Both the SF-12 and the HeartQoL questionnaires have a 4 week recall period.

Emotional and cognitive perceptions were measured by the Brief Illness Perception Questionnaire (B-IPQ). Higher scores (0-10) indicates stronger perceptions.¹⁸ No clear cut-offs for screening have been determined for B-IPQ. To reduce the degrees of freedom only the summary score was included for the main analyses in three categories based on the 25th and 75th quartile in the studied sample.

Loneliness was assessed by two ancillary questions, which have previously been used and tested in the Danish National Health Survey¹⁹. One question concerned whether patients experienced having someone to talk to if they needed support or were having problems, and the second question if they were alone, though preferring to be with others.

For health behavior, patients reported status of current or previous smoking behavior and alcohol intake during a typical week, as well as, current height and weight.

Outcomes

Mortality is a reliable and clinically important outcome in cardiac surgery; however, duration of hospitalization and stay in the ICU are common endpoints in cardiac surgical studies. The ICU stay is a standard component of the treatment and provides an indication of the patient's recovery profile and

is in effect a composite measure of the entire perioperative process.²⁰ Readmission is frequent, why it is an outcome with significant health and economic implications. Readmission rates are about 15% at 30 days after discharge^{21,22}, but varies greatly after 30 days from 19 to 56%^{22,23}. Thus, four outcomes were included, 1) death within 90 days of cardiac surgery, 2) prolonged stay in the ICU (≥ 72 hours), 3) prolonged hospital admission (≥ 10 days) and 4) readmission within 90 days from the time of cardiac surgery. Each outcome was evaluated in separate models.

Death

From the Danish Civil Registration System information on all-cause mortality within 90 days from cardiac surgery was obtained.

Prolonged length of stay

Length of stay was included as number of days in the ICU (LOS-ICU), as well as total length of hospital stay (LOS-HOSP). Length of hospital stay, and ICU stay were dichotomized to designate normal and prolonged length of stay. There is no consensus on the definition of prolonged length of stay in hospital following cardiac surgery. Previous studies have adopted the 75th percentile of the length of stay distribution, while others have defined prolonged length of stay as hospitalization for 10 or more days following cardiac surgery^{24,25}, which was used in this study.

In previous studies, prolonged length of stay in the ICU has been defined as from >24 to as much as >96 hours^{24,26–29}. For the present study, based on the existing literature and clinical framework, prolonged length of stay in the ICU was defined as >72 hours since a postoperative ICU stay lasting more than three days is most likely to include almost all patients suffering postoperative complications.

Readmission

Information on rehospitalization was obtained from the DNPR and was included as a dichotomous outcome of readmission within 90 days following cardiac surgery.

Statistical analysis

Baseline characteristics at time of admission were described using means and standard deviations (SD) for continuous measures and percentages for categorical measures.

Initially, logistic regression analyses were conducted to investigate the association between each candidate predictor variable and outcomes for both the total and DenHeart population. Using logistic regression models, we estimated odds ratio (OR) for death, readmission, LOS-ICU and LOS-HOSP adjusting for (1) age (10 years intervals) and sex, and (2) EuroSCORE I.

The number of missing values in the register-based data was low for educational level ($n = 110$ (3%)) and income ($n = 28$ (<1%)), however, to determine the best model based on variable selection, data were imputed, by assigning missing for educational level to basic education and missing for income to the median value. For the DenHeart population of 982 patients, 456 patients had missing data in one or more variables. However, for 18 (62%) of 29 items missingness was $\leq 4\%$. Thus, single mean imputation for each item was conducted for continuous variables whilst for categorical variables (smoking and loneliness), imputations were done by assigning missing to the category most frequently occurring, since missingness was <5% (see Supplementary Table 1).

To determine the incremental value of each candidate predictor variable, each of the predictor variables were excluded separately in a multiple regression model by using an automated backwards selection procedure with a set liberal significance level of 0.10. EuroSCORE was maintained in the models. The Receiver Operating Characteristic (ROC) curve including Area Under the Curve (AUC) and Brier score were used to determine discrimination and calibration, respectively ^{30,31}.

All analyses were conducted using SAS version 9.4.

Results

Demographic and outcome distribution

Total cohort

The study population comprised 3217 (82%) of a total of 3904 patients aged 18 years or above having undergone cardiac surgery from April 2013 to April 2014 (Figure 1). Median age was 68 (range 23 – 95), with almost 70% being 60-79 years, 76% were male and 2085 (65%) had a spouse or partner. Most patients (2340 (73%)) had an income of 50-150% of the median for the total population. Isolated CABG were performed on 1548 (48%) patients and 707 (22%) had non-isolated procedures of cardiac surgery (Table 1). Of the total population, 110 (3.4%) patients died within 90 days, 286 (9%) patients experienced prolonged ICU stay, 1653 (51%) patients prolonged hospital admission and 1249 (39%) patients were readmitted to hospital within 90 days after cardiac surgery. A total of 542 (14%) patients were excluded due to lack of follow-up data in the national registers. Furthermore, we had to exclude 145 (3%) patients for whom we did not have information on EuroSCORE (Figure 1, flow-chart).

DenHeart cohort

In the nested DenHeart sub-sample, including a total of 982 patients, 771 (79%) were male, the median age was 67, 320 (33%) lived alone and 705 (72%) had an income between 50-150% of the median (Table 1). Only 1 patient in the nested subsample died, 63 (6%) patients experienced prolonged LOS-ICU, 443 (45%) patients experienced prolonged LOS-HOSP and 348 (35%) patients were readmitted to hospital within 90 days after cardiac surgery. Since only one patient died in the DenHeart nested sub-sample, death was not included as an outcome in these analyses. In total 1576 (49%) did not participate in the DenHeart cross-sectional survey and were therefore excluded from the nested sub-sample (see flow-chart, Figure 1).

Individual candidate predictor associations in logistic models

Total cohort

Logistic analyses adjusted for EuroSCORE revealed that patients who were living alone (OR, 1.19; 95%CI, 1.02-1.38) and had a lower educational level (OR, 1.27; 95% CI, 1.04-1.54) were more likely to experience prolonged LOS-HOSP, whilst patients in the highest quartile for income were less likely to experience prolonged LOS-HOSP (OR, 0.83; 95% CI, 0.69-0.98) (Table 2).

Regarding prolonged LOS-ICU, patients who lived alone had an increased OR when adjusting for sex and age (OR, 1.33; 95% CI, 1.03-1.70), but the association was not present when adjusting for EuroSCORE (Table 2). The outcomes of readmission and death did not show any statistically significant associations.

DenHeart cohort

Logistic regression analyses in the nested DenHeart population revealed that a lower score on the physical component scale of the SF-12 was associated with LOS-HOSP when adjusting for age and sex, but not when adjusting for EuroSCORE. However, the mental component scale was associated with LOS-HOSP in both models (OR, 1.43; 95% CI, 1.02-2.01) and with LOS-ICU when adjusting for age and sex, but not when adjusted for EuroSCORE. Furthermore, a global score lower than the median on the HeartQoL questionnaire and a high score on the B-IPQ were associated with LOS-HOSP in both models (OR 1.35; 95% CI, 1.04-1.75 and OR, 1.58; 95% CI, 1.09-2.29, respectively). By contrast a medium score on the B-IPQ was found to be associated with LOS-ICU. Finally, being alone though preferring to be with others was associated with an increased OR for readmission within 90 days (OR, 1.41; 95% CI, 1.03-1.91) and not having someone to talk to was associated with an increased OR for LOS-HOSP (OR, 1.95; 95%CI, 1.25-3.04) when adjusting for EuroSCORE (Table 3).

Prediction equation of emotional, social, functional and behavioral factors

Total cohort

The multiple regression models based on comprehensive data revealed low educational level and living alone as predictors of prolonged LOS-HOSP (Table 4). The discriminative value was acceptable based on AUC of 0.625, however, the Brier score of 0.238 indicates a poor informative model (Table 4). Furthermore, living alone was found to be a predictor of LOS-ICU with an acceptable Brier score of 0.078 and AUC of 0.676 (Table 4), and death with a Brier score of 0.032 and AUC of 0.710 (Table 4). None of the candidate variables were predictive of readmission including EuroSCORE, which did not predict readmissions either (AUC, 0.53; Brier score, 0.24) (Table 4 and Supplementary figure 1).

DenHeart cohort

The multiple regression model for prolonged LOS-HOSP included a low global HeartQoL score and not having someone to talk to as predictors (Table 5). The discrimination was acceptable (AUC 0.62), however, the informative value of the model was poor (Brier score of 0.24) (Table 5). None of the candidate predictors in the DenHeart study were found to be predictive of prolonged LOS-ICU (Table 5), however, being alone though preferring to be with others predicted readmission (Table 5). Model fit was found to be poor (Brier score of 0.23), and discriminating ability was low (AUC of 0.56) (Table 5 and Supplementary figure 2).

All candidate predictor variables were tested for correlation with EuroSCORE. No or weak correlations were found (Gamma coefficients were between -0.19 and 0.20). Furthermore, correlation was tested among the included social factors. Educational level and income were found to be substantially correlated (Gamma coefficient, 0.511)

Discussion

In this cohort study emotional, social, functional and behavioral prognostic factors were tested for patients undergoing cardiac surgery. The principal findings were that 1) living alone predicted both

prolonged ICU stay and death for the total cohort of 3217 patients, and 2) low HRQoL and loneliness (not having someone to talk to) predicted prolonged hospital stay for the nested cohort of 982 patients undergoing cardiac surgery. Thus, information on cohabitation status may potentially be added to existing risk evaluation models due to its predictive value.

The predictive value of living alone is supported by Murphy & colleagues who found patients undergoing CABG surgery and living alone, were more than three times more likely to be readmitted to hospital (OR, 3.42; 95% CI, 1.38– 8.48) than those living with others ³². Being married, especially being in a highly satisfying marriage, has been found to offer a significant benefit to long-term survival after CABG (OR, 2.49; 95% CI, 1.47–4.24) ³³. The beneficial effect of cohabitation and relationship satisfaction on survival is likely multifactorial, which has been emphasized by earlier studies linking living alone with poor health outcomes. Patients who are socially isolated are more likely to smoke and have excessive alcohol intake ^{34,35}, delay seeking treatment ³⁶, and demonstrate non-compliance with medical regimens ³⁷, which may be due to a lack of emotional or practical support gained through living with another person ³².

In earlier studies a feeling of loneliness has been linked to several adverse health outcomes. For example, endorsing “yes” to “I feel lonely” was associated with increased 30-day (Rate Ratio (RR), 2.61; 95% CI, 1.15-5.95) and 5-year (RR, 1.78; 95% CI, 1.17-2.71) mortality among patients undergoing CABG ³⁸, and a response of “often” to the question “do you feel lonely” was associated with increased cardiovascular mortality among elderly Danish men (Hazard Ratio, 1.70; 95% CI, 1.03-2.81) ³⁹.

Several studies agree that HRQoL has become a necessary addition and key indicator of cardiac surgical outcomes ^{40–42}. This study found that reduced health-related quality of life predicts prolonged LOS-HOSP. The predictive value of HRQoL has been confirmed in earlier studies that have found low HRQoL to be predictive of both mortality following CABG with a 10 point lower SF-36 Physical

Component Summary score having an OR of 1.39; 95% CI, 1.11-1.77 ⁴³ and of one year cardiac functional status (OR, 2.73; 95% CI, 1.43–5.23) ⁴⁴.

For this study the intention was to investigate factors beyond the clinical indicators and physical health of the patients planned to undergo cardiac surgery. Traditional risk assessment in cardiac surgery has been a tool for patient selection and has been aimed at the perioperative patient pathways. With the proposed supplement the risk assessment can potentially be used to identify vulnerable groups of patients leading to improved patient management still with the overall aim to improve patient outcomes. Information on cohabitation status, loneliness and HRQoL could potentially be added to existing risk evaluation models in cardiac surgery. However, further research is warranted to validate the findings of the current study and to investigate interventions supporting the identified vulnerable groups of patients.

Strength and limitations

This study has several limitations. Firstly, we were restricted to the use of predictor variables based on existing data measured in previously collected data sets, which is a beneficial way to make full use of already collected data to address potentially important new research questions and avoid disturbing patients unnecessarily. However, we may not have included important prognostic variables (e.g. cognitive status and frailty), because they were not measured in the original studies. Secondly, the present study used corresponding datasets. When doing this there is a risk that the datasets differ in important aspects, such as baseline risk. However, in the current study a prediction model was developed for each dataset reducing bias due to this.

The risk score calculations available for this study are calculated based on EuroSCORE I and not the updated EuroSCORE II, why there is a risk of misclassification of high-risk patients due to use of the outdated model, however, EuroSCORE I has been found to over-estimate ⁴⁵ and EuroSCORE II to under-estimate mortality. ^{46–48} It is therefore reasonable to assume that the potential improvement of the model is not attributable to the choice of EuroSCORE model.

Non-response for the DenHeart study was high at 49% which might bias the results. Responders and non-responders of the DenHeart study has earlier been established to be similar regarding socio-demographics, however, the non-responders were more severely ill, had more comorbidity and thus a much higher mortality rate compared to responders ⁴⁹, which could have resulted in an underestimation of the associations between the predictor variables and the outcomes .

Imputations were utilized in the present study to maintain the sample size, assuming the missing values were missing at random. The use of mean imputations does not affect the estimate of the mean for the variable; however, it reduces the variance of the imputed variables. Furthermore, it assumes that the mean value of the respondents was a good estimate of the missing values, which may have resulted in conservative bias.

We used an automated stepwise approach to specify the models, principally due to its objectivity and that it generally results in smaller, clinically applicable models ⁵⁰, but stepwise methods have well-known limitations such as unstable variable selection ⁵¹ and biased coefficient estimation ⁵⁰. It is therefore conceivable that our choice to use stepwise selection may have reduced the predictive performance of the models. The overall model fit statistics indicate that the variance explained by our prediction models is at best modest. Perhaps some factors that are yet to be tested thoroughly in cardiac surgery, for example, frailty and mental state, explain additional variance in cardiac surgery. Despite the limitations of the study the models made informative predictions that should be externally validated in a similar population of patients undergoing cardiac surgery.

Finally, there was a possibility of interaction effects between candidate predictors, however, we assumed that the effect of each variable was persistent, independently of other factors.

Conclusion

We tested several emotional, social, functional and behavioural prognostic factors as a supplement to EuroSCORE and reported different aspects of model performance that can be interpreted for further research applications. Based on the cohorts included, living alone predicts death, prolonged hospital admission and prolonged ICU stay following cardiac surgery. Low educational level and impaired HRQoL were, furthermore found to be predictive of prolonged hospital admission.

Ethics approval

According to Danish legislation, surveys should only be approved by the Danish Data Protection agency (2007-58-0015/30-0937). Use of register data were permitted by The Danish national Board of health.

Funding

This work was supported by Helsefonden, [16-B-0205], the Danish Heart Foundation, [18-R124-A8454-22099] and Fabrikant Karl G. Andersens Fond, [9292].

Declaration of conflicting interests

None

Data availability

The information used in the analysis combines several Danish administrative registers (as described in the paper) and patient-reported questionnaires. The data use is subject to the European Union's General Data Protection Regulation (GDPR). The data were physically stored on computers at Statistics Denmark. Due to security considerations, the data cannot be transferred to computers outside Statistics Denmark. To obtain access to the register data employed in this paper researchers are required to submit a written application to gain approval from Statistics Denmark as well as the

DenHeart committee. The application must include a detailed description of the proposed project, including its purpose, and social contribution, as well as a description of the required datasets, variables, and analysis population. Applications can be submitted by researchers who are affiliated with Danish institutions accepted by Statistics Denmark, or by researchers outside of Denmark who collaborate with researchers affiliated with these institutions.

References

1. Prins C, de Villiers Jonker I, Botes L, et al. Cardiac surgery risk-stratification models. *Cardiovasc J Afr* 2012; 23: 160–4.
2. Nashef SA, Roques F, Michel P, et al. European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardiothorac Surg* 1999; 16: 9–13.
3. Nashef SAM, Roques F, Sharples LD, et al. Euroscore II. *Eur J Cardio-thoracic Surg* 2012; 41: 734–745.
4. Garcia-Valentin A, Mestres CA, Bernabeu E, et al. Validation and quality measurements for EuroSCORE and EuroSCORE II in the Spanish cardiac surgical population: a prospective, multicentre study†. *Eur J Cardiothorac Surg* 2015; 49: 399–405.
5. Borracci RA, Rubio M, Celano L, et al. Prospective validation of EuroSCORE II in patients undergoing cardiac surgery in Argentinean centres. *Interact Cardiovasc Thorac Surg* 2014; 18: 539–43.
6. Zhang G, Wang C, Wang L, et al. Validation of EuroSCORE II in Chinese Patients Undergoing Heart Valve Surgery. *Heart Lung Circ* 2013; 22: 606–611.
7. Cromhout PF, Moons P, Thygesen LC, et al. Time to expand risk evaluation systems for cardiac surgery? Looking beyond physiological parameters. *Eur J Cardiovasc Nurs* 2018; 17: 760–766.
8. Cromhout PF, Berg SK, Moons P, et al. Updating EuroSCORE by including emotional, behavioural, social and functional factors to the risk assessment of patients undergoing cardiac surgery: a study protocol. *BMJ Open* 2019; 9: 26745.
9. Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. *Scand J Public Health* 2011; 39: 30–3.

10. Pedersen CB. The Danish Civil Registration System. *Scand J Public Health* 2011; 39: 22–5.
11. Jensen VM, Rasmussen AW. Danish education registers. *Scand J Public Health* 2011; 39: 91–4.
12. Baadsgaard M, Quitzau J. Danish registers on personal income and transfer payments. *Scand J Public Health* 2011; 39: 103–105.
13. Berg SK, Svanholm J, Lauberg A, et al. Patient-reported outcomes at hospital discharge from Heart Centres, a national cross-sectional survey with a register-based follow-up: the DenHeart study protocol. *BMJ Open* 2014; 4: 4709.
14. Özcan C, Juel K, Lassen JF, et al. The Danish Heart Registry. *Clin Epidemiol* 2016; 8: 503–508.
15. Ware J, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey : Construction of Scales and Preliminary Tests of Reliability and Validity. *Med Care* 1996; 34: 220–233.
16. Gandek B, Ware JE, Aaronson NK, et al. Cross-validation of item selection and scoring for the SF-12 Health Survey in nine countries: results from the IQOLA Project. International Quality of Life Assessment. *J Clin Epidemiol* 1998; 51: 1171–8.
17. Oldridge N, Hofer S, McGee H, et al. The HeartQoL: Part I. Development of a new core health-related quality of life questionnaire for patients with ischemic heart disease. *Eur J Prev Cardiol* 2014; 21: 90–97.
18. Broadbent E, Petrie KJ, Main J, et al. The brief illness perception questionnaire. *J Psychosom Res* 2006; 60: 631–7.
19. Christensen AI, Ekholm O, Glümer C, et al. The Danish National Health Survey 2010. Study design and respondent characteristics. *Scand J Public Health* 2012; 40: 391–397.
20. Myles PS. Meaningful outcome measures in cardiac surgery. *J Extra Corpor Technol* 2014; 46: 23–7.

21. Hannan EL, Racz MJ, Walford G, et al. Predictors of readmission for complications of coronary artery bypass graft surgery. *JAMA* 2003; 290: 773–80.
22. Iribarne A, Chang H, Alexander JH, et al. Readmissions after cardiac surgery: Experience of the national institutes of health/canadian institutes of health research cardiothoracic surgical trials network. *Ann Thorac Surg* 2014; 98: 1274–80.
23. Sibilitz KL, Berg SK, Thygesen LC, et al. High readmission rate after heart valve surgery: A nationwide cohort study. *Int J Cardiol* 2015; 189: 96–104.
24. Almashrafi A, Alsabti H, Mukaddirov M, et al. Factors associated with prolonged length of stay following cardiac surgery in a major referral hospital in Oman: a retrospective observational study. *BMJ Open* 2016; 6: e010764.
25. Papachristofi O, Klein AA, Mackay J, et al. Effect of individual patient risk, centre, surgeon and anaesthetist on length of stay in hospital after cardiac surgery: Association of Cardiothoracic Anaesthesia and Critical Care (ACTACC) consecutive cases series study of 10 UK specialist centres. *BMJ Open* 2017; 7: e016947.
26. Azarfarin R, Ashouri N, Totonchi Z, et al. Factors influencing prolonged ICU stay after open heart surgery. *Res Cardiovasc Med* 2014; 3: e20159.
27. Atoui R, Ma F, Langlois Y, et al. Risk factors for prolonged stay in the intensive care unit and on the ward after cardiac surgery. *J Card Surg* 2008; 23: 99–106.
28. Ettema RGA, Peelen LM, Schuurmans MJ, et al. Prediction models for prolonged intensive care unit stay after cardiac surgery: systematic review and validation study. *Circulation* 2010; 122: 682–9.
29. Janssen DPB, Noyez L, Wouters C, et al. Preoperative prediction of prolonged stay in the intensive care unit for coronary bypass surgery. *Eur J Cardiothorac Surg* 2004; 25: 203–7.

30. Harrell FE, Califf RM, Pryor DB, et al. Evaluating the yield of medical tests. *JAMA* 1982; 247: 2543–6.
31. Gerds TA, Cai T, Schumacher M. The performance of risk prediction models. *Biometrical J* 2008; 50: 457–479.
32. Murphy BM, Elliott PC, Le Grande MR, et al. Living alone predicts 30-day hospital readmission after coronary artery bypass graft surgery. *Eur J Cardiovasc Prev Rehabil* 2008; 15: 210–215.
33. King KB, Reis HT. Marriage and long-term survival after coronary artery bypass grafting. *Heal Psychol* 2012; 31: 55–62.
34. Dickens CM, McGowan L, Percival C, et al. Lack of a close confidant, but not depression, predicts further cardiac events after myocardial infarction. *Heart* 2004; 90: 518–22.
35. Haustein K-O. Smoking and poverty. *Eur J Cardiovasc Prev Rehabil* 2006; 13: 312–8.
36. Schwarz B, Schoberberger R, Rieder A, et al. Factors delaying treatment of acute myocardial infarction. *Eur Heart J* 1994; 15: 1595–8.
37. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med* 2000; 160: 2101–7.
38. Herlitz J, Wiklund I, Caidahl K, et al. The Feeling of Loneliness Prior to Coronary Artery Bypass Grafting Might be a Predictor of Short-and Long-term Postoperative Mortality. *Eur J Vasc Endovasc Surg* 1998; 16: 120–125.
39. Olsen RB, Olsen J, Gunner-Svensson F, et al. Social networks and longevity. A 14 year follow-up study among elderly in Denmark. *Soc Sci Med* 1991; 33: 1189–1195.
40. Tully PJ. Quality-of-Life measures for cardiac surgery practice and research: a review and

primer. *J Extra Corpor Technol* 2013; 45: 8–15.

41. Noyez L, de Jager MJ, Markou ALP. Quality of life after cardiac surgery: underresearched research. *Interact Cardiovasc Thorac Surg* 2011; 13: 511–4.
42. Grady KL, Lee R, Subačius H, et al. Improvements in Health-Related Quality of Life Before and After Isolated Cardiac Operations. *Ann Thorac Surg* 2011; 91: 777–83.
43. Rumsfeld JS, MaWhinney S, McCarthy M, et al. Health-Related Quality of Life as a Predictor of Mortality Following Coronary Artery Bypass Graft Surgery. *JAMA* 1999; 281: 1298–303.
44. Falcoz P-E, Chocron S, Stoica L, et al. Open Heart Surgery: One-Year Self-Assessment of Quality of Life and Functional Outcome. *Ann Thorac Surg* 2003; 76: 1598–604.
45. Gogbashian A, Sedrakyan A, Treasure T. EuroSCORE: A systematic review of international performance. *European Journal of Cardio-thoracic Surgery* 2004; 25: 695–700.
46. Garcia-Valentin A, Mestres CA, Bernabeu E, et al. Validation and quality measurements for EuroSCORE and EuroSCORE II in the Spanish cardiac surgical population: a prospective, multicentre study†. *Eur J Cardiothorac Surg*. Epub ahead of print 11 March 2015. DOI: 10.1093/ejcts/ezv090.
47. Noyez L, Kievit PC, van Swieten HA, et al. Cardiac operative risk evaluation: The EuroSCORE II, does it make a real difference? *Neth Heart J* 2012; 20: 494–8.
48. Kalender M, Adademir T, Tasar M, et al. Validation of EuroSCORE II risk model for coronary artery bypass surgery in high-risk patients. *Kardiochirurgia i Torakochirurgia Pol* 2014; 11: 252–256.
49. Berg SK, Thorup CB, Borregaard B, et al. Patient-reported outcomes are independent predictors of one-year mortality and cardiac events across cardiac diagnoses: Findings from the national DenHeart survey. *Eur J Prev Cardiol* 2018; 204748731876976.

50. Steyerberg EW. *Clinical Prediction Models A Practical Approach to Development, Validation, and Updating*. New York: Springer Science & Business Media. Epub ahead of print 2005. DOI: 10.1016/B978-0-12-387667-6.00013-0.
51. Austin PC, Tu J V. Automated variable selection methods for logistic regression produced unstable models for predicting acute myocardial infarction mortality. *J Clin Epidemiol* 2004; 57: 1138–1146.

Total cohort			Nested cohort	
	N	%	N	%
<u>Sex</u>				
Women	777	24	211	21
Men	2440	76	771	79
<u>Age</u> (range 23-95)				
≤49	238	7	82	8
50-59	488	15	154	16
60-69	1118	35	361	37
70-79	1108	34	328	33
80+	265	8	57	6
Median (25%, 75% percentiles)	68 (60, 74)		67 (60, 73)	
Mean (SD)	66 (11)		65(11)	
IQR	14		13	
<u>Cohabitation status</u>				
Living with a partner	2085	65	662	67
Living alone	1132	35	320	33
<u>Educational level</u>				
Basic school	1190	37	342	35
Upper secondary or vocational education	1382	43	434	44
Higher education	645	20	206	21
<u>Equivalised disposable income</u> (median 177.577 DKK.)				
≤50% median	188	6	70	7
>50%-150% median	2340	73	705	72
>150% median	689	21	207	21
<u>Surgical procedure</u>				
CABG, isolated	1548	48	450	46
Aortic valve, isolated	597	19	186	19
Mitral valve, isolated	175	5	55	6
Aorta, isolated	142	4	37	4
Other, isolated	48	1	18	2
Non-isolated procedures	707	22	236	24
Reoperation	98	3	29	0.03
<u>EuroSCORE I</u>				
High	1518	47	413	42
Moderate	1124	35	197	20
Low	575	18	372	38

Table 1. Clinical and socio-demographic factors for the total and nested cohort.

		LOS-HOSP ⁱ		LOS-ICU ⁱⁱ		Readmissions		Death	
		Model 1 ^A	Model 2 ^B	Model 1 ^A	Model 2 ^B	Model 1 ^A	Model 2 ^B	Model 1 ^A	Model 2 ^B
		OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)
Educational level									
Basic school		1.27 (1.05-1.55)	1.27 (1.04-1.54)	0.82 (0.59-1.14)	0.81 (0.58-1.12)	0.96 (0.79-1.17)	0.93 (0.76-1.13)	0.93 (0.54-1.62)	1.04 (0.61-1.80)
Upper secondary / vocational		1.18 (0.98-1.43)	1.22 (1.01-1.48)	0.86 (0.62-1.18)	0.88 (0.64-1.22)	0.90 (0.74-1.09)	0.90 (0.74-1.09)	1.07 (0.62-1.82)	1.14 (0.67-1.95)
Higher education		1.00 (Reference)	#	#	#	#	#	#	#
Age group	≤49	1.07 (0.81-1.42)		1.04 (0.63-1.71)		1.53 (1.15-2.02)		0.32 (0.08-1.34)	
	50-59	0.81 (0.65-1.01)		0.57 (0.36-0.91)		1.22 (0.98-1.52)		0.74 (0.35-1.58)	
	60-69	1.00 (Reference)		#		#		#	
	70-79	1.21 (1.03-1.44)		1.17 (0.87-1.57)		0.93 (0.78-1.11)		1.45 (0.89-2.36)	
	≥80	2.12 (1.60-2.82)		1.83 (1.22-2.75)		1.18 (0.90-1.56)		4.37 (2.53-7.55)	
Sex	Female	1.10 (0.93-1.29)		1.36 (1.04-1.78)		0.97 (0.82-1.15)		1.81 (1.21-2.71)	
	Male	1.00 (Reference)		#		#		#	
EuroSCORE Low			0.70 (0.57-0.85)		0.52 (0.29-0.93)		0.86 (0.69-1.06)		0.36 (0.11-1.25)
Moderate			1.00 (Reference)		#		#		#
High			2.01 (1.72-2.36)		3.26 (2.40-4.44)		1.14 (0.97-1.33)		4.44 (2.59-7.61)
Equivalised disposable income									
≤50% median		1.06 (0.78-1.43)	1.03 (0.76-1.40)	1.19 (0.74-1.91)	1.18 (0.74-1.90)	1.22 (0.90-1.66)	1.20 (0.90-1.62)	0.93 (0.44-1.99)	1.03 (0.49-2.16)
>50%-150% median		1.00 (Reference)	#	#	#	#	#	#	#
≥150% median		0.77 (0.65-0.92)	0.83 (0.69-0.98)	0.75 (0.53-1.06)	0.84 (0.59-1.18)	1.01 (0.84-1.21)	1.10 (0.92-1.31)	0.64 (0.35-1.18)	0.65 (0.36-1.18)
Age group	≤49	1.07 (0.81-1.42)		1.05 (0.64-1.73)		1.53 (1.15-2.02)		0.32 (0.08-1.35)	
	50-59	0.83 (0.67-1.03)		0.59 (0.37-0.93)		1.22 (0.99-1.52)		0.76 (0.36-1.64)	
	60-69	1.00 (Reference)		#		#		#	
	70-79	1.20 (1.01-1.41)		1.13 (0.85-1.52)		0.93 (0.78-1.11)		1.39 (0.85-2.26)	
	≥80	2.10 (1.58-2.79)		1.75 (1.17-2.63)		1.18 (0.90-1.56)		4.17 (2.41-7.19)	
Sex	Female	1.07 (0.91-1.27)		1.28 (0.98-1.69)		0.96 (0.81-1.14)		1.71 (1.14-2.57)	
	Male	1.00 (Reference)		#		#		#	
EuroSCORE Low			0.71 (0.57-0.87)		0.53 (0.30-0.95)		0.86 (0.69-1.06)		0.38 (0.11-1.29)
Moderate			1.00 (Reference)		#		#		#
High			1.99 (1.70-2.33)		3.17 (2.33-4.31)		1.15 (0.98-1.34)		4.26 (2.49-7.30)

Cohabitation status								
Living with a partner	1.00 (Reference)	#	#	#	#	#	#	#
Living alone	1.24 (1.07-1.44)	1.19 (1.02-1.38)	1.33 (1.03-1.71)	1.26 (0.98-1.62)	0.95 (0.82-1.10)	0.95 (0.82-1.10)	1.36 (0.91-2.03)	1.42 (0.96-2.08)
Age group	≤49	1.05 (0.79-1.38)	1.02 (0.62-1.68)		1.54 (1.16-2.04)		0.31 (0.07-1.31)	
	50-59	0.80 (0.65-1.00)	0.57 (0.36-0.90)		1.23 (0.99-1.52)		0.73 (0.34-1.56)	
	60-69	1.00 (Reference)	#		#		#	
	70-79	1.23 (1.04-1.45)	1.16 (0.87-1.56)		0.93 (0.79-1.11)		1.43 (0.88-2.33)	
	≥80	2.12 (1.60-2.82)	1.76 (1.17-2.64)		1.19 (0.90-1.57)		4.18 (2.42-7.22)	
Sex	Female	1.06 (0.90-1.26)	1.26 (0.96-1.66)		0.99 (0.83-1.17)		1.64 (1.08-2.48)	
	Male	1.00 (Reference)	#		#		#	
EuroSCORE	Low	0.70 (0.57-0.86)		0.52 (0.29-0.93)		0.86 (0.70-1.06)		0.36 (0.11-1.26)
	Moderate	1.00 (Reference)		#		#		#
	High	2.00 (1.71-2.34)		3.16 (2.32-4.30)		1.14 (0.82-1.10)		4.29 (2.50-7.34)

Table 2. Associations between clinical and socio-demographic factors and all outcomes among 3217 patients having undergone cardiac surgery in Denmark, April 2013-April 2014.

ⁱ ≥ 10 days

ⁱⁱ ≥ 72 hours

^A Adjusted for age and sex

^B Adjusted for EuroSCORE

	LOS-HOSP ⁱ		LOS-ICU ⁱⁱ		Readmissions	
	Model 1 ^A	Model 2 ^B	Model 1 ^A	Model 2 ^B	Model 1 ^A	Model 2 ^B
	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)	OR (95%CI)
SF-12 (median, IQR, mean, SD)						
PCS (36.8, 13.2, 17.3, 9.6)						
≤mean-SD(28.2)	1.51 (1.07-2.12)	1.38 (0.98-1.95)	1.74 (0.94-3.22)	1.45 (0.78-2.68)	1.26 (0.89-1.79)	1.28 (0.91-1.81)
>mean-SD(28.2)	1.00 (Reference)	#	#	#	#	#
MCS (48.7, 16.4, 47.4, 11.2)						
≤mean-SD(37.3)	1.53 (1.09-2.14)	1.43 (1.02-2.01)	1.93 (1.06-3.54)	1.62 (0.88-2.95)	1.07 (0.75-1.52)	1.06 (0.75-1.49)
>mean-SD(37.3)	1.00 (Reference)	#	#	#	#	#
HeartQoL						
Global (1.5, 1.0, 1.5, 0.7)						
<median	1.43 (1.10-1.84)	1.35 (1.04-1.75)	1.51 (0.89-2.53)	1.29 (0.76-2.17)	0.99 (0.76-1.29)	1.01 (0.78-1.31)
≥median	1.00 (Reference)	#	#	#	#	#
Physical (1.3, 1.2, 1.3, 0.8)						
<median	1.22 (0.95-1.58)	1.18 (0.91-1.52)	1.15 (0.69-1.92)	1.00 (0.60-1.68)	1.07 (0.82-1.39)	1.08 (0.84-1.41)
≥median	1.00 (Reference)	#	#	#	#	#
Emotional (2.0, 1.3, 2.0, 0.7)						
<median	1.13 (0.87-1.45)	1.08 (0.83-1.40)	1.16 (0.69-1.95)	1.04 (0.62-1.74)	1.11 (0.85-1.45)	1.11 (0.85-1.44)
≥median	1.00 (Reference)	#	#	#	#	#
B-IPQ (46.0, 14.0, 45.5, 10.0)						
≤ 39 (q1)	1.00 (Reference)	#	#	#	#	#
40-53 (q2)	1.34 (0.99-1.81)	1.30 (0.96-1.76)	2.15 (1.08-4.28)	2.05 (1.02-4.09)	0.85 (0.62-1.15)	0.87 (0.64-1.19)
> 53 (q3)	1.61 (1.11-2.33)	1.58 (1.09-2.29)	1.82 (0.79-4.18)	1.70 (0.74-3.91)	0.90 (0.62-1.32)	0.98 (0.67-1.42)
Smoking						

Current	1.16 (0.70-1.92)	1.20 (0.72-2.00)	1.59 (0.64-4.00)	1.74 (0.69-4.39)	0.93 (0.55-1.59)	1.05 (0.62-1.78)
Former	1.00 (Reference)	#	#	#	#	#
Never	0.91 (0.69-1.20)	0.91 (0.68-1.20)	1.26 (0.72-2.20)	1.25 (0.72-2.19)	1.08 (0.81-1.45)	1.10 (0.83-1.47)
<u>Alcohol consumption</u>						
Heavy ^y	1.18 (0.74-1.90)	1.08 (0.67-1.74)	1.45 (0.63-3.34)	1.28 (0.55-2.97)	1.48 (0.91-2.39)	1.44 (0.89-2.31)
> recommendations	0.82 (0.56-1.20)	0.73 (0.52-1.03)	0.85 (0.38-1.91)	0.61 (0.29-1.29)	0.82 (0.55-1.22)	0.85 (0.60-1.22)
≤ recommendations	1.00 (Reference)	#	#	#	#	#
<u>Body Mass Index</u>						
< 30	1.00 (Reference)	#	#	#	#	#
≥ 30	0.95 (0.70-1.30)	1.02 (0.75-1.39)	1.39 (0.78-2.49)	1.58 (0.88-2.85)	0.95 (0.69-1.31)	0.96 (0.70-1.32)
≥30	1.05 (0.75-1.46)	1.11 (0.80-1.55)	1.64 (0.86-3.14)	1.85 (0.96-3.56)	1.01 (0.72-1.42)	1.03 (0.73-1.45)
25-29	1.00 (Reference)	#	#	#	#	#
< 25	1.27 (0.94-1.70)	1.24 (0.93-1.67)	1.47 (0.81-2.70)	1.43 (0.78-2.62)	1.17 (0.86-1.58)	1.19 (0.88-1.61)
<u>Loneliness 1</u>^x						
Yes	1.33 (0.98-1.80)	1.23 (0.91-1.68)	1.60 (0.91-2.82)	1.34 (0.76-2.36)	1.39 (1.02-1.09)	1.41 (1.03-1.91)
No	1.00 (Reference)	#	#	#	#	#
<u>Loneliness 2</u>^{xx}						
Yes	1.00 (Reference)	#	#	#	#	#
No	1.92 (1.24-2.98)	1.95 (1.25-3.04)	1.00 (0.42-2.39)	0.99 (0.41-2.38)	0.81 (0.51-1.29)	0.83 (0.53-1.32)

Table 3. Associations between patient-reported outcomes and all outcomes among 982 patients having undergone cardiac surgery in Denmark and participated in the DenHeart survey.

ⁱ ≥ 10 days

ⁱⁱ ≥ 72 hours

^A OR=odds ratio. Model 1, adjusted for sex and age

^B OR=odds ratio. Model 2, adjusted for EuroSCORE

^y >14 units/week (women); >21 units/week (men)

^x **Does it ever happen that you are alone, even though you would prefer to be with others?**

Yes = Yes, often / Yes, sometimes

No = Yes, but rarely / No

^{xx} **Do you have someone to talk to if you are having problems or need support?**

Yes = Yes, often / Yes, most of the time

No = Yes, sometimes / No, never or rarely

	LOS-HOSP ⁱ			LOS-ICU ⁱⁱ			DEATH			READMISSION		
	Number			Number			Number			Number		
	in	χ^2	P-value	in	χ^2	P-value	in	χ^2	P-value	in	χ^2	P-value
Equivalised disposable income	3	1.76	0.42	3	3.26	0.20	2	1.87	0.39	1	2.17	0.34
Educational level		5.48	0.06	2	1.93	0.38	3	0.47	0.79	3	0.92	0.63
Cohabitation		4.53	0.03		3.34	0.07		3.10	0.08	2	0.24	0.62
EuroScore		133.87	<.0001		86.55	<.0001		42.98	<.0001		8.31	0.02
Fit statistics	AUC	Brier score		AUC	Brier score		AUC	Brier score		AUC	Brier score	
	0.6245	0.2378		0.6759	0.0783		0.7099	0.0324		0.5273	0.2369	

Table 4. Model fitting and variable selection for all outcomes. 3217 patients having undergone cardiac surgery.

ⁱ ≥ 10 days


ⁱⁱ ≥ 72 hours


	LOS-HOSP ⁱ			LOS-ICU ⁱⁱ			READMISSION		
	Number			Number			Number		
	in	χ ²	P-value	in	χ ²	P-value	in	χ ²	P-value
BMI	11	0.01	0.91	1	2.30	0.1294	8	0.34	0.56
Loneliness 1	10	0.25	0.62	9	0.36	0.5462		4.72	0.03
Smoking	9	1.00	0.61	10	1.07	0.5866	10	0.42	0.81
SF-12 MCS	8	1.49	0.22	3	1.36	0.2440	11	0.003	0.95
SF-12 PCS	7	1.66	0.20	8	0.56	0.4530	3	1.76	0.18
HeartQol Physical	6	1.72	0.19	4	0.42	0.5168	6	1.33	0.25
HeartQol Emotional	5	1.27	0.26	6	1.44	0.2305	7	0.65	0.42
B-IPQ	4	3.49	0.17	2	4.56	0.1020	9	1.32	0.52
Alcohol	3	3.93	0.14	7	2.46	0.2924	2	3.66	0.16
HeartQoL Global		4.42	0.04	5	0.82	0.3656	5	0.19	0.66
Loneliness 2		7.72	0.01	11	0.11	0.7437	4	1.55	0.21
EuroScore		32.54	<.0001		19.26	<.0001		1.43	0.49
Fit statistics	AUC		Brier score	AUC		Brier score	AUC		Brier score
	0.6188		0.2357	0.6660		0.0586	0.5554		0.2264

Table 5. Model fitting and variable selection for all outcomes. 982 patients having undergone cardiac surgery and participated in the DenHeart study.


ⁱ ≥ 10 days

ⁱⁱ ≥ 72 hours





Supplementary Figure 1 ROC curves including AUC for discriminative performance for each model. 3217 patients having undergone cardiac surgery in Denmark, April 2013 to April 2014.



Supplementary figure 2. ROC curves including AUC for discriminative performance for each model. 982 patients having undergone cardiac surgery in Denmark April 2013 to April 2014 and who participated in the DenHeart study