



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Perspectives on the treatment of sequelae after cancer

protocol of an interview study of primary care physiotherapists

Toftdahl, Anne Katrine Skjølstrup; Hjoernholm, Laura Hvidaa; Simonsen, Mia; Stapelfeldt, Christina M; Thomsen, Janus Laust; Kongsgaard, Marianne; Riis, Allan

Published in:
BMJ Open

DOI (link to publication from Publisher):
[10.1136/bmjopen-2021-052378](https://doi.org/10.1136/bmjopen-2021-052378)

Creative Commons License
CC BY-NC 4.0

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Toftdahl, A. K. S., Hjoernholm, L. H., Simonsen, M., Stapelfeldt, C. M., Thomsen, J. L., Kongsgaard, M., & Riis, A. (2022). Perspectives on the treatment of sequelae after cancer: protocol of an interview study of primary care physiotherapists. *BMJ Open*, 12(4), e052378. Article e052378. <https://doi.org/10.1136/bmjopen-2021-052378>

General rights


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

BMJ Open Perspectives on the treatment of sequelae after cancer: protocol of an interview study of primary care physiotherapists

Anne Katrine Skjølstrup Toftdahl,¹ Laura Hvidaa Hjoernholm,² Mia Simonsen,³ Christina M Stapelfeldt,^{4,5} Janus Laust Thomsen,^{6,7} Marianne Kongsgaard,² Allan Riis ^{1,7}

To cite: Toftdahl AKS, Hjoernholm LH, Simonsen M, *et al*. Perspectives on the treatment of sequelae after cancer: protocol of an interview study of primary care physiotherapists. *BMJ Open* 2022;**12**:e052378. doi:10.1136/bmjopen-2021-052378

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-052378>).

Received 09 August 2021
Accepted 25 March 2022



© Author(s) or their employer(s) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Allan Riis; ariis@dcm.aau.dk

ABSTRACT

Introduction An increasing number of patients are living with fatigue, pain and other sequelae after cancer. About 30% of these patients express a need for physical rehabilitation and patients prefer to be involved in medical decision-making. This includes being offered individualised treatment with patient-defined goals and education during all aspects of treatment by physiotherapists with empathic skills. However, physiotherapists are often unaware of the relationship between cancer and its presenting symptoms. This can lead to inappropriate care and unrealistic goalsetting for rehabilitation. This calls for greater attention towards physiotherapist's treatment and their current perspectives on how treatment can be improved in physiotherapy clinics. The aim of this study is to explore physiotherapists perspectives on barriers, facilitators and potential solutions to improve the care of sequelae after cancer in physiotherapy clinics.

Methods and analysis This is a qualitative interview study using individual semistructured interviews with physiotherapists using a phenomenological approach to explore their lifeworld at work. We will recruit physiotherapists working in private clinics based on a variation in gender and years of working experience in private clinics. We will conduct between 10 and 16 interviews online via Microsoft Teams and thematically analyse data supported by NVivo software. Interviewing is scheduled to take place from April 2022.

Ethics and dissemination This is a non-intervention and qualitative study, and the local Ethics Committee in the North Denmark Region has stated that their approval is not needed. Informants will provide a written informed consent. Study information will be sent to the informant at least 3 days prior to the interview session and information will be summarised by the interviewer before the interview.

Aside from publication, results will be disseminated by two teaching institutions, a regional health care provider and DEFACTUM (a public Danish research institution focusing on increasing social equality in health).

INTRODUCTION

The number of people surviving cancer is increasing because of advances in early

Strengths and limitations of this study

- The interviewing is planned to take place online and this is a strength due to easing the availability for participation and thereby supporting a variation among participants.
- Online interviewing limits the building of trust and openness which might be more easily achieved during face-to-face interviewing.
- Transferring of results from this study requires careful thought about the similarity of contexts.

detection and treatment.¹ Among the people who survive cancer, 86% experience sequelae after cancer.² Sequelae are defined as problems arising due to the cancer or as side effects to the cancer treatment.³ Fatigue and pain are commonly reported, and other symptoms include tight surgical scars, lymphoedema, reduced functional capability, sexual dysfunction, depressive symptoms and anxiety.⁴

About 30% of patients with sequelae after cancer express a need for physical rehabilitation.⁵ In the literature, most physiotherapy-led interventions include exercise alone or as part of a combined treatment for the patient,⁶ and various types of exercise like yoga, aerobic training and resistance training can be effective in reducing the severity of the sequelae.⁷ Symptoms often present in a cluster with a combination of physical, psychological and social sequelae.^{8,9}

Patients with sequelae after cancer expect to be involved in medical decisions.¹⁰ In physiotherapy practices, this involvement includes an individualised treatment approach, working with patient-defined goals and offering education during all aspects of treatment by physiotherapists with empathic skills.¹¹ However, physiotherapists are often not aware of the relationship between cancer

and the presenting symptoms, and this can lead to an overestimation in goalsetting of the rehabilitation.¹² Regular physical activity reduces the sequelae and physical activity is associated with a decrease in mortality in patients with breast cancer.¹³ However, many patients do not believe that exercise improves their condition.¹⁴

Consequently, the care of sequelae provided by primary care physiotherapists has the potential to be significantly improved by increased attention to the presence of sequelae after cancer, the choice of treatment and goalsetting. This is an issue that the authors have discussed with a panel of patients with sequelae after cancer and a group of general practitioners in a workshop in Denmark.¹⁵ General practitioners suggested more educational material for patients and education of both general practitioners and primary care physiotherapists in identifying and treating sequelae. The workshop highlighted the need for enhanced qualification of the treatment of sequelae after cancer either during the medical education or implemented in the mandatory specialisation to become a general practitioner or introduced as a supplementary course following specialisation as general practitioner.¹⁵ However, we do not know physiotherapists' perspectives on barriers, facilitators and potential solutions to improve their primary care treatment of sequelae after cancer.

The aim of this study is therefore to explore physiotherapists perspectives on barriers, facilitators and potential solutions to improve the care of sequelae after cancer in physiotherapy clinics.

METHODS AND ANALYSIS

This study will be conducted as a qualitative survey with one-on-one interviews with physiotherapists following a semistructured interview guide. The development of this study protocol and the future reporting of the findings is informed by the consolidated criteria for reporting qualitative research (COREQ) checklist.¹⁶

Theoretical framework

To gain an insight into the subjective experiences and perceptions of the physiotherapists' treatment of sequelae after cancer, this study will use a phenomenological approach to data collection and thematic analysis, thereby allowing the phenomena to present themselves to the researchers while putting the researchers' preconceptions aside.¹⁷ Consequently, since the phenomenological approach requires a bracketing of the researchers' preunderstandings, AKST, LHH and AR will state their preunderstandings of the field and expectations to the findings in the interviews.¹⁸ This allows for later validation of the findings and transparency of reporting. This preunderstanding will be combined in a document and shared among the authors before the first interview (pilot interviewing not included). After finalising the analysis, we will revisit our preunderstanding and include this in the discussion section for the reporting on the findings.

Patient and public involvement

In 2020, a panel of four patients with sequelae after cancer was established. The panel has provided the authors with insight into their perspectives and perceived treatment in physiotherapists clinics. Furthermore, the conceptualisation of the overall project was discussed with them and they have pointed to a need for improving information given to patients but also a need for increasing physiotherapists knowledge about sequelae after cancer. The interview guide will be sent to the patient panel for feedback. The patient panel is not involved in recruitment to the study. Findings are presented to the National Patient Association for sequelae after cancer.¹⁹

Interviewer and research team

All interviews will be conducted by AKST. AKST is a female physiotherapist and Master of Science. AKST is currently a senior lecturer at the Department of Physiotherapy. AKST is experienced in interviewing and has previous experience from leading qualitative research. She has received extensive training in interview techniques such as open-ended questions, active listening and probing as part of her work as a physiotherapist and master's degree in Health Science. AKST will be supervised by LHH, MK and AR. Being a physiotherapist, AKST has knowledge about physiotherapy practice and, thereby, place her close to the field but she will have no prior knowledge of the informants.

Interview guide

The interview guide is developed from discussions among the research team, and the questions are informed by themes occurring from informal discussions with patients with sequelae after cancer, carried out by MK prior to the initiation of this study (online supplemental additional file 1). The interview guide will be further validated by feedback from the patient panel, two pilot interviews with physiotherapists, who have a special interest in treating cancer patients, giving them the opportunity to pinpoint relevant topics for further questioning. Finally, the interview guide will be pilot tested on MK, and any need for changes arising from feed-back and pilot testing will be made before the interviews are to be carried out. However, we will also allow for adaptations of the guide following the research interviews.

The interview guide will include open-ended questions that allow the participants to elaborate using examples from their clinical routines. Questions involving possible conflicts are saved for the last part of the interview.

Data collection

Interviewing is planned to take place between primo April and primo September 2022. During the interviews, the interviewer will continuously summarise the participants' accounts to ensure correct understanding and interpretation. All interviews will be audio-recorded and transcribed by LHH using the InqScribe (Inquirium LLC, <https://www.inqscribe.com/>) digital media transcription

software. A predefined transcription key will be used in the process, thus enabling the standardisation of transcription across the interviews. In the case of indistinguishable audio, another member of the research group will listen to the recording. If clarification of what was recorded is not possible, that specific part will not be transcribed or otherwise included in the analysis.

The transcriptions will be verified by AR, by listening to the audio recordings while reading the transcription of the spoken words and checking the quotes. Transcriptions or quotes will not be sent to the informants to be verified, as the interviewer will make a point of vocalising her understanding of the emerging themes during the interview, to verify whether the informant agrees with her understanding. Data will be pseudonymised. An identification key will be stored on a secured server. We will not make field notes.

Setting of data collection

Due to the COVID-19 situation, online interviews will be conducted using the Microsoft Teams online platform. To support and maintain the physiotherapists' role as health-care professionals in their natural working environment, interviews will be conducted in the physiotherapy clinics and preferably in their consultation rooms. During the interview, the informant will be encouraged to participate alone without patients or colleagues present. Likewise, the interviewer will be alone during the interview.

Participants

Physiotherapists working in Denmark in a private clinic will be eligible to participate. Eligible physiotherapists will be identified by AR and MK. The physiotherapists are to be selected using a purposive sampling technique²⁰ aiming for maximum variation in gender and years in clinical practice (in private physiotherapy clinics).

Inclusion criteria

We will include Danish speaking physiotherapists, working in private clinics in Denmark.

Exclusion criteria

We will exclude participants working in a clinic which already had provided one informant. Furthermore, to support a practice-oriented approach, physiotherapists who are not working at least one full day in clinical practice will be excluded from the sample.

Sampling

Based on a narrow research question, the expected similarity in practice organisation and the interviewers' experience with interview techniques, we are aiming for a sample size (n=10–16) until data saturation is reached.²¹ This saturation will be assessed during the data analysis, where new subjects and themes occurring at the end of the analysis will lead to the inclusion of new informants. Further interviews will be carried out, until data are considered to be sufficiently saturated. First, we conduct 10 interviews. If data saturation is not reached, additional

Table 1 Text analysis

Step	Primary action	Purpose
1	Listening to the recordings and reading the transcriptions	Becoming familiar with the data
2	Organising the data in meaning units	Creating initial codes
3	Organising the initial codes	Creating potential themes
4	Going through the coding for each potential theme	Ensuring a coherent pattern in the thematic placement of the codes
5	Identifying the essence of each theme	Themes are given their final labels/names
6	Statements are used to build up the argument for answering the research question	Writing the text and choosing quotes to support the text

The analysis is conducted in six steps according to Braun.²²

interviews in rounds of two will be conducted until saturation is reached.

Method of approach

Possible participants will be identified in national networks of physiotherapists working in private clinics and invited via LinkedIn and other social media. Where they will be invited to contact MK by e-mail or telephone. MK and AR will conduct the final identification of the sample to obtain maximum variation. MK will then send contact information to AKST, who will contact the participants, provide further study information and schedule an hour for the interview. Participants will not be paid or otherwise compensated for their participation. Participants will be aware of the overall purpose of the project but will not have personal knowledge of AKST.

Informed consent

Before participation in the interview, the informant will provide their written informed consent. Study information will be sent to the informant at least 3 days prior to the interview session.

Data analysis

The data will be analysed following a thematic text analysis as described by Braun and Clarke, as this strategy facilitates clear and systematic reflections on underlying themes occurring in the informants' statements.²² The thematic text analysis contains six phases (table 1).

In the initial phase, AKST will familiarise herself with the data by listening to the audio recordings and reading through the transcriptions of the interviews. The aim is to ensure an in-depth familiarity with the data prior to the analytical work. Therefore, the first phase will be concluded when the researcher can briefly summarise the apparent essence of each interview.²² In the second phase,



AKST and LHH will create initial codes and thereby organise the data into meaningful units. This organisation will be carried out using the quantitative coding programme NVivo (QSR International 2021, <https://www.qsrinternational.com/nvivo-qualitative-data-analysis-software/home>), and because of the phenomenological approach, the creation of initial codes will be done in an inductive manner. The inductive coding will focus on the themes appearing from the data and thereby ensure an open and reflexive approach enabling the phenomena to speak for themselves.

The third phase will consist of organising the initial codes in potential themes. Here, the researcher will consider which codes are fitting as possible overarching themes, and which codes constitute subthemes.²¹ The fourth phase will involve going through the coding for each theme to ensure a coherent pattern in the thematic placement of the codes. If the coded statements do not form a coherent pattern in the theme they are currently placed under, the deviating statements will be moved to other themes or the name of the theme will be changed. Furthermore, the depth and width of the coded themes will be discussed to consider whether some themes are considerable larger than others. If so, a possible division into smaller themes will be carried out to clarify what statement they represent.

In the fifth phase, the essence of each theme will be identified, with a focus on how each theme addresses the physiotherapists' perspectives on barriers, facilitators and potential solutions to improve primary care treatment of sequelae after cancer.²¹ Furthermore, the themes will be given their final label/names on the basis of their identified essence. In the sixth and final phase, the written analysis will be composed to enable the informants' statements to be used to build the argument for answering the research question. Quotes will be included to support essential perspectives. Data will be coded and analysed by the interviewer (AKST) with supervision from LHH and AR.

ETHICS AND DISSEMINATION

This is a non-intervention qualitative study and the local Ethics Committee in the North Denmark Region has stated that their approval is not needed (29 March 2021). Informants will provide written informed consent.

This study is conducted as part of a larger research project about developing interventions to improve the primary care treatment of sequelae after cancer in Denmark. The wider research project is conducted in collaboration between the Department of Physiotherapy at University College in Northern Denmark, The Research Unit for General Practice at the Department of Clinical Medicine in Aalborg, the Quality Unit for General Practice in the North Denmark Region and DEFACTUM (a public research institution in Denmark, focusing on increasing social equality in health). Thereby, we will be including practice, healthcare providers and

educational institutions for medical students and physiotherapy students in the project. This is believed to ensure the future implementation of findings from the project. Furthermore, the results will be disseminated via national and international conferences and will be published in a peer-reviewed international journal.

Author affiliations

¹Department of Physiotherapy, University College Northern Denmark, Aalborg, Denmark

²Quality Unit for General Practice in the North Denmark Region, Aalborg, Denmark

³Department of Public Health, Aarhus University, Aarhus, Denmark

⁴Public Health and Quality Improvement, Marselisborg Centret, Central Denmark Region, Aarhus, Denmark

⁵Department of Public Health/Section of Social Medicine and Rehabilitation, Aarhus University, Aarhus, Denmark

⁶Aalborg University Hospital, Aalborg University, Aalborg, Denmark

⁷Research Unit for General Practice in the North Denmark Region, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

Contributors The concept of this study was conceived by AKST, LHH, MS, CMS, JLT, MK and AR. LHH drafted the first version with supervision from AR. All authors commented during rounds of reviews. AKST wrote the last version. All authors read and approved the final version of the manuscript.

Funding This work was supported by the Cancer Group of Nord-KAP. Nord-KAP is the quality unit for the general practice in the North Denmark Region and the Cancer Group works to support a coherent treatment of patients with cancer. Furthermore, AKST's work is supported by internal funding from UCN.

Competing interests None declared.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Allan Riis <http://orcid.org/0000-0002-7009-3025>

REFERENCES

- 1 Miller KD, Siegel RL, Lin CC, *et al*. Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin* 2016;66:271–89.
- 2 Pearce A, Haas M, Viney R, *et al*. Incidence and severity of self-reported chemotherapy side effects in routine care: a prospective cohort study. *PLoS One* 2017;12:e0184360.
- 3 National cancer Institute at the National Institutes of health. USA. Available: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/long-term-side-effect> [Accessed 31 Mar 2021].
- 4 Gegechkori N, Haines L, Lin JJ. Long-Term and latent side effects of specific cancer types. *Med Clin North Am* 2017;101:1053–73.
- 5 Ohlsson-Nevo E, Alkebro I, Ahlgren J. Cancer patients' interest in participating in cancer rehabilitation. *Acta Oncol* 2019;58:1676–83.

- 6 Lin K-Y, Frawley HC, Denehy L, *et al.* Exercise interventions for patients with gynaecological cancer: a systematic review and meta-analysis. *Physiotherapy* 2016;102:309–19.
- 7 Hilfiker R, Meichtry A, Eicher M, *et al.* Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: a systematic review incorporating an indirect-comparisons meta-analysis. *Br J Sports Med* 2018;52:651–8.
- 8 Aziz NM. Cancer survivorship research: state of knowledge, challenges and opportunities. *Acta Oncol* 2007;46:417–32.
- 9 Harrington CB, Hansen JA, Moskowitz M, *et al.* It's not over when it's over: long-term symptoms in cancer survivors--a systematic review. *Int J Psychiatry Med* 2010;40:163–81.
- 10 Hahlweg P, Kriston L, Scholl I, *et al.* Cancer patients' preferred and perceived level of involvement in treatment decision-making: an epidemiological study. *Acta Oncol* 2020;59:967–74.
- 11 Wijma AJ, Bletterman AN, Clark JR, *et al.* Patient-centeredness in physiotherapy: what does it entail? A systematic review of qualitative studies. *Physiother Theory Pract* 2017;33:825–40.
- 12 Gupta E, Ng AH, Fu JB, *et al.* Cancer rehabilitation providers and oncology patient expectations for functional outcomes after inpatient rehabilitation. *Pm R* 2021;13:729–36. doi:10.1002/pmrj.12473
- 13 Ibrahim EM, Al-Homaidh A. Physical activity and survival after breast cancer diagnosis: meta-analysis of published studies. *Med Oncol* 2011;28:753–65.
- 14 Hirschev R, Docherty SL, Pan W, *et al.* Exploration of exercise outcome expectations among breast cancer survivors. *Cancer Nurs* 2017;40:E39–46.
- 15 Riis A, Kongsgaard M, Stapelfeldt CM. *Bedre Behandling AF senfølger efter kræft: Ideskabende workshop ud fra et primærsektorsperspektiv Danish]. PRACTICUS. The Danish College of General Practitioners. PRACTICUS no. 255. 9. Accepted in Press, 2021.*
- 16 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 17 Giorgi A. The theory, practice, and evaluation of the phenomenological method as a qualitative research procedure. *J Phenomenol Psychol* 1997;28:235–60.
- 18 Koch T. Interpretive approaches in nursing research: the influence of Husserl and Heidegger. *J Adv Nurs* 1995;21:827–36.
- 19 Sequele after cancer. Danish patient association. Available: <https://www.senfoelger.dk/om-senfoelger/>
- 20 Merriam SB, Tisdell EJ. *Qualitative research: a guide to design and implementation*. 4th ed, 2015.
- 21 Braun V, Clarke V. To saturate or not to saturate? Questioning data saturation as a useful concept for thematic analysis and sample-size rationales. *Qual Res Sport Exerc Heal* 2019:1–16.
- 22 Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.