



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Exploring the microbial influence on seasonal nitrous oxide concentration in a full-scale wastewater treatment plant using metagenome assembled genomes

Valk, Laura Christina; Peces, Miriam; Singleton, Caitlin Margaret; Laursen, Mads Dyring; Andersen, Mikkel Holmen; Mielczarek, Artur Tomasz; Nielsen, Per Halkjær

*Published in:*  
Water Research

*DOI (link to publication from Publisher):*  
[10.1016/j.watres.2022.118563](https://doi.org/10.1016/j.watres.2022.118563)

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2022

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Valk, L. C., Peces, M., Singleton, C. M., Laursen, M. D., Andersen, M. H., Mielczarek, A. T., & Nielsen, P. H. (2022). Exploring the microbial influence on seasonal nitrous oxide concentration in a full-scale wastewater treatment plant using metagenome assembled genomes. *Water Research*, 219, Article 118563. <https://doi.org/10.1016/j.watres.2022.118563>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Exploring the microbial influence on seasonal nitrous oxide concentration in a full-scale wastewater treatment plant using metagenome assembled genomes

Laura Christina Valk<sup>a,1</sup>, Miriam Peces<sup>a,1</sup>, Caitlin Margaret Singleton<sup>a</sup>, Mads Dyring Laursen<sup>a</sup>, Mikkel Holmen Andersen<sup>b</sup>, Artur Tomasz Mielczarek<sup>c</sup>, Per Halkjær Nielsen<sup>a,\*</sup>

<sup>a</sup> Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, 9220 Aalborg, Denmark

<sup>b</sup> Unisense Environment A/S, Aarhus, Denmark

<sup>c</sup> BIOFOS A/S, Copenhagen, Denmark

## ARTICLE INFO

### Keywords:

Nitrous oxide  
Full-scale WWTP  
Time-series  
Metagenomics  
Genomes

## ABSTRACT

Nitrous oxide is a highly potent greenhouse gas and one of the main contributors to the greenhouse gas footprint of wastewater treatment plants (WWTP). Although nitrous oxide can be produced by abiotic reactions in these systems, biological N<sub>2</sub>O production resulting from the imbalance of nitrous oxide production and reduction by microbial populations is the dominant cause. The microbial populations responsible for the imbalance have not been clearly identified, yet they are likely responsible for strong seasonal nitrous oxide patterns. Here, we examined the seasonal nitrous oxide concentration pattern in Avedøre WWTP alongside abiotic parameters, the microbial community composition based on 16S rRNA gene sequencing and already available metagenome-assembled genomes (MAGs). We found that the WWTP parameters could not explain the observed pattern. While no distinct community changes between periods of high and low dissolved nitrous oxide concentrations were determined, we found 26 and 28 species with positive and negative correlations to the seasonal N<sub>2</sub>O concentrations, respectively. MAGs were identified for 124 species (approximately 31% mean relative abundance of the community), and analysis of their genomic nitrogen transformation potential could explain this correlation for four of the negatively correlated species. Other abundant species were also analysed for their nitrogen transformation potential. Interestingly, only one full-denitrifier (*Candidatus* Dechloromonas phosphorivorans) was identified. 59 species had a *nosZ* gene predicted, with the majority identified as a clade II *nosZ* gene, mainly from the phylum Bacteroidota. A correlation of MAG-derived functional guilds with the N<sub>2</sub>O concentration pattern showed that there was a small but significant negative correlation with nitrite oxidizing bacteria and species with a *nosZ* gene (N<sub>2</sub>O reducers (DEN)). More research is required, specifically long-term activity measurements in relation to the N<sub>2</sub>O concentration to increase the resolution of these findings.

## 1. Introduction

Nitrous oxide (N<sub>2</sub>O) is a highly potent greenhouse gas, with a warming potential 300 times higher than CO<sub>2</sub> (Ravishankara et al., 2009; Terada et al., 2017). N<sub>2</sub>O is one of the main contributors to the greenhouse gas footprint of wastewater treatment plants (WWTP) (Daelman et al., 2013; Delre et al., 2019). Consequently, mitigating N<sub>2</sub>O-emissions would significantly reduce WWTP carbon footprints, but to create mitigation strategies we need to identify and understand the

N<sub>2</sub>O sources.

Emission of N<sub>2</sub>O in WWTPs is due to an imbalance in biotic and abiotic N<sub>2</sub>O production and complete consumption (Conthe et al., 2019). N<sub>2</sub>O is an intermediate of denitrification, where heterotrophic denitrifying bacteria (DEN) transform nitrate (NO<sub>3</sub><sup>-</sup>) to dinitrogen gas (N<sub>2</sub>) (Fig. 1) (Zumft, 1997). The accumulation of N<sub>2</sub>O can be due to incomplete denitrification by heterotrophic denitrifying bacteria, where the last transformation of N<sub>2</sub>O to dinitrogen gas (N<sub>2</sub>) via nitrous oxide reductase (NOS, EC 1.7.2.4; Fig. 1) is not completed (Kampschreur et al.,

\* Corresponding author.

E-mail address: [phn@bio.aau.dk](mailto:phn@bio.aau.dk) (P.H. Nielsen).


<sup>1</sup> These authors contributed equally to this work

<https://doi.org/10.1016/j.watres.2022.118563>

Received 11 February 2022; Received in revised form 5 May 2022; Accepted 6 May 2022

Available online 10 May 2022

0043-1354/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



**Fig. 1.** The nitrogen cycle with the dissimilatory conversions and their associated genes highlighted. The genes with quotation marks have not been included in the KEGG database (release 91, (Kanehisa, 2019)), and have not been included in the annotation. DEN, anoxic heterotrophic denitrifiers; AMX, anammox; CMX, comammox; AOB, ammonium oxidizing bacteria; NOB, nitrite oxidizing bacteria; DNRA, dissimilatory nitrate reduction to ammonium. Nitrifier denitrification was not specifically included for AOB. Adapted from Soler-Jofra et al. (2021).


2009). Additionally,  $N_2O$  is a by-product from ammonia oxidizing bacteria (AOB), converting the intermediate hydroxylamine to  $N_2O$  using cytochrome P460 (Caranto et al., 2016). Another  $N_2O$ -production pathway of AOB is nitrifier denitrification, where the  $NO_2^-$  produced by hydroxylamine dehydrogenase (HAO, EC 1.7.2.6) is reduced to  $N_2O$  via nitrite reductase (NIR, EC 1.7.2.1) and nitric oxide reductase (NOR, EC 1.7.2.5) (Fig. 1; (Caranto and Lancaster, 2017; Domingo-Félez and Smets, 2019; Wrage-Mönnig et al., 2018). Lastly,  $N_2O$  can also be produced via abiotic reactions between intermediates of nitrification and denitrification, or by metal-mediated reactions (Kampschreur et al., 2011; Soler-Jofra et al., 2016).

Although part of the  $N_2O$ -emission from WWTPs can originate from truncated-denitrifiers, lacking the nitrous oxide reductase gene (*nosZ*, K00376), there are also many non-denitrifying organisms that have *nosZ* genes predicted in their genome (Conthe et al., 2019; Hallin et al., 2018; Sanford et al., 2012). These organisms have the potential to reduce  $N_2O$ , and if their total NOS-activity is higher in the WWTP than the total combined activity of the nitric oxide reductase (NOR) and cytochrome P460, then the WWTP would be a  $N_2O$ -sink. The actual NOS-activity within a WWTP will depend on the genetic *nosZ* potential and factors influencing the NOS-activity. Post-translational and transcriptional regulation can cause an imbalance between  $N_2O$ -producing (NOR, cytochrome P460) and  $N_2O$ -consuming (NOS) enzymes (Liu et al., 2013; Lycus et al., 2017). Other factors that can influence the production-consumption balance are oxygen concentration,  $NO_2^-$ -accumulation, a low C/N ratio and storage polymer metabolism (Kampschreur et al., 2008; Law et al., 2012; Lu and Chandran, 2010; Otte et al., 1996; Wunderlin et al., 2012).

Short-term influences are not the only factors influencing  $N_2O$ -emissions. Seasonal abiotic parameters have been suggested to influence

long-term  $N_2O$ -emissions, such as temperature (for WWTPs located in temperate climates), accumulation of nitrogen-species, e.g.,  $NO_2^-$ ,  $NO_3^-$ ,  $NH_4^+$  and influent flow rate (Chen et al., 2019; Daelman et al., 2015; Gruber et al., 2021; Vasilaki et al., 2018). Seasonal community composition variation and dynamics have also been suggested to influence  $N_2O$ -emissions in full-scale WWTPs (Daelman et al., 2015; Gruber et al., 2021). However, most of the abundant organisms in WWTPs are undescribed and have unknown functions. A comprehensive database of WWTP-specific high quality metagenome-assembled genomes (HQ MAGs) was made available only recently; the MiDAS genome database (Singleton et al., 2021). This novel database might provide new insights into the (until now unknown) potential  $N_2O$  producers and consumers responsible for  $N_2O$  emissions.

Here we examine the  $N_2O$  production pattern and microbial community data of a full-scale Danish WWTP with nutrient removal spanning multiple years, combined with the MAG-derived nitrogen-transformation potential of the community. We searched for correlations between the relative abundance of specific species and the  $N_2O$ -production pattern. Species showing a statistically significant correlation were cross-referenced with the species representatives of the previously constructed MiDAS genome database of HQ MAGs (Singleton et al., 2021), and positive identity-hits were investigated for nitrogen-transformation potential. Community members with a substantial abundance (> 0.1% relative abundance) in the WWTP and microorganisms of special interest were also cross-referenced to the species representatives of the HQ MAGs, to ensure no key  $N_2O$ -producer or -consumer were omitted. The genomic  $N_2O$  reduction potential was investigated in more detail, based on clade-type and phylogeny. Additionally, the cumulative relative abundance of the species that were categorised into functional guilds, using metagenome-based



**Fig. 2.** The chemical and community characteristics of the Avedøre WWTP during the sampling period. A) The COD:N ratio, the concentration of nitrate in the effluent ( $\text{g N/m}^3$ ) and the  $\text{N}_2\text{O}$  concentration ( $\text{g N/m}^3$ ) of the Avedøre WWTP during the measured period. B) The growing population and the non-growing population. Species in blue have an identity-hit with a species in the MiDAS genome database, species in orange have no identity-hit.

classification, was compared to the  $\text{N}_2\text{O}$  concentration pattern over three years. This combined approach provided novel insights into the microbiology of nitrogen transformations in full-scale WWTPs, although a direct correlation to the  $\text{N}_2\text{O}$ -production pattern could not be established.

## 2. Methods

### 2.1. Plant process design and data collection

The Avedøre WWTP (Copenhagen, Denmark) has biological nitrogen removal and enhanced biological phosphate removal (EBPR) with a capacity to treat approximately  $70,000 \text{ m}^3$  of wastewater daily, under dry weather conditions, serving approximately 350,000 population equivalents (PE). Key parameters measured during the studied period are shown in Fig. 2A and other operational parameters are shown in Supplemental Data S1. Mainstream treatment consists of screening, grit removal, and primary settlement before biological treatment with activated sludge. Four lines, with two carousel reactors per line, work in parallel and have alternating aeration and anoxic periods. Aeration is controlled by STAR control and is based on the ammonia concentration. The average effluent total N was  $4.7 \pm 2.1 \text{ g N/m}^3$  during the period 2017 - 2020 (Supplemental Data S1). Nitrous oxide was measured in the liquid phase in Carrousel reactor 1 and 3 using two dissolved  $\text{N}_2\text{O}$  sensors (Unisense Environment A/S, Denmark) per line. Only line 3 was considered for this study, as line 1 changed conditions during the period studied.

### 2.2. 16S rRNA gene amplicon sequencing and bioinformatic analysis

Activated sludge biomass was sampled within the MiDAS project as previously described (McIlroy et al., 2015; Nierychlo et al., 2020). Briefly, the activated sludge was sampled every one-two weeks in the aerobic phase. DNA was extracted, prepared and amplified as previously described (Stokholm-Bjerregaard et al., 2017). The V1-V3 16S rRNA gene region was amplified using the 27F (AGAGTTTGATCCTGGCTCAG) (Lane, 1991) and 534R (ATTACCGCGGCTGCTGG) (Muyzer et al., 1993) primers. Forward reads were processed using usearch v11.0.667 (Edgar, 2010) and raw fastq files were filtered for phiX sequences using usearch -filter\_phiX, trimmed to 250 bp using usearch -fastx\_truncate -truncLen 250, and quality filtered using usearch -fastq\_filter with -fastq\_maxee


1.0. Dereplication of the sequences was done using usearch -fastx\_uniques with -sizeout. -unoise3 (Edgar, 2017), with standard settings to generate exact amplicon sequence variants (ASVs). The MiDAS 4 reference database (available at <https://www.midasfieldguide.org/guide/downloads>), combined with the SINTAX classifier with a confidence threshold of 0.8 (Edgar, 2018), was used to assign taxonomy at the species level (Nierychlo et al., 2020). For ASVs without species-level classification, the taxonomy was assigned at the lowest available taxonomic level (e.g. genus), and these ASVs were treated as separate species.

### 2.3. Classification of growing and non-growing species in activated sludge

The majority of species in activated sludge are found in the influent wastewater (Dottorini et al., 2021). Therefore, by means of mass-balances, it is possible to identify whether these species are expected to grow or die in the activated sludge reactors. In this study, we assigned the species into growing and non-growing groups based on the classification obtained by Dottorini et al., 2021. In Dottorini et al., 2021, the classification list is based on a mass-balance between paired influent and activated sludge samples collected from 11 municipal WWTPs across Denmark. Although Avedøre WWTP was not among the 11 WWTPs used to obtain the classification list, Avedøre WWTP has very similar taxa, process design and geographical location. Consequently, it can be assumed that species will follow the same growth groups as described in Dottorini et al., 2021. Species in WWTPs with different process designs may not follow the same growth groups, and mass-balances should be conducted between paired influent and activated sludge samples. The detailed mass-balance methodology can be found elsewhere (Dottorini et al., 2021).

### 2.4. MAG identification, annotation, and metabolic reconstruction

A list of the abundant species (relative read abundance of the 16S rRNA gene of  $>0.1\%$  at least twice in the sampling period) in the Avedøre WWTP was made, resulting in 367 16S rRNA gene defined species. This species list was cross-referenced with the 581 HQ MAGs species representatives previously recovered from Danish WWTPs ((Singleton et al., 2021); Supplemental Data S2). The 581 HQ MAG species representatives are defined as species at the whole genome level (95% ANI), and in cases where multiple genomes were recovered with



**Fig. 3.** Correlation of the relative read abundance (%) of species from the Avedøre WWTP with the seasonal  $\text{N}_2\text{O}$  concentration ( $\text{g N/m}^3$ ). The cumulative relative read abundance of A) positively correlated species, B) negatively correlated species.

>95% ANI to each other, the best genome (most complete, and least contaminated) was selected as the representative genome for the species group (Singleton et al., 2021). HQ MAGs that represented an abundant 16S rRNA gene defined species in Avedøre WWTP, with >98.6% average nucleotide identity (Yarza et al., 2014) across the full-length 16S rRNA gene, were annotated using EnrichM v0.5.0 (<https://github.com/geronimp/enrichM>) annotate with the EnrichM v10 database, incorporating a KO-annotated uniref100 database (Suzek et al., 2015). In some cases, one 16S rRNA gene defined MiDAS species is associated with multiple MAGs, for example, *Tetrasphaera midas\_5* is associated with 6 species representative MAGs. This occurs due to the discrepancy between 16S rRNA gene based and full-genome based species definitions (Singleton et al., 2022). In these instances we examined all the MAGs with a >98.6% identity-hit of their 16S rRNA genes to the 16S rRNA gene defined MiDAS species. R v4.0.5 (R Core Team, 2019, <https://www.R-project.org/>) and R-studio (Wickham et al., 2019) (RStudio Team, 2015) were used for downstream processing. Tidyverse (Wickham et al., 2019) was used to filter the KO-ids associated with nitrogen transformations and RColorBrewer v1.1–2 (<https://cran.r-project.org/web/packages/RColorBrewer/index.html>) was used to make Fig. 6. To reduce the probability of false positive annotations (Mise et al., 2021), a gene was assumed present if all KO-ids of the subunits were identified. Re-annotation of certain HQ MAGs (*Rhodanobacteraceae midas\_s\_2835*) was done with blastp (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). All HQ MAGs with predicted genes associated with ammonia oxidizing bacteria were additionally analysed for the presence of cytochrome p460 using HMMER v3.3.2 (<http://hmmmer.org/>) hmmbuild and hmmsearch with the PF16694 (seed sequence) from the Pfam database (El-Gebali et al., 2019). Sequences predicted by HMM with a sequence E-value of at least  $1 \times 10^{-8}$  were identified as positive hits. The key gene for nitrite oxidation, *nxrAB*, was predicted in five species (Fig. 4B, Supplemental Data S5). As the KEGG database does not distinguish between *narGH* and *nxrAB* (K00370, K00371), a protein tree was made (Supplemental Figure S3) comparing the putative nitrite oxidoreductase (NXR-A) sequence predicted in the HQ MAGs (Supplemental Data S5) with publicly available NXR-A and NAR-G sequences (Supplemental Data S3).


## 2.5. Nitrous oxide reductase and nitrite oxidoreductase protein trees

Fxtract v2.3 (<https://github.com/ctSkennerton/fxtract>) extracted all predicted NOS sequences (K00376) or NXR-A (K00370) from the HQ MAG species representatives. MUSCLE v3.8.31 (Edgar, 2004) was used to make a multisequence alignment as input for IQ-TREE version 2.0.6 (Minh et al., 2020). The best models for the NOS and NXR-A/NAR-G tree, LG + R7 and LG + F + R10 respectively, were identified using ModelFinder (Kalyaanamoorthy et al., 2017) and 1000 iterations were

made using ultrafast bootstrapping with UFBoot (Hoang et al., 2018). The NOS sequences were classified by clade type using GraftM v0.13.1 Graft (Boyd et al., 2018) and the nitrous oxide reductase package ([https://github.com/geronimp/graftm\\_gpkgs](https://github.com/geronimp/graftm_gpkgs)). iTol v5 (Letunic and Bork, 2021) was used to visualize and refine the tree. The positive control sequences for NOS were obtained from the uniprot database ([www.uniprot.org](http://www.uniprot.org)) Supplemental Table S1; A4IP44\_GEOTN, B8FZJ4\_DESHD, NO\_SZ\_PARDE, A0A3B7A9B0\_9RHO0, Q12M27\_SHEDE, I0AHV3\_IGNAJ, A0A0C5BY41\_METNJ, A0A3G6WI33\_RHOSH, NOSZ\_PSEST, NOSZ\_AC\_HCY, Q3SJ28\_THIDA, NOSZ\_BRADU, A0A069Q756\_PSEAI, Q5NZ01\_AROAE, Q6ZXZ8\_WOLSC, C6x281\_FLAB3. The positive control sequences for the NXR-A tree were obtained from the GraftM nitrite oxidoreductase and nitrate reductase package ([https://github.com/geronimp/graftm\\_gpkgs](https://github.com/geronimp/graftm_gpkgs)) Supplemental Data S3.

## 2.6. Data processing of 16S rRNA gene amplicon time-series

Downstream processing of 16S rRNA gene time-series was carried out in R v4.0.5 (R Core Team, 2019, <https://www.R-project.org/>) and R-studio (RStudio Team, 2015) using the following packages: ampvis2 v.2.6.1 (Andersen et al., 2018), tidyverse v.1.3.0 (Wickham et al., 2019) and Hmisc v. 4.5.0 (<https://CRAN.R-project.org/package=Hmisc>). Prior to data analysis, samples with less than 10,000 reads were discarded and duplicate samples from the same sampling point were combined by averaging the relative abundance of each ASV. The total reads per sample ranged from 32,481 to 124,155. The relative abundance of species was visualised with boxplots using the mean relative abundance for growing and non-growing species. Differences in microbial community beta-diversity were explored by non-metric multidimensional scaling (NMDS), using Bray-Curtis dissimilarity distance in ampvis2. Based on the  $\text{N}_2\text{O}$  concentration profile during the 2017–2020 period, two grouping periods have been described *ad-hoc* using the mean  $\text{N}_2\text{O}$  concentration as a threshold value. The two groups are defined as “high” ( $\text{N}_2\text{O}$  concentration was higher than  $0.138 \text{ g N/m}^3$ ) and “low” ( $\text{N}_2\text{O}$  concentration was lower than  $0.138 \text{ g N/m}^3$ ). Pearson correlations between species time-series and  $\text{N}_2\text{O}$  concentration were assessed with the Hmisc package. To approximate the data to a normal distribution, species time-series were transformed using a robust-centered log-ratio transformation and  $\text{N}_2\text{O}$  concentration was squared-root transformed prior correlation. P-values of the correlations were adjusted using the Benjamini & Hochberg method to control the false discovery rate in multiple comparisons (Benjamini and Hochberg, 1995).




**Fig. 4.** The mean abundance (%) and genetic nitrogen transformation potential of bacterial species with a corresponding HQ MAG (Singleton et al., 2021) in the Avedøre WWTP. A) the positive (green dot) and negative (red dot) associated species with N<sub>2</sub>O-production pattern and B) the top 33 most abundant species with a corresponding HQ MAG. All species had a relative abundance of at least 0.1% twice in the sampling period. Heatmap strips indicate the mean relative abundance (%). Genes were considered present if all the subunits of the gene were predicted. (Supplemental Data S2).

### 3. Results

#### 3.1. Avedøre WWTP showed a distinct seasonal N<sub>2</sub>O concentration pattern

The full-scale Avedøre WWTP with biological nitrogen and phosphorus removal showed a distinct seasonal N<sub>2</sub>O-production pattern with

a recurring peak in the period between March and June (Fig. 2A). This was not concurrent with other operational parameters, as the COD:N ratio in the influent and nitrate in the effluent (g N/m<sup>3</sup>) were relatively constant over time (Fig. 2A). Although there was a nitrate peak in the effluent in the winter of 2018/2019, it did not appear to influence the N<sub>2</sub>O concentration during this period. The contrast between the relatively stable abiotic parameters and the strong N<sub>2</sub>O-production pattern



**Fig. 5.** Relative read abundance and nitrogen transformation potential of the ammonia oxidizing bacteria present in the Avedøre WWTP. A) The relative abundance (%) of all *Nitrosomonas* species over time, with the  $N_2O$ -concentration ( $g\ N/m^3$ ) depicted in red. B) Predicted nitrogen transformation genes associated with ammonium oxidation and nitrifier denitrification identified in the HQ MAGs of the Avedøre WWTP. No HQ MAG is available for *Nitrosomonas midas\_s\_723*.

could indicate that the trend was caused by the microbial community present in the WWTP. However, the top 25 growing and non-growing species (Fig. 2B) did not show many known  $N_2O$ -consumers or -producers. The growing species mainly consisted of known process-critical species in WWTPs with nutrient removal, including biological P-removal, and many undescribed species (Nierychlo et al., 2020). The non-growing species were only present because of continuous addition via the influent wastewater (Dottorini et al., 2021) and likely not very active in the transformation processes.


### 3.2. The relative abundance of a subset of species was significantly positively or negatively correlated to the $N_2O$ concentration pattern

We hypothesized that the strong  $N_2O$ -production pattern would be reflected by the relative read abundance of microbial community members of the Avedøre WWTP. We identified a total of 54 species (8.7% mean cumulative relative read abundance), 39 growing-species and 15 ambiguous-species, with a significant correlation to the seasonal  $N_2O$  concentration pattern (Supplemental data S4). 26 species had a significant positive correlation ( $\rho > 0.4$ , p-value  $< 0.01$  (Supplemental Data S4, Fig. 3)). The most abundant positively correlated species was SJA-25 midas\_s\_31 (Bacteroidota), with an average relative abundance between 0.1 and 0.4%, but two peaks around 1.2% relative abundance, (Supplemental Figure S1). The mean relative abundance of all other positively correlated species was below 1%, with most species around 0.2% (Supplemental Figure S1). Five of the 26 species (0.8% mean cumulative relative read abundance) had an identity-hit ( $>98.6\%$  nucleotide identity) with a species representative in the MiDAS genome database, and were subsequently investigated for their nitrogen transformation potential. While all five species had nitrogen transformation genes predicted (Fig. 4A, green dot), only *Terrimonas midas\_s\_743* encoded denitrification genes directly associated with  $N_2O$  production and reduction, *norBC* (K04561, K02305) and *nosZ* (K00376), respectively. Interestingly, two other positively correlated species, *Geothrix midas\_s\_443* and *Rhodiferax midas\_s\_320*, also had the predicted genes *nrfAH* and *nirBD*, respectively, which have been suggested to be involved in  $N_2O$  production (Heo et al., 2020). 28 species were

identified with a significant negative correlation ( $\rho < -0.4$ , p-value  $< 0.01$ ) (Supplemental Data S4), with *Ca. Amarolinea midas\_s\_1* (Supplemental Figure S2) showing the highest relative read abundance. *Ca. Amarolinea midas\_s\_1* had a read abundance of approximately 2.5% throughout the sampling period, except for a peak of 15% around the end of September 2018 and a peak of 5% around February 2020 (Supplemental Figure S2). Of the 23 species, nine (2.9% mean cumulative relative read abundance) had an identity-hit with a species representative of the MiDAS genome database. Analysis of these genomes showed that all species had nitrogen transformation genes predicted (Fig. 4A, red dot). Four species, *Ca. Amarolinea midas\_s\_1*, *Saprosiraceae midas\_s\_321*, *Saprosiraceae midas\_s\_862* and 37-13 midas\_s\_1223 (Bacteroidota), had a *nosZ* gene predicted without the *norBC* genes, indicating those species as potential  $N_2O$  reducers. Of those four species, only *Ca. Amarolinea midas\_s\_1* had denitrification gene predicted (*narGHI*) in combination with the *nosZ*.

### 3.3. The most abundant microorganisms had a highly diverse nitrogen-transformation potential

The nitrogen transformation potential of all abundant species with an identity hit with a HQ MAG from the MiDAS genome database (Singleton et al., 2021) was additionally investigated. Species considered significantly abundant had a relative read abundance of at least 0.1% twice a year. Of the 367 abundant species (80.3% of the total cumulative relative read abundance), 124 species had an identity-hit with a species representative of the MiDAS genome database. Together the species represented by a HQ MAG had a mean cumulative relative abundance of 31.4%. Of the 156 species representatives, 108 had one or more nitrogen transformation genes (Supplemental Data S5). An overview of the predicted nitrogen-transformation genes of the top 33 most abundant species with an associated HQ MAG, based on the mean relative abundance, showed a varied nitrogen transformation potential (Fig. 4B). Interestingly, only one potential full-denitrifier was predicted in the Avedøre WWTP community, *Ca. Dechloromonas phosphorivorans* (Fig. 4B). Although full denitrification potential was rare, partial-denitrification was not. There were 81 species with at least



**Fig. 6. Bootstrapped phylogenetic nitrous oxide reductase (NOS, EC 1.7.2.4) tree.** The tree represents 88 NOS sequences, showing the diversity of the predicted NOS protein sequences in the de-replicated HQ MAGs of the most abundant species in the Avedøre WWTP. Branches with bootstraps >70% are indicated with black circles. Experimentally confirmed and database-derived NOS sequences are indicated by blue dots (Supplemental Table S1). The different NOS clades are depicted with light blue (clade I) or pink (clade II). The mean abundance (%), based on the 16S rRNA gene amplicon sequencing, of the species that had reached a relative abundance of at least 0.1% twice in the sampling period in the Avedøre WWTP, is indicated by the bar chart. The scale bar at the lower left corresponds to the mean number of amino acid substitutions per site.

one of the denitrification genes predicted, including 11 potential  $N_2O$  producers (*norBC* predicted, but no *nosZ*) and 59 potential  $N_2O$  reducers (*nosZ* predicted, but no *norBC*). An interesting observation was that the HQ MAG of *Rhodanobacteraceae* midas\_s\_2835 had both a *nirK* and a *nirS* predicted, suggested to be mutually exclusive (Jones et al., 2013). Indeed, re-annotation of the *nirK* and *nirS* genes with BLAST could only replicate this predicted function for *nirK*. The closest hit for *nirS* was a c-type cytochrome (Supplemental Data S6). Further study is required to identify the function of this predicted gene.


To investigate if the potential NOB were correctly predicted, a NXR-A protein tree was made (Supplementary Figure S3). *Nitrotoga* species are well-known NOB (Boddicker and Mosier, 2018), but the HQ-MAG of *Nitrotoga* midas\_s\_181 had only one of the two subunits predicted, *nrxB* (K00371) (Supplemental Data S5), and was therefore not included in the protein tree. The NXR-A sequences of the two NS9 marine group species clustered with the cytoplasmic NXR-A and NAR-G sequences. OLB14 midas\_s\_849 clustered most closely to the periplasmic nitrite oxidoreductase sequences, and *Nitrospira defluvii* clustered most closely to

NXR-A sequences of other *Nitrospira* species. Lastly, *Stenotrophobacter* midas\_s\_1188 clustered near branches with a variety of functions. Although most sequences clustered near potential NXR-A proteins, experimental confirmation is required to functionally characterize these predicted proteins and determine if these species are actual NOB.

#### 3.4. Two novel *Nitrosomonas* species might be important $N_2O$ producers, with multiple $N_2O$ production pathways predicted

Although no AOB were identified within the top 33 most abundant species represented by a HQ MAG (Singleton et al., 2021) (Fig. 4B), they were present in lower abundance. Three *Nitrosomonas* species were identified, *Nitrosomonas* midas\_s\_139, *Nitrosomonas* midas\_s\_717 and *Nitrosomonas* midas\_s\_723, with a maximum relative abundance of 0.45%, 0.21% and 0.23%, respectively (Fig. 5). HQ MAGs are available for *Nitrosomonas* midas\_s\_139 and *Nitrosomonas* midas\_s\_717. Both species had the genes predicted of the canonical ammonia oxidation pathway, ammonia monooxygenase (*amoABC*; K10944, K10945,





**Fig. 7.** Non-metric multidimensional scaling analysis (NMDS) between growing and non-growing species, and the N<sub>2</sub>O-concentration measured (g N/m<sup>3</sup>). The analysis was based on the Bray-Curtis dissimilarity, with A) the individual community members labelled according to their correlation with the N<sub>2</sub>O concentration (g N/m<sup>3</sup>), and B) grouped according to the period in which they correlate, with red the high N<sub>2</sub>O-concentration period and blue the low N<sub>2</sub>O-concentration period.

K10946) and hydroxylamine dehydrogenase (*hao*; K10535). Additionally, nitrifier denitrification was predicted in both AOB, with *nirS/K* and *norBC* identified. One other species, *Ottowia midas\_s\_482* (family *Comamonadaceae*), had hydroxylamine dehydrogenase, but no ammonia monooxygenase, predicted. All three species were subsequently investigated for the potential to produce N<sub>2</sub>O from hydroxylamine with cytochrome p460 (Caranto et al., 2016; Kozłowski et al., 2014). Only the two *Nitrosomonas* species were predicted to encode the cytochrome p460 (Fig. 5). Based on the genomic prediction of *Ottowia midas\_s\_482*, it appears that although this bacteria might be capable of converting hydroxylamine and/or nitric oxide to nitrite via *hao*, and nitrite to nitric oxide via *nirS*, it does not appear to have the potential to produce N<sub>2</sub>O (Fig. 1 and 5B).

### 3.5. The majority of the *nosZ* genes were predicted in HQ MAGs from the Bacteroidota phylum

Of the 124 species with an associated HQ MAG, 59 had a *nosZ* gene predicted in their genome (Supplemental Data S5). However, comparing the predicted NOS sequences with NOS sequences from known N<sub>2</sub>O reducers (Fig. 6), we found a large evolutionary distance between the NOS sequences of known N<sub>2</sub>O reducers (blue dots) and the predicted NOS sequences from the species in the Avedøre WWTP. The known N<sub>2</sub>O reducers were predominantly species ( $n = 16$ ) from the Proteobacteria phylum, with 12 out of 16 clustering together and all encoding a clade I NOS. It appeared that the protein tree was dominated by NOS clade II sequences, with the majority of sequences from Bacteroidota species (53 out of the 76). Only one species (*Rhodobacter midas\_s\_63*) clustered with *nosZ* sequences of known N<sub>2</sub>O reducers. To support the annotation of the HQ MAGs, a NOS sequence from *Ignavibacterium album* strain DSM 19864 (Sanford et al., 2012), and from a *Flavobacterium* species (Kim et al., 2020) from the Bacteroidota phylum was included in the tree.


These sequences clustered with the other sequences from Bacteroidota, showing that the *nosZ* sequences of the Bacteroidota were closely related to the characterised NOS of *I. album* DSM 19864.

Of the bacteria predicted to have a *nosZ* gene, only five had a mean abundance higher than 0.5% during the period sampled (Fig. 6). The microorganisms with the highest abundance were a *Ca. Amarolinea midas\_s\_1* (Chloroflexota), *Rhodobacter midas\_s\_63* (Proteobacteria), *Ca. Dechloromonas phosphorivorans* (Proteobacteria), *Ca. Promineofilaceae midas\_s\_461* (Chloroflexota) and *Terrimonas midas\_s\_390* (Bacteroidota), with 1.4%, 0.9%, 0.9%, 0.6% and 0.5% mean abundance, respectively.

### 3.6. The N<sub>2</sub>O-production pattern could not be explained by the relative abundance of species, individually or grouped into functional guilds

No individual member seemed to be solely responsible for the seasonal N<sub>2</sub>O concentration pattern (Fig. 3). Non-growing species are not expected to take an active role in the process. As expected, there was a large overlap of the community between high and low N<sub>2</sub>O-production periods (Fig. 7B) and an absence of a correlation between the non-growing microbial community and the N<sub>2</sub>O concentration (g N/m<sup>3</sup>) (Fig. 7A). Although there was an overlap between high and low N<sub>2</sub>O-production periods for the growing communities (Fig. 7B), a slight gradient was observed between the beta-diversity of growing microbial communities and N<sub>2</sub>O concentration (Fig. 7A). Additionally, as shown in Section 3.2, a number of individual species were positively correlated with N<sub>2</sub>O concentration-level.

However, it was possible that the cumulative relative abundance of a functional guild of microorganisms could show a better correlation to the N<sub>2</sub>O-production pattern. The functional guilds were based on the presence of key nitrogen-transforming or N<sub>2</sub>O-associated genes predicted in the HQ MAGs. The three AOB and the full denitrifier (*Ca.*



**Fig. 8.** Time-series of the cumulative relative read abundance (%) of the 16S rRNA gene amplicons of the HQ MAG-determined functional guilds in Avedøre WWTP. The line shows the N<sub>2</sub>O-concentration (g N/m<sup>3</sup>). AOB (*amoABC* and *hao* predicted), NOB (*nrxAB* predicted), N<sub>2</sub>O-producers (DEN) (*norBC* predicted, but no *nosZ*), N<sub>2</sub>O-reducers (DEN) (*nosZ*, but no *norBC*) predicted and full-denitrifiers (complete DEN) (*narGHI* or *napAB*, *nirK* or *nirS*, *norBC* and *nosZ* predicted).

*Dechloromonas phosphorivorans*) were grouped according to their functionality, AOB and complete DEN, respectively. The other species were grouped according to the presence of the *nosZ* gene (N<sub>2</sub>O-reducers (DEN)) or the presence of the *norBC* gene without a *nosZ* gene (N<sub>2</sub>O-producers (DEN)). The NOB without a *nosZ* gene, but with a *nrxAB*, were

grouped together as well (NOB). There was a small but significant negative correlation ( $\rho = -0.37$ ,  $p$ -value =  $2.3 \times 10^{-3}$ , Fig. 8, Supplemental Data S7) with species grouped as (potential) NOB ( $n = 3$ , *nrxAB*, but no *nosZ*, Supplemental Data S7). Additionally, species grouped as DEN—N<sub>2</sub>O-reducers ( $n = 59$ ) ( $\rho = -0.48$ ,  $p$ -value =  $5.6 \times 10^{-5}$ , Fig. 8, Supplemental Data S7; encoding *nosZ*) had a significant negative correlation with the N<sub>2</sub>O concentration. The relative read abundance of the DEN—N<sub>2</sub>O producers (encoding *norBC*, but not *nosZ*,  $n = 8$ ) was dominated by one species from the *Saprospiraceae* family, OLB8 midas\_s\_29 (Fig. 8, Supplemental Data S7). While at first glance there was a similar pattern in relative read abundance of this specific group and the seasonal N<sub>2</sub>O concentration, the large time difference makes a direct metabolic correlation unlikely. This was also reflected in the statistical analysis as no correlation ( $\rho = 3.0 \times 10^{-3}$ ,  $p$ -value = 0.98, Fig. 8, Supplemental Data S7) in relative read abundance of the N<sub>2</sub>O-producers was identified. The full-denitrifiers ( $n = 1$ ), consisting only of the *Ca. D. phosphorivorans* (Fig. 8, Supplemental Data S7), showed an increased relative abundance during high N<sub>2</sub>O periods. However, the specific pattern in increase was not reflected in the N<sub>2</sub>O concentration pattern as it was not significantly correlated ( $\rho = 3.2 \times 10^{-3}$ ,  $p$ -value = 0.98, Supplemental Data S7). Lastly, the relative read abundance of the other functional group able to produce N<sub>2</sub>O, the AOB (*amoABC* and *hao* predicted,  $n = 3$ ), had no significant correlation to the N<sub>2</sub>O concentration pattern ( $\rho = -9.6 \times 10^{-2}$ ,  $p$ -value = 0.45, Supplemental Data S7) (Fig. 8).

#### 4. Discussion

##### 4.1. The N<sub>2</sub>O concentration pattern appeared to be microbially mediated by several candidate species

The abiotic and biotic influences on the N<sub>2</sub>O-emissions in full-scale WWTPs have been studied at multiple locations across Europe (Chen et al., 2019; Daelman et al., 2015; Gruber et al., 2020, 2021). These four studies also report the highest N<sub>2</sub>O-emission between March and May and additionally indicated a correlation between N<sub>2</sub>O, seasonality and accumulation of nitrite and/or nitrate. These studies also acknowledge the relationship between the accumulation of intermediates and the microbial conversions. Such annual patterns in microbial conversions may also be determined by variations in the abundance of specific bacterial taxa, a change in the activity of certain taxa due to environmental variations, or a combination of the two. We identified 54 species showing a strong recurrent seasonality and a significant correlation to the N<sub>2</sub>O concentration, indicating that the abundance of certain taxa could be important. The majority (39) of these taxa also belonged to the growing fraction, which is assumed to be metabolically active and not only present due to continuous transportation by the influent wastewater (Dottorini et al., 2021). Of these growing species, 15 had an identity-hit with a HQ MAG, so it was possible to investigate their genomic potential in greater detail.

*Terrimonas midas\_s\_743* was the only positively correlated species with a HQ MAG that had denitrification genes predicted directly associated with both N<sub>2</sub>O production (*norBC*) and reduction (*nosZ*). Although species of this genus have been identified in (partial) denitrification communities, it has not been directly associated with N<sub>2</sub>O emissions (Zhang et al., 2021). Four of the negatively associated species with a HQ MAG had the potential to reduce N<sub>2</sub>O. The filamentous bacteria *Ca. Amarolinea midas\_s\_1*, the most abundant of the species, has previously been suggested to have the potential for N<sub>2</sub>O reduction (Andersen et al., 2019). Additionally, other species from the *Ca. Amarolinea* genus have been reported to actively transcribe *nosZ* in an anammox bioreactor community (Suenaga et al., 2021). The other three species were *Bacteroidota* (*Saprospiraceae* midas\_s\_321, *Saprospiraceae* midas\_s\_862 and 37–13 midas\_s\_1223). While *Saprospiraceae* has been associated with N<sub>2</sub>O emissions in soils and anammox bioreactors (Jones et al., 2013, 2014; Speth et al., 2016), this is the first instance that

members of the *Saprosiraceae* family have been predicted as N<sub>2</sub>O-reducers in a full-scale WWTP community. Another closely related genus within the Bacteroidota, *Flavobacterium*, has also been associated with N<sub>2</sub>O production in WWTP, suggesting that Bacteroidota species might have a more important role than previously thought (Kim et al., 2020). All four species are interesting candidates to investigate in more detail for their N<sub>2</sub>O-reduction potential.

The majority (39 of the 54) of the significantly correlated species did not have an identity-hit with a HQ MAG in the MiDAS genome database (Singleton et al., 2021). It is difficult to investigate the possible genetic traits these species had to explain a significant correlation and possible effect on the seasonal N<sub>2</sub>O concentration. It also shows that many of the species in the WWTP remain functionally elusive and that we need to increase the number of HQ MAGs to improve the resolution and increase our insight into species functionality.

However, some species correlating with the seasonal N<sub>2</sub>O concentration, but without a HQ MAG available, had a genus-level association with known (partial) denitrifiers. Although functional traits are often species-specific (Petriglieri et al., 2021), these species could be of special interest to study in more detail. Among the species with a significant correlation to the seasonal N<sub>2</sub>O concentration was a *Rhodobacter*, a genus harbouring known denitrifiers and with species previously correlated with N<sub>2</sub>O emissions in WWTPs (Vieira et al., 2019). Other microbial lineages that significantly correlated with N<sub>2</sub>O emission (*Ca. Competibacter*, *Ca. Accumulibacter* and *Dechloromonas* species) have been shown to produce and reduce N<sub>2</sub>O in lab-scale enrichment experiments (Roy et al., 2021; Vieira et al., 2018; Yoon et al., 2016). While multiple *Ca. Accumulibacter* and *Dechloromonas* species had a HQ MAG available, only *Ca. D. phosphorivorans* had genes predicted directly associated with N<sub>2</sub>O production and reduction. *Dechloromonas* species have not been attributed with high N<sub>2</sub>O emissions, under optimal growth conditions, due to their high affinity to N<sub>2</sub>O (Suenaga et al., 2019; Yoon et al., 2016). However, non-ideal conditions, e.g. salt or alkaline stress, may stimulate N<sub>2</sub>O-production by this species (Han et al., 2019).

The species grouped as NOB consisting of *Nitrospira defluvii*, *Nitrotoga midas\_s\_181* and two predicted species with at least one copy of the *nrxAB* gene, had a statistically significant negative correlation with the seasonal N<sub>2</sub>O concentration. This correlation was also observed by Gruber et al. (2021). Although NOB have no direct genes able to produce or reduce N<sub>2</sub>O, they can be involved in complex symbiosis with AOB or heterotrophic bacteria (Daims et al., 2016) potentially affecting seasonal N<sub>2</sub>O concentration. More research is required to elucidate the relationship between abundance of NOB and the seasonal N<sub>2</sub>O concentration.

AOB have been suggested as mainly responsible for N<sub>2</sub>O production in WWTPs in several studies (Dias et al., 2022; Law et al., 2012; Wunderlin et al., 2012). Although the AOB in the Avedøre WWTP had the potential to produce N<sub>2</sub>O via both the hydroxylamine pathway and nitrifier denitrification (Caranto et al., 2016; Wrage-Mönnig et al., 2018), no significant correlation was identified between their abundance and the N<sub>2</sub>O concentration peak. Such stable abundance, in contrast to strongly changing seasonal N<sub>2</sub>O concentration, was also observed by Gruber et al. (2021). More detailed research is required to determine if this discrepancy demonstrates that AOB do not significantly contribute to the seasonal N<sub>2</sub>O-production peak.

#### 4.2. Denitrification seemed dominated by partial denitrifiers

Full-denitrifiers are often assumed to dominate the denitrification process in WWTP (Nielsen and McMahon, 2014). However, in our study of Avedøre WWTP we only identified one full-denitrifier (NO<sub>3</sub><sup>-</sup> to N<sub>2</sub>, (Lu et al., 2014; Zumft, 1997)), *Ca. D. phosphorivorans*. This is also reflected in the MiDAS genome database of species from WWTPs all over Denmark. While Singleton et al. (2021) identified 21 full-denitrifiers, they identified manifold more HQ MAGs with at least one gene of the pathway predicted. The variety in denitrification gene combinations is

well established and seen in many natural and engineered environments (Baker et al., 2015; Graf et al., 2014; Lu et al., 2014). Division of labor between different species has additionally been shown in denitrifying enrichment cultures (Vieira et al., 2018; Wang et al., 2021), suggesting that the denitrifying potential, and as such the N<sub>2</sub>O production or reduction potential, is likely impacted by the community structure.

#### 4.3. The high N<sub>2</sub>O reduction potential, mainly identified in Bacteroidota species, was dominated by clade II *nosZ*

The majority of the *nosZ* genes were predicted in members of the Bacteroidota. The importance of this phylum in N<sub>2</sub>O reduction has been observed in soils (Jones et al., 2014). While functionally verified N<sub>2</sub>O-reducers are predominantly Proteobacteria (Conthe et al., 2018; Hallin et al., 2018; Suenaga et al., 2018), the many undiscovered N<sub>2</sub>O-reducers from other phyla might play an important role in natural and engineered environments (Jones et al., 2013; Sanford et al., 2012; Singleton et al., 2021).

The majority of clade I *nosZ* gene sequences have been identified in Proteobacteria, and are normally associated with denitrifying organisms (Hallin et al., 2018; Sanford et al., 2012; Yoon et al., 2016). Interestingly, this was not observed in our analysis. Although there were three HQ MAGs with a denitrifier clade I *nosZ* gene predicted, none of these HQ MAGs had the full denitrification pathway predicted. Additionally, none of these species had a significant correlation with the seasonal N<sub>2</sub>O concentration. The majority (69 of the 72) of the *nosZ* sequences predicted within the Avedøre WWTP community were clade II. Although this has been observed in other environments (Jones et al., 2014; Kim et al., 2020; Orellana et al., 2014; Suenaga et al., 2021), this is the first study showing this overwhelming dominance of clade II over clade I (ratio = 25.7). It could be that the proposed higher oxygen tolerance and affinity for N<sub>2</sub>O of the clade II NOS was selective for this community (Yoon et al., 2016).

#### 4.4. Other factors than the genomic N<sub>2</sub>O production or reduction potential predict the N<sub>2</sub>O concentration in the Avedøre WWTP

The community in Avedøre WWTP had a large genetic potential to reduce N<sub>2</sub>O during the period studied. Other studies have shown that genomic N<sub>2</sub>O reduction potential and N<sub>2</sub>O emission level are correlated (Jones et al., 2014; Vieira et al., 2018), with high *nosZ* values relating to low N<sub>2</sub>O emissions. In our dataset there were a high number of HQ MAGs with a *nosZ* predicted (approximately 45% of the HQ MAGs available), suggesting a high genomic potential to reduce N<sub>2</sub>O. This would imply that other factors than the genetic potential determine the N<sub>2</sub>O concentration in the WWTP. However, as we do not cover the entire community with the MiDAS genome database, long term monitoring of the *nosZ* gene in the WWTP will be required to examine the *nosZ* potential of the community without a HQ MAG.

Many factors can influence the actual N<sub>2</sub>O reduction activity in full-scale WWTPs, with dissolved oxygen concentrations often suggested as one of the key-inhibitors for N<sub>2</sub>O reduction (Law et al., 2012; Pan et al., 2013; Suenaga et al., 2018). The seasonal change in temperature can affect the dissolved oxygen concentration in the WWTP, as well as the microbial conversion rates, but its effect should be expected to be observed in both spring and fall. Additionally, previous studies have identified only weak to no correlation between temperature as a parameter and N<sub>2</sub>O emissions (Ahn et al., 2010; Chen et al., 2019), suggesting that temperature was not the dominant influencing factor. Another key-parameter in N<sub>2</sub>O reduction is the electron availability, as all enzymatic conversions in the denitrification pathway require electrons, electron competition within the pathway can decrease in N<sub>2</sub>O reduction activity (Conthe et al., 2018; Felgate et al., 2012). However, the COD:N ratio in the influent and the COD quality in the effluent remained stable throughout the studied period (Supplementary Data S1). This does not exclude the possibility that the type of electron donor

**Table 1**

**KEGG orthology identifier (KO-id) used in this study.** Functional guilds, the three-letter abbreviation of their associated genes, the KEGG orthology identifier (KO-id), and the nitrogen-transformation performed by the protein encoded by the gene. DEN; denitrifier, AOB; ammonia oxidizing bacteria, NOB; nitrite oxidizing bacteria, CMX; comammox, DNRA; dissimilatory nitrite reducing bacteria, AMX; anammox bacteria.

Functional guild	Gene	KO-id	Transformation
DEN	<i>narGHI<sup>d</sup></i>	K00370, K00371, K00374	$\text{NO}_3^- > \text{NO}_2^-$
	<i>napAB</i>	K02567, K02568	$\text{NO}_3^- > \text{NO}_2^-$
	<i>nirK</i>	K00368	$\text{NO}_2^- > \text{NO}$
	<i>nirS</i>	K15864	$\text{NO}_2^- > \text{NO}$
	<i>norBC</i>	K04561, K02305	$\text{NO} > \text{N}_2\text{O}$
AOB	<i>nosZ</i>	K00376	$\text{N}_2\text{O} > \text{N}_2$
	<i>amoABC</i>	K10944, K10945, K10946	$\text{NH}_3^+ > \text{NH}_2\text{OH}$
	<i>hao</i>	K10535	$\text{NH}_2\text{OH} > \text{NO}$
	*		$\text{NO} > \text{NO}_2^-$
	<i>Cyt-p460</i>		$\text{NH}_2\text{OH} > \text{N}_2\text{O}$
NOB	<i>nrxAB<sup>a</sup></i>	K00370, K00371	$\text{NO}_2^- > \text{NO}_3^-$
CMX	<i>amoABC</i>	K10944, K10945, K10946	$\text{NH}_3^+ > \text{NH}_2\text{OH}$
	<i>hao</i>	K10535	$\text{NH}_2\text{OH} > \text{NO}$
	*		$\text{NO} > \text{NO}_2^-$
DNRA	<i>nrxAB<sup>a</sup></i>	K00370, K00371	$\text{NO}_2^- > \text{NO}_3^-$
	<i>narGHI<sup>d</sup></i>	K00370, K00371, K00374	$\text{NO}_3^- > \text{NO}_2^-$
	<i>napAB</i>	K02567, K02568	$\text{NO}_3^- > \text{NO}_2^-$
	<i>nrfAH</i>	K03385, K15876	$\text{NO}_2^- > \text{NH}_4^+$
	<i>nirBD</i>	K00362, K00363	$\text{NO}_2^- > \text{NH}_4^+$
AMX	<i>nirS</i>	K15864	$\text{NO}_2^- > \text{NO}$
	<i>hszABC</i>	K20932, K20933, K20934	$\text{NO} + \text{NH}_4^+ > \text{N}_2\text{H}_4$
	<i>hdh</i>	K20935	$\text{N}_2\text{H}_4 > \text{N}_2$

\*No gene has been conclusively identified for this conversion (Kuypers et al., 2018).

<sup>a</sup> The NXR and NAR are homologs and cannot be distinguished by KEGG IDs. A protein tree is used to resolve potential function as NXR or NAR.

or carbon source, shown to severely impact the  $\text{N}_2\text{O}$  reductase activity (Itokawa et al., 2001; Pan et al., 2013; Qi et al., 2022; Ribera-Guardia et al., 2014, 2016), changed over the year. Table 1

$\text{N}_2\text{O}$  production can occur in both the aerobic and in the anoxic phase, by AOB and (truncated) denitrifiers (Song et al., 2020). As  $\text{N}_2\text{O}$  reduction only occurs during the anoxic phase (Kampschreur et al., 2009; Suenaga et al., 2018), part of the produced  $\text{N}_2\text{O}$  will directly be sparged out of the liquid during the aerobic phase. Although aerobic denitrification may occur (Robertson et al., 1989, 1995) and produce  $\text{N}_2\text{O}$ , the majority of the aerobically produced  $\text{N}_2\text{O}$  is produced by AOB (Tallec et al., 2008). Low dissolved oxygen concentrations and high nitrite concentrations have been shown to stimulate  $\text{N}_2\text{O}$  production by AOB (Kampschreur et al., 2009; Kozłowski et al., 2016). As stated above, if the seasonal temperature effect on dissolved oxygen affected  $\text{N}_2\text{O}$  production by AOB, this would be expected to occur in both spring and fall. While no nitrite concentrations were measured in the period studied, it might be an interesting additional parameter to measure in future studies.

Future work is required to elucidate the factors influencing seasonal (under-)utilization of the high genetic  $\text{N}_2\text{O}$  reduction potential of the community. Although this work provides a list of interesting candidates that could act as a  $\text{N}_2\text{O}$  sink within Avedøre WWTP, many species are novel with little to no knowledge of their (potential) function. Metagenomics data, combined with  $\text{N}_2\text{O}$  reductase activity assays measured with activated sludge throughout the year(s), could provide more insight into the physiology of  $\text{N}_2\text{O}$  sink species. This could result in mitigation strategies controlling species composition and  $\text{N}_2\text{O}$  production and reduction activity.

## 5. Conclusions

- We found a recurrent strong seasonal  $\text{N}_2\text{O}$  concentration pattern, with spring peaks and autumn lows, from Avedøre WWTP over 3 years.

- The relative read abundance of 54 species was significantly correlated to the  $\text{N}_2\text{O}$  concentrations, either positively or negatively.
- Partial denitrifiers, instead of complete denitrifiers, appeared to dominate the genetic denitrification potential of the community.
- The cumulative abundances of  $\text{N}_2\text{O}$  reducers and NOB had a significant negative correlation to  $\text{N}_2\text{O}$  concentrations.
- Clade II *nosZ* genes, primarily found in Bacteroidota, were more common than clade I *nosZ* genes in the HQ MAGs.
- Multiple novel potential  $\text{N}_2\text{O}$  sink species have been identified for further study.
- Overall, there was some correlation between community structure and  $\text{N}_2\text{O}$  concentration, but activity measurements of specific bacteria are needed to understand the correlation in greater detail.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The authors wish to thank the Avedøre WWTP for sharing the operational data, and all the student helpers and operators for taking the activated sludge samples. The authors additionally wish to thank Nick Green for help with the HQ MAG annotation. The project has been funded by the Villum Foundation (Dark Matter, grant 13351).

## Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.watres.2022.118563.

## References

- Ahn, J.H., Kim, S., Park, H., Rahm, B., Pagilla, K., Chandran, K., 2010.  $\text{N}_2\text{O}$  Emissions from Activated Sludge Processes, 2008–2009: results of a National Monitoring Survey in the United States. *Environ. Sci. Technol.* 44 (12), 4505–4511. <https://doi.org/10.1021/es903845y>.
- Andersen, K.S., Kirkegaard, R.H., Karst, S.M., Albertsen, M., 2018. *ampvis2: an R package to analyse and visualise 16S rRNA amplicon data* [Preprint]. *Bioinformatics*. <https://doi.org/10.1101/299537>.
- Andersen, M.H., McIlroy, S.J., Nierychlo, M., Nielsen, P.H., Albertsen, M., 2019. Genomic insights into *Candidatus Amarolinea aalborgensis* gen. nov., sp. nov., associated with settleability problems in wastewater treatment plants. *Syst. Appl. Microbiol.* 42 (1), 77–84. <https://doi.org/10.1016/j.syapm.2018.08.001>.
- Baker, B.J., Lazar, C.S., Teske, A.P., Dick, G.J., 2015. Genomic resolution of linkages in carbon, nitrogen, and sulfur cycling among widespread estuary sediment bacteria. *Microbiome* 3 (1), 14. <https://doi.org/10.1186/s40168-015-0077-6>.
- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)* 57 (1), 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>.
- Boddicker, A.M., Mosier, A.C., 2018. Genomic profiling of four cultivated *Candidatus Nitrotoga* spp. predicts broad metabolic potential and environmental distribution. *ISME J.* 12 (12), 2864–2882. <https://doi.org/10.1038/s41396-018-0240-8>.
- Caranto, J.D., Lancaster, K.M., 2017. Nitric oxide is an obligate bacterial nitrification intermediate produced by hydroxylamine oxidoreductase. *Proc. Natl. Acad. Sci.* 114 (31), 8217–8222. <https://doi.org/10.1073/pnas.1704504114>.
- Caranto, J.D., Vilbert, A.C., Lancaster, K.M., 2016. *Nitrosomonas europaea* cytochrome P460 is a direct link between nitrification and nitrous oxide emission. *Proc. Natl. Acad. Sci.* 113 (51), 14704–14709. <https://doi.org/10.1073/pnas.1611051113>.
- Chen, X., Mielczarek, A.T., Habicht, K., Andersen, M.H., Thornberg, D., Sin, G., 2019. Assessment of full-scale  $\text{N}_2\text{O}$  emission characteristics and testing of control concepts in an activated sludge wastewater treatment plant with alternating aerobic and anoxic phases. *Environ. Sci. Technol.* 53 (21), 12485–12494. <https://doi.org/10.1021/acs.est.9b04889>.
- Conthe, M., Lycus, P., Arntzen, M.Ø., Ramos da Silva, A., Frostegård, Å., Bakken, L.R., Kleerebezem, R., van Loosdrecht, M.C.M., 2019. Denitrification as an  $\text{N}_2\text{O}$  sink. *Water Res.* 151, 381–387. <https://doi.org/10.1016/j.watres.2018.11.087>.
- Conthe, M., Wittorf, L., Kuenen, J.G., Kleerebezem, R., van Loosdrecht, M.C.M., Hallin, S., 2018. Life on  $\text{N}_2\text{O}$ : deciphering the ecophysiology of  $\text{N}_2\text{O}$  respiring bacterial communities in a continuous culture. *ISME J.* 12 (4), 1142–1153. <https://doi.org/10.1038/s41396-018-0063-7>.

- Daelman, M.R.J., van Voorhuizen, E.M., van Dongen, U.G.J.M., Volcke, E.I.P., van Loosdrecht, M.C.M., 2013. Methane and nitrous oxide emissions from municipal wastewater treatment—Results from a long-term study. *Water Sci. Technol.* 67 (10), 2350–2355. <https://doi.org/10.2166/wst.2013.109>.
- Daelman, M.R.J., van Voorhuizen, E.M., van Dongen, U.G.J.M., Volcke, E.I.P., van Loosdrecht, M.C.M., 2015. Seasonal and diurnal variability of N<sub>2</sub>O emissions from a full-scale municipal wastewater treatment plant. *Sci. Total Environ.* 536, 1–11. <https://doi.org/10.1016/j.scitotenv.2015.06.122>.
- Daims, H., Lückler, S., Wagner, M., 2016. A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol.* 24 (9), 699–712. <https://doi.org/10.1016/j.tim.2016.05.004>.
- Delre, A., ten Hoeve, M., Scheutz, C., 2019. Site-specific carbon footprints of Scandinavian wastewater treatment plants, using the life cycle assessment approach. *J. Clean Prod.* 211, 1001–1014. <https://doi.org/10.1016/j.jclepro.2018.11.200>.
- Dias, D.F.C., Marques, R., Martins, C., Martins, A., Oehmen, A., 2022. The impact of a seasonal change in loading rate on the nitrous oxide emissions at the WWTP of a tourist region. *Sci. Total Environ.* 804, 149987 <https://doi.org/10.1016/j.scitotenv.2021.149987>.
- Domingo-Félez, C., Smets, B.F., 2019. Regulation of key N<sub>2</sub>O production mechanisms during biological water treatment. *Curr. Opin. Biotechnol.* 57, 119–126. <https://doi.org/10.1016/j.copbio.2019.03.006>.
- Dottorini, G., Michaelsen, T.Y., Kucheryavskiy, S., Andersen, K.S., Kristensen, J.M., Peces, M., Wagner, D.S., Nierychlo, M., Nielsen, P.H., 2021. Mass-immigration determines the assembly of activated sludge microbial communities. *Proc. Natl. Acad. Sci.* 118 (27), e2021589118 <https://doi.org/10.1073/pnas.2021589118>.
- Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic. Acids. Res.* 32 (5), 1792–1797. <https://doi.org/10.1093/nar/gkh340>.
- Edgar, R.C., 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26 (19), 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>.
- Edgar, R.C., 2017. Accuracy of microbial community diversity estimated by closed- and open-reference OTUs. *PeerJ* 5, e3889. <https://doi.org/10.7717/peerj.3889>.
- Edgar, R.C., 2018. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics* 34 (14), 2371–2375. <https://doi.org/10.1093/bioinformatics/bty113>.
- El-Gebali, S., Mistry, J., Bateman, A., Eddy, S.R., Luciani, A., Potter, S.C., Qureshi, M., Richardson, L.J., Salazar, G.A., Smart, A., Sonnhammer, E.L.L., Hirsh, L., Paladin, L., Piovesan, D., Tosatto, S.C.E., Finn, R.D., 2019. The Pfam protein families database in 2019. *Nucleic. Acids. Res.* 47 (D1), D427–D432. <https://doi.org/10.1093/nar/gky995>.
- Felgate, H., Giannopoulos, G., Sullivan, M.J., Gates, A.J., Clarke, T.A., Baggs, E., Rowley, G., Richardson, D.J., 2012. The impact of copper, nitrate and carbon status on the emission of nitrous oxide by two species of bacteria with biochemically distinct denitrification pathways: denitrification and nitrous oxide release. *Environ. Microbiol.* 14 (7), 1788–1800. <https://doi.org/10.1111/j.1462-2920.2012.02789.x>.
- Graf, D.R.H., Jones, C.M., Hallin, S., 2014. Intergenomic comparisons highlight modularity of the denitrification pathway and underpin the importance of community structure for N<sub>2</sub>O emissions. *PLoS One* 9 (12), e114118. <https://doi.org/10.1371/journal.pone.0114118>.
- Gruber, W., Niederdorfer, R., Ringwald, J., Morgenroth, E., Bürgmann, H., Joss, A., 2021. Linking seasonal N<sub>2</sub>O emissions and nitrification failures to microbial dynamics in a SBR wastewater treatment plant. *Water Research X* 11, 100098. <https://doi.org/10.1016/j.wroa.2021.100098>.
- Gruber, W., Villez, K., Kipf, M., Wunderlin, P., Siegrist, H., Vogt, L., Joss, A., 2020. N<sub>2</sub>O emission in full-scale wastewater treatment: proposing a refined monitoring strategy. *Sci. Total Environ.* 699, 134157 <https://doi.org/10.1016/j.scitotenv.2019.134157>.
- Hallin, S., Philippot, L., Löffler, F.E., Sanford, R.A., Jones, C.M., 2018. Genomics and ecology of novel N<sub>2</sub>O-reducing microorganisms. *Trends Microbiol.* 26 (1), 43–55. <https://doi.org/10.1016/j.tim.2017.07.003>.
- Han, H., Song, B., Song, M.J., Yoon, S., 2019. Enhanced nitrous oxide production in denitrifying *Dechloromonas aromatica* strain RCB under salt or alkaline stress conditions. *Front Microbiol.* 10, 1203. <https://doi.org/10.3389/fmicb.2019.01203>.
- Heo, H., Kwon, M., Song, B., Yoon, S., 2020. Involvement of NO<sub>3</sub><sup>-</sup> in ecophysiological regulation of dissimilatory nitrate/nitrite reduction to ammonium (DNRA) is implied by physiological characterization of soil DNRA bacteria isolated via a colorimetric screening method. *Appl. Environ. Microbiol.* 86 (17) <https://doi.org/10.1128/AEM.01054-20> e01054-20.
- Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., Vinh, L.S., 2018. UFBoot2: improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* 35 (2), 518–522. <https://doi.org/10.1093/molbev/msx281>.
- Itokawa, H., Hanaki, K., Matsuo, T., 2001. Nitrous oxide production in high-loading biological nitrogen removal process under low cod/n ratio condition. *Water Res.* 35 (3), 657–664. [https://doi.org/10.1016/S0043-1354\(00\)00309-2](https://doi.org/10.1016/S0043-1354(00)00309-2).
- Jones, C.M., Graf, D.R., Bru, D., Philippot, L., Hallin, S., 2013. The unaccounted yet abundant nitrous oxide-reducing microbial community: a potential nitrous oxide sink. *ISME J.* 7 (2), 417–426. <https://doi.org/10.1038/ismej.2012.125>.
- Jones, C.M., Spor, A., Brennan, F.P., Breuil, M.-C., Bru, D., Lemanceau, P., Griffiths, B., Hallin, S., Philippot, L., 2014. Recently identified microbial guild mediates soil N<sub>2</sub>O sink capacity. *Nat. Clim. Chang.* 4 (9), 801–805. <https://doi.org/10.1038/nclimate2301>.
- Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermini, L.S., 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* 14 (6), 587–589. <https://doi.org/10.1038/nmeth.4285>.
- Kampschreur, M.J., Kleerebezem, R., de Vet, W.W.J.M., van Loosdrecht, M.C.M., 2011. Reduced iron induced nitric oxide and nitrous oxide emission. *Water Res.* 45 (18), 5945–5952. <https://doi.org/10.1016/j.watres.2011.08.056>.
- Kampschreur, M.J., Tan, N.C.G., Kleerebezem, R., Picioreanu, C., Jetten, M.S.M., Loosdrecht, M.C.M., 2008. Effect of dynamic process conditions on nitrogen oxides emission from a nitrifying culture. *Environ. Sci. Technol.* 42 (2), 429–435. <https://doi.org/10.1021/es071667p>.
- Kampschreur, M.J., Temmink, H., Kleerebezem, R., Jetten, M.S.M., van Loosdrecht, M.C.M., 2009. Nitrous oxide emission during wastewater treatment. *Water Res.* 43 (17), 4093–4103. <https://doi.org/10.1016/j.watres.2009.03.001>.
- Kanehisa, M., 2019. Toward understanding the origin and evolution of cellular organisms. *Protein Sci.* 28 (11), 1947–1951. <https://doi.org/10.1002/pro.3715>.
- Kim, D.D., Park, D., Yoon, H., Yun, T., Song, M.J., Yoon, S., 2020. Quantification of *nosZ* genes and transcripts in activated sludge microbiomes with novel group-specific qPCR methods validated with metagenomic analyses. *Water Res.* 185, 116261 <https://doi.org/10.1016/j.watres.2020.116261>.
- Kozłowski, J.A., Kits, K.D., Stein, L.Y., 2016. Comparison of nitrogen oxide metabolism among diverse Ammonia-Oxidizing Bacteria. *Front Microbiol.* 7. <https://doi.org/10.3389/fmicb.2016.01090>.
- Kozłowski, J.A., Price, J., Stein, L.Y., 2014. Revision of N<sub>2</sub>O-producing pathways in the ammonia-oxidizing bacterium *Nitrosomonas europaea* ATCC 19718. *Appl. Environ. Microbiol.* 80 (16), 4930–4935. <https://doi.org/10.1128/AEM.01061-14>.
- Kuyper, M.M.M., Marchant, H.K., Kartal, B., 2018. The microbial nitrogen-cycling network. *Nat. Rev. Microbiol.* 16 (5), 263–276. <https://doi.org/10.1038/nrmicro.2018.9>.
- Lane, D., 1991. 16S/23S rRNA sequencing. *Nucleic Acid Techn. Bacterial Systematics* 115–175.
- Law, Y., Ye, L., Pan, Y., Yuan, Z., 2012. Nitrous oxide emissions from wastewater treatment processes. *Philos. Trans. R. Soc. B: Biol. Sci.* 367 (1593), 1265–1277. <https://doi.org/10.1098/rstb.2011.0317>.
- Letunic, I., Bork, P., 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic. Acids. Res.* 49 (W1), W293–W296. <https://doi.org/10.1093/nar/gkab301>.
- Liu, B., Mao, Y., Bergaust, L., Bakken, L.R., Frostegård, Å., 2013. Strains in the genus *Thauera* exhibit remarkably different denitrification regulatory phenotypes: denitrification regulatory phenotypes in *T. hausera*. *Environ. Microbiol.* <https://doi.org/10.1111/1462-2920.12142> n/a-n/a.
- Lu, H., Chandran, K., 2010. Factors promoting emissions of nitrous oxide and nitric oxide from denitrifying sequencing batch reactors operated with methanol and ethanol as electron donors. *Biotechnol. Bioeng.* <https://doi.org/10.1002/bit.22704> n/a-n/a.
- Lu, H., Chandran, K., Stensel, D., 2014. Microbial ecology of denitrification in biological wastewater treatment. *Water Res.* 64, 237–254. <https://doi.org/10.1016/j.watres.2014.06.042>.
- Lycus, P., Lovise Bothun, K., Bergaust, L., Peele Shapleigh, J., Reier Bakken, L., Frostegård, Å., 2017. Phenotypic and genotypic richness of denitrifiers revealed by a novel isolation strategy. *ISME J.* 11 (10), 2219–2232. <https://doi.org/10.1038/ismej.2017.82>.
- McIlroy, S.J., Awata, T., Nierychlo, M., Albertsen, M., Kindaichi, T., Nielsen, P.H., 2015. Characterization of the *in situ* ecophysiology of novel phylogenies in nutrient removal activated sludge treatment plants. *PLoS One* 10 (9), e0136424. <https://doi.org/10.1371/journal.pone.0136424>.
- Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., Lanfear, R., 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* 37 (5), 1530–1534. <https://doi.org/10.1093/molbev/msaa015>.
- Mise, K., Masuda, Y., Senoo, K., Itoh, H., 2021. Undervalued pseudo-*nifH* sequences in public databases distort metagenomic insights into biological nitrogen fixers. *mSphere* 6 (6). <https://doi.org/10.1128/msphere.00785-21> e00785-21.
- Muyzer, G., de Waal, E.C., Uitterlinden, A.G., 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* 59 (3), 695–700. <https://doi.org/10.1128/aem.59.3.695-700.1993>.
- Nielsen, P.H., McMahon, K.D., 2014. Microbiology and microbial ecology of the activated sludge process. *Activated Sludge—100 Years and Counting*. IWA Publishing, pp. 53–76.
- Nierychlo, M., Andersen, K.S., Xu, Y., Green, N., Jiang, C., Albertsen, M., Dueholm, M.S., Nielsen, P.H., 2020. MiDAS 3: an ecosystem-specific reference database, taxonomy and knowledge platform for activated sludge and anaerobic digesters reveals species-level microbiome composition of activated sludge. *Water Res.* 182, 115955 <https://doi.org/10.1016/j.watres.2020.115955>.
- Orellana, L.H., Rodriguez-R, L.M., Higgins, S., Chee-Sanford, J.C., Sanford, R.A., Ritalahti, K.M., Löffler, F.E., Konstantinidis, K.T., 2014. Detecting nitrous oxide reductase (*nosZ*) genes in soil metagenomes: method development and implications for the nitrogen cycle. *MBio* 5 (3). <https://doi.org/10.1128/mBio.01193-14>.
- Otte, S., Grobden, N.G., Robertson, L.A., Jetten, M.S., Kuenen, J.G., 1996. Nitrous oxide production by *Alcaligenes faecalis* under transient and dynamic aerobic and anaerobic conditions. *Appl. Environ. Microbiol.* 62 (7), 2421–2426. <https://doi.org/10.1128/aem.62.7.2421-2426.1996>.
- Pan, Y., Ni, B.-J., Bond, P.L., Ye, L., Yuan, Z., 2013. Electron competition among nitrogen oxides reduction during methanol-utilizing denitrification in wastewater treatment. *Water Res.* 47 (10), 3273–3281. <https://doi.org/10.1016/j.watres.2013.02.054>.
- Petriglieri, F., Singleton, C., Peces, M., Petersen, J.F., Nierychlo, M., Nielsen, P.H., 2021. *Candidatus* *Dechloromonas phosphoritropha* and “*Ca. D. phosphorivorans*”, novel polyphosphate accumulating organisms abundant in wastewater treatment systems. *ISME J.* 15 (12), 3605–3614. <https://doi.org/10.1038/s41396-021-01029-2>.

- Qi, C., Zhou, Y., Suenaga, T., Oba, K., Lu, J., Wang, G., Zhang, L., Yoon, S., Terada, A., 2022. Organic carbon determines nitrous oxide consumption activity of clade I and II *nosZ* bacteria: genomic and biokinetic insights. *Water Res.* 209, 117910. <https://doi.org/10.1016/j.watres.2021.117910>.
- Ravishankara, A.R., Daniel, J.S., Portmann, R.W., 2009. Nitrous Oxide (N<sub>2</sub>O): the dominant ozone-depleting substance emitted in the 21st century. *Science* 326 (5949), 123–125. <https://doi.org/10.1126/science.1176985>.
- Ribera-Guardia, A., Kassotaki, E., Gutierrez, O., Pijuan, M., 2014. Effect of carbon source and competition for electrons on nitrous oxide reduction in a mixed denitrifying microbial community. *Process Biochem.* 49 (12), 2228–2234. <https://doi.org/10.1016/j.procbio.2014.09.020>.
- Ribera-Guardia, A., Marques, R., Arangio, C., Carvalho, M., Oehmen, A., Pijuan, M., 2016. Distinctive denitrifying capabilities lead to differences in N<sub>2</sub>O production by denitrifying polyphosphate accumulating organisms and denitrifying glycogen accumulating organisms. *Biotechnol.* 219, 106–113. <https://doi.org/10.1016/j.biortech.2016.07.092>.
- Robertson, L.A., Cornelisse, R., De Vos, P., Hadiotomoto, R., Kuenen, J.G., 1989. Aerobic denitrification in various heterotrophic nitrifiers. *Antonie Van Leeuwenhoek* 56 (4), 289–299. <https://doi.org/10.1007/BF00443743>.
- Robertson, L.A., Dalsgaard, T., Revsbech, N.-P., Kuenen, J.G., 1995. Confirmation of 'aerobic denitrification' in batch cultures, using gas chromatography and <sup>15</sup>N mass spectrometry. *FEMS Microbiol. Ecol.* 18 (2), 113–120. <https://doi.org/10.1111/j.1574-6941.1995.tb00168.x>.
- Roy, S., Nirakar, P., Yong, N.G.H., Stefan, W., 2021. Denitrification kinetics indicates nitrous oxide uptake is unaffected by electron competition in *Accumulibacter*. *Water Res.* 189, 116557. <https://doi.org/10.1016/j.watres.2020.116557>.
- Sanford, R.A., Wagner, D.D., Wu, Q., Chee-Sanford, J.C., Thomas, S.H., Cruz-Garcia, C., Rodriguez, G., Massol-Deya, A., Krishnani, K.K., Ritalahti, K.M., Nissen, S., Konstantinidis, K.T., Löffler, F.E., 2012. Unexpected nondenitrifier nitrous oxide reductase gene diversity and abundance in soils. *Proc. Natl. Acad. Sci.* 109 (48), 19709–19714. <https://doi.org/10.1073/pnas.1211238109>.
- Singleton, C.M., Petriglieri, F., Kristensen, J.M., Kirkegaard, R.H., Michaelsen, T.Y., Andersen, M.H., Kondrotaitė, Z., Karst, S.M., Dueholm, M.S., Nielsen, P.H., Albertsen, M., 2021. Connecting structure to function with the recovery of over 1000 high-quality metagenome-assembled genomes from activated sludge using long-read sequencing. *Nat. Commun.* 12 (1), 2009. <https://doi.org/10.1038/s41467-021-22203-2>.
- Singleton, C.M., Petriglieri, F., Wasmund, K., Nierychlo, M., Kondrotaitė, Z., Petersen, J.F., Peces, M., Dueholm, M.S., Wagner, M., Nielsen, P.H., 2022. The novel genus, '*Candidatus* Phosphoribacter', previously identified as *Tetrasphaera*, is the dominant polyphosphate accumulating lineage in EBPR wastewater treatment plants worldwide. *ISME J.* <https://doi.org/10.1038/s41396-022-01212-z>.
- Soler-Jofra, A., Pérez, J., van Loosdrecht, M.C.M., 2021. Hydroxylamine and the nitrogen cycle: a review. *Water Res.* 190, 116723. <https://doi.org/10.1016/j.watres.2020.116723>.
- Soler-Jofra, A., Stevens, B., Hoekstra, M., Picioreanu, C., Sorokin, D., van Loosdrecht, M.C.M., Pérez, J., 2016. Importance of abiotic hydroxylamine conversion on nitrous oxide emissions during nitrification of reject water. *Chem. Eng. J.* 287, 720–726. <https://doi.org/10.1016/j.cej.2015.11.073>.
- Song, M.J., Choi, S., Bae, W.B., Lee, J., Han, H., Kim, D.D., Kwon, M., Myung, J., Kim, Y.M., Yoon, S., 2020. Identification of primary effectors of N<sub>2</sub>O emissions from full-scale biological nitrogen removal systems using random forest approach. *Water Res.* 184, 116144. <https://doi.org/10.1016/j.watres.2020.116144>.
- Speth, D.R., in 't Zandt, M.H., Guerrero-Cruz, S., Dutilh, B.E., Jetten, M.S.M., 2016. Genome-based microbial ecology of anammox granules in a full-scale wastewater treatment system. *Nat. Commun.* 7 (1), 11172. <https://doi.org/10.1038/ncomms11172>.
- Stokholm-Bjerregaard, M., McIlroy, S.J., Nierychlo, M., Karst, S.M., Albertsen, M., Nielsen, P.H., 2017. A critical assessment of the microorganisms proposed to be important to enhanced biological phosphorus removal in full-scale wastewater treatment systems. *Front. Microbiol.* 8, 718. <https://doi.org/10.3389/fmicb.2017.00718>.
- Suenaga, T., Hori, T., Riya, S., Hosomi, M., Smets, B.F., Terada, A., 2019. Enrichment, isolation, and characterization of high-affinity N<sub>2</sub>O-reducing bacteria in a gas-permeable membrane reactor. *Environ. Sci. Technol.* 53 (20), 12101–12112. <https://doi.org/10.1021/acs.est.9b02237>.
- Suenaga, T., Ota, T., Oba, K., Usui, K., Sako, T., Hori, T., Riya, S., Hosomi, M., Chandran, K., Lackner, S., Smets, B.F., Terada, A., 2021. Combination of <sup>15</sup>N tracer and microbial analyses discloses N<sub>2</sub>O sink potential of the anammox community. *Environ. Sci. Technol.* 55 (13), 9231–9242. <https://doi.org/10.1021/acs.est.1c00674>.
- Suenaga, T., Riya, S., Hosomi, M., Terada, A., 2018. Biokinetic characterization and activities of N<sub>2</sub>O-reducing bacteria in response to various oxygen levels. *Front. Microbiol.* 9, 697. <https://doi.org/10.3389/fmicb.2018.00697>.
- Suzek, B.E., Wang, Y., Huang, H., McGarvey, P.B., Wu, C.H., the UniProt Consortium, 2015. UniProt clusters: a comprehensive and scalable alternative for improving sequence similarity searches. *Bioinformatics* 31 (6), 926–932. <https://doi.org/10.1093/bioinformatics/btu739>.
- Tallec, G., Garnier, J., Billen, G., Goussailles, M., 2008. Nitrous oxide emissions from denitrifying activated sludge of urban wastewater treatment plants, under anoxia and low oxygenation. *Biotechnol. Technol.* 99 (7), 2200–2209. <https://doi.org/10.1016/j.biortech.2007.05.025>.
- Terada, A., Sugawara, S., Hojo, K., Takeuchi, Y., Riya, S., Harper, W.F., Yamamoto, T., Kuroiwa, M., Isobe, K., Katsuyama, C., Suwa, Y., Koba, K., Hosomi, M., 2017. Hybrid nitrous oxide production from a partial nitrifying bioreactor: hydroxylamine interactions with nitrite. *Environ. Sci. Technol.* 51 (5), 2748–2756. <https://doi.org/10.1021/acs.est.6b05521>.
- Vasilaki, V., Volcke, E.I.P., Nandi, A.K., van Loosdrecht, M.C.M., Katsou, E., 2018. Relating N<sub>2</sub>O emissions during biological nitrogen removal with operating conditions using multivariate statistical techniques. *Water Res.* 140, 387–402. <https://doi.org/10.1016/j.watres.2018.04.052>.
- Veira, A., Galinha, C.F., Oehmen, A., Carvalho, G., 2019. The link between nitrous oxide emissions, microbial community profile and function from three full-scale WWTPs. *Sci. Total Environ.* 651, 2460–2472. <https://doi.org/10.1016/j.scitotenv.2018.10.132>.
- Veira, A., Ribera-Guardia, A., Marques, R., Barreto Crespo, M.T., Oehmen, A., Carvalho, G., 2018. The link between the microbial ecology, gene expression, and biokinetics of denitrifying polyphosphate-accumulating systems under different electron acceptor combinations. *Appl. Microbiol. Biotechnol.* 102 (15), 6725–6737. <https://doi.org/10.1007/s00253-018-9077-3>.
- Wang, Y., Gao, H., Wells, G., 2021. Integrated omics analyses reveal differential gene expression and potential for cooperation between denitrifying polyphosphate and glycogen accumulating organisms. *Environ. Microbiol.* 23 (6), 3274–3293. <https://doi.org/10.1111/1462-2920.15486>.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., ..., Yutani, H., 2019. Welcome to the Tidyverse. *J. Open Source Software* 4 (43), 1686. <https://doi.org/10.21105/joss.01686>.
- Wrage-Mönnig, N., Horn, M.A., Well, R., Müller, C., Velthof, G., Oenema, O., 2018. The role of nitrifier denitrification in the production of nitrous oxide revisited. *Soil Biol. Biochem.* 123, A3–A16. <https://doi.org/10.1016/j.soilbio.2018.03.020>.
- Wunderlin, P., Mohn, J., Joss, A., Emmenegger, L., Siegrist, H., 2012. Mechanisms of N<sub>2</sub>O production in biological wastewater treatment under nitrifying and denitrifying conditions. *Water Res.* 46 (4), 1027–1037. <https://doi.org/10.1016/j.watres.2011.11.080>.
- Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F.O., Ludwig, W., Schleifer, K.-H., Whitman, W.B., Euzéby, J., Amann, R., Rosselló-Móra, R., 2014. Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat. Rev. Microbiol.* 12 (9), 635–645. <https://doi.org/10.1038/nrmicro3330>.
- Yoon, S., Nissen, S., Park, D., Sanford, R.A., Löffler, F.E., 2016. Nitrous oxide reduction kinetics distinguish bacteria harboring clade I *NosZ* from those harboring clade II *NosZ*. *Appl. Environ. Microbiol.* 82 (13), 3793–3800. <https://doi.org/10.1128/AEM.00409-16>.
- Zhang, M., Gao, J., Liu, Q., Fan, Y., Zhu, C., Liu, Y., He, C., Wu, J., 2021. Nitrite accumulation and microbial behavior by seeding denitrifying phosphorus removal sludge for partial denitrification (PD): the effect of COD/NO<sub>3</sub> ratio. *Biotechnol.* 323, 124524. <https://doi.org/10.1016/j.biortech.2020.124524>.
- Zumft, W.G., 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* 61, 84. <https://doi.org/10.1128/mmb.61.4.533-616.1997>.