



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

The HLS19-COM-P, a New Instrument for Measuring Communicative Health Literacy in Interaction with Physicians

Development and Validation in Nine European Countries

Finbråten, Hanne Sørberg; Nowak, Peter; Griebler, Robert; Bíró, Éva; Vrđelja, Mitja; Charafeddine, Rana; Griese, Lennert; Bøggild, Henrik; Schaeffer, Doris; Link, Thomas; Kucera, Zdenek; Mancini, Julien; Pelikan, Jürgen M.

Published in:

International Journal of Environmental Research and Public Health

DOI (link to publication from Publisher):

[10.3390/ijerph191811592](https://doi.org/10.3390/ijerph191811592)

Creative Commons License
CC BY 4.0

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Finbråten, H. S., Nowak, P., Griebler, R., Bíró, É., Vrđelja, M., Charafeddine, R., Griese, L., Bøggild, H., Schaeffer, D., Link, T., Kucera, Z., Mancini, J., & Pelikan, J. M. (2022). The HLS19-COM-P, a New Instrument for Measuring Communicative Health Literacy in Interaction with Physicians: Development and Validation in Nine European Countries. *International Journal of Environmental Research and Public Health*, 19(18), Article 11592. <https://doi.org/10.3390/ijerph191811592>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -



Article

The HLS₁₉-COM-P, a New Instrument for Measuring Communicative Health Literacy in Interaction with Physicians: Development and Validation in Nine European Countries

Hanne Søberg Finbråten ^{1,*}, Peter Nowak ², Robert Griebler ², Éva Bíró ³, Mitja Vrdelja ⁴, Rana Charafeddine ⁵, Lennert Griese ⁶, Henrik Bøggild ⁷, Doris Schaeffer ⁶, Thomas Link ⁸, Zdenek Kucera ⁹, Julien Mancini ¹⁰ and Jürgen M. Pelikan ¹¹

¹ Department of Health and Nursing Sciences, Faculty of Social and Health Sciences, Inland Norway University of Applied Sciences, P.O. Box 400, 2418 Elverum, Norway

² Competence Centre for Health Promotion and Health System, Austrian National Public Health Institute, A-1010 Vienna, Austria

³ Department of Public Health and Epidemiology, Faculty of Medicine, University of Debrecen, H-4028 Debrecen, Hungary

⁴ Communication Unit, National Institute of Public Health, Trubarjeva 2, 1000 Ljubljana, Slovenia

⁵ Department of Public Health and Epidemiology, Sciensano, 1050 Brussels, Belgium

⁶ Interdisciplinary Centre for Health Literacy Research [ICHL], School of Public Health, Bielefeld University, 33615 Bielefeld, Germany

⁷ Public Health and Epidemiology, Department of Health Science and Technology, Aalborg University, 9220 Aalborg, Denmark

⁸ Department of Quality Measurement and Patient Survey, Austrian National Public Health Institute, A-1010 Vienna, Austria

⁹ Czech Health Literacy Institute, Sokolská 490/31, 120 00 Prague, Czech Republic

¹⁰ Aix Marseille University, APHM INSERM, IRD, ISSPAM, SESSTIM, Cancer, Biomedicine & Society Group, Ligue 2019 Accredited Team, F-13009 Marseille, France

¹¹ WHO-CC Health Promotion in Hospitals and Health Care, Austrian National Public Health Institute, A-1010 Vienna, Austria

* Correspondence: hanne.finbraten@inn.no



Citation: Finbråten, H.S.; Nowak, P.; Griebler, R.; Bíró, É.; Vrdelja, M.; Charafeddine, R.; Griese, L.; Bøggild, H.; Schaeffer, D.; Link, T.; et al. The HLS₁₉-COM-P, a New Instrument for Measuring Communicative Health Literacy in Interaction with Physicians: Development and Validation in Nine European Countries. *Int. J. Environ. Res. Public Health* **2022**, *19*, 11592. <https://doi.org/10.3390/ijerph191811592>

Academic Editor: Paul B. Tchounwou

Received: 6 July 2022

Accepted: 12 August 2022

Published: 14 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Background: Sufficient communicative health literacy (COM-HL) is important for patients actively participating in dialogue with physicians, expressing their needs and desires for treatment, and asking clarifying questions. There is a lack of instruments combining communication and HL proficiency. Hence, the aim was to establish an instrument with sufficient psychometric properties for measuring COM-HL. Methods: The HLS₁₉-COM-P instrument was developed based on a conceptual framework integrating HL with central communicative tasks. Data were collected using different data collection modes in nine countries from December 2019 to January 2021 ($n = 18,674$). Psychometric properties were assessed using Rasch analysis and confirmatory factor analysis. Cronbach's alpha and Person separation index were considered for reliability. Results: The 11-item version (HLS₁₉-COM-P-Q11) and its short version of six items (HLS₁₉-COM-P-Q6) fit sufficiently the unidimensional partial credit Rasch model, obtained acceptable goodness-of-fit indices and high reliability. Two items tend to under-discriminate. Few items displayed differential item functioning (DIF) across person factors, and there was no consistent pattern in DIF across countries. All items had ordered response categories. Conclusions: The HLS₁₉-COM-P instrument was well accepted in nine countries, in different data collection modes, and could be used to measure COM-HL.

Keywords: Calgary-Cambridge Guide framework (C-CG); communicative health literacy; confirmatory factor analysis; data collection modes; HLS₁₉; measurement; physician-patient communication; Rasch analysis

1. Introduction

In recent decades, research has highlighted the importance of health literacy (HL) as a relevant determinant of health. Among others, low HL is found to be associated with poorer health [1–3], unhealthier behaviour and a higher number of visits to healthcare services [3–6]. There are relatively high proportions of people with low HL, and there is also a social gradient for HL [3,5].

By communicating in a “health literate” way addressing patients’ needs, health professionals can play a critical role in taking the patients’ HL into account and strengthening their HL. Adapting health communication to the patients’ HL might also enable them to play an active role as co-producer in healthcare and be able to make sound health decisions.

The quality of communication between patients and health professionals is one of the patients’ main concerns and the most important reason for (dis-)satisfaction with healthcare [7–9]. It has an important impact on trust in healthcare providers and the healthcare system. Successful communication in healthcare is associated with a wide range of improved healthcare outcomes and does also contribute to the workplace satisfaction of health professionals [10–12]. This indicates the importance of communication proficiency and responsibility for good communication skills on the side of health professionals, who must be regularly trained in this regard.

Patients (the term is used here as a synonym also for healthcare users, clients, citizens, individuals, and people) strongly rely on personal communication with health professionals [13] to understand and manage their health issues. Thus, health(care) communication and patient participation in healthcare have been recognized as a decisive part of HL, as well as a critical determinant of successful disease management and health outcomes [14,15]. However, patients with low HL tend to report poorer communication skills in dialogue with health professionals than those with higher HL [16,17]. Furthermore, patients with lower literacy are less likely to ask questions to health professionals [18]. Hence, sufficient communicative skills of health professionals are especially necessary to actively involve patients with low HL in the dialogue, so they can use health information to manage their health situation properly. Information about patients’ communicative HL (COM-HL) is, thus, needed to better adapt health communication accordingly. In addition, personal communication seems to be an important support for patients to navigate the healthcare system. Smith et al. [19] claim that interactive HL, or COM-HL, comprises the skills to interact with both the healthcare system and the providers. Communication is also related to the theoretical framework and definition of navigational HL [20]. Hence, COM-HL is seen as very relevant for “navigational HL” [20,21].

Nutbeam [22–24] stresses the importance of interactive processes by defining communicative/interactive HL as one of three relevant types of HL and linking it to greater autonomy and personal empowerment. Expanding Nutbeam’s conceptualization of communicative/interactive HL by integrating the HL framework of the European Health Literacy Survey (HLS-EU) Consortium [25,26], the following definition of COM-HL in healthcare is proposed: “Communicative health literacy refers to patients’ communicative and social skills that enable them to actively engage in face-to-face encounters with healthcare professionals, to give and seek information, derive meaning from it, and apply this information in decision making and in co-producing their health care” [27] (p. 235).

So far, there is no instrument measuring COM-HL as an independent construct [28]. Nutbeam [23] highlights the need for a specific measurement tool for these “oral literacy and social skills”. Based on Nutbeam’s concept, Ishikawa et al. [29] developed the “Functional, Communicative and Critical HL” scale, where the dimension intending to measure COM-HL consists of five items. However, these items are more about processing health information in general and do not comprise patients’ ability to interact with healthcare professionals. On the other hand, Chinn & McCarthy [30] suggested three items on COM-HL in relation to doctor or nurse interaction. However, these items were a part of a scale intending to also measure functional and critical HL. Hence, the COM-HL items were not considered or tested as a stand-alone scale. The 47-item version of the European HL-Survey

Questionnaire (HLS-EU-Q47) [26] also includes a few items concerning COM-HL but without systematic reference to the broader research on healthcare communication. O'Hara et al. [31] published the concept of the “Conversational Health Literacy Assessment Tool (CHAT)” to provide a short actionable survey tool for the clinical context to assess patients' ability to interact with health professionals, but the 10 items mainly focus on general health information-seeking behaviour and health promotion activities. Only one item focuses on interactive behaviour.

In summary, previous instruments meant to measure COM-HL were either developed to measure only certain communicative tasks or outcomes or to capture only certain aspects of healthcare communication. As far as we know, no previous instrument integrated systematic findings from communication research and HL research into one instrument. We would also like to consider COM-HL as a relational proficiency. Hence, there is a need for a new instrument that covers the COM-HL skills necessary for actively participating in health communication with healthcare professionals, especially physicians.

Based on this background, this article aims to establish an international instrument with sufficient psychometric properties, intending to measure communicative health literacy in patient–physician communication.

2. Materials and Methods

2.1. Development of the Instrument for Measuring Communicative Health Literacy in Patient–Physician Communication

The new COM-HL with physicians in healthcare instrument, HLS₁₉-COM-P, was developed in the context of the Health Literacy Population Survey Project 2019–2021 (HLS₁₉), a European HL survey within 17 participating countries from the WHO-Europe region [3]. The HLS₁₉-COM-P is based on a comprehensive theoretical framework that integrates Nutbeam's [22] idea of COM-HL, the basic competencies of information processing according to the HL framework of the HLS-EU Consortium [25,26], and the main communicative tasks of the Calgary-Cambridge Guide framework (C-CG) [32]. The C-CG framework has been developed over the last 25 years and integrates the results of different research traditions to serve as a guide to teach health professionals in patient-centred communication skills. This framework is also used as a framework for assessments of communicative skills of health professionals [33]. The C-CG describes 56 single communicative practices of a health professional in six main phases of a routine interaction in healthcare. Within these six main phases, the C-CG identifies the communicative tasks of patients that need to be considered in the conceptual framework for COM-HL (see Figure 1).

C-CG main phases	Patients' communicative tasks	Participating actively
1. Initiating the session	1. Opening the session and giving initial information	
2. Gathering information	2. Giving full information	
3. Providing structure	3. Understanding and following the agenda	
4. Building relationship	4. Expressing own views / feelings and trusting the health professionals (e.g., physician)	
5. Explanation and planning	5. Understanding and decision making (extract information & derive meaning)	
6. Closing the session	6. Final understanding, agreement (and application)	

Figure 1. Overview of the main communicative practices of health professionals of the Calgary-Cambridge Guide to the Medical Interview (C-CG) and main communicative tasks of patients, constituting the Conceptual Framework for Communicative Health Literacy.

Based on our conceptual framework and definition of COM-HL (see above), the HLS₁₉-COM-P was developed in a multistage process by a working group of representatives from HLS₁₉ countries interested in COM-HL (see Figure 2). After creating the conceptual framework in the first step, the HLS-EU-Q47 [26,34] instrument for measuring general HL was reviewed for suitable items in the second step. Five items on communication in

healthcare were identified (Q5, Q8, Q9, Q13 and Q16), but, according to the C-CG, these items only measure aspects of two main phases of interactions between patients and health professionals (explanation and planning; closing the session). In particular, key patient communication tasks were not captured by the HLS-EU-Q47, e.g., presenting their concerns and preferences, and asking questions. Therefore, as a third step, a targeted literature search in English or German language was conducted to identify existing instruments and possible items for measuring COM-HL. In addition to the HLS-EU-Q47, a total of 20 different instruments were found. Since none of these instruments covers all relevant aspects of the underlying conceptual framework and definition of COM-HL, the working group decided to develop a set of items inspired by these instruments. Hence, in the fourth step, the most relevant items were selected from the pool of 183 items and tested in an expert panel and in two focus group interviews. The aim was to identify at least one item per C-CG main phase and the related main communicative task (see Figure 1 and Table 1) to capture the main HL-related challenges in healthcare communication. A preliminary set of 15 items (including 5 items from the HLS-EU-Q47) was selected, which was adapted to the question format of the HLS₁₉-HL instruments. In accordance with the HLS₁₉ methodology [3], the items are formulated as direct questions (see Table 1) and are rated using a four-point Likert response scale: very easy (4), easy (3), difficult (2), very difficult (1). The comprehensibility and importance of the individual items were assessed in two focus groups involving potential survey participants in Austria (using a German preliminary version of the instrument). Of the focus group participants ($n = 14$), four were men, they were aged 18–54, four had a university degree, eight had high school graduation, and two participants had compulsory school as their highest completed education. Three of the participants were chronically ill. In general, the 15 items were well received and understood. However, the focus group interviews revealed that the term “health professional” was not well accepted by participants because their experiences varied by type of health professional. The term was, therefore, perceived as too vague, making it difficult to form opinions and respond to the items. These general insights were included in creating the original English version. In addition, the set of items and item wording were discussed in several feedback loops within the working group, with the HLS₁₉ International Coordination Center, and the HLS₁₉ Consortium to ensure transferability to different national contexts. The discussion in the working group indicated that the status of different health professions varies widely in the participating countries, while the status of physicians seems to be quite similar and comparable. Based on these considerations the COM-HL instrument focuses exclusively on physician–patient communication. To create an independent instrument for measuring COM-HL, the five items from the HLS-EU-Q47 were excluded, constituting a set of 11 items (HLS₁₉-COM-P-Q11) reflecting the conceptual framework. The HLS₁₉-COM-P instrument measures all six main communicative phases of physician–patient interactions according to the C-CG and can be used to analyze the dimensions of COM-HL in accordance with Nutbeam [22] and the basic competencies of information processing according to the conceptual model of HL developed by the HLS-EU Consortium [25,26].

In addition, a 6-item short form (HLS₁₉-COM-P-Q6) (see Table 1) was suggested. The short form was proposed based on content considerations, and the shorter length allowed more countries to include HLS₁₉-COM-P in their national survey. The short form might also be included more easily in future studies with patients.




Figure 2. Steps in the development of the instrument to measure communicative health literacy.

Table 1. Overview of items included in the instrument for measuring communicative health literacy. Items constituting the short version are in italics.

Patients' Communicative Tasks	Items	
	On a Scale from Very Easy to Very Difficult, How Easy Would You Say It Is for You ...	
1. Opening the session and giving initial information	COM1	... to describe to your doctor your reasons for coming to the consultation?
2. Giving full information	COM2	... to make your doctor listen to you without being interrupted?
	COM3	... <i>to explain your health concerns to your doctor?</i>
	COM4	... <i>to get enough time in the consultation with your doctor?</i>
3. Understanding and following the agenda	COM5	... <i>to express your personal views and preferences to your doctor?</i>
	COM6	... to get the information you need from your doctor?
4. Expressing one's own views and trusting	COM7	... to understand the words used by your doctor?
	COM8	... <i>to ask your doctor questions in the consultation?</i>
	COM9	... <i>to be involved in decisions about your health in dialogue with your doctor?</i>
5. Understanding and decision making	COM10	... <i>to recall the information you get from your doctor?</i>
	COM11	... to use the information from your doctor to take care of your health?
6. Final understanding and agreement		

2.2. Translation Process

The HLS₁₉-COM-P instrument was developed in English and translated into eight languages. In six countries (Austria [AT], Belgium [BE] (Dutch translation), Germany [DE], Denmark [DK], Hungary [HU] and Slovenia [SI]), the translation process followed a two-step procedure: first, two forward translations were prepared, one by the National Study Centre (NSC) and one by the data collection agency (DCA), and, second, a comparison of the two translations was carried out by the NSC, with the most appropriate translation being selected in consensus with the DCA in case of differences. The AT and DE versions were also aligned. In three countries (BE (French translation), Bulgaria (BG) and Czech Republic (CZ)), only one forward translation was performed. Back-translation was conducted in CZ and SI. In France [FR], the BE French translation was used with very minor adaptations.

2.3. Data Collection

The HLS₁₉-COM-P instrument was included as an optional package in the HLS₁₉ survey [3]. Countries that chose this optional package could either use the 11-item version (HLS₁₉-COM-P-Q11) or the 6-item short version (HLS₁₉-COM-P-Q6). Data on COM-HL were collected in nine countries (AT, BE, BG, CZ, DE, DK, FR, HU, and SI). The 11-item version was applied in three of these (AT, DE, and SI). Data were collected using different modes of data collection (Table 2) from December 2019 to June 2021 based on multi-stage random sampling or quota sampling procedures in most countries. A mix of survey methods was used in three countries (BG, CZ, and SI). Except for DE, all surveys took place during the COVID-19 pandemic, which had an impact on possible data collection modes. The sample size in the countries varied from 865 (BG) to 3602 (DK) respondents and data on COM-HL were collected from a total of 18,674 respondents.

Table 2. Main characteristics of data collection in countries measuring communicative health literacy.

Country	Item Set	Language	Mode of Data Collection	Sampling Procedure	Period of Data Collection	Number of Respondents ⁱ
AT	Q11	German	CATI	Multi-stage random sampling	16 March–26 May 2020	2954
BE	Q6	Dutch, French	CAWI	Quota sampling	30 January–28 February 2020; 1–26 October 2020	1000
BG	Q6	Bulgarian	CAPI, CAWI	Proportional stratified sampling (CAPI) and random quota sampling (CAWI)	15 August–30 November 2020 (CAPI); 1 April–1 June 2021 (CAWI)	859
CZ	Q6	Czech	CATI, CAWI	Random digital procedure (CATI) and random quota sampling (CAWI)	10–24 November 2020	1597
DE	Q11	German	PAPI	Multi-stage random and quota sampling combined	13 December 2019–27 January 2020	2133
DK	Q6	Danish	CAWI	Multi-stage random sampling	11 December 2020–5 February 2021	3600
FR	Q6	French	CAWI	Quota sampling	27 May–5 June 2020; 8–18 January 2021	2003
HU	Q6	Hungarian	CATI	Multi-stage random sampling	2–20 December 2020	1186
SI	Q11	Slovenian	CAPI, self-administered paper and pencil ⁱⁱ , CAWI	Multi-stage random sampling	9–15 March 2020; 9 June 2020–10 August 2020	3342

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; HU = Hungary; PAPI = paper-assisted personal interviews; SI = Slovenia. ⁱ The number of respondents who have answered one or more HLS₁₉-COM-P items. ⁱⁱ Only 12 individuals responded using paper and pencil. These records were excluded from the analyses.

2.4. Analyses

At the overall level, the psychometric properties of the HLS₁₉-COM-P-Q11 and its short version HLS₁₉-COM-P-Q6 were assessed using Rasch analysis and by using one-factorial confirmatory factor analysis (CFA). CFA provides detailed information about the overall fit of the model [35] but has some shortcomings (e.g., the results are sample- and scale-dependent and the standard error of measurement is constant) [36,37]. Rasch models are considered as parsimonious models meeting the requirements of fundamental measurement [38]. Rasch analysis also provides detailed information about the items. Hence,

Rasch analysis was performed at both overall and item levels [39]. As mentioned above, the HLS₁₉-COM-P items are interpreted as ordinal scaled. In the International Report on the Methodology, Results, and Recommendations of the European Health Literacy Population Survey 2019–2021 (HLS₁₉) of M-POHL [3], the analyses were based on dichotomous data. In this article, the assessment of psychometric properties is conducted on both the polytomous version of the COM-HL items (very difficult—difficult—easy—very easy) and on a dichotomized version of the COM-HL items (very easy/easy versus difficult/very difficult) to explore which one might be preferable. The internal consistency and reliability were assessed using Cronbach's alpha and the Person Separation Index (PSI). If conclusions are to be drawn at the individual or group level, the indexes are recommended to exceed 0.85 or 0.65, respectively [40]. Omega for categorical data was used as an index for composite reliability [41]. In addition, the average variance extracted (AVE) was evaluated. An AVE value of ≥ 0.5 could be considered as acceptable [42]. The analyses were conducted for each country and separately for different modes of data collection if used within a country.

In terms of Rasch analysis, data were tested against the partial-credit parameterization [43] of the unidimensional Rasch model [44]. Analyses at the overall level included data-model fit, targeting (mean person location), and dimensionality [39]. Chi-square statistics were applied to assess the data-model fit. The targeting of the HLS₁₉-COM-P-Q11 and the HLS₁₉-COM-P-Q6 was assessed by comparing the item and person location distributions on the same metric. An instrument could be deemed as well-targeted if the mean person location values are around zero [45]. Graphical displays for targeting were also inspected. Dimensionality was assessed using the combined procedure of principal component analysis (PCA) of residuals and paired *t*-tests [46–48]. Based on the PCA of residuals, two subsets of items were made, and paired *t*-tests were used to examine whether the subsets provided significantly different person location estimates. A scale could be considered sufficiently unidimensional if the proportion of individuals with significantly different person location estimates on the pair of compared subscales does not exceed 5% (or if the lower bound of the binomial 95% confidence interval (CI) does not exceed 5%) [46,48].

CFA was performed for a one-factor model of the HLS₁₉-COM-P-Q11 and the HLS₁₉-COM-P-Q6 using a WLSMV estimator with diagonally weighted least squares [49–51]. The following goodness-of-fit (GOF) indices were considered: standardized root mean square residual (SRMR), root mean square error of approximation (RMSEA), comparative fit index (CFI), Tucker–Lewis index (TLI), goodness-of-fit index (GFI) and adjusted goodness-of-fit index (AGFI). Schumacker and Lomax [52] recommend SRMR < 0.05, RMSEA ≤ 0.05 –0.08, CFI, TLI, GFI and AGFI values close to 0.90 or 0.95, whereas Hu and Bentler [53] claim that SRMR values close to or below 0.08 indicate sufficient overall fit. An overview of GOF indices with reference values could also be found in Table S1: Fit indices considered in confirmatory factor analysis.

Rasch analyses at a finer level included assessing item fit, response dependence, ordering of response categories and differential item functioning (DIF) [39]. Chi-square probability values above a Bonferroni-adjusted *p*-value of 5% and fit residuals within the range of ± 2.5 indicate adequate item fit [45]. In addition, the mean square residual fit statistic (MNSQ) infit was used to assess item fit. For measuring COM-HL at the population level, infit between 0.7 and 1.3 was considered as sufficient [54]. A residual correlation of <0.3 was applied as an indicator of response dependency. In addition, residual correlations were assessed relative to each other [55]. For ordinal data, the threshold ordering was inspected both statistically and graphically to examine whether the response categories could be considered working as intended. A key requirement of measurement is that items measure invariantly across levels of different person factors, such as gender, age, and education. Lack of invariance in measurement across person factors is called differential item functioning (DIF) [56,57]. Uniform DIF means that there are consistent systematic differences in responses across person factor levels, whereas nonuniform DIF is present if the DIF varies along the latent trait, i.e., the persons factor interacts with the latent

trait [56]. Items were inspected for DIF across different levels of person factors (gender, age, educational level, status of employment, ability to pay bills, self-perceived level in society and self-reported general health status), both statistically using two-way analysis of variance of standardized residuals and graphically by inspecting item characteristic curves [58]. Statistical significance was assumed at a Bonferroni-adjusted p -value $\leq 5\%$. An overview of the tests performed with reference values is also found in Table S2: Analyses and tests with reference values considered in Rasch analyses.

Since chi-square statistics are sensitive to sample size, there is a risk of drawing false conclusions due to large sample sizes [59]. Therefore, the amend sample size function in the software RUMM2030 was used to draw a random sub-sample for analyses concerning data-model fit, item fit and DIF. As recommended, the sample size was calculated by multiplying the number of items (11/6) by the number of thresholds (3 for polytomous items), with 10–30 persons per threshold [58], indicating that a sample size of 330–990/180–540 ($11/6 \times 3 \times 10/30$) can be deemed as adequate for these analyses.

Items with a negative item location estimate could be considered as relatively easy to endorse, whereas the opposite is the case for items with a positive item location estimate. A higher value indicates that the item is having a higher difficulty level and is, consequently, harder to endorse [58].

Convergent and discriminant validity are also facets of construct validity [60]. As the HLS₁₉-COM-P intends to measure an aspect of HL, COM-HL, it would be expected that the COM-HL score is positively (moderately) correlated with general HL (GEN-HL) and with navigational HL (HL-NAV; convergent validity) but is still a separate construct (discriminant validity). For convergent validity, Pearson's correlations were used to assess the associations between COM-HL and GEN-HL and HL-NAV. A positive moderate correlation between these scores would be expected, as the instruments intend to capture different aspects of HL. To assess discriminant validity, the combined procedure of PCA of residuals and paired sample t -test was applied to investigate whether the different HL instruments could be considered measuring distinctive constructs. GEN-HL and HL-NAV were measured using the HLS₁₉-Q12 and the HLS₁₉-NAV instruments, each consisting of 12 items [3].

Rasch analyses were performed using the software RUMM2030Plus [61] and ACER ConQuest 5 [62], the lavaan package [51] for R [63] was applied for CFA, and the correlation analyses were conducted using R.

2.5. Missing

On average, there were few missing values. In most countries, the number of missing values varied between 0 and 2 or 3%. In BG data, COM11 had 11% missing values. Conducting Rasch analysis, missing data were handled through full information maximum likelihood estimation (FIML), whereas the other analyses included respondents that had at least 80% of valid responses.

3. Results

Table 3 provides details regarding the main characteristics of the samples, including the key demographics, socioeconomic variables, and health status.

Table 3. Sample characteristics (in percentages) for the participating countries in HLS₁₉, measuring HLS₁₉-COM-P, divided by data collection mode.

Characteristic		AT	BE	BG CAPI	BG CAWI	CZ CATI	CZ CAWI	DE	DK	FR	HU	SI CAPI	SI CAWI	
<i>n</i>		2954	1000	402	457	531	1066	2133	3600	2003	1186	1855	1487	
Gender	male	44.2	49.6	29.6	24.5	40.7	51.3	49.6	43.9	49.2	47.8	47.0	45.5	
	female	55.8	50.4	70.4	75.5	59.3	48.7	50.2	56.1	50.8	52.2	53.0	54.5	
	missing	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	
Age #	Dichotomized	≤45	34.1	43.9	60.5	66.5	18.6	57.0	38.3	21.7	48.6	33.6	28.4	51.8
		≥46	65.8	56.1	37.8	33.5	81.4	43.0	60.8	78.3	51.4	66.4	71.6	48.2
		missing	0.1	0.0	1.7	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0
	Categorized (1st version)	18 to 25 years	6.8	9.0	18.9	20.5	1.7	12.7	9.4	4.4	12.0	7.2	5.3	11.5
		26 to 65 years	70.4	74.7	73.2	75.3	49.7	76.3	65.2	60.7	74.8	65.4	61.9	75.6
		66 years and older	22.7	16.3	6.2	4.2	48.6	11.0	24.5	34.9	13.2	27.4	32.8	12.9
		missing	0.1	0.0	1.7	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0
	Categorized (2nd version)	18 to 45 years	34.1	43.9	60.5	66.5	18.6	57.0	38.3	21.7	48.6	33.6	28.4	51.8
		46 to 75 years	56.3	53.3	37.1	32.4	68.0	41.4	50.0	69.1	51.4	57.5	57.9	44.2
76 years and older		9.5	2.8	0.7	1.1	13.4	1.6	10.8	9.2	0.0	8.9	13.7	4.0	
missing		0.1	0.0	1.7	0.0	0.0	0.0	0.9	0.0	0.0	0.0	0.0	0.0	
Highest level of completed education	Upper secondary school (ISCED 0 to 3)	61.9	14.7	32.3	22.1	86.6	71.8	54.4	15.1	17.9	70.4	80.9	54.3	
	above	38.1	84.1	66.2	77.5	13.2	28.2	43.5	84.8	82.1	29.6	19.1	45.7	
	missing	0.0	1.2	1.5	0.4	0.2	0.0	2.1	0.1	0.0	0.0	0.0	0.0	
Status of employment	employed	59.7	57.7	81.4	87.8	34.5	72.5	60.5	55.2	67.5	56.9	44.8	70.3	
	unemployed or retired	40.0	38.4	14.4	9.4	65.2	27.2	38.1	40.9	32.5	42.5	54.9	28.4	
	missing	0.3	3.9	4.2	2.8	0.3	0.3	1.4	3.9	0.0	0.6	0.3	1.3	
Ability to pay bills	easy	85.8	62.4	61.5	63.0	81.4	67.4	73.7	92.8	74.6	67.4	56.2	61.2	
	difficult	13.3	37.6	33.3	35.2	18.3	32.6	22.5	6.9	25.4	31.3	42.5	38.7	
	missing	0.9	0.0	5.2	1.8	0.4	0.0	3.8	0.3	0.0	1.3	1.3	0.1	
Self-perceived level in society (1 to 10)	level 4 or lower ⁱ	6.8	10.6	11.7	9.0	13.9	17.4	17.3	11.4	20.3	26.1	25.9	20.6	
	level 5 or higher	87.2	89.4	77.9	79.0	84.0	82.6	80.0	88.3	79.7	72.6	71.1	79.1	
	missing	6.0	0.0	10.4	12.0	2.1	0.0	2.7	0.3	0.0	1.3	3.0	0.3	
Self-reported general health	good or fair	97.0	92.1	96.3	96.3	85.5	91.5	93.0	92.6	92.6	91.1	90.2	96.3	
	bad	2.9	7.9	3.5	3.3	14.3	8.5	6.9	7.3	7.4	8.8	9.7	3.6	
	missing	0.1	0.0	0.2	0.4	0.2	0.0	0.1	0.1	0.0	0.1	0.1	0.1	

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; HU = Hungary; SI = Slovenia. [#] We used different categorizations of age when conducting analyses of differential item functioning to explore if the categorization had an impact on the results in these analyses. Patients aged ≥ 76 are perceived as a vulnerable subpopulation. In FR, the sample was collected among people aged below 75. ⁱ People with a score of ≤4 are considered at a low level in society [64,65].

3.1. Rasch Analyses at the Overall Level

At the overall level, the polytomous scored HLS₁₉-COM-P-Q11 displayed misfit in all countries when a sample size of $n = 660$ (20 persons for each of the 33 thresholds) (11 items \times 3 thresholds) was considered (Table 4, [39]). Reducing the sample size to 330 in each country, the HLS₁₉-COM-P-Q11 displayed acceptable overall data-model fit. Applying a sample size of $n = 360$, the polytomous scored HLS₁₉-COM-P-Q6 displayed acceptable overall data-model fit in AT and DE data. Reducing the sample size to 180 (6 items \times 3 thresholds \times 10) the short version also displayed acceptable data-model fit in the other countries. The proportion of significant different person location estimates across subtests for the HLS₁₉-COM-P-Q11 varied between 4.8% (SI; CAPI) and 7.9% (DE; PAPI), and between 3.0% (HU; CATI and SI; CAPI) and 7.5% (DK; CAWI) for HLS₁₉-COM-P-Q6, indicating that the scales could be considered sufficiently unidimensional in all countries. The targeting of both the long and the short version could have been better, as the items, on average, were quite easy to endorse (mean person location varying between 1.38 (DE; PAPI) and 2.73 (SI; CAWI), and 1.21 (DE; PAPI) and 2.47 (SI; CAWI) for HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6, respectively (Table 4, [39]).

In countries applying different modes of data collection, higher mean person location was observed for data obtained from CAWI than from CAPI (SI (long and short version) and BG). A higher mean person location was also observed in data obtained from CATI compared with CAWI (CZ; Figure 3a,b).




Figure 3. (a) Person item distribution for HLS₁₉-COM-P-Q6 in the Czech Republic based on CATI data. (b) Person item distribution for HLS₁₉-COM-P-Q6 in the Czech Republic based on CAWI data.

Assessing fit to the Rasch model based on dichotomized items indicated low power of analysis of fit and a decreased PSI. The results based on dichotomous data also led to a sharp increase in the number of records with extreme scores (4–12 times more extreme records). Reasonable power of analyses of fit was observed only for the DE version of the dichotomized HLS₁₉-COM-P-Q11. In DE data, the number of extreme records increased from 81 when analyses were based on polytomous data to 642 when based on dichotomous data. The PSI decreased from 0.89 to 0.62. Hence, the following results from Rasch analyses provided in this paper are based on polytomous data.

3.2. Confirmatory Factor Analysis

Regardless of considering polytomous or dichotomous data, most goodness-of-fit indices for both the HLS₁₉-COM-P-Q11 and the HLS₁₉-COM-P-Q6 could be considered as acceptable when using a one-factor model (Table 5). In DE and SI data, the RMSEA for HLS₁₉-COM-P-Q11 was above the recommended reference value. In countries applying different data collection modes, the goodness-of-fit indices were approximately the same, except for the SI HLS₁₉-COM-P-Q6 data, where data collected using CAPI had somewhat better fit than CAWI data. Comparing goodness of fit indices based on dichotomous versus polytomous data, the SRMR was either equivalent or lower when analyses were based on polytomous data, whereas the opposite was the case considering RMSEA.

Table 4. Overall fit for HLS₁₉-COM-P-Q11 (left) and HLS₁₉-COM-P-Q6 (right) to the partial credit parametrization of the unidimensional polytomous Rasch model.

	Q11				Q6											
	AT n = 2954	DE n = 2133	SI n = 1856	SI n = 1487	AT n = 2952	BE n = 1000	BG n = 402	BG n = 457	CZ n = 531	CZ n = 1066	DE n = 2133	DK n = 3600	FR n = 2003	HU n = 1186	SI n = 1855	SI n = 1487
mode	CATI	PAPI	CAPI	CAWI	CATI	CAWI	CAPI	CAWI	CATI	CAWI	PAPI	CAWI	CAWI	CATI	CAPI	CAWI
χ^2, p	81.5, <0.001	84.5, <0.001	94.2, <0.001	108.1, <0.001	33.2, 0.1	57.3, <0.001	62.9, <0.001	93.2, <0.001	84.4, <0.001	51.3, 0.001	34.6, 0.07	86.7, <0.001	44.5, 0.01	52.1, 0.001	45.8, 0.005	47.7, 0.003
Mean person location	2.57	1.38	2.55	2.73	2.39	2.20	1.34	1.58	2.13	1.54	1.21	1.97	1.85	1.88	2.36	2.47
Dimensionality, % (lower 95% CI proportion)	6.1 (5.3)	7.9 (7.0)	4.8 (3.8)	6.9 (5.8)	5.3 (4.5)	4.4 (3.0)	6.5 (4.4)	5.3 (3.3)	5.8 (4.0)	5.1 (3.8)	5.0 (4.1)	7.5 (6.7)	3.6 (2.6)	3.0 (1.7)	3.0 (2.0)	4.4 (3.3)

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CI = confidence interval; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; HU = Hungary; PAPI = paper-assisted personal interviews; SI = Slovenia. Chi-square (χ^2) is based on $n = 660$ and 360 for HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6, respectively [39].

Table 5. Fit indices for the one-factor model of the HLS₁₉-COM-P-Q11 (left) and HLS₁₉-COM-P-Q6 (right), for each country, divided by data collection mode. Analyses are based on both polytomous and dichotomous (marked in grey) data.

Fit-Indices	mode	Q11				Q6											
		AT n = 2766	DE n = 2064	SI n = 1781	SI n = 1471	AT n = 2827	BE n = 1000	BG n = 333	BG n = 394	CZ n = 504	CZ n = 1066	DE n = 2101	DK n = 3574	FR n = 2003	HU n = 1125	SI n = 1788	SI n = 1477
		CATI	PAPI	CAPI	CAWI	CATI	CAWI	CAPI	CAWI	CATI	CAWI	PAPI	CAWI	CAWI	CATI	CAPI	CAWI
SRMR	polytomous data	0.04	0.07	0.04	0.05	0.03	0.02	0.03	0.03	0.02	0.02	0.03	0.01	0.02	0.03	0.01	0.03
	dichotomous data	0.06	0.07	0.06	0.07	0.03	0.04	0.06	0.06	0.07	0.02	0.03	0.02	0.03	0.05	0.02	0.03
RMSEA	polytomous data	0.07	0.11	0.10	0.10	0.06	0.05	0.07	0.07	0.03	0.03	0.07	0.03	0.05	0.07	0.03	0.07
	dichotomous data	0.02	0.05	0.03	0.04	0.00	0.02	0.02	0.04	0.00	0.00	0.03	0.01	0.02	0.02	0.00	0.01
RMSEA; CI, lower	polytomous data	0.06	0.11	0.10	0.09	0.05	0.03	0.04	0.04	0.00	0.01	0.05	0.02	0.04	0.05	0.01	0.05
	dichotomous data	0.02	0.05	0.03	0.03	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00
RMSEA; CI, upper	polytomous data	0.07	0.12	0.11	0.11	0.07	0.07	0.11	0.10	0.06	0.05	0.08	0.04	0.06	0.08	0.04	0.08
	dichotomous data	0.03	0.06	0.04	0.04	0.02	0.05	0.07	0.07	0.05	0.03	0.04	0.02	0.04	0.05	0.01	0.03
RMSEA; p-value	polytomous data	0.00	0.00	0.00	0.00	0.06	0.52	0.12	0.18	0.79	0.91	0.02	1.00	0.52	0.06	1.00	0.02
	dichotomous data	1.00	0.16	1.00	1.00	1.00	0.98	0.80	0.71	0.95	1.00	1.00	1.00	0.98	1.00	1.00	1.00
CFI	polytomous data	1.00	0.98	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	dichotomous data	0.99	0.99	1.00	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
TLI	polytomous data	0.99	0.98	1.00	0.99	1.00	1.00	1.00	1.00	1.00	0.99	1.00	1.00	1.00	1.00	1.00	1.00
	dichotomous data	0.99	0.98	0.99	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
GFI	polytomous data	1.00	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	dichotomous data	1.00	0.99	1.00	0.99	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
AGFI	polytomous data	0.99	0.98	0.99	0.99	0.99	1.00	0.99	0.99	1.00	1.00	0.99	1.00	1.00	0.99	1.00	0.99
	dichotomous data	0.99	0.98	0.99	0.99	1.00	1.00	0.99	0.99	1.00	1.00	0.99	1.00	1.00	1.00	1.00	1.00

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CFI = comparative fit index; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; GFI = goodness-of-fit index; HU = Hungary; PAPI = paper-assisted personal interviews; RMSEA = root mean square error of approximation; SI = Slovenia. SRMR = standardized root mean square residual; TLI = Tucker–Lewis index.

3.3. Reliability

Both the HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6 obtained acceptable to high reliability indices (Table 6). Based on polytomous data, the PSI, Cronbach's alpha, and omega for the HLS₁₉-COM-P-Q11 varied from 0.86 (AT) to 0.89 (DE), from 0.91 (AT) to 0.94 (SI, CAPI and CAWI), and 0.92 (AT) to 0.95 (SI, CAPI and CAWI), respectively. PSI, Cronbach's alpha, and omega for the HLS₁₉-COM-P-Q6 based on polytomous data varied from 0.75 (AT) to 0.83 (CZ, CATI and CAWI, DK and FR), 0.84 (DE) to 0.90 (BE, DK, SI, CAPI) and from 0.85 (DE) to 0.90 (BE), respectively. A decrease in Cronbach's alpha and omega was observed when analyses were based on dichotomous data. Both the HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6 obtained acceptable values of AVE.

3.4. Fit at the Item Level

Using a sample of 990 from each country, the HLS₁₉-COM-P-Q11 item COM1 ("describe to your doctor your reasons for coming to the consultation") displayed significant misfit ($p < 0.001$) in all three countries but had acceptable infit and fit residual. This was also the case for DE (PAPI) and SI (CAPI and CAWI) when reducing the sample size to 660. The item misfit was not significant at Bonferroni 5% for AT considering a sample size of 660. In data from AT (CATI), items COM4 ("get enough time in the consultation with your doctor"; fit residual of 3.87 and infit of 1.21) and COM7 ("understand the words used by your doctor"; fit residual of 3.21 and infit of 1.18) tend to under-discriminate (Table S3: Item fit statistics for HLS₁₉-COM-P-Q11 for each country, [39]). The other items displayed acceptable fit.

Most HLS₁₉-COM-P-Q11 items worked invariantly across different levels of person factors. However, in DE data, item COM7 ("understand the words used by your doctor") displayed significant DIF for education, where those having maximum upper secondary school as the highest completed education (ISCED 0 to 3) scored significantly lower than those having higher education despite the same location on the latent trait. This was also evident in SI CAWI data. In addition, item COM7 ("understand the words used by your doctor") did also display significant DIF for paying bills in SI CAWI data. In SI CAWI data, item COM6 ("get the information you need from your doctor") displayed DIF for age and education. However, the DIF was not evident when reducing the sample size to 660. The same was the case for age in item COM4 ("get enough time in the consultation with your doctor") in AT data. In SI CAPI data, item COM10 ("recall the information you get from your doctor") displayed DIF for age depending on how the variable was categorized (Table S3: Item fit statistics for HLS₁₉-COM-P-Q11 for each country, [39]). None of the items displayed DIF when it comes to self-reported health.

Response dependency was observed between items COM1 ("describe to your doctor your reasons for coming to the consultation") and COM3 ("explain your health concerns to your doctor") ($r = 0.35$) in the DE data (not reported in the Table). The response categories worked well for all items in all countries. Applying the HLS₁₉-COM-P-Q11, item COM1 ("describe to your doctor your reasons for coming to the consultation") was the easiest to endorse in all countries, whereas items COM4 ("get enough time in the consultation with your doctor"), COM7 ("understand the words used by your doctor"), COM5 ("express your personal views and preferences to your doctor") and COM9 ("be involved in decisions about your health in dialogue with your doctor") were the hardest in AT, DE, SI CAWI and Slovenian CAPI data, respectively (Table S3: Item fit statistics for HLS₁₉-COM-P-Q11 for each country, [39]).

Table 6. Reliability indices for the HLS₁₉-COM-P-Q11 (left) and HLS₁₉-COM-P-Q6 (right), for each country, divided by data collection mode. Calculation of Person separation index is based on polytomous data, whereas Cronbach’s alpha, omega and average variance extracted are displayed for both polytomous and dichotomous (marked in grey) data.

		Q11				Q6											
		AT CATI	DE PAPI	SI CAPI	SI CAWI	AT CATI	BE CAWI	BG CAPI	BG CAWI	CZ CATI	CZ CAWI	DE PAPI	DK CAWI	FR CAWI	HU CATI	SI CAPI	SI CAWI
Person separation index	polytomous data ⁱ	0.86	0.89	0.88	0.88	0.75	0.82	0.80	0.82	0.83	0.83	0.81	0.83	0.83	0.77	0.78	0.79
	dichotomous data ⁱⁱ	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cronbach’s alpha	polytomous data	0.91	0.90	0.94	0.94	0.86	0.90	0.87	0.88	0.87	0.88	0.84	0.90	0.89	0.87	0.90	0.89
	dichotomous data	0.78	0.84	0.87	0.86	0.68	0.80	0.81	0.80	0.70	0.81	0.74	0.78	0.80	0.78	0.80	0.79
Omega	polytomous data	0.92	0.93	0.95	0.95	0.86	0.90	0.87	0.89	0.87	0.88	0.85	0.89	0.89	0.87	0.89	0.89
	dichotomous data	0.80	0.86	0.91	0.90	0.71	0.81	0.84	0.83	0.74	0.82	0.76	0.80	0.81	0.81	0.82	0.81
Average variance extracted	polytomous data	0.63	0.58	0.74	0.71	0.64	0.71	0.67	0.67	0.67	0.66	0.57	0.71	0.70	0.67	0.73	0.70
	dichotomous data	0.55	0.57	0.73	0.68	0.57	0.67	0.72	0.69	0.62	0.67	0.56	0.67	0.67	0.68	0.73	0.70

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; HU = Hungary; PAPI = paper-assisted personal interviews; SI = Slovenia. ⁱ [39]; ⁱⁱ Due to low power, it could not be calculated.

For the HLS₁₉-COM-P-Q6, most items worked well in most countries. However, item COM4 (“get enough time in the consultation with your doctor”) under-discriminated in BG (CAPI data: fit residual of 3.02 and infit of 1.36, CAWI data: fit residual of 2.01 and infit of 1.27) and DK data (fit residual of 7.21 and infit of 1.36), while item COM10 (“recall the information you get from your doctor”) tend to under-discriminate in BE (fit residual of 3.02 and infit of 1.29), CZ (CATI data: fit residual of 1.98 and infit of 1.39, CAWI data: fit residual of 3.48 and infit of 1.34), DK (fit residual of 4.87 and infit of 1.37) and HU (fit residual of 1.58 and infit of 1.33) data (Table S4: Item fit statistics for HLS₁₉-COM-P-Q6 for each country, [39]). The item COM4 (“get enough time in the consultation with your doctor”) also displayed significant DIF across age categories (depending on categorization), level of education and employment status in BG CAPI data. In addition, the item displayed DIF for self-perceived social level in society and self-reported general health status, but this was not significant when reducing the sample size to 360 (Table S4: Item fit statistics for HLS₁₉-COM-P-Q6 for each country, [39]). Significant uniform and nonuniform DIF across age categories (depending on categorization) was also observed for item COM3 (“explain your health concerns to your doctor”) in BG CAWI data. For the other countries, no significant DIF was observed when considering a sample size of 360.

No response dependency nor unordered response categories were observed for HLS₁₉-COM-P-Q6. In most countries, item COM3 (“explain your health concerns to your doctor”) was the easiest to endorse (in BE data and SI CAWI data, COM8 (“ask your doctor questions in the consultation”) was the easiest and, in HU data, COM10 (“recall the information you get from your doctor”) was the easiest), whereas item COM4 (“get enough time in the consultation with your doctor”) was the hardest in most countries (except for CZ CATI data, CZ CAWI data, FR and SI CAPI data, where COM10 (“recall the information you get from your doctor”), COM9 (“be involved in decisions about your health in dialogue with your doctor”), COM5 (“express your personal views and preferences to your doctor”) and COM9 (“be involved in decisions about your health in dialogue with your doctor”) were the hardest to endorse, respectively; Table S4: Item fit statistics for HLS₁₉-COM-P-Q6 for each country).

3.5. Invariance across Modes and Countries

In SI data, the HLS₁₉-COM-P-Q11 items COM4 (“get enough time in the consultation with your doctor”), COM5 (“express your personal views and preferences to your doctor”; CAPI > CAWI) and COM7 (“understand the words used by your doctor”; CAPI < CAWI) displayed DIF across mode ($n = 990$). When the sample size was reduced to $n = 660$, DIF was only evident in item COM5 (F-ratio: 12.75, $p < 0.001$; Figure 4). Item COM5 also displayed DIF across mode in data from the SI six-items version, but this was not significant when the sample size was reduced to 540. Using approximately equal sample sizes from CZ CATI and CAWI HLS₁₉-COM-P-Q6 data, item COM10 (“recall the information you get from your doctor”) displayed DIF across modes. Item COM3 (“explain your health concerns to your doctor”) did display DIF across mode in BG data. These DIFs were not significant at Bonferroni adjusted 5% level when applying a sample size of 540.

Using equal sample sizes ($n = 500$) from countries applying CATI (AT, CZ, and HU), items COM3 (“explain your health concerns to your doctor”), COM4 (“get enough time in the consultation with your doctor”) and COM10 (“recall the information you get from your doctor”) displayed DIF across the countries. The same items displayed DIF when drawing a random sample of 400 for each country applying CAPI/PAPI (BG, DE, and SI). Using random samples of 500 from countries applying CAWI (BE, BG, CZ, DK, FR and SI), items COM3 (“explain your health concerns to your doctor”), COM4 (“get enough time in the consultation with your doctor”), COM9 (“be involved in decisions about your health in dialogue with your doctor”) and COM10 (“recall the information you get from your doctor”) displayed DIF. The mean person location for the random samples of HLS₁₉-COM-P-Q6 data collected using CATI, CAPI/PAPI and CAWI were 2.08, 1.60 and 1.89, respectively.




Figure 4. Graphical comparison between means of CAPI and CAWI in Slovenian data for item COM5 (“express your personal views and preferences to your doctor”).

3.6. Convergent and Discriminant Validity

The long and the short version of the instrument were, as expected, highly correlated; r varied from 0.97 (AT) to 0.98 (SI).

The scores obtained from HLS₁₉-COM-P-Q11 and the HLS₁₉-COM-P-Q6 were moderately to highly correlated with scores of GEN-HL (measured using HLS₁₉-Q12) and HL-NAV (measured using HLS₁₉-NAV) when analyses were based on polytomous data (Table 7). When conducting analyses based on dichotomous data, the correlation between COM-HL and related HL scores could be considered as small to large (Table 7). Lower correlation coefficients were observed in all countries when analyses were based on dichotomous data compared to polytomous.

Applying the combined PCA of residuals and paired sample t -test procedure, the HLS₁₉-Q12 (GEN-HL) and HLS₁₉-COM-P-Q11/Q6 item residuals loaded on separate components in all countries. The same was the case for the HLS₁₉-COM-P-Q11/Q6 and HLS₁₉-NAV items. The proportions of significant t -tests varied from 21.8% (DE) to 27.7% (SI, CAWI) for HLS₁₉-Q12 and HLS₁₉-COM-P-Q11, and between 15.1% (CZ, CATI) and 21.9% (BE) for HLS₁₉-Q12 and HLS₁₉-COM-P-Q6. The proportion of significant t -tests between HLS₁₉-COM-P-Q11 and HLS₁₉-NAV varied between 27.4% (DE) and 36.9% (SI, CAWI), and for HLS₁₉-COM-Q6 and HLS₁₉-NAV between 18.5% (AT) and 33.2% (BE).

3.7. Distribution of COM-HL Score

The score based on dichotomous items shows a left-skewed distribution with a clear ceiling effect in all countries, for both the long and short versions, regardless of the survey method (see Figure S1a: Distribution of HLS₁₉-COM-P-Q11 dichotomous score by country and survey mode; and Figure S1b: Distribution of HLS₁₉-COM-P-Q6 dichotomous score by country and survey mode). The score based on polytomous items is rather normally distributed in most countries, both for the long and short version, although, in some countries, the positive extreme values are disproportionately represented in the distribution (Figure S2a: Distribution of HLS₁₉-COM-P-Q11 polytomous score by country and survey mode; and Figure S2b: Distribution of HLS₁₉-COM-P-Q6 polytomous score by country and survey mode).

Table 7. Correlation between COM-HL scores (based on HLS₁₉-COM-P-Q11 to the left and HLS₁₉-COM-P-Q6 to the right) and general (GEN-HL) and navigational (HL-NAV) health literacy scores, based on polytomous and dichotomous (marked in grey) data. Results are divided by country and data collection mode.

		Q11				Q6											
		AT n = 2954	DE n = 2133	SI n = 1856	SI n = 1487	AT n = 2952	BE n = 1000	BG n = 402	BG n = 457	CZ n = 531	CZ n = 1066	DE n = 2133	DK n = 3600	FR n = 2003	HU n = 1186	SI n = 1855	SI n = 1487
mode		CATI	PAPI	CAPI	CAWI	CATI	CAWI	CAPI	CAWI	CATI	CAWI	PAPI	CAWI	CAWI	CATI	CAPI	CAWI
GEN-HL	polytomous data	0.54	0.59	0.59	0.53	0.52	0.35	0.56	0.65	0.49	0.49	0.56	0.55	0.60	0.47	0.58	0.51
	dichotomous data	0.37	0.54	0.51	0.44	0.34	0.27	0.46	0.53	0.39	0.45	0.50	0.47	0.52	0.36	0.49	0.41
HL-NAV	polytomous data	0.57	0.55	0.55	0.53	0.56	0.44	-	-	0.44	0.49	0.54	-	0.51	-	0.54	0.52
	dichotomous data	0.49	0.48	0.48	0.42	0.46	0.36	-	-	0.38	0.45	0.45	-	0.44	-	0.48	0.41

AT = Austria; BE = Belgium; BG = Bulgaria; CAPI = computer-assisted personal interviews; CATI = computer-assisted telephone interviews; CAWI = computer-assisted web interviews; CZ = Czech Republic; DE = Germany; DK = Denmark; FR = France; Gen-HL = general health literacy; HL-NAV = navigational health literacy; HU = Hungary; PAPI = paper-assisted personal interviews; SI = Slovenia. PAPI: paper-assisted personal interviews. BG, DK, and HU did not measure HL-NAV.

4. Discussion

Based on our theoretical framework that integrates the idea of COM-HL of Nutbeam [22], the basic competencies of information processing according to the HL framework of the HLS Consortium [3,25,26] and the main communicative tasks of the C-CG framework [32] we succeeded in developing a brief international instrument with acceptable psychometric properties and strong reliability for measuring COM-HL in patient–physician interaction.

4.1. Construct Validity and Reliability

The HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6 data display acceptable fit to the unidimensional Rasch model (considering a reduced sample size) and acceptable goodness-of-fit indices in CFA. Both HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6 gave sufficient unidimensional data, implying that it could be statistically defensible to calculate a total score [66] of COM-HL based on these instruments. Sufficiently high reliability indices do also allow for drawing conclusions about COM-HL both at group and individual levels [40].

However, the targeting of both HLS₁₉-COM-P-Q11 and HLS₁₉-COM-P-Q6 could have been better. Overall, the items were quite easy to endorse, implying a ceiling effect. Mistargeting might bring decreased reliability, as the precision of the instrument becomes poorer [56]. Hence, the instrument could benefit from adding items that are harder to endorse. On the other hand, for identifying groups with difficulties in HL-COM, the instrument performs well.

Most items displayed acceptable fit to the Rasch model. However, item COM10 (“recall the information you get from your doctor”) under-discriminated in CZ, DK, and HU HLS₁₉-COM-P-Q6 data. Under-discriminating items tend also to measure something else that is not positively correlated with the latent trait [67], here, COM-HL. The item is about recalling health information, which might be dependent on other cognitive processes than HL. Hence, in future studies, one might consider replacing this item with item COM11 in the short version of the instrument. This item might also be more in line with the cognitive domain to “apply” health information in the conceptual model of Sørensen et al. [25]. In addition, item COM4 (“get enough time in the consultation with your doctor”) under-discriminated in BG (CAPI) and DK HLS₁₉-COM-P-Q6 data. In AT data, the fit residual was also somewhat elevated, but the infit could be deemed as acceptable. Experiences of having sufficient time in consultation with a physician may depend on other things in addition to COM-HL, such as the number and type of health issues that the patient would like to discuss and the patients’ age [68]. In our conceptual framework, there are no other items covering this dimension. However, item COM4 might be replaced by item COM6 (“get the information you need from your doctor”), which could also be an indicator for understanding and following the agenda. In the short version, there are no items covering the dimension opening the session and giving initial information. However, these communicative tasks might be somewhat overlapping, which was confirmed in DE data, as response dependency was observed between items COM1 and COM3.

Few items showed DIF across different levels of person factors and, where DIF was present, there was no consistent pattern across countries. This indicates that the instrument works quite invariantly. However, the HLS₁₉-COM-P-Q11 item COM7 (“understand words used by your doctor”) displayed DIF for education in DE and SI CAWI data. The source of DIF might be that patients with low education are less familiar with medical jargon than those with higher education. In BG HLS₁₉-COM-P-Q6 CAPI data, COM4 (“get enough time in the consultation with your doctor”) displayed DIF for several person factors. As mentioned above, there might be several reasons that some patients perceive a need for more time in consultation with physicians.

On one hand, the COM-HL score was moderate to highly correlated with GEN-HL and HL-NAV scores, indicating that the instruments measure something common and, consequently, ensure convergent validity. The scores are all based on related instruments, all intending to measure certain aspects of HL. On the other hand, the combined PCA

of residuals and *t*-test procedure show that they are measuring distinctive constructs (discriminant validity). Hence, we conclude that the instruments for measuring COM-HL, GEN-HL and HL-NAV are measuring different aspects but could be considered parts of the family of HL instruments. Content and face validity are also ensured by using the theory-based model and definition of communicative HL with physicians in healthcare for selecting and operationalizing the included indicators. Concurrent predictive validity should be further explored in future studies.

4.2. Using Dichotomous or Polytomous Scores

Due to a change in the labelling of the response categories from the HLS-EU [26] to the HLS₁₉ survey, the HLS₁₉ Consortium decided to use dichotomized scores when reporting on HL in the international report [3] to ease the comparison between the surveys. However, no items of the HLS₁₉-COM-P-Q11 or HLS₁₉-COM-P-Q6 displayed unordered response categories, indicating that the four-point response categories used in the HLS₁₉-survey worked well at least for the HLS₁₉-COM-P instruments. Conducting Rasch analyses based on dichotomous items did result in an increased number of extreme records. The Cronbach's alpha values based on polytomous data were also higher than those reported based on dichotomized data, which would also be expected as more response categories yield more scoring points. The correlation between COM-HL and other HL scores was also stronger when analyses were based on polytomous scores. This is in line with Jiao et al. [69] who also found that results based on polytomous scores have slightly higher measurement precision compared to results from dichotomous scoring. Dichotomized scoring might be easier to understand but yields a loss of information and a loss of power [70]. However, dichotomization might reduce the effect of outliers [71].

4.3. Data Collection Mode

Different data collection modes were applied across countries and, for some countries, also within the country. The advantage is that the instrument was evaluated for different modes. However, according to Bowling [72], using different modes might bring more response bias than within a single mode. For countries that have used multiple modes, we also found that the mean person location differed across modes. Especially in CZ data, the difference in mean person location estimates between CATI and CAWI data was significant. In CZ, there was a predominance of younger people who responded to the CAWI version, whereas the CATI responses were dominated by older people, as these were hard to reach in CAWI sampling. Hence, the CAWI and CATI samples in CZ are also incomparable due to including different age groups. This was the case also in SI data. Some items did also display DIF across different data collection modes, implying that people responding to questionnaires operationalized by different data collection modes might interpret the items differently. However, the DIF across modes was marginal.

Even though the results should be interpreted with caution, due to DIF across countries, the mean person location was, on average, highest for countries that collected data by CATI and lowest in countries that used CAPI/PAPI. Both might be affected by response bias, but CAPI/PAPI brings less cognitive burden and is usually the most preferred data collection mode for the respondents [72]. Braekman et al. [73] did also find that, in a health survey, responses collected via self-administered modes (web versus PAPI) were more comparable than responses collected via self- and interviewer-administrated modes (web versus CAPI). However, the authors also found that simple and factual questions (such as healthcare use) are less prone to mode differences when comparing self- and interviewer-administered modes. Future studies which intend to compare scores across countries and modes of data collection should take actions to minimize mode effects, such as providing instructions for different modes in order to provide the same perceived stimuli to respondents. To reduce DIF across languages or countries, the translated versions of the instrument should also be assessed to ensure that items are interpreted in the same way across countries.

4.4. How to Use the Instrument

The HLS₁₉-COM-P intends to measure COM-HL in general adult populations and comprise skills that are necessary to actively participate in an interaction with physicians within a healthcare setting. The COM-HL score is standardized in the range of 0 to 100. Scores are only computed for respondents who have answered at least 80% of the HLS₁₉-COM-P items. If less than 80% of the items contain valid responses, the score is set to “missing”. A higher score value signifies a higher level of COM-HL. The score should be interpreted in light of contextual factors related to the health system of the present country.

The instrument belongs to the HLS₁₉ Consortium. The use of the instrument is free, but any use of it needs a contractual agreement between the nonprofit applicant and the HLS₁₉ Consortium. Further information can be found here: <https://m-pohl.net/tools> (accessed on 1 June 2022).

4.5. Strengths and Limitations

The psychometric properties of the instrument were assessed in large country representative samples from nine countries and are assessed in different data collection modes. The development of the instrument relies on a theory-based definition and conceptual framework of COM-HL.

A limitation is that the instrument only measures patients’ COM-HL in interaction with physicians. However, other healthcare professionals, such as nurses, are also providing health communication in healthcare settings. Hence, a version of the instrument comprising COM-HL in interacting with nurses should also be piloted. As the instrument intends to measure COM-HL in interaction with physicians in a healthcare setting, the instrument should be further tested among patients in clinical settings, especially with relevant indicators for predictive validity of the instrument. The validity and reliability of the HLS₁₉-COM-P should also be further explored in people with chronic illnesses.

In most countries, data were collected during the COVID-19 pandemic, which could have had an impact on the responses, as face-to-face encounters in this period were restricted to some extent. However, in DE, data were collected before the pandemic, and the psychometric properties of the HLS₁₉-COM-P do not differ much from DE to the other countries.

As the analyses are based on self-reported data, and some also from interviews, there might be a risk of response bias, such as social desirability. There is also a risk of recall misclassification as the experiences of physician–patient interactions might vary because of diverse factors, such as time since the last interaction, frequency of interactions, individual dependence on healthcare, cognitive skills, etc. Selection bias might also have occurred.

5. Conclusions

The HLS₁₉-COM-P-Q11 and the short version HLS₁₉-COM-P-Q6 worked quite well in the nine countries and across different data collection modes, even though misfit was found in a few items in some countries. The scale was also well accepted in all countries, with few missing values. Hence, this instrument could be used for identifying COM-HL in populations, and results from this instrument can be used to give recommendations for policy, practice and for COM-HL interventions (e.g., communication training for physicians). To our understanding, this is the first instrument to measure COM-HL as a separate construct in the family of related HLS instruments. However, the HLS₁₉-COM-P should be further evaluated in clinical settings and should be adapted to measure COM-HL also in relation to other health professions.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph191811592/s1>, Figure S1a. Distribution of HLS₁₉-COM-P-Q11 dichotomous score by country and survey mode, Figure S1b. Distribution of HLS₁₉-COM-P-Q6 dichotomous score by country and survey mode, Figure S2a. Distribution of HLS₁₉-COM-P-Q11 polytomous score by country and survey mode, Figure S2b. Distribution of HLS₁₉-COM-P-Q6 polytomous score by country and survey mode, Table S1: Fit indices considered in confirmatory factor analysis, Table S2: Analyses and tests with reference values considered in Rasch analyses, Table S3: Item fit statistics for HLS₁₉-COM-P-Q11 for each country, Table S4: Item fit statistics for HLS₁₉-COM-P-Q6 for each country.

Author Contributions: Conceptualization, H.S.F., P.N., R.G., É.B., M.V., R.C., L.G., J.M., H.B., Z.K., D.S. and J.M.P.; formal analysis, T.L. and H.S.F.; writing—original draft preparation, H.S.F., P.N., R.G., É.B. and M.V.; writing—review and editing, all authors. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding, but the data collection was supported either by ministries of health, university, public health institutes or insurance funds in the different countries. AT: The Austrian Health Literacy Survey was commissioned and financed by the Austrian Federal Health Agency and the Federation of Austrian Social Insurance Institutions. BE: There was no support for data collection in Belgium from any organisation. BG: Medical University—Sofia, Faculty of Public Health and AGREEMENT No. D-125/24.06.2020, PROJECT No. 8418/21.11.2019, “GRANT-2020”, MU-Sofia. CZ: Data collection was funded by (all seven) Czech health insurance funds. DE: The German study was funded by the German Federal Ministry of Health, grant number: Kapitel 1504 Titel 54401, ZMV I 1-2518 004 (HLS-GER 2). DK: The distribution of the questionnaire was funded by Aalborg University. FR: The research was supported by the National Public Health agency (Santé Publique France, 21DPPA040-0) and by Ligue Contre le Cancer (LIGUE2019). HU: The research was supported by the Ministry of Human Capacities, Hungary (IV/956-4/2020/EKF). SI: The national survey of Health literacy in Slovenia took place within the framework of the project *Increasing health literacy in Slovenia—ZaPiS*, which is co-financed by the Republic of Slovenia in the amount of 20% of the value and the European Union from the European Social Fund in the amount of 80% of the value.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki. In each country, it has been ensured that ethical requirements have been met. For more information about ethical considerations, data protection, and informed consent by country, please see *International Report on the methodology, results and recommendations of the European Health Literacy Population Survey 2019–2021 (HLS₁₉) of M-POHL* [3].

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Information about data supporting reported results can be found on the M-POHL webpage, https://m-pohl.net/Design_Methods (accessed on 1 June 2022).

Acknowledgments: The authors would like to acknowledge The International Coordination Centre (ICC) of the HLS₁₉ Project, especially Christa Straßmayr for project coordination. We will also thank all national HLS₁₉ team members who contributed to the HLS₁₉ Project, and to the Bulgarian principal investigators and policy representatives for organizing the HLS₁₉ project in their country and providing data for our article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Berkman, N.D.; Sheridan, S.L.; Donahue, K.E.; Halpern, D.J.; Crotty, K. Low health literacy and health outcomes: An updated systematic review. *Ann. Intern. Med.* **2011**, *155*, 97. [CrossRef]
2. Levin-Zamir, D.; Baron-Epel, O.B.; Cohen, V.; Elhayany, A. The association of health literacy with health behavior, socioeconomic indicators, and self-assessed health from a national adult survey in Israel. *J. Health Commun.* **2016**, *21*, 61–68. [CrossRef] [PubMed]
3. The HLS₁₉ Consortium of the WHO Action Network M-POHL. *International Report on the Methodology, Results and recommendations of the European Health Literacy Population Survey 2019–2021 (HLS₁₉) of M-POHL*; Austrian National Public Health Institute: Vienna, Austria, 2021.
4. Vandenbosch, J.; Van den Broucke, S.; Vancorenland, S.; Avalosse, H.; Verniest, R.; Callens, M. Health literacy and the use of healthcare services in Belgium. *J. Epidemiol. Community Health* **2016**, *70*, 1032–1038. [CrossRef] [PubMed]

5. Sørensen, K.; Pelikan, J.M.; Rothlin, F.; Ganahl, K.; Slonska, Z.; Doyle, G.; Fullam, J.; Kondilis, B.; Agrafiotis, D.; Uiters, E.; et al. Health literacy in Europe: Comparative results of the European health literacy survey (HLS-EU). *Eur. J. Public Health* **2015**, *25*, 1053–1058. [CrossRef] [PubMed]
6. Schaeffer, D.; Berens, E.-M.; Gille, S.; Griese, L.; Klinger, J.; de Sombre, S.; Vogt, D.; Hurrelmann, K. *Gesundheitskompetenz der Bevölkerung in Deutschland vor und während der Corona Pandemie: Ergebnisse des HLS-GER 2*; Universität Bielefeld: Bielefeld, Germany, 2021. [CrossRef]
7. Clever, S.L.; Jin, L.; Levinson, W.; Meltzer, D.O. Does doctor–patient communication affect patient satisfaction with hospital care? Results of an analysis with a novel instrumental variable. *Health Serv. Res.* **2008**, *43*, 1505–1519. [CrossRef]
8. Langewitz, W.; Conen, D.; Nübling, M.; Weber, H. Kommunikation ist wesentlich-Defizite der Betreuung im Krankenhaus aus der Sicht von Patienten und Patientinnen. *PPmP-Psychother. Psychosom. Med. Psychol.* **2002**, *52*, 348–354. [CrossRef]
9. Stahl, K.; Nadj-Kittler, M. *Zentrale Faktoren der Patienten-und Mitarbeiterzufriedenheit*; Picker Institut Deutschland gGmbH: Hamburg, Germany, 2013.
10. Mead, N.; Bower, P. Patient-centred consultations and outcomes in primary care: A review of the literature. *Patient Educ. Couns.* **2002**, *48*, 51–61. [CrossRef]
11. Sator, M.; Nowak, P.; Menz, F. *Verbesserung der Gesprächsqualität in der Krankenversorgung–Praxismodelle und Entwicklungsinitiativen*; Gesundheit Österreich GmbH: Vienna, Austria, 2015.
12. Street, R.L.; Makoul, G.; Arora, N.K.; Epstein, R.M. How does communication heal? Pathways linking clinician–patient communication to health outcomes. *Patient Educ. Couns.* **2008**, *74*, 295–301. [CrossRef]
13. Chen, X.; Hay, J.L.; Waters, E.A.; Kiviniemi, M.T.; Biddle, C.; Schofield, E.; Li, Y.; Kaphingst, K.; Orom, H. Health literacy and use and trust in health information. *J. Health Commun.* **2018**, *23*, 724–734. [CrossRef]
14. Paasche-Orlow, M.K.; Wolf, M.S. The causal pathways linking health literacy to health outcomes. *Am. J. Health Behav.* **2007**, *31*, S19–S26. [CrossRef]
15. Van der Heide, I.; Poureslami, I.; Mitic, W.; Shum, J.; Rootman, I.; FitzGerald, J.M. Health literacy in chronic disease management: A matter of interaction. *J. Clin. Epidemiol.* **2018**, *102*, 134–138. [CrossRef] [PubMed]
16. Castro, C.M.; Wilson, C.; Wang, F.; Schillinger, D. Babel babble: Physicians’ use of unclarified medical jargon with patients. *Am. J. Health Behav.* **2007**, *31*, S85–S95. [CrossRef] [PubMed]
17. Schillinger, D.; Bindman, A.; Wang, F.; Stewart, A.; Piette, J. Functional health literacy and the quality of physician–patient communication among diabetes patients. *Patient Educ. Couns.* **2004**, *52*, 315–323. [CrossRef]
18. Katz, M.G.; Jacobson, T.A.; Veledar, E.; Kripalani, S. Patient literacy and question-asking behavior during the medical encounter: A mixed-methods analysis. *J. Gen. Intern. Med.* **2007**, *22*, 782–786. [CrossRef] [PubMed]
19. Smith, S.K.; Nutbeam, D.; McCaffery, K.J. Insights into the concept and measurement of health literacy from a study of shared decision-making in a low literacy population. *J. Health Psychol.* **2013**, *18*, 1011–1022. [CrossRef] [PubMed]
20. Griese, L.; Berens, E.-M.; Nowak, P.; Pelikan, J.M.; Schaeffer, D. Challenges in Navigating the Health Care System: Development of an Instrument Measuring Navigation Health Literacy. *Int. J. Environ. Res. Public Health* **2020**, *17*, 5731. [CrossRef]
21. Griese, L.; Schaeffer, D.; Berens, E.-M. Navigational health literacy among people with chronic illness. *Chronic Illn.* **2022**, 1–12. [CrossRef]
22. Nutbeam, D. Health literacy as a public health goal: A challenge for contemporary health education and communication strategies into the 21st century. *Health Promot. Int.* **2000**, *15*, 259–267. [CrossRef]
23. Nutbeam, D. The evolving concept of health literacy. *Soc. Sci. Med.* **2008**, *67*, 2072–2078. [CrossRef]
24. Nutbeam, D. Defining and measuring health literacy: What can we learn from literacy studies? *Int. J. Public Health* **2009**, *54*, 303–305. [CrossRef]
25. Sørensen, K.; Van den Broucke, S.; Fullam, J.; Doyle, G.; Pelikan, J.; Slonska, Z.; Brand, H. Health literacy and public health: A systematic review and integration of definitions and models. *BMC Public Health* **2012**, *12*, 80. [CrossRef] [PubMed]
26. HLS-EU Consortium. Comparative Report of Health Literacy in Eight EU Member States. *The European Health Literacy Survey HLS-EU (First Revised and Extended Version. Date 22 July 2014)*. 2012. Available online: <http://www.health-literacy.eu> (accessed on 17 June 2022).
27. Nowak, P.; Finbråten, H.S.; Biro, E.; Bøggild, H.; Charafeddine, R.; Mancini, J.; Griebler, R.; Griese, L.; Kucera, Z.; Link, T.; et al. Communicative Health Literacy with physicians in health care services. In *International Report on the Methodology, Results, and Recommendations of the European Health Literacy Population Survey 2019–2021 (HLS19) of M-POHL*; The HLS19 Consortium of the WHO Action Network M-POHL, Ed.; Austrian National Public Health Institute: Vienna, Austria, 2021; pp. 233–274.
28. Nouri, S.S.; Rudd, R.E. Health literacy in the “oral exchange”: An important element of patient–provider communication. *Patient Educ. Couns.* **2015**, *98*, 565–571. [CrossRef] [PubMed]
29. Ishikawa, H.; Takeuchi, T.; Yano, E. Measuring functional, communicative, and critical health literacy among diabetic patients. *Diabetes Care* **2008**, *31*, 874–879. [CrossRef] [PubMed]
30. Chinn, D.; McCarthy, C. All Aspects of Health Literacy Scale (AAHLS): Developing a tool to measure functional, communicative and critical health literacy in primary healthcare settings. *Patient Educ. Couns.* **2013**, *90*, 247–253. [CrossRef] [PubMed]
31. O’Hara, J.; Hawkins, M.; Batterham, R.; Dodson, S.; Osborne, R.H.; Beauchamp, A. Conceptualisation and development of the conversational health literacy assessment tool (CHAT). *BMC Health Serv. Res.* **2018**, *18*, 199. [CrossRef] [PubMed]
32. Silverman, J.; Kurtz, S.; Draper, J. *Skills for Communication with Patients*, 3rd ed.; Radcliffe Publishing Ltd.: London, UK, 2013.

33. Iversen, E.D.; Wolderslund, M.O.; Kofoed, P.-E.; Gulbrandsen, P.; Poulsen, H.; Cold, S.; Ammentorp, J. Codebook for rating clinical communication skills based on the Calgary-Cambridge Guide. *BMC Med. Educ.* **2020**, *20*, 140. [CrossRef]
34. Sørensen, K.; Van den Broucke, S.; Pelikan, J.M.; Fullam, J.; Doyle, G.; Slonska, Z.; Kondilis, B.; Stoffels, V.; Osborne, R.H.; Brand, H. Measuring health literacy in populations: Illuminating the design and development process of the European Health Literacy Survey Questionnaire (HLS-EU-Q). *BMC Public Health* **2013**, *13*, 948. [CrossRef]
35. Gebhardt, E. Latent Path Models within an IRT Framework. Ph.D. Thesis, University of Melbourne, Melbourne, Australia, 2016.
36. Petrillo, J.; Cano, S.J.; McLeod, L.D.; Coon, C.D. Using classical test theory, item response theory, and Rasch measurement theory to evaluate patient-reported outcome measures: A comparison of worked examples. *Value Health* **2015**, *18*, 25–34. [CrossRef]
37. Wilson, M.; Allen, D.D.; Li, J.C. Improving measurement in health education and health behavior research using item response modeling: Comparison with the classical test theory approach. *Health Educ. Res.* **2006**, *21* (Suppl. 1), i19–i32. [CrossRef]
38. Kreiner, S. The Rasch model for dichotomous items. In *Rasch Models in Health*; Christensen, K.B., Kreiner, S., Mesbah, M., Eds.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2013; pp. 5–25.
39. Guttersrud, Ø.; Le, C.; Pettersen, K.S.; Finbråten, H.S. Rasch Analyses of Data Collected in 17 Countries—A Technical Report to Support Decision-Making within the M-POHL Consortium, Oslo, Norway. 2021, p. 37. Available online: https://m-pohl.net/sites/m-pohl.net/files/inline-files/Guttersrud%20et%20al_Rasch%20analyses%20of%20data%20collected%20in%2017%20countries_2021_0.pdf (accessed on 1 July 2022).
40. Frisbie, D.A. Reliability of Scores From Teacher-Made Tests. *Educ. Meas. Issues Pract.* **1988**, *7*, 25–35. [CrossRef]
41. Green, S.B.; Yang, Y. Reliability of summed item scores using structural equation modeling: An alternative to coefficient alpha. *Psychometrika* **2009**, *74*, 155–167. [CrossRef]
42. Ahmad, S.; Zulkurnain, N.N.A.; Khairushalimi, F.I. Assessing the validity and reliability of a measurement model in Structural Equation Modeling (SEM). *Br. J. Math. Comput. Sci.* **2016**, *15*, 1–8. [CrossRef] [PubMed]
43. Masters, G.N. A Rasch model for partial credit scoring. *Psychometrika* **1982**, *47*, 149–174. [CrossRef]
44. Rasch, G. *Probabilistic Models for Some Intelligence and Attainment Tests*, Expanded ed.; University of Chicago Press: Chicago, IL, USA, 1960.
45. Tennant, A.; Conaghan, P.G. The Rasch measurement model in rheumatology: What is it and why use it? When should it be applied, and what should one look for in a Rasch paper? *Arthritis Care Res.* **2007**, *57*, 1358–1362. [CrossRef]
46. Hagell, P. Testing rating scale unidimensionality using the principal component analysis (PCA)/*t*-test protocol with the Rasch model: The primacy of theory over statistics. *Open J. Stat.* **2014**, *4*, 456–465. [CrossRef]
47. Horton, M.; Marais, I.; Christensen, K.B. Dimensionality. In *Rasch Models in Health*; Christensen, K.B., Kreiner, S., Mesbah, M., Eds.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2013; pp. 137–157.
48. Smith, E.V. Detecting and evaluating the impact of multidimensionality using item fit statistics and principal component analysis of residuals. *J. Appl. Meas.* **2002**, *3*, 205–231.
49. Beaujean, A.A. *Latent Variable Modeling Using R: A Step-by-Step Guide*; Routledge: New York, NY, USA, 2014.
50. Kline, R.B. *Principles and Practice of Structural Equation Modeling*; Guilford Publications: New York, NY, USA, 2015.
51. Rosseel, Y. lavaan: An R package for structural equation modeling. *J. Stat. Softw.* **2012**, *48*, 1–36. [CrossRef]
52. Schumacker, R.E.; Lomax, R.G. *A beginner's Guide to Structural Equation Modeling*, 3rd ed.; Routledge: New York, NY, USA, 2010.
53. Hu, L.T.; Bentler, P.M. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct. Equ. Modeling Multidiscip. J.* **1999**, *6*, 1–55. [CrossRef]
54. Wright, B.D.; Linacre, J.M. Reasonable mean-square fit values. *Rasch Meas. Trans.* **1994**, *8*, 370.
55. Marais, I. Local Dependence. In *Rasch Models in Health*; Christensen, K.B., Kreiner, S., Mesbah, M., Eds.; John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2013; pp. 111–130.
56. Hagquist, C.; Bruce, M.; Gustavsson, J.P. Using the Rasch model in nursing research: An introduction and illustrative example. *Int. J. Nurs. Stud.* **2009**, *46*, 380–393. [CrossRef]
57. Hagquist, C. Explaining differential item functioning focusing on the crucial role of external information—An example from the measurement of adolescent mental health. *BMC Med. Res. Methodol.* **2019**, *19*, 185. [CrossRef] [PubMed]
58. Andrich, D.; Marais, I. *A Course in Rasch Measurement Theory: Measuring in the Educational, Social and Health Sciences*; Springer: Singapore, 2019.
59. Lantz, B. The large sample size fallacy. *Scand. J. Caring Sci* **2013**, *27*, 487–492. [CrossRef] [PubMed]
60. Cohen, L.; Manion, L.; Morrison, K. *Research Methods in Education*; Taylor & Francis Group: Oxford, UK, 2018.
61. Andrich, D.; Sheridan, B. *RUMM2030Plus*; Rumm Laboratory Pty Ltd.: Duncraig, Australia, 2019.
62. Adams, R.J.; Wu, M.L.; Cloney, D.; Wilson, M.R. *ACER ConQuest: Generalised Item Response Modelling Software [Computer software], Version 5*; Australian Council for Educational Research: Camberwell, Australia, 2020.
63. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2013.
64. Adler, N.E.; Epel, E.S.; Castellazzo, G.; Ickovics, J.R. Relationship of subjective and objective social status with psychological and physiological functioning: Preliminary data in healthy, White women. *Health Psychol.* **2000**, *19*, 586. [CrossRef]
65. Lampert, T.; Hoebel, J.; Kuntz, B.; Müters, S.; Kroll, L.E. *Messung des Sozioökonomischen Status und des Subjektiven Sozialen Status in KiGGs Welle 2*; Robert Koch-Institut: Berlin, Germany, 2018.
66. Andrich, D. *Rasch Models for Measurement*; SAGE: Newsbury Park, CA, USA, 1988; Volume 68.

67. Masters, G.N. Item discrimination: When more is worse. *J. Educ. Meas.* **1988**, *25*, 15–29. [[CrossRef](#)]
68. Deveugele, M.; Derese, A.; van den Brink-Muinen, A.; Bensing, J.; De Maeseneer, J. Consultation length in general practice: Cross sectional study in six European countries. *BMJ* **2002**, *325*, 472. [[CrossRef](#)] [[PubMed](#)]
69. Jiao, H.; Liu, J.; Haynie, K.; Woo, A.; Gorham, J. Comparison between dichotomous and polytomous scoring of innovative items in a large-scale computerized adaptive test. *Educ. Psychol. Meas.* **2012**, *72*, 493–509. [[CrossRef](#)]
70. Fedorov, V.; Mannino, F.; Zhang, R. Consequences of dichotomization. *Pharm. Stat. J. Appl. Stat. Pharm. Ind.* **2009**, *8*, 50–61. [[CrossRef](#)]
71. DeCoster, J.; Iselin, A.-M.R.; Gallucci, M. A conceptual and empirical examination of justifications for dichotomization. *Psychol. Methods* **2009**, *14*, 349. [[CrossRef](#)]
72. Bowling, A. Mode of questionnaire administration can have serious effects on data quality. *J. Public Health* **2005**, *27*, 281–291. [[CrossRef](#)]
73. Braekman, E.; Charafeddine, R.; Demarest, S.; Driekens, S.; Berete, F.; Gisle, L.; Van der Heyden, J.; Van Hal, G. Comparing web-based versus face-to-face and paper-and-pencil questionnaire data collected through two Belgian health surveys. *Int. J. Public Health* **2020**, *65*, 5–16. [[CrossRef](#)]