



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Characterization of the Structural and Mechanical Changes of the Biceps Brachii and Gastrocnemius Muscles in the Subacute and Chronic Stage after Stroke

García-Bernal, María Isabel; González-García, Paula; Madeleine, Pascal; Casuso-Holgado, María Jesús; Heredia-Rizo, Alberto Marcos

Published in:
International Journal of Environmental Research and Public Health

DOI (link to publication from Publisher):
[10.3390/ijerph20021405](https://doi.org/10.3390/ijerph20021405)

Creative Commons License
CC BY 4.0

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
García-Bernal, M. I., González-García, P., Madeleine, P., Casuso-Holgado, M. J., & Heredia-Rizo, A. M. (2023). Characterization of the Structural and Mechanical Changes of the Biceps Brachii and Gastrocnemius Muscles in the Subacute and Chronic Stage after Stroke. *International Journal of Environmental Research and Public Health*, 20(2), Article 1405. <https://doi.org/10.3390/ijerph20021405>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Article

Characterization of the Structural and Mechanical Changes of the Biceps Brachii and Gastrocnemius Muscles in the Subacute and Chronic Stage after Stroke

María Isabel García-Bernal ^{1,2}, Paula González-García ^{1,2,*}, Pascal Madeleine ³,
María Jesús Casuso-Holgado ^{1,4} and Alberto Marcos Heredia-Rizo ^{1,4}

¹ Departamento de Fisioterapia, Facultad de Enfermería, Fisioterapia y Podología, Universidad de Sevilla, 41009 Sevilla, Spain

² Instituto de Biomedicina de Sevilla (IBIS), 41013 Sevilla, Spain

³ Sport Sciences—Performance and Technology, Department of Health Science and Technology, Aalborg University, 9220 Aalborg, Denmark

⁴ Uncertainty, Mindfulness, Self, Spirituality (UMSS) Research Group, Universidad de Sevilla, 41009 Sevilla, Spain

* Correspondence: pgonzalez@us.es; Tel.: +34-954486528

Abstract: The objective of this study was to characterize the changes of muscle tone, stiffness, and thickness of upper and lower limb muscles in stroke survivors. Forty patients with subacute or chronic stroke and 31 controls were included and measured using myotonometry (MyotonPRO), with multiple site assessments at muscle belly (MB) and musculotendinous (MT) locations of the biceps brachii and gastrocnemius muscles. Muscle thickness (ultrasonography) was obtained for each muscle. Upper and lower limb motor performance was evaluated with the Fugl–Meyer Assessment for Upper Extremity and the Functional Ambulance Category. Overall, muscle tone and stiffness were significantly higher at MT than at MB sites. Among stroke patients, differences between the paretic and nonparetic limb were found for the biceps brachii, with lower muscle tone, stiffness, and thickness of the paretic side (all, $p < 0.05$). There were weak to moderate correlations between mechanical (myotonometry) and structural (ultrasound) muscular changes, regardless of the post-stroke stage. This suggests that myotonometry and ultrasonography assess similar, although different, constructs and can be combined in the clinical setting. Their discriminative ability between the paretic and nonparetic sides and between participants with and without stroke differs depending on the muscle, the functional level, and the stroke stage.

Keywords: stroke; spasticity; muscle; myotonometry; outcomes assessment; ultrasound



Citation: García-Bernal, M.I.; González-García, P.; Madeleine, P.; Casuso-Holgado, M.J.; Heredia-Rizo, A.M. Characterization of the Structural and Mechanical Changes of the Biceps Brachii and Gastrocnemius Muscles in the Subacute and Chronic Stage after Stroke. *Int. J. Environ. Res. Public Health* **2023**, *20*, 1405. <https://doi.org/10.3390/ijerph20021405>

Academic Editor: Guglielmo Duranti

Received: 21 November 2022

Revised: 5 January 2023

Accepted: 10 January 2023

Published: 12 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Post-stroke spasticity (PSS) is a major sensorimotor disorder that can be present from the early stages after stroke [1] and frequently involves the flexor muscles of the upper limb and the extensor muscles of the lower limb [2]. Structural changes in muscle and tendon appear as a result of PSS, leading to impaired motor control, disuse [3], worse recovery, and disability [4]. However, recent systematic reviews suggest that there is little, and conflicting, evidence about the specific structural and mechanical adaptations that occur in the upper and lower limb muscles after stroke [5,6]. Understanding this could help to guide clinical decisions aimed at targeting focal spasticity of specific muscles and locations within the muscles (i.e., botulin toxin) [7] or at restoring muscle function during physical rehabilitation [8].

Monitoring structural and mechanical soft-tissue properties in PSS is a complex task. Clinical measures are indirect, suboptimal, and show limited reliability and reproducibility [9]. Alternatively, direct measurement tools, such as ultrasound elastography and

myotonometry, have emerged as promising procedures to provide quantitative useful information in patient populations [10–12]. Current literature suggests a non-uniform distribution of the muscle mechanical properties after stroke, with different adaptations in the upper and lower limbs [8,13]. However, it has been recommended to measure several spots within the muscle and tendon to better characterize PSS [13]. Topographical mapping, based on multiple site assessments within an area of interest, has been used extensively for surface electromyography (SEMG) [14] and quantitative sensory testing [15] and, more recently, to report the spatial distribution of muscle stiffness in people with chronic pain [16].

The objective of this study was to characterize the topographical maps of tone and stiffness of the biceps brachii and gastrocnemius muscles in stroke patients, compared with healthy controls, and observe the possible association between structural (muscle thickness) and mechanical (muscle tone and stiffness) changes. We hypothesized that: (1) among persons with stroke, musculotendinous sites would show higher tone and stiffness than muscle belly sites, structural and mechanical changes would be correlated, and muscle adaptations would differ between the upper and the lower limb; (2) myotonometry and ultrasound measurements would be able to discriminate between participants with and without stroke.

2. Materials and Methods

2.1. Design

This cross-sectional study was approved by the Institutional Review Board (Junta de Andalucía Ethical Committee for Biomedical Research, code number CI 1222-N-16) and followed the STROBE guidelines for observational studies. All participants provided written informed consent prior to data collection.

2.2. Participants

This is a secondary analysis. The sample was the same as in a previous study investigating the influence of muscle position during assessment (relaxed vs. stretched) on the mechanical properties of spastic muscles [13]. Stroke participants were recruited from public and private settings and assigned to either group according to time after stroke. Twenty stroke patients in the subacute stage (1.5 to 9 months after the event) and 20 patients in the chronic stage (more than 9 months after the event) [17] participated in the study. A total of 31 healthy participants, from the same population-based cohort, were included in the control group. Patients with a first-ever stroke were selected if they had a score ≥ 1 in the Modified Ashworth Scale for the biceps brachii and gastrocnemius muscles [18] and showed a normal cognitive status (score ≥ 24 in the Mini-Mental State Examination) [19]. The exclusion criteria were: severe upper or lower limb injury, a diagnosed behavioral disorder, recent changes in medication for PSS, having received botulinum toxin injections within 3 months, and any other neurological disease. Stroke participants were undergoing regular care, mostly involving physical therapy and medication intake.

2.3. Outcome Measures

Tone and stiffness of the biceps brachii and gastrocnemius muscles were measured with the MyotonPRO (Myoton AS, Tallinn, Estonia), using several site assessments. The MyotonPRO is a noninvasive hand-held device that applies consecutive short mechanical impulses on the surface of the skin to induce a dynamic soft-tissue response (Figure S1). Using an accelerometer, muscle tone (tension) is measured as the frequency of the signal and muscle stiffness is the resultant natural oscillation that characterizes the resistance to the external force [20]. For the biceps brachii, participants lay supine, with 45° elbow flexion and neutral forearm. First, we measured the distance d between the coracoid process of the scapula and the muscle insertion at the radial tuberosity. Then, using a wax pencil, a 13-point grid was drawn, with adjacent points separated longitudinally by $1/8$ of the d value and transversally by $1/6$ of d , except for point 13, which was separated by $1/12$ of

d. Points 1, 2, 3, 4 and 13 corresponded to musculotendinous (MT) sites, whereas points 5, 6, 7, 8, 9, 10, 11, and 12 were located at the muscle belly (MB). For the gastrocnemius, participants were prone, with the knee in 45° flexion. We calculated the distance d between the upper edge of the calcaneus and the lower edge of the gastrocnemius medialis, and e between the lower edge of the gastrocnemius medialis and the midpoint between the medial and lateral condyle of the femur. Points 1 to 8, corresponding to MB sites, were separated longitudinally by $1/5$ of e and transversally by $1/11$ of e . Points 9 to 12 were located at MT sites, with a distance between adjacent points of $1/5$ of d . Assessment sites are described in detail in Figure 1. All measures were taken twice for each site bilaterally, and the average score was used for analysis [16]. The topographical maps were generated with the Matlab software 9.1 (The Mathworks, Natick, MA, USA). For that purpose, muscle tone and stiffness data from the MyotonPRO were used, applying an inverse distance weighted interpolation method.




Figure 1. Measurement sites for the biceps brachii (A) and gastrocnemius (B) muscles.

Resting measures of muscle thickness were obtained bilaterally at points 7 and 8 for the biceps brachii and at points 5 and 6 for the gastrocnemius muscle. We used B-mode ultrasound imaging (4.2 to 13 MHz) coupled with a multifrequency linear array transducer, model 12L-RS (GE Logiq V2, GE Health Care, Chicago, IL, USA). The average of two consecutive measures was taken for analysis. The muscle thickness was measured as the vertical distance between the superficial and deep aponeurosis for the gastrocnemius and between the adipose tissue–muscle interface and the muscle–bone interface for the biceps brachii [21].

Upper and lower limb motor performance was evaluated with the Fugl–Meyer Assessment for Upper Extremities (FMA-UE) [22] and the Functional Ambulation Category (FAC) [23], respectively. The same examiners, who remained blinded to participants' allocation group, collected all demographic and clinical measures.

2.4. Statistical Analysis

Sample size was estimated using the software G*Power (v. 3.1.9.2, Heinrich-Heine University, Düsseldorf, Germany). We considered an alpha level of 0.05, an 80% power, and a large effect size ($n^2 = 0.15$) for differences in muscle tone and stiffness between groups. This analysis revealed that at least 19 participants were required per study group.

The PASW Advanced Statistics (SPSS Inc., Chicago, IL, USA), version 26.0, was used for statistical processing. Normal distribution of the data was analyzed with the Shapiro–Wilk test. A three-way analysis of variance (ANOVA) was used to detect differences in tone and

stiffness of the biceps brachii and gastrocnemius, using sites (MT vs. MB) and sides (paretic vs. nonparetic) of testing as the within-participant factors, and group as the between-participant factor. Associations between muscle thickness and myotonometry were tested with the Pearson product-moment correlation coefficient analysis or the Spearman rank test. The level of significance was set to $p < 0.05$.

3. Results

A total of 71 adults (31 controls, 20 subacute stroke, and 20 chronic stroke) were included (Figure 2).




Figure 2. Flowchart diagram of the study participants.

The baseline clinical and demographic features of participants can be seen in Table 1. Upper limb motor performance was severely compromised among stroke survivors (overall score, 19 ± 20.8 points), whereas 15% of them (6/30) could be categorized as nonfunctional ambulators, according to the FAC (FAC = 0).

Table 1. Baseline clinical and demographic features of participants.

| | Subacute Stroke (n = 20) | Chronic Stroke (n = 20) | Control Group (n = 31) | p Value |
|--|-----------------------------|------------------------------|---------------------------|---------|
| Age (years) | 60.2 ± 9.7 | 61.45 ± 9.7 | 60.8 ± 10.6 | 0.926 |
| Sex, female, % (n) | 35% (7) | 35% (7) | 45.2% (14) | 0.689 |
| Time after stroke (weeks) | 17 (6–34) | 242.5 (58–1108) | N/A | <0.001 |
| Affected side, left, % (n) | 55% (11) | 75% (15) | N/A | 0.289 |
| Hand dominance, right, left, ambidextrous, % (n) | 100% (20) | 85% (17); 5% (1); 10% (2) | 80.6% (25); 19.4% (6) | 0.131 |
| Leg dominance, right, left, ambidextrous, % (n) | 95% (19); 5% (1); | 80% (16); 5% (1); 15% (3) | 83.9% (26); 16.1% (5) | 0.319 |

The topographical maps of tone and stiffness of the biceps brachii and gastrocnemius muscles are shown in Figures 3 and 4.




Figure 3. Muscle tone (Hz; **A**) and stiffness (N/m; **B**) maps based on average values of the assessed locations from the biceps brachii muscle in individuals with subacute and chronic stroke, and control participants.




Figure 4. Muscle tone (Hz; **A**) and stiffness (N/m; **B**) maps based on average values of the assessed locations from the gastrocnemius muscle in individuals with subacute and chronic stroke, and control participants.

Tables 2 and 3 include the mean ± SD values for tone and stiffness at the different sites, sides, and groups. Table 4 shows the data for muscle thickness.

Table 2. Mean ± SD biceps brachii stiffness (N/m) and tone (Hz) over assessed sites (points 1 to 13, MB and MT sites) in stroke patients and controls.

| Site | Measure | Control Participants (n = 31) | | Subacute Stroke (n = 20) | | Chronic Stroke (n = 20) | |
|-----------------------------|-----------|-------------------------------|-------------------|--------------------------|-------------------|-------------------------|-------------------|
| | | Dominant Side | Non-Dominant Side | Affected Side | Non-Affected Side | Affected Side | Non-Affected Side |
| Point 1 | Stiffness | 311.1 ± 82.7 | 343.9 ± 78.7 | 302.9 ± 101.9 | 355.5 ± 112.8 | 322.4 ± 94.3 | 434.1 ± 139.2 |
| | Tone | 17.2 ± 2.7 | 18.2 ± 2.5 | 16.7 ± 3.1 | 18.1 ± 3.4 | 17.3 ± 3.0 | 20.6 ± 3.9 |
| Point 2 | Stiffness | 359.5 ± 55.9 | 372.9 ± 58.1 | 357.7 ± 59.8 | 400.5 ± 97.1 | 388.9 ± 85.6 | 478.2 ± 111.0 |
| | Tone | 19.0 ± 2.4 | 19.3 ± 2.3 | 18.5 ± 2.2 | 19.5 ± 4.1 | 19.5 ± 3.5 | 22.9 ± 4.2 |
| Point 3 | Stiffness | 264.9 ± 49.8 | 283.4 ± 49.1 | 260.9 ± 48.0 | 293.4 ± 71.9 | 254.6 ± 46.4 | 315.7 ± 61.1 |
| | Tone | 15.1 ± 2.0 | 15.6 ± 1.9 | 15 ± 2.0 | 15.7 ± 2.5 | 14.5 ± 1.9 | 16.7 ± 2.2 |
| Point 4 | Stiffness | 335.7 ± 52.7 | 346.5 ± 43.2 | 335.7 ± 55.3 | 370.8 ± 69.8 | 332.5 ± 51.5 | 407.4 ± 85.4 |
| | Tone | 17.6 ± 2.2 | 18.0 ± 1.7 | 17.3 ± 2.6 | 17.9 ± 3.2 | 17.1 ± 2.6 | 19.8 ± 3.2 |
| Point 5 | Stiffness | 250.5 ± 47.7 | 256.9 ± 52.2 | 227.4 ± 41.3 | 267.3 ± 63.3 | 223.0 ± 47.2 | 265.6 ± 67.1 |
| | Tone | 14.1 ± 2.0 | 14.2 ± 1.9 | 13.5 ± 1.6 | 15.0 ± 2.5 | 13.2 ± 2.0 | 14.5 ± 2.6 |
| Point 6 | Stiffness | 319.1 ± 49.7 | 340.6 ± 57.6 | 339.7 ± 82.9 | 369.7 ± 89.3 | 318.5 ± 67.6 | 380.1 ± 96.0 |
| | Tone | 16.6 ± 2.2 | 17.4 ± 2.3 | 17.1 ± 3.2 | 17.8 ± 3.4 | 16.6 ± 2.9 | 18.6 ± 3.5 |
| Point 7 | Stiffness | 235.2 ± 26.4 | 240.2 ± 29.8 | 220.4 ± 35.6 | 242.9 ± 30.7 | 228.0 ± 29.7 | 238.9 ± 31.8 |
| | Tone | 13.4 ± 1.6 | 13.2 ± 1.4 | 13.0 ± 1.5 | 13.7 ± 1.5 | 13.5 ± 1.3 | 13.4 ± 1.3 |
| Point 8 | Stiffness | 307.2 ± 78.7 | 312.4 ± 67.5 | 297.8 ± 75.5 | 381.6 ± 131.3 | 315.3 ± 70.4 | 364.1 ± 101.2 |
| | Tone | 15.9 ± 2.9 | 16.1 ± 2.5 | 15.5 ± 2.5 | 17.8 ± 4.0 | 16.1 ± 2.2 | 17.7 ± 3.4 |
| Point 9 | Stiffness | 237.3 ± 23.9 | 235.3 ± 19.8 | 223.9 ± 31.4 | 245.4 ± 29.1 | 242.6 ± 31.9 | 245.3 ± 23.7 |
| | Tone | 13.3 ± 1.3 | 13.1 ± 1.0 | 12.9 ± 1.6 | 14.1 ± 1.9 | 13.7 ± 1.5 | 13.7 ± 1.2 |
| Point 10 | Stiffness | 262.1 ± 38.9 | 264.9 ± 37.1 | 260.1 ± 41.5 | 294.1 ± 47.4 | 267.9 ± 30.7 | 302.2 ± 44.1 |
| | Tone | 14.4 ± 1.8 | 14.4 ± 1.9 | 14.2 ± 1.6 | 15.2 ± 1.9 | 14.9 ± 1.5 | 15.9 ± 2.1 |
| Point 11 | Stiffness | 242.5 ± 20.8 | 246.9 ± 26.7 | 233.9 ± 26.7 | 251.8 ± 37.8 | 257.2 ± 32.6 | 263.3 ± 25.7 |
| | Tone | 13.7 ± 1.0 | 13.9 ± 1.2 | 13.6 ± 1.5 | 14.5 ± 1.8 | 14.6 ± 1.9 | 14.2 ± 1.6 |
| Point 12 | Stiffness | 247.9 ± 44.9 | 252.9 ± 37.8 | 243.3 ± 32.5 | 276.1 ± 43.1 | 248.2 ± 23.2 | 280.3 ± 37.3 |
| | Tone | 14.2 ± 1.8 | 14.0 ± 1.6 | 13.5 ± 1.2 | 14.8 ± 1.6 | 14.1 ± 1.4 | 14.9 ± 1.9 |
| Point 13 | Stiffness | 253.4 ± 39.7 | 258.8 ± 48.4 | 245.6 ± 33.2 | 251.9 ± 54.2 | 275.7 ± 75.2 | 259.9 ± 39.1 |
| | Tone | 14.3 ± 1.4 | 14.5 ± 1.9 | 14.3 ± 1.3 | 14.4 ± 2.5 | 15.1 ± 2.9 | 14.2 ± 1.7 |
| Muscle belly (MB) sites | Stiffness | 256.8 ± 27.1 | 262.2 ± 28.1 | 248.1 ± 36.3 | 277.7 ± 35.8 | 254.8 ± 27.6 | 280.9 ± 35.4 |
| | Tone | 14.3 ± 1.4 | 14.3 ± 1.3 | 13.9 ± 1.3 | 15.1 ± 1.5 | 14.4 ± 1.2 | 15.0 ± 1.5 |
| Musculotendinous (MT) sites | Stiffness | 295.3 ± 41.2 | 310.2 ± 36.0 | 288.7 ± 37.8 | 318.1 ± 58.1 | 302.4 ± 49.9 | 351.9 ± 50.7 |
| | Tone | 16.4 ± 1.7 | 16.8 ± 1.5 | 16.1 ± 1.5 | 16.7 ± 2.3 | 16.4 ± 2.1 | 18.1 ± 2.1 |

Table 3. Mean ± SD gastrocnemius muscle stiffness (N/m) and tone (Hz) over assessed sites (points 1 to 12, MB and MT sites) in stroke patients and controls.

| Site | Measure | Control Participants (n = 31) | | Subacute Stroke (n = 20) | | Chronic Stroke (n = 20) | |
|---------|-----------|-------------------------------|-------------------|--------------------------|-------------------|-------------------------|-------------------|
| | | Dominant Side | Non-Dominant Side | Affected Side | Non-Affected Side | Affected Side | Non-Affected Side |
| Point 1 | Stiffness | 255.4 ± 22.1 | 271.5 ± 37.6 | 253.9 ± 29.6 | 273.5 ± 47.4 | 283.4 ± 52.5 | 295.1 ± 65.8 |
| | Tone | 14.6 ± 1.1 | 15.3 ± 1.8 | 14.4 ± 1.5 | 15.2 ± 2.3 | 15.4 ± 2.2 | 15.8 ± 2.5 |
| Point 2 | Stiffness | 259.3 ± 28.1 | 267.2 ± 33.1 | 263.1 ± 39.2 | 273.3 ± 51.6 | 292.8 ± 43.3 | 291.5 ± 73.4 |
| | Tone | 14.9 ± 1.2 | 15.2 ± 1.4 | 14.5 ± 1.8 | 14.9 ± 2.4 | 15.8 ± 1.8 | 15.7 ± 2.8 |
| Point 3 | Stiffness | 277.5 ± 24.8 | 280.7 ± 23.2 | 265.0 ± 25.2 | 282.8 ± 40.5 | 298.1 ± 43.3 | 300.8 ± 59.4 |
| | Tone | 14.9 ± 1.4 | 15.0 ± 1.3 | 14.2 ± 1.4 | 15.0 ± 2.2 | 15.4 ± 2.5 | 15.9 ± 2.6 |
| Point 4 | Stiffness | 279.1 ± 19.0 | 283.4 ± 26.4 | 269.9 ± 19.6 | 281.4 ± 30.8 | 289.9 ± 40.7 | 290.7 ± 66.2 |
| | Tone | 14.7 ± 1.4 | 15.3 ± 1.6 | 14.0 ± 1.3 | 14.8 ± 1.8 | 15.6 ± 2.3 | 15.6 ± 2.9 |

Table 3. Cont.

| Site | Measure | Control Participants (n = 31) | | Subacute Stroke (n = 20) | | Chronic Stroke (n = 20) | |
|-----------------------------|-----------|-------------------------------|-------------------|--------------------------|-------------------|-------------------------|-------------------|
| | | Dominant Side | Non-Dominant Side | Affected Side | Non-Affected Side | Affected Side | Non-Affected Side |
| Point 5 | Stiffness | 288.8 ± 26.6 | 290.7 ± 28.1 | 278.7 ± 31.3 | 289.3 ± 29.7 | 317.4 ± 51.7 | 312.9 ± 58.4 |
| | Tone | 15.7 ± 1.4 | 15.9 ± 1.5 | 14.9 ± 1.8 | 15.7 ± 2.0 | 16.8 ± 2.8 | 17.1 ± 3.0 |
| Point 6 | Stiffness | 282.3 ± 18.7 | 281.6 ± 26.2 | 278.0 ± 26.7 | 284.3 ± 30.2 | 293.5 ± 40.7 | 295.7 ± 66.5 |
| | Tone | 14.9 ± 1.5 | 15.3 ± 1.7 | 14.4 ± 1.7 | 14.8 ± 2.0 | 15.8 ± 2.3 | 16.2 ± 2.6 |
| Point 7 | Stiffness | 306.8 ± 26.3 | 308.1 ± 25.4 | 290.4 ± 25.1 | 299.2 ± 31.7 | 334.0 ± 58.3 | 327.0 ± 66.2 |
| | Tone | 16.0 ± 1.4 | 16.2 ± 1.5 | 15.3 ± 1.5 | 15.7 ± 2.2 | 17.9 ± 3.1 | 17.7 ± 2.8 |
| Point 8 | Stiffness | 314.7 ± 29.2 | 310.7 ± 27.9 | 305.4 ± 26.8 | 308.8 ± 37.3 | 332.3 ± 47.1 | 329.4 ± 57.2 |
| | Tone | 16.2 ± 1.7 | 16.5 ± 1.6 | 15.7 ± 1.8 | 16.4 ± 2.5 | 17.5 ± 2.7 | 17.6 ± 3.2 |
| Point 9 | Stiffness | 416.4 ± 40.4 | 421.6 ± 40.9 | 380.6 ± 49.1 | 386.9 ± 44.5 | 428.9 ± 81.9 | 400.8 ± 71.4 |
| | Tone | 21.1 ± 2.2 | 21.1 ± 2.2 | 19.3 ± 2.3 | 19.7 ± 2.4 | 20.9 ± 3.2 | 20.8 ± 3.1 |
| Point 10 | Stiffness | 483.2 ± 58.6 | 482.5 ± 56.5 | 435.3 ± 59.8 | 440.2 ± 60.8 | 482.3 ± 89.5 | 445.7 ± 77.6 |
| | Tone | 23.9 ± 3.3 | 23.5 ± 2.6 | 21.8 ± 2.7 | 22.0 ± 2.5 | 23.4 ± 3.6 | 22.4 ± 3.5 |
| Point 11 | Stiffness | 562.1 ± 90.1 | 571.3 ± 98.3 | 521.7 ± 98.9 | 509.5 ± 66.6 | 572.1 ± 117.5 | 510.4 ± 98.2 |
| | Tone | 26.9 ± 3.9 | 26.3 ± 3.6 | 25.2 ± 3.9 | 24.4 ± 3.1 | 26.8 ± 4.5 | 24.0 ± 4.0 |
| Point 12 | Stiffness | 679.4 ± 91.1 | 683.8 ± 119.5 | 670.1 ± 164.1 | 663.8 ± 109.8 | 651.4 ± 121.5 | 663.6 ± 164.9 |
| | Tone | 29.3 ± 2.8 | 29.4 ± 3.8 | 29.2 ± 4.6 | 28.6 ± 3.5 | 28.4 ± 3.5 | 28.2 ± 5.0 |
| Muscle belly (MB) sites | Stiffness | 280.3 ± 13.7 | 284.2 ± 17.4 | 273.3 ± 19.8 | 284.0 ± 29.4 | 301.3 ± 35.5 | 302.3 ± 61.1 |
| | Tone | 15.1 ± 0.9 | 15.5 ± 1.1 | 14.6 ± 1.0 | 15.2 ± 1.7 | 16.0 ± 1.9 | 16.3 ± 2.5 |
| Musculotendinous (MT) sites | Stiffness | 515.9 ± 56.1 | 475.2 ± 55.3 | 476.4 ± 70.5 | 478.2 ± 55.4 | 518.1 ± 98.4 | 484.2 ± 84.2 |
| | Tone | 24.8 ± 2.4 | 23.1 ± 2.4 | 23.2 ± 2.8 | 23.1 ± 2.3 | 24.4 ± 3.4 | 23.4 ± 3.3 |

Table 4. Muscle thickness (cm) of the biceps brachii and gastrocnemius muscles in stroke survivors.

| Muscle | Site | Subacute Stroke (n = 20) | | Chronic Stroke (n = 20) | | Control Group (n = 31) | |
|----------------|---------|--------------------------|-------------------|-------------------------|-------------------|------------------------|-------------------|
| | | Affected Side | Non-Affected Side | Affected Side | Non-Affected Side | Dominant Side | Non-Dominant Side |
| Biceps brachii | Point 7 | 1.97 ± 0.52 | 2.27 ± 0.66 | 1.83 ± 0.39 | 2.33 ± 0.45 | 2.41 ± 0.51 | 2.37 ± 0.59 |
| | Point 8 | 1.41 ± 0.36 | 1.78 ± 0.59 | 1.24 ± 0.28 | 1.66 ± 0.51 | 1.75 ± 0.72 | 1.57 ± 0.48 |
| Gastrocnemius | Point 5 | 1.47 ± 0.42 | 1.60 ± 0.26 | 1.25 ± 0.51 | 1.45 ± 0.47 | 1.70 ± 0.41 | 1.75 ± 0.40 |
| | Point 6 | 1.34 ± 0.45 | 1.29 ± 0.46 | 1.12 ± 0.41 | 1.35 ± 0.44 | 1.41 ± 0.29 | 1.41 ± 0.28 |

3.1. Mechanical and Structural Muscular Adaptations among Stroke Survivors

For the biceps brachii, the ANOVA revealed a significant sites*sides interaction for muscle tone ($F = 1.937; p = 0.020; \eta^2 = 0.023$) and stiffness ($F = 2.728; p = 0.001; \eta^2 = 0.032$), and a side effect for muscle thickness ($F = 19.367; p < 0.001; \eta^2 = 0.110$). For the gastrocnemius muscle, a sides*group interaction was found for stiffness ($F = 4.269; p = 0.039; \eta^2 = 0.004$). Tone and stiffness were significantly higher at MT than at MB sites, and in patients in the chronic stage compared with the subacute stroke group (all, $p < 0.001$). There were differences between sides for the biceps brachii, with lower tone, stiffness, and thickness of the affected upper limb ($p < 0.001$). Muscle thickness was correlated with tone ($r = 0.355, p < 0.001$) and stiffness ($r = 0.353, p < 0.001$) for the biceps brachii, and with stiffness for the gastrocnemius ($r = 0.237, p = 0.003$).

3.2. Discriminative Ability between Stroke Survivors and Healthy Controls

A significant group*site interaction was demonstrated for the biceps brachii stiffness ($F = 1.732; p = 0.010; \eta^2 = 0.023$) and for the gastrocnemius muscle tone ($F = 1.942; p = 0.003; \eta^2 = 0.025$) and stiffness ($F = 1.742; p = 0.012; \eta^2 = 0.023$). Muscle tone and stiffness were higher in chronic stroke patients than in controls, and lower in the subacute stroke group compared with healthy participants (only for the gastrocnemius). Stroke survivors showed

lower biceps brachii and gastrocnemius muscle thickness than did those in the control group (all, $p < 0.05$).

4. Discussion

The structural and mechanical properties of the biceps brachii and gastrocnemius muscles were heterogeneously distributed among stroke survivors. Tone and stiffness were higher at MT than at MB sites, and the biceps brachii muscle tone, stiffness, and thickness were lower at the affected side. Myotonometry and ultrasound measures were significantly correlated, and both techniques could discriminate between the paretic and nonparetic upper limb and between participants with and without stroke.

4.1. Mechanical and Structural Muscular Adaptations among Stroke Survivors

Current literature has characterized the muscle adaptations after stroke with SEMG [14], mechanomyography [24], and ultrasonography [6,8]. Myotonometry uses superficial mechanical deformation and represents a convenient approach that is considerably less costly than elastography [25] and has good psychometric properties [11]. Hence, it can be of high clinical value to track muscle mechanical changes [26], especially in stroke populations [9]. Previous research with myotonometry has measured a single site, often at the midportion of the MB, which does not reflect the spatial distribution [11]. This is the first study in which myotonometry is used to quantify multiple sites within the MB and tendon to image the differences in tone and stiffness between upper and lower limb muscles in people with subacute or chronic stroke.

In both biceps brachii and gastrocnemius muscles, tone and stiffness were significantly higher at MT than at MB sites, in line with findings in adults with spinal cord injury [27] and Parkinson's disease [28]. These differences can be explained by the structural and functional adaptations that occur after stroke, such as variations in the number and length of sarcomeres in the skeletal muscles [29]. Additionally, reduced muscle thickness [6], together with increased tendon compliance and muscle pennation angle [30], leads to muscle atrophy, which is usually observed following prolonged disuse. In fact, disuse is considered the main factor involved in most of these muscular changes [6].

Differences in the level of function may help to understand the distinct behaviors of the upper and lower limb muscles. Participants with stroke reported a severe upper limb motor impairment (FMA-UE < 19) that denotes a restricted ability to function (e.g., bring the arm into the body, extend the elbow, or relax the fingers) [31]. The involvement of the paretic upper limb in daily activities is related to the extent of motor restriction [32]. The most natural way to respond to that is to rely on the nonaffected upper limb [33]. This would explain the decreased tone, stiffness, and thickness of the biceps brachii of the affected side. As regards the lower limb motor performance, 45% of stroke participants (18/40) were able to walk without physical assistance (FAC ≥ 3) [23]. Despite the fact that their walking ability was somehow preserved, stroke survivors tend to be inactive and sedentary [34]. Several studies conclude that changes in muscle architecture after stroke may not be limited to the spastic side [35], and that the nonparetic lower limb also adapts [5,6]. For example, the bilateral overuse of lower limb muscles to walk or support body weight [36] may account for the lack of differences between sides for the gastrocnemius muscle tone, stiffness, and thickness. Overall, some of the adaptations of the lower limb have been described as muscle specific [37], which makes it difficult to reach a definite conclusion. These findings seem to suggest the need to involve both lower extremities during rehabilitation, although further research is necessary to support this claim.

Weak to moderate correlations were observed between myotonometry and ultrasound measures [21,36,38], with inconsistent results across muscles. According to these findings, both techniques assess similar, although different, constructs and, therefore, can be combined for the clinical assessment of PSS [21].

4.2. Discriminative Ability between Stroke Survivors and Healthy Controls

Changes in muscle architecture can take place as early as 3 weeks after stroke [39]. Muscle thickness decreases in the first months of recovery [40], whereas the evolution to the chronic stage leads to increased muscle tone and stiffness. This has been shown in the upper [41,42] and lower [36,38] limb in chronic stroke survivors. Neural and morphological changes can support these findings [43]. However, evidence is still conflicting about how the structural and mechanical muscle properties may evolve with time. For example, Mirbagheri et al. described two temporal patterns of change for muscle stiffness over the first 12 months: a progressive increase or a slow decline after the first four weeks in patients with mild motor impairment [44]. Again, the level of motor performance represents a key aspect in most of these changes. The current results indicate that treatment strategies must be carefully chosen according to the stroke stage. Additional research may help to understand the best therapeutic approach at each stage. Previous studies concluded that task-oriented repetitive training can be recommended during the acute and subacute stages to speed recovery and prevent disuse [45]. For the chronic stage, multimodal rehabilitation programs including localized soft-tissue therapies may be of more interest. Longitudinal studies monitoring the structural and mechanical muscular changes in larger cohorts are warranted to improve the clinical management of PSS.

4.3. Study Limitations

Chronic stage was defined as more than 9 months post-stroke [17], although new standards may recommend otherwise. Assessments were conducted in the relaxed position to prevent fatigue. There is a high heterogeneity among studies in the measurement protocols using myotonometry and ultrasonography, which may negatively impact the clinical interpretation of the results [11,21]. Future studies could also combine SEMG and mechanomyography as an index of muscle performance with myotonometry or ultrasonography, as recently performed in healthy individuals [46]. Stroke patients were involved in different treatment routines. Since physical activity and medication intake can modulate muscle tone, the possible impact of the physical rehabilitation programs on the results needs to be considered.

5. Conclusions

Topographical maps of the biceps brachii and gastrocnemius muscles revealed a heterogeneous distribution of structural and mechanical properties, with lower muscle tone, stiffness, and thickness of the paretic upper limb, and increasing tone and stiffness in MT compared with MB locations and from subacute to chronic stroke stage. Among stroke survivors, the discriminative ability of myotonometry and ultrasonography can be influenced by the assessed muscle, the stroke stage, and the level of motor performance.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph20021405/s1>, Figure S1: MyotonPRO device in use.

Author Contributions: Conceptualization, M.I.G.-B., P.G.-G. and A.M.H.-R.; methodology, M.I.G.-B., P.G.-G., P.M., M.J.C.-H. and A.M.H.-R.; software, P.M. and A.M.H.-R.; formal analysis, M.I.G.-B., P.G.-G., P.M. and A.M.H.-R.; writing—original draft preparation, M.I.G.-B., P.G.-G., P.M., M.J.C.-H. and A.M.H.-R.; writing—review and editing, M.I.G.-B., P.G.-G., P.M., M.J.C.-H. and A.M.H.-R.; supervision, M.I.G.-B.; project administration, M.I.G.-B.; funding acquisition, M.I.G.-B. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Ilustre Colegio Profesional de Fisioterapeutas de Andalucía, grant number 03729/19D/MA.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the Institutional Review Board of Junta de Andalucía Ethical Committee for Biomedical Research (protocol code CI 1222-N-16 and 24/03/2019) for studies involving humans.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: We wish to thank the participants who took part in the study, and the stroke units at CRECER and DACE for helping us during the recruitment process.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Wissel, J.; Verrier, M.; Simpson, D.M.; Charles, D.; Guinto, P.; Papapetropoulos, S.; Sunnerhagen, K.S. Post-Stroke Spasticity: Predictors of Early Development and Considerations for Therapeutic Intervention. *PM R* **2015**, *7*, 60–67. [[CrossRef](#)] [[PubMed](#)]
2. Trompetto, C.; Marinelli, L.; Mori, L.; Pelosin, E.; Currà, A.; Molfetta, L.; Abbruzzese, G. Pathophysiology of Spasticity: Implications for Neurorehabilitation. *BioMed Res. Int.* **2014**, *2014*, 354906. [[CrossRef](#)] [[PubMed](#)]
3. Raghavan, P. Upper Limb Motor Impairment Post Stroke. *Phys. Med. Rehabil. Clin. N. Am.* **2016**, *26*, 599–610. [[CrossRef](#)]
4. Pundik, S.; McCabe, J.; Skelly, M.; Tatsuoka, C.; Daly, J.J. Association of Spasticity and Motor Dysfunction in Chronic Stroke. *Ann. Phys. Rehabil. Med.* **2019**, *62*, 397–402. [[CrossRef](#)] [[PubMed](#)]
5. Hunnicutt, J.L.; Gregory, C.M. Skeletal Muscle Changes Following Stroke: A Systematic Review and Comparison to Healthy Individuals. *Top. Stroke Rehabil.* **2017**, *24*, 463–471. [[CrossRef](#)]
6. Faturi, F.M.; Lopes Santos, G.; Ocamoto, G.N.; Russo, T.L. Structural Muscular Adaptations in Upper Limb after Stroke: A Systematic Review. *Top. Stroke Rehabil.* **2019**, *26*, 73–79. [[CrossRef](#)]
7. Del Prete, C.M.; Viva, M.G.; de Trane, S.; Brindisino, F.; Barassi, G.; Specchia, A.; di Iorio, A.; Pellegrino, R. An Observational Cross-Sectional Study of Gender and Disability as Determinants of Person-Centered Medicine in Botulinum Neurotoxin Treatment of Upper Motoneuron Syndrome. *Toxins* **2022**, *14*, 246. [[CrossRef](#)]
8. Lehoux, M.C.; Sobczak, S.; Cloutier, F.; Charest, S.; Bertrand-Grenier, A. Shear Wave Elastography Potential to Characterize Spastic Muscles in Stroke Survivors: Literature Review. *Clin. Biomech.* **2020**, *72*, 84–93. [[CrossRef](#)]
9. Luo, Z.; Lo, W.L.A.; Bian, R.; Wong, S.; Li, L. Advanced Quantitative Estimation Methods for Spasticity: A Literature Review. *J. Int. Med. Res.* **2019**, *48*, 0300060519888425. [[CrossRef](#)]
10. Miller, T.; Ying, M.; Sau Lan Tsang, C.; Huang, M.; Pang, M.Y.C. Reliability and Validity of Ultrasound Elastography for Evaluating Muscle Stiffness in Neurological Populations: A Systematic Review and Meta-Analysis. *Phys. Ther.* **2021**, *101*, pzaa188. [[CrossRef](#)]
11. García-Bernal, M.I.; Heredia-Rizo, A.M.; González-García, P.; Cortés-Vega, M.D.; Casuso-Holgado, M.J. Validity and Reliability of Myotonometry for Assessing Muscle Viscoelastic Properties in Patients with Stroke: A Systematic Review and Meta-Analysis. *Sci. Rep.* **2021**, *11*, 5062. [[CrossRef](#)]
12. Murillo, C.; Falla, D.; Sanderson, A.; Rushton, A.; Heneghan, N.R. Shear Wave Elastography Investigation of Multifidus Stiffness in Individuals with Low Back Pain. *J. Electromyogr. Kinesiol.* **2019**, *47*, 19–24. [[CrossRef](#)]
13. García-Bernal, M.; González-García, P.; Casuso-Holgado, M.; Cortés-Vega, M.; Heredia-Rizo, A. Measuring Mechanical Properties of Spastic Muscles after Stroke. Does Muscle Position during Assessment Really Matter? *Arch. Phys. Med. Rehabil.* **2022**, *103*, 2368–2374. [[CrossRef](#)]
14. Munneke, M.A.M.; Bakker, C.D.; Goverde, E.A.; Pasman, J.W.; Stegeman, D.F. On the Electrode Positioning for Bipolar EMG Recording of Forearm Extensor and Flexor Muscle Activity after Transcranial Magnetic Stimulation. *J. Electromyogr. Kinesiol.* **2018**, *40*, 23–31. [[CrossRef](#)]
15. Alburquerque-Sendín, F.; Madeleine, P.; Fernández-De-Las-Peñas, C.; Camargo, P.R.; Salvini, T.F. Spotlight on Topographical Pressure Pain Sensitivity Maps: A Review. *J. Pain Res.* **2018**, *11*, 215–225. [[CrossRef](#)]
16. Heredia-Rizo, A.M.; Petersen, K.K.; Arendt-Nielsen, L.; Madeleine, P. Eccentric Training Changes the Pressure Pain and Stiffness Maps of the Upper Trapezius in Females with Chronic Neck-Shoulder Pain: A Preliminary Study. *Pain Med.* **2020**, *21*, 1936–1946. [[CrossRef](#)]
17. Sarasso, S.; Määttä, S.; Ferrarelli, F.; Poryazova, R.; Tononi, G.; Small, S.L. Plastic Changes Following Imitation-Based Speech and Language Therapy for Aphasia: A High-Density Sleep EEG Study. *Neurorehabil. Neural Repair* **2014**, *28*, 129–138. [[CrossRef](#)]
18. Kumar, R.T.S.; Pandyan, A.D.; Sharma, A.K. Biomechanical Measurement of Post-Stroke Spasticity. *Age Ageing* **2006**, *35*, 371–375. [[CrossRef](#)]
19. Moon, S.H.; Choi, J.H.; Park, S.E. The Effects of Functional Electrical Stimulation on Muscle Tone and Stiffness of Stroke Patients. *J. Phys. Ther. Sci.* **2017**, *29*, 238–241. [[CrossRef](#)]

20. Ilahi, S.; Masi, A.T.; White, A.; Devos, A.; Henderson, J.; Nair, K. Quantified Biomechanical Properties of Lower Lumbar Myofascia in Younger Adults with Chronic Idiopathic Low Back Pain and Matched Healthy Controls. *Clin. Biomech.* **2020**, *73*, 78–85. [[CrossRef](#)]
21. Miller, T.; Ying, M.T.C.; Chung, R.C.K.; Pang, M.Y.C. Convergent Validity and Test-Retest Reliability of Multimodal Ultrasonography and Related Clinical Measures in People With Chronic Stroke. *Arch. Phys. Med. Rehabil.* **2022**, *103*, 459–472.e4. [[CrossRef](#)]
22. Barbosa, N.E.; Forero, S.M.; Galeano, C.P.; Hernández, E.D.; Landinez, N.S.; Sunnerhagen, K.S.; Murphy, M.A. Translation and Cultural Validation of Clinical Observational Scales—The Fugl-Meyer Assessment for Post Stroke Sensorimotor Function in Colombian Spanish. *Disabil. Rehabil.* **2019**, *41*, 2317–2323. [[CrossRef](#)] [[PubMed](#)]
23. Mehrholz, J.; Wagner, K.; Rutte, K.; Meißner, D.; Pohl, M. Predictive Validity and Responsiveness of the Functional Ambulation Category in Hemiparetic Patients After Stroke. *Arch. Phys. Med. Rehabil.* **2007**, *88*, 1314–1319. [[CrossRef](#)] [[PubMed](#)]
24. Hu, X.L.; Tong, K.Y.; Li, L. The Mechanomyography of Persons after Stroke during Isometric Voluntary Contractions. *J. Electromyogr. Kinesiol.* **2007**, *17*, 473–483. [[CrossRef](#)] [[PubMed](#)]
25. Kelly, J.P.; Koppenhaver, S.L.; Michener, L.A.; Proulx, L.; Bisagni, F.; Cleland, J.A. Characterization of Tissue Stiffness of the Infraspinatus, Erector Spinae, and Gastrocnemius Muscle Using Ultrasound Shear Wave Elastography and Superficial Mechanical Deformation. *J. Electromyogr. Kinesiol.* **2018**, *38*, 73–80. [[CrossRef](#)]
26. Liu, C.L.; Feng, Y.N.; Zhang, H.Q.; Li, Y.P.; Zhu, Y.; Zhang, Z.J. Assessing the Viscoelastic Properties of Upper Trapezius Muscle: Intra- and Inter-Tester Reliability and the Effect of Shoulder Elevation. *J. Electromyogr. Kinesiol.* **2018**, *43*, 226–229. [[CrossRef](#)]
27. Ge, J.S.; Chang, T.T.; Zhang, Z.J. Reliability of Myotonometric Measurement of Stiffness in Patients with Spinal Cord Injury. *Med. Sci. Monit.* **2020**, *26*, e924811-1. [[CrossRef](#)]
28. Marusiak, J.; Jaskólska, A.; Budrewicz, S.; Koszewicz, M.; Jaskólski, A. Increased Muscle Belly and Tendon Stiffness in Patients with Parkinson’s Disease, as Measured by Myotonometry. *Mov. Disord.* **2011**, *26*, 2119–2122. [[CrossRef](#)]
29. Son, J.; Rymer, W.Z.; Lee, S.S.M. Limited Fascicle Shortening and Fascicle Rotation May Be Associated with Impaired Voluntary Force-Generating Capacity in Pennate Muscles of Chronic Stroke Survivors. *Clin. Biomech.* **2020**, *75*, 105007. [[CrossRef](#)]
30. Li, L.; Tong, K.Y.; Hu, X. The Effect of Poststroke Impairments on Brachialis Muscle Architecture as Measured by Ultrasound. *Arch. Phys. Med. Rehabil.* **2007**, *88*, 243–250. [[CrossRef](#)]
31. Woodbury, M.L.; Velozo, C.A.; Richards, L.G.; Duncan, P.W. Rasch Analysis Staging Methodology to Classify Upper Extremity Movement Impairment after Stroke. *Arch. Phys. Med. Rehabil.* **2013**, *94*, 1527–1533. [[CrossRef](#)]
32. Narai, E.; Hagino, H.; Komatsu, T.; Togo, F. Accelerometer-Based Monitoring of Upper Limb Movement in Older Adults with Acute and Subacute Stroke. *J. Geriatr. Phys. Ther.* **2016**, *39*, 171–177. [[CrossRef](#)]
33. Jones, T.A.; Allred, R.P.; Jefferson, S.C.; Kerr, A.L.; Woodie, D.A.; Cheng, S.Y.; Adkins, D.A.L. Motor System Plasticity in Stroke Models Intrinsically Use-Dependent, Unreliably Useful. *Stroke* **2013**, *44*, 104–106. [[CrossRef](#)] [[PubMed](#)]
34. Fini, N.A.; Holland, A.E.; Keating, J.; Simek, J.; Bernhardt, J. How Physically Active Are People Following Stroke? Systematic Review and Quantitative Synthesis. *Phys. Ther.* **2017**, *97*, 707–717. [[CrossRef](#)]
35. Berenpas, F.; Martens, A.M.; Weerdesteyn, V.; Geurts, A.C.; van Alfen, N. Bilateral Changes in Muscle Architecture of Physically Active People with Chronic Stroke: A Quantitative Muscle Ultrasound Study. *Clin. Neurophysiol.* **2017**, *128*, 115–122. [[CrossRef](#)]
36. Wang, J.-S.; Lee, S.-B.; Moon, S.-H. The Immediate Effect of PNF Pattern on Muscle Tone and Muscle Stiffness in Chronic Stroke Patient. *J. Phys. Ther. Sci.* **2016**, *28*, 967–970. [[CrossRef](#)]
37. Jakubowski, K.L.; Terman, A.; Santana, R.V.C.; Lee, S.S.M. Passive Material Properties of Stroke-Impaired Plantarflexor and Dorsiflexor Muscles. *Clin. Biomech.* **2017**, *49*, 48–55. [[CrossRef](#)]
38. Fröhlich-Zwahlen, A.K.; Casartelli, N.C.; Item-Glatthorn, J.F.; Maffiuletti, N.A. Validity of Resting Myotonometric Assessment of Lower Extremity Muscles in Chronic Stroke Patients with Limited Hypertonia: A Preliminary Study. *J. Electromyogr. Kinesiol.* **2014**, *24*, 762–769. [[CrossRef](#)]
39. Scherbakov, N.; von Haehling, S.; Anker, S.D.; Dirnagl, U.; Doehner, W. Stroke Induced Sarcopenia: Muscle Wasting and Disability after Stroke. *Int. J. Cardiol.* **2013**, *170*, 89–94. [[CrossRef](#)]
40. Beckwée, D.; Cuyppers, L.; Lefeber, N.; de Keersmaecker, E.; Scheys, E.; van Hees, W.; Perkisas, S.; de Raedt, S.; Kerckhofs, E.; Bautmans, I.; et al. Skeletal Muscle Changes in the First Three Months of Stroke Recovery: A Systematic Review. *J. Rehabil. Med.* **2022**, *54*, jrm00308. [[CrossRef](#)]
41. Gao, J.; He, W.; Du, L.J.; Chen, J.; Park, D.; Wells, M.; Fowlkes, B.; O’Dell, M. Quantitative Ultrasound Imaging to Assess the Biceps Brachii Muscle in Chronic Post-Stroke Spasticity: Preliminary Observation. *Ultrasound Med. Biol.* **2018**, *44*, 1931–1940. [[CrossRef](#)]
42. Lee, S.S.M.; Jakubowski, K.L.; Spear, S.C.; Rymer, W.Z. Muscle Material Properties in Passive and Active Stroke-Impaired Muscle. *J. Biomech.* **2019**, *83*, 197–204. [[CrossRef](#)]
43. Stubbs, P.W.; Nielsen, J.F.; Sinkjær, T.; Mrachacz-Kersting, N. Short-Latency Crossed Spinal Responses Are Impaired Differently in Sub-Acute and Chronic Stroke Patients. *Clin. Neurophysiol.* **2012**, *123*, 541–549. [[CrossRef](#)]
44. Mirbagheri, M.M.; Tsao, C.; Rymer, W.Z. Natural History of Neuromuscular Properties after Stroke: A Longitudinal Study. *J. Neurol. Neurosurg. Psychiatry* **2009**, *80*, 1212–1217. [[CrossRef](#)]

45. Jeon, B.J.; Kim, W.H.; Park, E.Y. Effect of Task-Oriented Training for People with Stroke: A Meta-Analysis Focused on Repetitive or Circuit Training. *Top. Stroke Rehabil.* **2015**, *22*, 34–43. [[CrossRef](#)]
46. Tier, L.; Salomoni, S.E.; Hug, F.; Besomi, M.; Hodges, P.W. Shear Modulus of Multifidus and Longissimus Muscles Measured Using Shear Wave Elastography Correlates with Muscle Activity, but Depends on Image Quality. *J. Electromyogr. Kinesiol.* **2021**, *56*, 102505. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.