



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Factors influencing rectal hypersensitivity in irritable bowel syndrome: A systematic review and meta-analysis

Roberts, Christopher; Albusoda, Ahmed; Farmer, Adam D.; Aziz, Qasim

Published in:
Neurogastroenterology and Motility

DOI (link to publication from Publisher):
[10.1111/nmo.14515](https://doi.org/10.1111/nmo.14515)

Creative Commons License
CC BY 4.0

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Roberts, C., Albusoda, A., Farmer, A. D., & Aziz, Q. (2023). Factors influencing rectal hypersensitivity in irritable bowel syndrome: A systematic review and meta-analysis. *Neurogastroenterology and Motility*, 35(4), Article e14515. Advance online publication. <https://doi.org/10.1111/nmo.14515>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW

Factors influencing rectal hypersensitivity in irritable bowel syndrome: A systematic review and meta-analysis

Christopher Roberts^{1,2}  | Ahmed Albusoda¹  | Adam D. Farmer^{1,3,4,5}  | Qasim Aziz¹ 

¹Centre for Neuroscience, Surgery and Trauma, Blizard Institute, Wingate Institute of Neurogastroenterology, Barts and the London School of Medicine & Dentistry, Queen Mary University of London, London, UK

²Department of Paediatric Gastroenterology, University Hospital Southampton, Southampton, UK

³Mech-Sense, Department of Gastroenterology, Aalborg University Hospital, Aalborg, Denmark

⁴Institute of Applied Clinical Sciences, University of Keele, Keele, UK

⁵Department of Gastroenterology, University Hospitals of North Midlands NHS Trust, Stoke on Trent, UK

Correspondence

Qasim Aziz and Christopher Roberts, The Wingate Institute of Neurogastroenterology, Barts and the London School of Medicine and Dentistry, 26 Ashfield Street, Whitechapel, London E1 2AJ, UK.

Email: q.aziz@qmul.ac.uk and chrisrobertsberkhamsted@gmail.com

Funding information

AA was funded by People Programme of the European Union's Seventh Framework Programme under REA, Grant/Award Number: grant agreement no.607652 (NeuroGu); ADF was funded by West Midlands Clinical Research Scholarship

Abstract

Background: A frequent, although not universal, feature of irritable bowel syndrome (IBS) is heightened sensitivity to mechanical stimulation of the rectum, termed rectal hypersensitivity (RH). Differences in RH-based on sex, IBS subtype, IBS diagnostic criteria and age of population studied are incompletely understood. We aimed to determine whether IBS population had lower pain thresholds than healthy controls.

Methods: We searched MEDLINE and EMBASE databases (1970–2021). Prospective studies that compared pain/discomfort thresholds to mechanical rectal stimuli in IBS and healthy controls were included. Data were pooled for meta-analyses and effect sizes were calculated with 95% confidence interval (CIs).

Results: Our search strategy identified 809 studies of which 32 studies met the inclusion criteria. Reduced rectal pain thresholds was more common in IBS patients compared to healthy controls with an effect size of 1.00 95% CIs (0.77–1.24) ($p < 0.0001$) ($I^2 = 78.6\%$). The pediatric IBS population had lower pain thresholds than adult IBS populations ($p = 0.05$) but no difference based on IBS diagnostic criteria, subtype or sex.

Conclusion & Inferences: The results suggest that reduced rectal pain threshold to experimental stimulation is far more common in IBS patients than healthy controls. Further research is required to understand the pathophysiological and therapeutic implications of rectal sensitivity such as its role in measuring response to treatment and prognosis in IBS.

KEYWORDS

abdominal pain, irritable bowel syndrome, visceral hypersensitivity

1 | INTRODUCTION

Rectal hypersensitivity is defined as increased sensitivity to experimental stimuli applied to the gastrointestinal (GI) tract.¹ It

can arise due to a combination of either heightened sensitivity to noxious stimuli (hyperalgesia) and/or non-noxious stimuli (allodynia) due to factors such as peripheral and central sensitisation.² Additional mechanisms include alterations in central factors such

Adam D. Farmer and Qasim Aziz joint senior authors.

Study Registration: PROSPERO Reference - CRD42018095687.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Neurogastroenterology & Motility* published by John Wiley & Sons Ltd.

as aberrant brain processing³ and abnormal descending inhibitory control of pain pathways.^{4,5}

Irritable bowel syndrome (IBS) is characterized by visceral pain and altered bowel habits. IBS is thought to be a disorder of a dysfunctional gut-brain axis where symptoms are present in the absence of demonstrable organic disease.⁶ IBS is subtyped according to the predominant bowel habit into IBS with constipation (IBS-C), IBS with diarrhea (IBS-D), mixed bowel habit IBS (IBS-M) and IBS unclassified (IBS-U).⁷ IBS has a reported prevalence of around 4.5% and is associated with a large reduction in quality of life.⁸

In IBS, the cause of visceral pain is incompletely understood but it is considered that rectal hypersensitivity exerts an important effect.^{9,10} Rectal pain thresholds can be evaluated using mechanical (manual or automated using a barostat), nutrient, chemical, thermal or electrical stimuli to discriminate whether pain thresholds are higher or lower in different groups. The intensity of pain when using such techniques is most commonly measured using a self-report visual analogue scale (VAS). In IBS patients' there is a correlation between symptom severity and pain sensitivity to mechanical rectal distension.^{11,12} Provocation tests suffer from significant heterogeneity as distension protocols and definitions for a painful stimulus vary from study to study, although recent international efforts have sought to improve standardization.¹³ This variation in testing conditions has prevented the development of standardized normal values from which hyper, normo and hyposensate individuals can be identified. Repeated exposure to experimental provocation stimuli can normalize rectal sensation probably due to habituation.¹⁴ However, mechanical stimulation is currently regarded as the most reliable instrument to assess rectal sensitivity.¹⁵ At present rectal provocation testing is rarely used outside of GI physiology units in the clinical setting.¹⁶ Rectal sensation testing can be considered in the evaluation of rectal hyposensitivity when assessing for megarectum and in the consideration of fecal incontinence when anal sphincter function may be impaired.¹⁶ Current standard of practice is for balloon distension either using phasic or ramp distension techniques, barostat is mainly reserved for research environments.

The primary aims of this study were to assess if pain thresholds to mechanical rectal stimulation were different in the IBS population compared to healthy controls. Secondary aims were to ascertain if there were differences in rectal pain thresholds based on IBS diagnostic criteria used, IBS subtype, sex or age. The reason for this is that there is some evidence that Rome III and IV represent a more severe phenotype of the disease than the earlier Rome iterations as reported prevalence is higher using previous versions of the Rome criteria^{8,17} compared to using Rome IV. In addition, there is some evidence that some patients, particularly those with IBS-C may demonstrate hyposensitivity^{18,19} but this is not a universal conclusion.²⁰ Therefore we aimed to see if there were differences in pain thresholds based on IBS subtype.

Despite the large number of published studies on rectal provocation testing in IBS patients, to the best of our knowledge, there has been no meta-analysis on the topic.

Key Points

Background

- Reduced rectal pain thresholds are considered to be a feature of irritable bowel syndrome (IBS) although this epiphenomenon is variably reported.

Findings

- In 32 studies with 1452 individuals with IBS, compared to 567 healthy controls, reduced rectal pain thresholds were more common with a large effect size.
- Pediatric IBS patients demonstrated larger reduction in pain thresholds than adults.
- There were no differences based on diagnostic criteria, subtype, or sex.

What is the impact on clinical practice in the foreseeable future?

- Reduced rectal pain thresholds are frequently seen in patients with IBS and may impact clinical presentation. Therefore this phenomenon requires physician awareness.
- Rectal hypersensitivity should be considered an important mechanism of pain in IBS and future studies should focus on understanding its pathophysiology and importance as a marker of response to treatment.

2 | MATERIALS AND METHODS

2.1 | Study population and study design

The systematic review and meta-analyses were conducted according to the PRISMA recommendations and were registered with PROSPERO (Reference CRD42018095687).²¹ The search of the literature was performed using MEDLINE and EMBASE (1970–2021). This was carried out using the set search strategies outlined in the Table S1. There were no language restrictions. Eligibility criteria are shown in Box 1. The bibliographies of all relevant studies and available meeting abstracts were screened to identify studies that were missed by the original search criteria. Senior authors were contacted to provide additional information where required. Articles were assessed independently by two reviewers (CR and AA) using the predetermined eligibility criteria. Disagreements were resolved by consensus.

2.2 | Data extraction

The name of the first author, year of publication, location of study, IBS population size, control population size, IBS criteria and primary

BOX 1 Eligibility criteria

Inclusion criteria

- Diagnosis of IBS according to ROME criteria
- Assessing of IBS and a healthy control population
- Measurement of pain/ discomfort thresholds using mechanical rectal distension
- Prospective study.

Exclusion criteria

- If testing pain/discomfort thresholds by means other than mechanical such as electrical.
- Retrospective studies were rejected due to the risk of repeat data.

outcome data which were recorded in means and standard deviations were extracted into an Excel Spreadsheet (Excel 2016, Microsoft).

The primary outcome was to see if there was a difference in pain/discomfort threshold in IBS and control populations. Secondary outcomes were assessed to see if pain thresholds differed between IBS subtype, IBS diagnostic criteria used, sex and age studied. Paediatric populations were defined as being less than 18 years old. To perform secondary analysis, data were pooled from the studies that provided the necessary information to perform the subgroup analyses. Standard deviations were calculated according to the Cochrane Collaboration guidelines.²²

2.3 | Quality assessment and risk of bias

Two investigators performed a biased assessment independently for all studies included in the meta-analysis. Bias was scored in six areas using a modified checklist for case-control studies.²³ These areas were: (1) blinding of assessors, (2) use of international criteria to diagnose IBS, (3) use of aged-matched controls, (4) use of sex-matched controls, (5) exclusion of other chronic pain disorders and (6) controlling for other known factors that affect pain sensation such as anxiety and depression.

2.4 | Data analysis

Data were pooled for meta-analysis and a random effect model using the Hartung-Knapp-Sidik-Jonkman method was chosen. Heterogeneity was assessed using the I^2 statistical test which gives values between 0% and 100%, with 0% representing no observed heterogeneity. Outcomes were assessed using Hedges' g effect sizes and are reported with 95% confidence intervals (CI). A pre-specified secondary analysis was performed to determine if the effect size was modified in various subgroups. Meta-regression was performed to determine if rectal compliance played a role in the

development of reduced rectal pain thresholds using studies that provided data on rectal compliance. The statistical criterion was $p < 0.05$. Evidence of publication bias was assessed by using a funnel plot and Egger's Test. Propriety software was used to perform the meta-analysis and generate the plots (Comprehensive Meta-Analysis Version 2, Biostat, Version 2) and (R, R Foundation for Statistical Computing).

3 | RESULTS

3.1 | Search results

The search generated 809 citations of which 81 were classed as relevant and 32 met the inclusion criteria comprising 1452 individuals with IBS and 567 controls, see [Figure 1](#). In total, 49 studies were rejected. The characteristics of the included studies are shown in [Table S2](#).

3.2 | Rectal hypersensitivity in IBS

Irritable bowel syndrome participants had lower pain thresholds in comparison to healthy controls with a large effect size, 1.00 95% CIs (0.77–1.24) ($p < 0.0001$) ($I^2 = 78.6\%$), see [Figure 2](#).

3.3 | IBS effect size based on IBS subtype, diagnostic criteria, sex and age

Of the 31 studies included in the meta-analysis, nine provided data for the different IBS subtypes. Compared to healthy controls, the effect size was strongly positive for both IBS-C and IBS-D. There was a significant difference between the subgroups ($p = 0.001$), however, when IBS-M subtype along with studies that did not differentiate between subtypes were removed from the analysis there was no significant difference in pain thresholds ($p = 0.40$). The effect size for IBS-C is 0.98 (0.39–1.58) and for IBS-D 1.37 (0.36–2.37). A forest plot is shown in [Figure 3 plot A](#).

Of the 31 included studies, nine used Rome I, 18 used Rome II, two used Rome III and two used Rome IV. There was no statistical difference found between the groups ($p = 0.21$). A forest plot is in [Figure 3 plot B](#).

Of the 31 studies included eight provided data about sexes covering 242 women and 83 men with IBS. There was no statistical difference identified between the sexes ($p = 0.13$). A forest plot is in [Figure 3 plot C](#).

Three studies examined 39 pediatric individuals with IBS. The effect size for adult IBS patients compared to adult controls was 0.94 95% CI (0.70–1.17) whereas, in the pediatric IBS patients compared to pediatric controls was 1.85 95% CI (–0.07–3.77). There was a statistically significant difference between adult and pediatric populations ($p = 0.05$). A forest plot is in [Figure 3 plot D](#).





FIGURE 1 Flow diagram of the assessment of studies included in the IBS meta-analysis.

3.4 | Heterogeneity

Heterogeneity was high as shown by $I^2 = 78.6\%$ which is regarded as being a high level of heterogeneity, the reasons for this high level were unclear, a number of statistical tests were used to look for the reasons including detecting outliers, an influence analysis and a Baujat plot which all revealed that the Camilleri et al.²⁰ and Bouin et al.²⁴ studies provided a significant amount of the observed heterogeneity. With these studies removed $I^2 = 61\%$ which is regarded as medium heterogeneity and the effect size remained relatively unaffected at 1.07 (0.85–1.28). See Figure S1.

Another cause of possible heterogeneity was the variability in rectal provocation testing using either rectal balloon distension or rectal barostat. Despite this, when we adjusted for the device used heterogeneity remained unchanged at $I^2 = 73\%$. When the different distension protocols using ramp or phasic distension protocols were corrected for, then heterogeneity improved to $I^2 = 65\%$. There was variability in the terms used. When we assessed studies that used pain and discomfort thresholds then heterogeneity did improve to $I^2 = 61\%$. Interestingly, there was no difference in pain threshold effect sizes between studies that use pain or discomfort as their definition ($p = 0.67$).

FIGURE 2 A forest plot of the effect sizes for pain thresholds in an IBS vs a control population. The pooled effect size is 0.97 95% CI (0.77–1.17) ($p < 0.0001$).



3.5 | Meta-regression

Results on rectal compliance were provided in seven studies covering 413 individuals with IBS and 160 healthy controls. Lower levels of rectal compliance were found to be associated with a lower rectal pain threshold ($p = 0.006$).

3.6 | IBS study quality assessment

The quality of the studies included in the meta-analysis is summarized in Table S3. All the papers were scored out of 12 with zero indicating the least risk of bias. In most studies, the assessors were not blinded and hence most scored two points for this section although the majority did use a single-blind protocol. A funnel plot was performed to assess for the presence of publication bias which is included in the Figure S2 and an Egger's test was carried out which

showed the presence of publication bias. When the publication bias was corrected the resulting effect size remained highly positive at 0.66 95% CI (0.38–0.93) $p < 0.0001$.

4 | DISCUSSION

This meta-analysis demonstrates average rectal pain thresholds were lower in IBS patients in comparison to healthy controls. This observation is common to IBS subtypes and is not modified by the diagnostic criteria or sex. However, pain thresholds were lower in pediatric IBS populations compared to adults.

Chronic visceral pain is a central defining factor in IBS and contributes to healthcare seeking and reduces the quality of life.^{25,26} The absolute cause of chronic visceral pain in IBS is incompletely understood but visceral hypersensitivity remains the germane hypothesis.²⁷ Although rectal provocation testing is not considered to




FIGURE 3 Forest Plots showing effect size in different subgroups. Each subgroup is present with the p value for the difference between all subgroups. (A) IBS subtype – IBS-D and IBS-C ($p = 0.4$), (B) Rome criteria used to diagnose IBS ($p = 0.21$), (C) Sex ($p = 0.13$) and (D) Age group studied ($p = 0.05$).

display the prerequisite receiver operator characteristics for diagnosis of IBS, provocation testing can be used in the clinical setting in the assessment of evacuation disorders or when a megarectum may be suspected.¹⁶ Our results demonstrate, at least in the participants included in the meta-analysis, that reduced pain thresholds are a defining feature in people with IBS compared to healthy controls.

All included studies used the four different versions of the Rome criteria and although there have been changes to the diagnostic criteria for IBS, our findings indicate that this does not impact the degree of rectal hypersensitivity. In some way, this is surprising as the Rome IV classification of IBS requires the presence of weekly pain and prevalence of IBS drops between the different Rome versions.^{9,17} It is difficult to make firm conclusions here as there were only four studies that looked at either Rome III or IV and so further studies are required to confirm whether rectal hypersensitivity is more common in the newer iterations of Rome. There was a trend towards Rome IV having a greater reduction in pain thresholds than individuals who were identified by other Rome criteria but this did not reach statistical significance. This suggests that reduced rectal pain thresholds may be more common in individuals who have been diagnosed using Rome IV compared to the earlier versions of the criteria. If reduced rectal pain thresholds were more common in the Rome IV population this would indicate that the amount of reduction of rectal pain thresholds may co-relate with disease severity,

this does conform with previous work.¹¹ However further work is required to confirm this hypothesis.

Interestingly, no difference in rectal pain thresholds was observed between IBS-C and IBS-D. Previous studies have suggested there may be a proportion of patients particularly those with IBS-C who may be hyposensitive due to alterations in rectal sensorimotor function and compliance.^{18,20} Conversely, in IBS-D, sensory defecatory urge threshold and rectal compliance are diminished in comparison to IBS-C and healthy controls.²⁸ Given the lack of overall effect between subtypes, our data suggest that whilst there are differences in individual studies, an overall effect is not present and reduced rectal pain thresholds are common to all IBS subtypes. From our data, it is far more likely for IBS patients to demonstrate reduced rectal pain thresholds but in many of our studies there were subgroups of IBS patients who had higher pain rectal pain thresholds than controls. These individuals may be hyposensitive.

Pediatric IBS patients were more likely to have lower rectal pain thresholds than adult IBS patients. The reason for this difference is unclear and it could be reasoned that pediatric IBS may represent a more severe phenotype. However, other reasons may also explain this difference. For instance, even though both pediatric²⁹ and adult^{30,31} IBS populations often demonstrate hypervigilance as a significant number reported pain before any stimulus was applied. It is plausible that this hypervigilance is more common in the pediatric

IBS cohort. To date, there has been no comparison in the prevalence of hypervigilance between adult and pediatric populations. Similarly, anxiety, stress and hormonal factors may also play a role in the differences observed between adult and pediatric cohorts. Further research is required to understand the reasons for the difference in rectal sensitivity between adult and pediatric patients. Greater anxiety before the procedure may play a role in explaining the difference in rectal pain thresholds between pediatric and adult patients. This could explain the significantly increased levels of hypervigilance in the pediatric populations. Unfortunately, in the studies that assessed anxiety this was in the form of generalized anxiety rather than anxiety towards the distension procedure itself.

The results of the meta-regression show that reduced rectal compliance plays a key role in the lower pain thresholds seen in patients with IBS compared to healthy controls; however, these results do not fully explain the lower pain thresholds seen in IBS patients compared to controls. Reduced rectal compliance is linked with reduced pain thresholds in provocation testing as the rectum is less able to expand as distension increases and therefore a pain stimulus is likely to occur at a lower threshold.³²

The cause of reduced pain thresholds and likely rectal hypersensitivity is unclear, there is a significant thought that subtle levels of inflammation and immune activation seen in IBS may play a key role.^{33,34} In vitro testing shows that high burdens of inflammation may lead to the sensitizing effect of inflammation on afferent neurons.³⁴ In this regard, some neurotrophic factors, such as nerve growth factor, and heightened expression of the transient receptor potential cation channel subfamily V member 1 (TRPV1) and the purinergic P2X3 receptor measured in the mucosa have been implicated.^{35–37} TRPV1 channels are further shown to play a key role in rectal hypersensitivity as the reversal of their activation in a mouse model was able to normalize pain response to rectal distension.³⁸

However, in some patient's inflammation may be required as a triggering event such as in post-infectious IBS. In post-infectious IBS occurs in around 10% of episodes of acute gastroenteritis (AG).³⁹ In AG there is a burden of inflammation which should then improve. The cause of why these symptoms persist is unclear but some genetic and environmental factors have been identified relating to both the episode of AG itself and some prior risk factors such as female, younger age and previous anxiety/depression.^{40,41} Interestingly these risk factors are also risk factors for IBS in non-AG patients.¹⁷ A similar phenomenon is seen in patients with quiescent IBD where most patients do not experience long-term abdominal patients, but a subset does which is regarded as being IBD-IBS crossover. The risk factors for development of IBS-IBD crossover seem to be similar to those of IBS and post-infectious IBS further confirming that co-existing psychological comorbidities appear to play a key role in the development of IBS.

Another major factor that has been implicated in the development of IBS is hypervigilance.

Hypervigilance is already known to play a key role in rectal hypersensitivity as experimental stimulation is likely to represent a 'threat' to the patient with repeated exposure reducing this 'threat'

through habituation. Habituation has been demonstrated to normalize perceptual ratings to rectal stimulation in IBS patients and this is associated with decreased activity in brain regions and networks associated with pain processing.¹⁵ This indicates that psychological factors play a role in pain processing. In IBS, abdominal pain-related fear learning and memory processes are altered, which may contribute to central pain amplification and hypervigilance which may be enhanced in those with comorbid anxiety and depression.^{41,42} Experimentally induced negative emotions during painful rectal distension even in healthy volunteers can lead to increased brain activity in the left thalamus and right dorsal posterior cingulate gyrus.⁴³ Besides, alterations in descending pain modulatory pathways also contribute to rectal hypersensitivity in IBS.⁴⁴ Indeed, altered brain processing to rectal stimulation is seen in IBS patients compared to healthy controls in areas involved in descending pain modulation.^{3,5,45} The above mechanisms are likely to be more prominent in IBS than the inflammation solely. In post-infectious IBS the symptoms persist well after the inflammation improves and is thought to be due to central sensitisation.⁴⁶

This study is subject to several limitations. There was significant heterogeneity in the study although this was improved after outliers were removed. There was though still some heterogeneity that persisted. This is likely because of the differences in the site of experimental stimulation and methods used for determining pain given that it is a subjective experience with marked intra-and inter-individual variability.⁴⁷ There was an attempt to correct the difference in study methodology by only evaluation studies that assessed rectal as opposed to colonic sensation. All the studies included took place in tertiary care settings so may represent a more severe phenotype than what is seen in other settings.

There is an established link between the presence of depression and anxiety reporting pain at lower pain intensities, so it was surprising that studies did not try and correct for these confounding factors.^{48,49} The fact that only some did will have likely increased the observed heterogeneity and unfortunately, a meta-regression was not possible given the lack of data. However, how important such factors are in IBS is unclear as van der Veek et al.¹² did not demonstrate differential rectal sensitivity when IBS participants were stratified according to levels of anxiety, depression or somatization.¹²

Suggestions for further research include using a standardized rectal provocation testing¹³ in healthy individuals to identify accurate normal values, which would allow for identification of hyper, hypo and normosensate individuals with IBS. Clinical trials could then be designed focusing on these groups to determine what treatments would be the most efficacious within the different populations.

This meta-analysis indicates that reduced pain thresholds are an epiphenomenon strongly associated with both adult and pediatric IBS populations in comparison to healthy controls. The mechanism causing reduced pain thresholds is not fully understood but is likely due to a combination of peripheral and central factors and is likely to vary in different populations and diseases. Standardized rectal provocation testing, given that it is cheap and widely available, may facilitate the identification of hypersensate individuals

who may benefit from personalisation of treatment strategies that aim to reduce hypersensitivity such as in the use of pain modulators and/ or psychological therapies. However further work is required to suggest that serial testing could be used as a biomarker for success in the treatment of IBS and as a prognostic indicator of long-term outcomes. Rectal barostat could be used in the clinical setting to identify possible responses to pharmacological^{50,51} and psychological⁵² treatments.

AUTHOR CONTRIBUTIONS

Christopher Roberts & Ahmed Albusoda: Designed search protocol; performed data collection; manuscript preparation; statistical analysis; critical revision of the manuscript for important intellectual content. Joint first authors. Adam D Farmer & Qasim Aziz: Pioneered study concept and design, technical support; critical revision of the manuscript for important intellectual content, project supervision. Joint senior authors.

ACKNOWLEDGEMENT

AA was funded by the People Programme of the European Union's Seventh Framework Programme under REA grant agreement no.607652 (NeuroGu). ADF was supported by the West Midlands Clinical Research Network Scholarship.

CONFLICT OF INTEREST

None of the authors have any conflict of interest to declare.

ORCID

Christopher Roberts  <https://orcid.org/0000-0002-6215-8044>

Ahmed Albusoda  <https://orcid.org/0000-0001-9733-4776>

Adam D. Farmer  <https://orcid.org/0000-0003-1902-2640>

Qasim Aziz  <https://orcid.org/0000-0002-2718-2065>

REFERENCES

- Camilleri M, Coulie B, Tack JF. Visceral hypersensitivity: facts, speculations, and challenges. *Gut*. 2001;48(1):125-131. doi:10.1136/gut.48.1.125
- Knowles CH, Aziz Q. Basic and clinical aspects of gastrointestinal pain. *Pain*. 2009;141(3):191-209. doi:10.1016/j.pain.2008.12.011
- Tillisch K, Mayer EA, Labus JS. Quantitative meta-analysis identifies brain regions activated during rectal distension in irritable bowel syndrome. *Gastroenterology*. 2011;140(1):91-100. doi:10.1053/j.gastro.2010.07.053
- Wilder-Smith OHG. Chronic pain and surgery: a review of new insights from sensory testing. *J Pain Palliat Care Pharmacother*. 2011;25(2):146-159. doi:10.3109/15360288.2010.505256
- Albusoda A, Ruffle JK, Friis KA, et al. Systematic review with meta-analysis: conditioned pain modulation in patients with the irritable bowel syndrome. *Aliment Pharmacol Ther*. 2018;48(8):797-806. doi:10.1111/apt.14965
- Chey WD, Kurlander J, Eswaran S. Irritable bowel syndrome: a clinical review. *JAMA*. 2015;313(9):949-958. doi:10.1001/jama.2015.0954
- Lacy BE, Mearin F, Chang L, et al. Bowel disorders. *Gastroenterology*. 2016;150(6):1393-1407. doi:10.1053/j.gastro.2016.02.031
- Palsen OS, Whitehead W, Törnblom H, Sperber AD, Simren M. Prevalence of Rome IV functional bowel disorders among adults in the United States, Canada, and the United Kingdom. *Gastroenterology*. 2020;158(5):1262-1273.e3. doi:10.1053/j.gastro.2019.12.021
- Farmer AD, Aziz Q. Gut pain & visceral hypersensitivity. *Br J Pain*. 2013;7(1):39-47. doi:10.1177/2049463713479229
- Mertz H, Naliboff B, Munakata J, Niazi N, Mayer EA. Altered rectal perception is a biological marker of patients with irritable bowel syndrome. *Gastroenterology*. 1995;109(1):40-52. doi:10.1016/0016-5085(95)90267-8
- van der Veek PPJ, van Rood YR, Masclee AAM. Symptom severity but not psychopathology predicts visceral hypersensitivity in irritable bowel syndrome. *Clin Gastroenterol Hepatol*. 2008;6(3):321-328. doi:10.1016/j.cgh.2007.12.005
- Kanazawa M, Palsen OS, Thiwan SIM, et al. Contributions of pain sensitivity and colonic motility to IBS symptom severity and predominant bowel habits. *Am J Gastroenterol*. 2008;103(10):2550-2561. doi:10.1111/j.1572-0241.2008.02066.x
- Boeckxstaens GE, Drug V, Dumitrascu D, et al. Phenotyping of subjects for large scale studies on patients with IBS. *Neurogastroenterol Motil*. 2016;28(8):1134-1147. doi:10.1111/nmo.12886
- Naliboff BD, Berman S, Suyenobu B, et al. Longitudinal change in perceptual and brain activation response to visceral stimuli in irritable bowel syndrome patients. *Gastroenterology*. 2006;131(2):352-365. doi:10.1053/j.gastro.2006.05.014
- Mujagic Z, Keszthelyi D, Aziz Q, et al. Systematic review: instruments to assess abdominal pain in irritable bowel syndrome. *Aliment Pharmacol Ther*. 2015;42(9):1064-1081. doi:10.1111/apt.13378
- Carrington E, Heinrich H, Knowles CH, et al. The international anorectal physiology working group (IAPWG) recommendations: standardized testing protocol and the London classification for disorders of anorectal function. *Neurogastroenterol Motil*. 2020;32(1):e13679. doi:10.1111/nmo.13679
- Lovell RM, Ford AC. Global prevalence of and risk factors for irritable bowel syndrome: a meta-analysis. *Clin Gastroenterol Hepatol*. 2012;10(7):712-721.e4. doi:10.1016/j.cgh.2012.02.029
- Burgell RE, Mark SS. Rectal hyposensitivity. *J Neurogastroenterol Motil*. 2012;18(4):384. doi:10.5056/jnm.2012.18.4.373
- Harraf F, Schmulson M, Saba L, et al. Subtypes of constipation predominant irritable bowel syndrome based on rectal perception. *Gut*. 1998;43(3):388-394. doi:10.1136/gut.43.3.388
- Camilleri M, McKinzie S, Busciglio I, et al. Prospective study of motor, sensory, psychologic, and autonomic functions in patients with irritable bowel syndrome. *Clin Gastroenterol Hepatol*. 2008;6(7):772-781.e5. doi:10.1016/j.cgh.2008.02.060
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;339:b2700. doi:10.1136/bmj.b2700
- Chandler J, Higgins J, Deeks J, Davenport C, Clarke M. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.2.0 (Updated February 2017)*. The Cochrane Collaboration; 2017.
- Whiting PF, Rutjes AWS, Westwood ME, et al. Quadas-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann Intern Med*. 2011;155(8):529. doi:10.7326/0003-4819-155-8-201110180-00009
- Bouin M, Plourde V, Boivin M, et al. Rectal distention testing in patients with irritable bowel syndrome: sensitivity, specificity, and predictive values of pain sensory thresholds. *Gastroenterology*. 2002;122(7):1771-1777. doi:10.1053/gast.2002.33601
- Akehurst RL, Brazier JE, Mathers N, et al. Health-related quality of life and cost impact of irritable bowel syndrome in a UK primary care setting. *Pharmacoeconomics*. 2002;20(7):455-462. doi:10.2165/00019053-200220070-00003
- Gralnek IM, Hays RD, Kilbourne AA, Naliboff B, Mayer EA. The impact of irritable bowel syndrome on health-related

- quality of life. *Gastroenterology*. 2000;119(3):654-660. doi:[10.1053/gast.2000.16484](https://doi.org/10.1053/gast.2000.16484)
27. Farmer AD, Aziz Q. Visceral pain hypersensitivity in functional gastrointestinal disorders. *Br Med Bull*. 2009;91(1):123-136. doi:[10.1093/bmb/ldp026](https://doi.org/10.1093/bmb/ldp026)
 28. Zar S, Benson MJ, Kumar D. Rectal afferent hypersensitivity and compliance in irritable bowel syndrome: differences between diarrhoea-predominant and constipation-predominant subgroups. *Eur J Gastroenterol Hepatol*. 2006;18(2):151-158. doi:[10.1097/00042737-200602000-00007](https://doi.org/10.1097/00042737-200602000-00007)
 29. van Ginkel R, Voskuil WP, Benninga MA, Taminiau JAJM, Boeckstaens GE. Alterations in rectal sensitivity and motility in childhood irritable bowel syndrome. *Gastroenterology*. 2001;120(1):31-38. doi:[10.1053/gast.2001.20898](https://doi.org/10.1053/gast.2001.20898)
 30. Naliboff BD, Munakata J, Fullerton S, et al. Evidence for two distinct perceptual alterations in irritable bowel syndrome. *Gut*. 1997;41(4):505-512. doi:[10.1136/gut.41.4.505](https://doi.org/10.1136/gut.41.4.505)
 31. Dorn SD, Palsos OS, Thiwan SIM, et al. Increased colonic pain sensitivity in irritable bowel syndrome is the result of an increased tendency to report pain rather than increased neurosensory sensitivity. *Gut*. 2007;56(9):1202-1209. doi:[10.1136/gut.2006.117390](https://doi.org/10.1136/gut.2006.117390)
 32. Cremon C, Gargano L, Morselli-Labate AM, et al. Mucosal immune activation in irritable bowel syndrome: gender-dependence and association with digestive symptoms. *Am J Gastroenterol*. 2009;104(2):392-400. doi:[10.1038/ajg.2008.94](https://doi.org/10.1038/ajg.2008.94)
 33. Bercik P, Verdu EF, Collins SM. Is irritable bowel syndrome a low-grade inflammatory bowel disease? *Gastroenterol Clin North Am*. 2005;34(2):235-245. doi:[10.1016/j.gtc.2005.02.007](https://doi.org/10.1016/j.gtc.2005.02.007)
 34. di Mola FF, Friess H, Zhu ZW, et al. Nerve growth factor and Trk high affinity receptor (TrkA) gene expression in inflammatory bowel disease. *Gut*. 2000;46(5):670-679. doi:[10.1136/gut.46.5.670](https://doi.org/10.1136/gut.46.5.670)
 35. Yiangou Y, Facer P, Baecker PA, et al. ATP-gated ion channel P2X3 is increased in human inflammatory bowel disease. *Neurogastroenterol Motil*. 2001;13(4):365-369. doi:[10.1046/j.1365-2982.2001.00276.x](https://doi.org/10.1046/j.1365-2982.2001.00276.x)
 36. Yiangou Y, Facer P, Dyer NHC, et al. Vanilloid receptor 1 immunoreactivity in inflamed human bowel. *Lancet*. 2001;357(9265):1338-1339. doi:[10.1016/S0140-6736\(00\)04503-7](https://doi.org/10.1016/S0140-6736(00)04503-7)
 37. Perna E, Aguilera-Lizarraga J, Florens MV, et al. Effect of resolvins on sensitisation of TRPV1 and visceral hypersensitivity in IBS. *Gut*. 2021;70(7):1275-1286. doi:[10.1136/gutjnl-2020-321530](https://doi.org/10.1136/gutjnl-2020-321530)
 38. Klem F, Wadhwa A, Prokop LJ, et al. Prevalence, risk factors, and outcomes of irritable bowel syndrome after infectious enteritis: a systematic review and meta-analysis. *Gastroenterology*. 2017;152(5):1042-1054.e1. doi:[10.1053/j.gastro.2016.12.039](https://doi.org/10.1053/j.gastro.2016.12.039)
 39. Villani AC, Lemire M, Thabane M, et al. Genetic risk factors for post-infectious irritable bowel syndrome following a waterborne outbreak of gastroenteritis. *Gastroenterology*. 2010;138(4):1502-1513. doi:[10.1053/j.gastro.2009.12.049](https://doi.org/10.1053/j.gastro.2009.12.049)
 40. Jonefjäll B, Strid H, Öhman L, Svedlund J, Bergstedt A, Simren M. Characterization of IBS-like symptoms in patients with ulcerative colitis in clinical remission. *Neurogastroenterol Motil*. 2013;25(9):756-e578. doi:[10.1111/nmo.12163](https://doi.org/10.1111/nmo.12163)
 41. Icenhour A, Langhorst J, Benson S, et al. Neural circuitry of abdominal pain-related fear learning and reinstatement in irritable bowel syndrome. *Neurogastroenterol Motil*. 2015;27(1):127. doi:[10.1111/nmo.12489](https://doi.org/10.1111/nmo.12489)
 42. Rosenberger C, Elsenbruch S, Scholle A, et al. Effects of psychological stress on the cerebral processing of visceral stimuli in healthy women. *Neurogastroenterol Motil*. 2009;21(7):740-e45. doi:[10.1111/j.1365-2982.2009.01295.x](https://doi.org/10.1111/j.1365-2982.2009.01295.x)
 43. Wilder-Smith CH, Robert-Yap J. Abnormal endogenous pain modulation and somatic and visceral hypersensitivity in female patients with irritable bowel syndrome. *World J Gastroenterol*. 2007;13(27):3699-3704. doi:[10.3748/wjg.v13.i27.3699](https://doi.org/10.3748/wjg.v13.i27.3699)
 44. Elsenbruch S. Abdominal pain in irritable bowel syndrome: a review of putative psychological, neural and neuro-immune mechanisms. *Brain Behav Immun*. 2011;25(3):386-394. doi:[10.1016/j.bbi.2010.11.010](https://doi.org/10.1016/j.bbi.2010.11.010)
 45. Balemans D, Mondelaers SU, Cibert-Goton V, et al. Evidence for long-term sensitization of the bowel in patients with post-infectious-IBS. *Sci Rep*. 2017;7(1):13606. doi:[10.1038/s41598-017-12618-7](https://doi.org/10.1038/s41598-017-12618-7)
 46. Farmer AD, Coen SJ, Kano M, et al. Psychophysiological responses to pain identify reproducible human clusters. *Pain*. 2013;154(11):2266-2276. doi:[10.1016/j.pain.2013.05.016](https://doi.org/10.1016/j.pain.2013.05.016)
 47. Grinsvall C, Törnblom H, Tack J, van Oudenhove L, Simrén M. Psychological factors selectively upregulate rectal pain perception in hypersensitive patients with irritable bowel syndrome. *Neurogastroenterol Motil*. 2015;27(12):1772-1782. doi:[10.1111/nmo.12689](https://doi.org/10.1111/nmo.12689)
 48. Grinsvall C, Törnblom H, Tack J, van Oudenhove L, Simrén M. Relationships between psychological state, abuse, somatization and visceral pain sensitivity in irritable bowel syndrome. *United European Gastroenterol J*. 2018;6(2):300-309. doi:[10.1177/2050640617715851](https://doi.org/10.1177/2050640617715851)
 49. Thua NM, Murray CDR, Winchester WJ, et al. Amitriptyline modifies the visceral hypersensitivity response to acute stress in the irritable bowel syndrome. *Aliment Pharmacol Ther*. 2009;29(5):552-560. doi:[10.1111/j.1365-2036.2008.03918.x](https://doi.org/10.1111/j.1365-2036.2008.03918.x)
 50. Houghton LA, Fell C, Whorwell PJ, Jones I, Sudworth DP, Gale JD. Effect of a second-generation $\alpha 2\delta$ ligand (pregabalin) on visceral sensation in hypersensitive patients with irritable bowel syndrome. *Gut*. 2007;56(9):1218-1225. doi:[10.1136/gut.2006.110858](https://doi.org/10.1136/gut.2006.110858)
 51. Edebol-Carlman H, Ljótsson B, Linton SJ, et al. Face-to-face cognitive-behavioral therapy for irritable bowel syndrome: the effects on gastrointestinal and psychiatric symptoms. *Gastroenterol Res Pract*. 2017;2017:1-9. doi:[10.1155/2017/8915872](https://doi.org/10.1155/2017/8915872)

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Roberts C, Albusoda A, Farmer AD, Aziz Q. Factors influencing rectal hypersensitivity in irritable bowel syndrome: A systematic review and meta-analysis. *Neurogastroenterology & Motility*. 2023;35:e14515. doi:[10.1111/nmo.14515](https://doi.org/10.1111/nmo.14515)