

Aalborg Universitet

A short update on equipackable graphs

Vestergaard, Preben D.

Publication date: 2007

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA):

Vestergaard, P. D. (2007). A short update on equipackable graphs. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2007-06

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

AALBORG UNIVERSITY

A short update on equipackable graphs by Preben Dahl Vestergaard R-2007-06 February 2007

DEPARTMENT OF MATHEMATICAL SCIENCES AALBORG UNIVERSITY Fredrik Bajers Vej 7 G • DK-9220 Aalborg Øst • Denmark Phone: +45 96 35 80 80 • Telefax: +45 98 15 81 29 URL: www.math.auc.dk/research/reports/reports.htm



A short update on equipackable graphs

P.D. Vestergaard

Department of Mathematical Sciences, Aalborg University Fredrik Bajers Vej 7G, DK-9220 Aalborg Ø, Denmark e-mail: pdv@math.aau.dk

To be published in **Discrete Mathematics**

Keywords: packing, decomposition, randomly packable, randomly decomposable, path 2000 Mathematics Subject Classification: 05C70, 05C38

Abstract

A graph is called equipackable if every maximal packing in it is also maximum. This generalizes randomly packable graphs. We survey known results both on randomly packable graphs and on equipackable graphs. As a new result is given a characterization of P_3 -equipackable graphs with all valencies at least two.

1 Definitions

Let H be a subgraph of G. A collection H_1, H_2, \ldots, H_k of edge disjoint subgraphs of G, each isomorphic to H, is called an H-packing of G, and it is maximal if $G - \bigcup_{i=1}^k E(H_i)$ contains no subgraph isomorphic to H. It is maximum if no H-packing in G with more than k copies of H exists. Bosák

[3] wrote a book on packing, he used the equivalent term *decomposition*, and there are many papers on the topic. G is called *H*-packable if there exists an *H*-packing of G which uses all edges in G and G is called *randomly H*-packable if every maximal *H*-packing in G uses all edges in G, i.e., if every *H*-packing can be extended to a decomposition of the edges of G into copies of H. Note that not every graph H produces a family of randomly *H*-packable graphs, if e.g. H is the disjoint union of K_2 and K_3 , no other graph than H itself is randomly *H*-packable.

As a relaxation of random *H*-packability we define *G* to be *H*-equipackable if every maximal *H*-packing is also a maximum *H*-packing. So the randomly *H*-packable graphs is contained as a subclass in the class of *H*-equipackable graphs. This paper focuses on $H = P_3$.

2 Notation

A graph G has order |V(G)| and size |E(G)|. The path and circuit on k vertices is denoted by P_k and C_k , respectively. By $C_m \bullet C_n$ we denote the graph of order n+m-1 obtained from two circuits C_m and C_n by identifying one vertex from each. $S_{2k+1}^{(r)}$ denotes the graph obtained from r paths P_{2k+1} by identifying their center vertices. The corona $H \circ K_1$ on H is is the graph of order 2|H| obtained by adding for each vertex x of H one new vertex x' and a new edge xx'. By H + G we denote the graph obtained from two disjoint graphs H and G by adding edges joining each vertex of H to each vertex of G. A matching in the graph G is a set of independent edges in G, it is perfect if it covers all vertices of G. By M_t , $t \ge 1$, we denote a matching having t edges. The union of k disjoint copies of a graph G is denoted by nG, e.g. $M_t = tK_2$.

3 Results

An early result by Caro and Schönheim is

Lemma 1 ([4]) A connected graph G is P_3 -packable if and only if G has even size.

Observation A connected graph G of odd size contains an edge whose deletion leaves a connected graph, which necessarily is of even size and therefore by Lemma 1 is P_3 -packable.

It is clear that a maximum P_3 -packing in a connected P_3 -equipackable graph contains either all or all but one edge of the graph.

It follows that a P_3 -equipackable connected graph of even size is also randomly P_3 -packable. Another useful observation is that if there is a maximal P_3 -packing of a connected graph G which omits at least two edges, then Gis not P_3 -equipackable.

Ruiz characterized randomly P_3 -packable graphs.

Theorem 1 ([11]) A connected graph G is randomly P_3 -packable if and only if $G \cong C_4$ or $G \cong K_{1,2k}, k \ge 1$.

Thus P_3 -equipackable graphs of even size are quadrilaterals or stars. It remains to characterize P_3 -equipackable graphs of odd size. Hartnell and Vestergaard did that for graphs of girth at least five.

Theorem 2 ([5]) A connected graph G of girth 5 or more is P_3 -equipackable if and only if G satisfies one of the following:

(i) G is a tree consisting of a single star (i.e., $K_{1,n}$)

(ii) G is a tree which has two stems that are at distance 3, where the vertices on this shortest path are w_1 and w_2 . Furthermore the stems are of odd parity and have no neighbours other than leaves and w_1 or w_2 . In addition w_1 and w_2 are of degree two.

(iii) G is a tree which has two stems that are at distance two where w is the common neighbour of the stems. The two stems must be of different parity and neither stem has other neighbours than its leaves and w. Furthermore, the vertex w must be of degree two.

(iv) G is a tree which has two stems that are adjacent where these stems are of the same parity and these stems have only each other and their leaves as neighbours.

(v) G is either C_7 , C_5 or has 5+2m vertices where G consists of a circuit of length 5 along with 2m leaves attached to exactly one node on the 5-cycle.

The remaining problem is to characterize P_3 -equipackable graphs of girth 3 and 4. We shall consider graphs with $\delta(G) \geq 2$.

Theorem 3 Let G be a graph with $\delta(G) \geq 2$, which is connected and has a cutvertex. Then G is P_3 -equipackable if and only if either (1) G is of order eight and is obtained from two vertex disjoint quadrilaterals together with an edge joining a vertex in one quadrilateral to a vertex in the other or (2) G is a $C_3 \bullet C_4$.

Proof. The two graphs described can be checked to be P_3 -equipackable. Conversely, let x be a cutvertex in the P_3 -equipackable graph G and let A'_1, \ldots, A'_m be the components of G - x, while A_1, \ldots, A_m are the graphs spanned in G by $A'_i \cup \{x\}$. We first observe that $|E(A_i)|$ can be odd for at most one i. Assume namely A'_i is a component with $|E(A_i)|$ odd. We shall by P_3 -removals isolate a subgraph of A'_i having odd size. If there is an even number of edges from x to A'_i we P_3 -remove them in pairs. This isolates A'_i with an odd number of edges. If there is an odd number of edges from x to A'_i we can P_3 -remove all but one of them, say xy, and as $\delta(G) \geq 2$ we have that $y \in V(A'_i)$ is incident with an edge $yz, z \in A'_i \setminus \{y\}$. By removal of $\{xy, yz\}$ we also in this case have isolated a subgraph of A'_i with an odd number of edges and $|E(A_i)|$ can by the observation after Lemma 1 be odd for at most one index i.

If all $|E(A_i)|$ are even, then G has even size and, by the remark after Lemma 1, G is randomly P_3 -packable and hence by Theorem 1 is C_4 or a star, but C_4 has no cutvertex and the star violates $\delta(G) \geq 2$, so this case cannot occur.

Thus $|E(A_1)|$, say, is odd and all other $|E(A_i)|, i \ge 2$, are even.

Subcase 1. Assume a bridge xy of G joins x to A'_1 . P_3 -remove yx, xz, $z \in N(x) \cap V(A'_2)$. Then A'_1 is isolated and as a connected graph of even size, it is therefore randomly P_3 -packable and by Theorem 1 isomorphic to C_4 as $\delta(G) \geq 2$. The graph spanned in G by the union of A_2, \ldots, A_m is connected, of even size and it is also randomly P_3 -packable. By Theorem 1 it is isomorphic to C_4 and G consists of two disjoint C_4 's joined by an edge as claimed in (1).

Subcase 2. Assume x is joined to A'_1 by more than one edge. Again the graph spanned by the union of A_2, \ldots, A_m is connected and of even size. It is randomly P_3 -packable because we can by P_3 -removals inside A_1 make sure that the unique non-removed edge is isolated inside A'_1 . Thus all of A_2, \ldots, A_m spans a C_4 . We can deduce that A_1 is isomorphic to C_3 , because A_1 minus an edge must be randomly P_3 -packable and C_4 cannot be fitted in to ensure P_3 -equipackability of G, so A_1 minus an edge must be a star and only P_3 with its two ends joined to x will do. In this case G is the disjoint union of a quadrilateral and a triangle with one vertex from each identified as claimed in (2). This proves Theorem 3.

Theorem 4 A connected graph G with $\delta(G) \geq 2$ is P_3 -equipackable if and only if G is one of of the graphs listed in Figure 1.

Proof. We can by inspection verify that the graphs in Fig. 1 all are P_3 -equipackable. If G has a cutvertex we know by Theorem 3 that $G \cong C_3 \bullet C_4$ or G can be obtained by joining two quadrilaterals by an edge. So assume that G is 2-connected and P_3 -equipackable, we must prove that G is one of the remaining graphs listed on Fig. 1. Let C with length ℓ be a longest circuit in G. If $\ell = 3$ necessarily $G \cong C_3$, a graph in the family on top of Fig. 1. If $\ell = 4$ and C has no diagonal the only possibility is $G \cong C_4$ since by 2-connectivity any x in $V(G) \setminus V(C)$ must be joined to C by two independent paths. Each path must be an edge, otherwise G would contain a circuit longer than four. Also x must have valency 2. Thus G is of even size, that implies that G is randomly P_3 -packable and hence by Theorem 1 must be a C_4 . If $\ell = 4$ and C has a diagonal we can by a similar argument obtain that G is a graph in the family on top of Fig. 1. If $\ell = 5$ and G has order 5 we obtain the four graphs of Fig. 1. If $\ell = 5$ and G has order > 5 each x in $V(G) \setminus V(C)$ is by 2-connectivity joined by two independent paths to C.

Again, not to produce a longer circuit, x is joined by edges to two vertices at distance two on C, and x has valency two. But now it is easy to see that we by P_3 -deletions can isolate two edges, so this case cannot occur. If $\ell = 6$ we can find that C must contain a triangular diagonal. If G has order 6, either Gis this graph, included in Fig. 1, or we can by P_3 -removals isolate two edges, a contradiction. In the remaining cases either G is C_7 , included in Fig. 1, or we can for each vertex x on C pairwise P_3 -remove its adjacent edges in $E(G) \setminus E(C)$, so at most one edge besides the two circuit edges remain at x. If the end vertex y of such an edge has a neighbour z such that $yz \notin E(C)$, we P_3 -remove xy, yz. We have thus isolated a component consisting of C and possibly at some C-vertices one other edge in $E(G) \setminus E(C)$, a pendent edge or a diagonal. At most one vertex on C has a pendent edge, otherwise we could by P_3 -removals on C isolate the two pendent edges, a contradiction. We can now see that we by further P_3 -removals can isolate two edges, a contradiction. This proves the theorem.



Figure 1: All P_3 -equipackable graphs which are connected and have all valencies at least two.

4 Randomly packable graphs

Equipackability is a relaxation of random packability, so let us mention a few results from packings, all have potential for generalizations to equipackability.

Theorem 1 by Ruiz [11] was later generalized from P_3 to $K_{1,r}$ by Barrientos, Bernasconi, Jeltsch, Tronisco and Ruiz:

Theorem 5 ([1]) For $r \ge 2$ a connected graph is randomly $K_{1,r}$ -packable if and only if it is $K_{r,r}$ or it is bipartite with all valencies in one partite set being multiples of r and all valencies in the other set being less than r.

4.1 Randomly path-packable graphs

Beineke, Goddard, Hamburger [2] generalized Theorem 1 from P_3 to P_k , $3 \le k \le 6$, and Molina with coauthors [9, 10] extended to $k \le 10$:

Characterization of connected, randomly P_k -packable graphs		
[11]	Randomly P_2 -packable	Trivially every graph
[11]	Randomly P_3 -packable	C_4 and stars of even size
[2]	Randomly P_4 -packable	$P_4, K_4, K_{2,3}, C_6, C_3 \bullet C_3$
[9, 10]		
[2]	Randomly P_5 -packable	$P_5, K_{2,4}, C_4 \bullet C_4, C_8, S_5^{(k)}, k \ge 2$
[9, 10]		
	Randomly P_6 -packable	$P_6, C_{10}, C_5 \bullet C_5, K_4 \circ K_1$ and
[2]		the graph obtained by joining two
[9, 10]		new vertices by an edge to
		the same valency 2 vertex of a $K_{2,4}$
	Randomly P_7, P_8, P_9 ,- and P_{10} -packable	Families of graphs whose descriptions
[9, 10]		become increasingly complex
		with growing k

4.2 Randomly matching-packable graphs

Ruiz characterized randomly M_2 -packable graphs:

Theorem 6 ([11]) A graph is randomly M_2 -packable if and only if it one of the following: $C_4, K_4, 2K_3, K_3 \cup K_{1,3}, 2K_{1,n}$ or $2nK_2, n \ge 1$.

This was generalized to randomly M_t -packable graphs by Beineke, Goddard, Hamburger, but only for graphs with sufficiently many edges:

Theorem 7 ([2] Th. 2.5) For a given integer $t \ge 2$, a graph with at least $2t^3 - t^2$ edges is randomly M_t -packable if and only if it is isomorphic to tH, where H is either nK_2 or $K_{1,n}$ for some $n \ge 1$.

Those graphs in which each matching can be extended to a perfect matching are called *randomly matchable* and if each matching extends to a maximum matching, which is not necessecarily perfect, they are called *equimatchable*. Sumner characterized randomly matchable graphs:

Theorem 8 ([12]) A connected graph G is randomly matchable if and only if $G \cong K_{2n}$ or $G \cong K_{n,n}$, where n is a positive integer.

Lesk, Plummer and Pulleyblank [6] gave a characterization of equimatchable graphs in terms of the Gallai-Edmonds structure theorem (described in [7]). Define a *total matching* to be a subset X of $E(G) \cup V(G)$ such that no two elements of X are adjacent or incident. Topp and Vestergaard characterized totally equimatchable graphs:

Theorem 9 ([13]) A connected graph G is totally equimatchable if and only if G is $K_n, K_{n,n}$ or $K_1 + \bigcup_{i=1}^n K_{2m_i}$, where n and m_1, m_2, \ldots, m_n are any positive integers.

4.3 Randomly K_n -packable graphs

Beineke, Hamburger and Goddard proved

Theorem 10 ([2] Th. 3.1) A graph G is randomly K_n -packable if and only if every edge is in precisely one copy of K_n in G.

McSorley and Porter ([8]) have considered a vertex variant. They let

 $(*) \{G_{\alpha n}\}_{\alpha=1}^{\infty} = G_n, G_{2n}, \dots, G_{\alpha n}, \dots$

be a sequence of graphs such that $G_1 \stackrel{\sim}{=} K_n$, $G_{\alpha n}$ has order αn and $G_{\alpha n} - K_n \stackrel{\sim}{=} G_{(\alpha-1)n}$ for any complete subgraph K_n in $G_{\alpha n}$, $\alpha \ge 2$.

For $0 \leq \lambda \leq n$ they call (*) a (K_n, λ) -removable sequence if $G_{\alpha n}$ is $(n-1) + (\alpha - 1)\lambda$ -regular, and they prove that for a fixed $n \geq 2$ there is a unique (K_n, λ) -removable sequence for $\lambda = 0, n-1$ or n.

Theorem 11 ([8]) For a fixed $n \geq 2$,

 $\{\alpha K_n\}_{\alpha=1}^{\infty}$ is the unique $(K_n, 0)$ -removable sequence. $\{K_{\underbrace{\alpha, \alpha, \alpha, \ldots, \alpha}_n}\}_{\alpha=1}^{\infty}$ is the unique $(K_n, n-1)$ -removable sequence.

 $\{K_{\alpha n}\}_{\alpha=1}^{\infty}$ is the unique (K_n, n) -removable sequence.

5 Open problems

Molina and coauthors in [10] posed a still unsolved problem: Does the characterizations become easier if only 2-connected graphs are considered? Their examples of 2-connected randomly P_k -packable graphs contain only two copies of P_k and with three exceptions have a vertex of odd degree, and they asked whether that holds generally.

References

- C. Barrientos, A. Bernasconi, E. Jeltsch, C. Tronosco, S. Ruiz: "Randomly star-decomposable graphs", Proc. 250th Anniversary Conf. on Graph Theory, Congr. Numr. 64 (1988), 193-195.
- [2] L.W. Beineke, P. Hamburger, W.D. Goddard, "Random packings of graphs", Discrete Mathematics 125(1994), 45-54.

- [3] Juraj Bosák, "Decompositions of graphs". Kluwer Academic Publishers. Dordrecht 1990. ISBN 0-7923-0747-X.
- [4] Y. Caro, J. Schönheim, "Decompositions of trees into isomorphic subtrees", Ars Combinatoria 9 (1980),119-130.
- [5] B.L. Hartnell, P.D. Vestergaard: "Equipackable graphs", to appear in Bulletin of ICA.
- [6] M. Lesk, M.D. Plummer and W.R. Pulleyblank, "Equi-matchable graphs", Graph Theory and Combinatorics (Cambridge, 1983), 239-254.
- [7] L. Lovász, M.D. Plummer, "Matching Theory", North-Holland Mathematics Studies 121, Annals of Discrete Mathematics 29, Elsevier Science Publishers, Amsterdam 1986.
- [8] John P. McSorley, Thomas D. Porter: "K_p-removable sequences of graphs". J. Combin. Math. Comput. 45 (2003), 43-62.
- [9] Robert Molina, "On randomly P_k -decomposable graphs", Congressus Numerantium 148 (2001), 207-221.
- [10] Robert Molina, Myles McNally, Ken Smith: "Characterizing randomly P_k -decomposable graphs for $k \leq 9$ ", Congressus Numerantium 156 (2002), 211-221.
- [11] Sergio Ruiz, "Randomly decomposable graphs", Discrete Mathematics 57(1985), 123-128.
- [12] D.P. Sumner: "Randomly matchable graphs", Journal of Graph Theory 3 (1979), 183-186.
- [13] J. Topp, P.D. Vestergaard, "Totally equimatchable graphs", Discrete Mathematics 164 (1997), 285 – 290.