



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Comparison of metagenomes from EBPR wastewater treatment plants to other environments shows a conserved functional potential

Albertsen, Mads; Saunders, Aaron Marc; Nielsen, Kåre Lehmann; Nielsen, Per Halkjær

*Publication date:*  
2012

*Document Version*  
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Albertsen, M., Saunders, A. M., Nielsen, K. L., & Nielsen, P. H. (2012). *Comparison of metagenomes from EBPR wastewater treatment plants to other environments shows a conserved functional potential*. Poster presented at Danish Conference on Biotechnology and Molecular Biology, Vejle, Denmark.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Comparison of metagenomes from EBPR wastewater treatment plants to other environments shows a conserved functional potential

Mads Albertsen, Aaron M. Saunders, Kåre L. Nielsen and Per H. Nielsen

Department of Biotechnology, Chemistry and Environmental Engineering, Aalborg University, Sohngaardsholmsvej 49, DK-9000 Aalborg, Denmark  
Email: ma@bio.aau.dk





## Introduction

Metagenomics can provide comprehensive information about the entire microbial community of a certain ecosystem, e.g. a wastewater treatment plant. Such studies on full-scale plants are needed to improve the understanding of key microorganisms in the EBPR process and the entire community in order to improve the process for efficient phosphorus removal and recovery.

## Objectives

- Compare the functional potential of two full-scale EBPR plants to other environments by the use of metagenomics.
- Investigate, in detail, similarities and differences between selected available reference genomes and the bacteria present in the full-scale EBPR plants.

## Methods




**Figure 1.** Principle coordinate analysis (PCA) of the functional potential of the two EBPR metagenomes and 39 metagenomes selected to represent a wide range of environments.

The EBPR metagenomes form a distinct cluster when compared to the other environments. However, no single factor could be identified that explained the groupings, underlining the complex selection pressures that form the different microbial communities.

**Figure 2.** Identification of selection pressures on one of the key P-removal bacteria by mapping metagenome reads to the genome.

It can be seen that several parts of the genome have low coverage in both the Aalborg West and East metagenomes (examples highlighted by inserts). Low coverage indicates that these genes are not present in the genomes of the species in the wastewater treatment plants. Many genes involved in exopolymer production and CRISPR phage defense had very low coverage. This indicates that not only CRISPR genes are involved in the direct phage defense, but also the structure of the extracellular matrix is altered in response to phage attacks.



## Conclusions and Further Work

- EBPR wastewater treatment plants show a conserved functional potential when compared to a wide range of environments. However, no single factors could be attributed to the clustering. More metagenomes will enable more detailed investigations.
- The hyper-variability of genes involved in exopolymer production indicates a very active phage defence. Phages might play a large role in shaping the microbial community and the dynamic interplay of bacteria and phages in full-scale EBPR wastewater treatment plants will be interesting to investigate in further details.
- More reference genomes from wastewater treatment plants are needed to make in-depth analysis. We are currently generating the needed genomes by a combination of single cell genomics and metagenomics.