



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

On quasi-greedy bases associated with unitary representations of countable groups

Nielsen, Morten

Publication date:
2013

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Nielsen, M. (2013). *On quasi-greedy bases associated with unitary representations of countable groups*. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2013-05

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

**On quasi-greedy bases associated with
unitary representations of countable groups**

by

Morten Nielsen

R-2013-05

June 2013

DEPARTMENT OF MATHEMATICAL SCIENCES
AALBORG UNIVERSITY

Fredrik Bajers Vej 7G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 99 40 99 40 ■ Telefax: +45 99 40 35 48

URL: <http://www.math.aau.dk>



ON QUASI-GREEDY BASES ASSOCIATED WITH UNITARY REPRESENTATIONS OF COUNTABLE GROUPS

MORTEN NIELSEN

ABSTRACT. We consider the natural generating system for a cyclic subspace of a Hilbert space generated by a dual integrable unitary representation of a countable abelian group. We prove, under mild hypothesis, that whenever the generating system is a quasi-greedy basis it must also be an unconditional Riesz basis. A number of applications to Gabor systems and to general Vilenkin systems are considered. In particular, we show that any Gabor Schauder basis that also forms a quasi-greedy system in $L_2(\mathbb{R})$ is in fact a Riesz basis, and therefore satisfies the classical Balian-Low theorem.

1. INTRODUCTION

Let G be a discrete countable abelian group, and suppose that $g \rightarrow T_g$ is a unitary representation of G on a Hilbert space \mathcal{H} . For fixed $\psi \in \mathcal{H}$, we are interested in properties of the system $\mathcal{Q} := \{T_g\psi : g \in G\}$ in the cyclic subspace

$$(1.1) \quad \mathcal{S} := \langle \psi \rangle = \overline{\text{Span}\{T_g\psi : g \in G\}}^{\mathcal{H}}.$$

Cyclic subspaces are important in many areas of both pure and applied mathematics. One example is in harmonic analysis where systems of the type \mathcal{Q} form the foundation for generating wavelet and Gabor systems, see [1, 5].

For representation systems such as wavelet and Gabor systems, the stability and approximation properties of \mathcal{Q} are important, see e.g. [2, 6, 12, 13]. In this paper we study approximation properties of systems \mathcal{Q} given that \mathcal{Q} has some stability properties. More precisely, we suppose that \mathcal{Q} , with some fixed ordering, forms a Schauder basis for \mathcal{S} with \mathcal{Q} having the additional property that decreasing rearrangements of expansions relative to \mathcal{Q} are norm convergent. Put another way, we assume that the approximants obtained from \mathcal{Q} by thresholding an expansion in the system converge. Thresholding is a very natural way to build approximants, and for Riesz bases it is known to be (up to a constant) the optimal way to build m -terms approximants.

Systems for which decreasing rearrangements are norm convergent are known as quasi-greedy bases. Every unconditional basis is also a quasi-greedy basis, but

2000 Mathematics Subject Classification. 42C10, 41A45.

Key words and phrases. Quasi-greedy bases, dual integrable representation, Gabor systems, integer translates, Vilenkin system.

[†]The author is supported by the Danish Council for Independent Research. Natural Sciences, Grant 12-124675, "Mathematical and Statistical Analysis of Spatial Data".

it is known that conditional quasi-greedy bases exist not only in Banach spaces but also in infinite dimensional Hilbert spaces, see [10] and [16]. However, our main result shows that under mild hypothesis there are no conditional quasi-greedy bases of the type \mathcal{Q} generated by a unitary representation of a countable group. Put another way, if thresholding approximation is convergent, then \mathcal{Q} is an unconditional Riesz basis for \mathcal{S} .

One special case related to this setup has been studied earlier. For a cyclic subspace of $L^2(\mathbb{R}^d)$ generated by integer translates of a single function, it was shown in [13] that no conditional quasi-greedy bases of integer translates exists. However, the abstract approach in the present paper will show that the result in [13] is just one instance of a more general theory that applies to many other “popular” systems in harmonic analysis such as Gabor and Vilenkin systems. Our general result relies on the notion of a dual integrable representation introduced in [9].

The structure of the paper is as follows. In Section 2 we review some needed background material on quasi-greedy systems in a Hilbert space and on dual integrable representations of locally compact groups. Section 3 contains our main result, Theorem 3.3. Finally, Section 4 and Section 5 contain examples related to Gabor and Vilenkin systems, respectively.

2. QUASI-GREEDY SYSTEMS IN A HILBERT SPACE

The present section reviews some needed background material on quasi-greedy systems and on dual integrable representations. The dual integrable representations provide a suitable abstract framework for our study in Section 3 of quasi-greedy systems in cyclic subspaces of the type given by Eq. (1.1).

2.1. Quasi-greedy systems in a Hilbert space. An ordered family $\mathcal{B} = \{x_n : n \in \mathbb{N}\}$ of vectors in a Hilbert space \mathcal{H} is a *Schauder basis* for \mathcal{H} if for every $x \in \mathcal{H}$ there exists a unique sequence $\{\alpha_n := \alpha_n(x)\}_{n \in \mathbb{N}}$ of scalars such that

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N \alpha_n x_n = x$$

in the norm topology of \mathcal{H} . The unique choice of scalars implies that $x \mapsto \alpha_n(x)$ is a linear functional on \mathcal{H} , and by the Riesz representation theorem there exists a unique vector y_n such that $\alpha_n(x) = \langle x, y_n \rangle$. It follows that

$$(2.1) \quad \langle x_m, y_n \rangle = \delta_{m,n}, \quad m, n \in \mathbb{N}.$$

and that there exists a smallest constant $C = C(\mathcal{B}) \geq 1$ such that, for every $n \in \mathbb{N}$,

$$(2.2) \quad 1 \leq \|x_n\| \cdot \|y_n\| \leq C.$$

A pair of sequences $(\{u_n\}_{n \in \mathbb{N}}, \{v_n\}_{n \in \mathbb{N}})$ in \mathcal{H} is a *bi-orthogonal system* if $\langle u_m, v_n \rangle = \delta_{m,n}$, $m, n \in \mathbb{N}$. We say that $\{v_n\}_{n \in \mathbb{N}}$ is a *dual sequence* to $\{u_n\}_{n \in \mathbb{N}}$, and vice versa. A dual sequence is not necessarily uniquely defined. In fact, it is unique if and only if the original sequence is complete in \mathcal{H} (i.e., if the span of the original sequence is dense in \mathcal{H}). A complete sequence $\{x_n : n \in \mathbb{N}\}$ with dual sequence $\{y_n\}$ is a Schauder basis for \mathcal{H} if and only if the partial sum operators

$S_N(x) = \sum_{n=1}^N \langle x, y_n \rangle x_n$ are uniformly bounded on \mathcal{H} . Eq. (2.1) shows that every Schauder basis $\{x_n : n \in \mathbb{N}\}$ for \mathcal{H} has an associated bi-orthogonal system $(\{x_n\}, \{y_n\})$ with a uniquely determined dual sequence. Furthermore, the dual sequence $\{y_n\}$ is also a Schauder basis for \mathcal{H} .

We now fix a biorthogonal system $(x_n, x_n^*)_{n \in \mathbb{N}}$ with $\{x_n\}$ complete in \mathcal{H} . We assume that the system is quasi-normalized, i.e., $\inf_n \|x_n\|_{\mathcal{H}} > 0$ and $\sup_n \|x_n^*\|_{\mathcal{H}} < \infty$, see (2.2). For each $x \in \mathcal{H}$ and $m \in \mathbb{N}$, we define

$$\mathcal{G}_m(x) := \sum_{n \in A} x_n^*(x) x_n,$$

where A is a set of cardinality m satisfying $|x_n^*(x)| \geq |x_k^*(x)|$ whenever $n \in A$ and $k \notin A$. Whenever A is not uniquely defined, we arbitrarily pick any such set. The definition of \mathcal{G}_m leads directly to the definition of a quasi-greedy system, see [10].

Definition 2.1. A quasi-normalized biorthogonal system $(x_n, x_n^*)_{n \in \mathbb{N}} \subset \mathcal{H} \times \mathcal{H}$, with $\text{span}_n(x_n)$ dense in \mathcal{H} , is called a quasi-greedy system if there exists a constant Q such that

$$(2.3) \quad \|\mathcal{G}_m(x)\|_{\mathcal{H}} \leq Q \|x\|_{\mathcal{H}}, \quad \forall x \in \mathcal{H}.$$

If the system is also a Schauder basis for \mathcal{H} , we will use the term quasi-greedy basis.

Remark 2.2. It was proved by Wojtaszczyk in [16] that a system is quasi-greedy if and only if for each $x \in \mathcal{H}$, the sequence $\mathcal{G}_m(x)$ converges to x in norm.

2.2. Dual integrable representations of LCA groups. We now introduce a suitable setting to study the quasi-greedy property in cyclic subspaces of the type given by Eq. (1.1).

We will use the terminology and notation presented in [9]. Suppose that G is a locally compact abelian (LCA) group. We shall use additive notation for G . A character of G is a continuous map $\chi : G \rightarrow \{z \in \mathbb{C} : |z| = 1\}$ for which

$$\chi(g_1 + g_2) = \chi(g_1) \cdot \chi(g_2), \quad g_1, g_2 \in G.$$

The character group of G , denoted by \widehat{G} , is the multiplicative group of all characters of G .

A representation of a LCA group G on a Hilbert space \mathcal{H} is a strongly continuous map $g \rightarrow T_g$ from G into $\mathcal{L}(H, H)$ of bounded linear operators on \mathcal{H} with bounded inverses such that $T_g \circ T_h = T_{g+h}$ for all $g, h \in G$. The representation is called unitary if all the operators T_g are unitary, i.e., $\langle T_g \phi, T_g \psi \rangle = \langle \phi, \psi \rangle$ for all $\phi, \psi \in \mathcal{H}$ and $g \in G$.

Fix a Haar measure χ on \widehat{G} , and let $g \rightarrow T_g$ be a unitary representation of G on a Hilbert space \mathcal{H} . This representation is called dual integrable if there exists a function

$$[\cdot, \cdot] : \mathcal{H} \times \mathcal{H} \rightarrow L^1(\widehat{G}, d\chi)$$

such that

$$(2.4) \quad \langle \phi, T_g \psi \rangle = \int_{\widehat{G}} [\phi, \psi] \overline{\chi(g)} d\chi.$$

An abstract way to interpret dual integrability, which was pointed out in [9], is through Stone's theorem, see e.g. [3, Theorem 4.44]. According to Stone's theorem there is a regular measure P on \widehat{G} with values in the set of self-adjoint projections on \mathcal{H} such that

$$T_g = \int_{\widehat{G}} \chi(g) dP(\chi), \quad g \in G.$$

For $\phi, \psi \in \mathcal{H}$, the function $\mu_{\phi, \psi}(S) := \langle P(S)\phi, P(S)\psi \rangle$ defines a complex valued measure on \widehat{G} such that

$$\langle \phi, T_g \psi \rangle = \int_{\widehat{G}} \overline{\chi(g)} d\mu_{\phi, \psi}(\chi).$$

Comparing this formula to (2.4), we see that dual integrability happens exactly when $d\mu_{\phi, \psi}$ is absolutely continuous wrt. the Haar measure $d\chi$ in which case $[\phi, \psi]$ is the Radon-Nikodym derivative of $d\mu_{\phi, \psi}$.

It follows from [9, Corollary 2.4] that $[\cdot, \cdot]$ is always a sesqui-linear form and a Hermitian symmetric map. In particular, $[T_g \phi, \psi](\chi) = \chi(g)[\phi, \psi](\chi) = [\phi, T_{-g} \psi](\chi)$ a.e. \widehat{G} .

For $\psi \in \mathcal{H} \setminus \{0\}$ one defines $\Omega_\psi = \{\chi \in \widehat{G} : [\psi, \psi](\chi) > 0\}$. Consider the cyclic space $\langle \psi \rangle = \overline{\{T_g \psi : g \in G\}}^{\mathcal{H}}$. It is proved in [9] that S_ψ defined by

$$(2.5) \quad S_\psi(\phi) := \chi_{\Omega_\psi} \frac{[\phi, \psi]}{[\psi, \psi]}$$

is a one-to-one isometry from $\langle \psi \rangle$ onto the weighted space

$$L^2(\Omega_\psi, [\psi, \psi](\chi) d\chi).$$

2.3. Countable abelian groups. We now focus on the restricted case where G is a countable abelian group with the discrete topology. We assume that G is ordered according to $G = \{g_n\}_{n=1}^\infty$, and we let $g \rightarrow T_g$ be a dual integrable representation of G on a Hilbert space \mathcal{H} .

For fixed $\psi \in \mathcal{H}$, we consider the system $\mathcal{Q} := \{T_{g_n} \psi : n \in \mathbb{N}\}$ in the cyclic subspace

$$\mathcal{S} := \langle \psi \rangle = \overline{\{T_{g_n} \psi : n \in \mathbb{N}\}}^{\mathcal{H}}.$$

Notice that \mathcal{Q} is automatically complete in \mathcal{S} . We now suppose \mathcal{Q} forms a Schauder basis for \mathcal{S} . It is easy to verify that completeness is equivalent to $[\psi, \psi](\chi) > 0$ a.e. $[\widehat{G}]$. The Schauder basis property is more delicate. We will discuss the special setup where we have either a Gabor system or a shift invariant system in Section 4.

Now we use the isometry S_ψ given by (2.5) to translate properties of the system \mathcal{Q} to properties of characters.

Lemma 2.3. *Suppose $G = \{g_n\}_{n \in \mathbb{N}}$ is a countable abelian group with $g \rightarrow T_g$ a dual integrable representation of G on a Hilbert space \mathcal{H} . Let S_ψ be defined by (2.5). For any*

finite scalar sequence $\{a_n\}_{n=1}^N$, we have

$$S_\psi \left(\sum_{n=1}^N a_n T_{g_n} \psi \right) (\chi) = \sum_{n=1}^N a_n \chi(g_n).$$

Proof. According to (2.4), we have

$$S_\psi(T_{g_k} \psi)(\chi) = \chi(g_k) \frac{[\psi, \psi]}{[\psi, \psi]} = \chi(g_k),$$

and the result now follows from sesqui-linearity of the bracket $[\cdot, \cdot]$. \square

Since S_ψ is a one-to-one isometry it follows immediately that \mathcal{Q} is a Schauder basis for \mathcal{S} if and only if $\tilde{\mathcal{Q}} := \{\chi(g_n)\}_{n=1}^\infty$ is a Schauder basis for $L^2(\widehat{G}, [\psi, \psi](\chi) d\chi)$. Moreover, \mathcal{Q} is quasi-greedy in \mathcal{S} if and only if $\tilde{\mathcal{Q}}$ is quasi-greedy in

$$L^2(\widehat{G}, [\psi, \psi](\chi) d\chi).$$

The Dirichlet kernel associated with $\tilde{\mathcal{Q}}$ is given by

$$D_N(\chi) = \sum_{n=1}^N \chi(g_n),$$

and it follows that the partial sum operator \widetilde{S}_N associated to $\tilde{\mathcal{Q}}$ is given by

$$\widetilde{S}_N(f)(\alpha) = \int_{\widehat{G}} D_N(\alpha - \chi) f(\chi) d\chi.$$

The following property will be essential to prove Theorem 3.3.

Definition 2.4. With the notation considered in Section 2.3, we say that $\tilde{\mathcal{Q}}$ has the *Fejér property* provided that the kernels $\{F_N(\chi) := |D_N(\chi)|^2/N\}_{N \in \mathbb{N}}$ satisfy the following condition. There exists an increasing sequence $\{N_j\}_{j=1}^\infty \subset \mathbb{N}$ such that for every $n \in \mathbb{N}$ and for a.e. $\alpha \in \widehat{G}$,

$$(2.6) \quad \lim_{j \rightarrow \infty} \int_{\widehat{G}} F_{N_j}(\alpha - \chi) \chi(g_n) d\chi = \alpha(g_n).$$

Notice that since F_N is non-negative with $\int_{\widehat{G}} F_N d\chi = 1$, and the characters $\{\chi(g_n)\}_n$ form a complete system in $L^1(\widehat{G}, d\chi)$, the Fejér property implies that $\{F_{N_j}\}_j$ is an approximate identity for $L^1(\widehat{G}, d\chi)$.

3. PROPERTIES OF QUASI-GREEDY SYSTEMS FOR CYCLIC SUBSPACES

This section contains our main result, Theorem 3.3. The main idea behind Theorem 3.3 is to obtain pointwise information on \widehat{G} for the function $[\psi, \psi]$ by “probing” the function by a suitable approximation of the identity. This idea was introduced in this context in [13]. For another application of this approach see [8, Theorem 4.7]. We first consider some general properties of quasi-greedy system that will be used later to obtain information about the function $[\psi, \psi]$ associated with the cyclic

subspace \mathcal{S} given by (1.1). Quasi-greedy bases are known to be unconditional relative to sign-changes in the following sense.

Lemma 3.1 ([16]). *Suppose $\{e_k\}_{k \in \mathbb{N}}$ is a quasi-greedy system in a Hilbert space \mathcal{H} . Then there exist constants $0 < c_1 \leq c_2 < \infty$ such that for every choice of signs $\epsilon_k = \pm 1$ and any finite subset $A \subset \mathbb{N}$ we have*

$$(3.1) \quad c_1 \left\| \sum_{k \in A} e_k \right\|_{\mathcal{H}} \leq \left\| \sum_{k \in A} \epsilon_k e_k \right\|_{\mathcal{H}} \leq c_2 \left\| \sum_{k \in A} e_k \right\|_{\mathcal{H}},$$

where c_1 and c_2 depend only on the quasi-greedy constant for the system.

For our purpose, Lemma 3.1 is not quite enough. When we consider translates of the Dirichlet kernel, we need to be able to handle arbitrary unimodular complex coefficients and not only ± 1 as covered by Lemma 3.1. For that purpose, we consider the following proposition, which will be essential for the proof of Theorem 3.3.

Proposition 3.2. *Suppose that $\{e_k\}_{k \in \mathbb{N}}$ is a quasi-greedy system in a Hilbert space \mathcal{H} . Then there exist constants $0 < c_1 \leq c_2 < \infty$ such that for every finite unimodular sequence $\{\alpha_k\}_{k=1}^N \subset \mathbb{C}$, we have*

$$(3.2) \quad c_1 N^{1/2} \leq \left\| \sum_{k=1}^N \alpha_k e_k \right\|_{\mathcal{H}} \leq c_2 N^{1/2}.$$

Proof. For technical reasons we define a new scalar sequence $\{\beta_k\}_{k \in \mathbb{N}}$ by

$$\beta_k = \begin{cases} \alpha_k & 1 \leq k \leq N \\ 1, & k > N. \end{cases}$$

Now observe that $\{\beta_k e_k\}_{k \in \mathbb{N}}$ is also a quasi-greedy system in \mathcal{H} . By inspection, we see that the greedy approximation operator $\tilde{\mathcal{G}}_m$ for $\{\beta_k e_k\}_{k \in \mathbb{N}}$ is identical to the approximation operator \mathcal{G}_m for $\{e_k\}_{k \in \mathbb{N}}$. This follows from the trivial observation that if f_k is the dual element to e_k , then $\beta_k f_k$ is the dual element to $\beta_k e_k$, since $|\beta_k| = 1$.

Next we recall that the Hilbert space \mathcal{H} has Rademacher type and cotype 2, see [15, §III.A]. Hence, for any sequence $\{f_\ell\}_{\ell \in \mathbb{N}} \subset \mathcal{H}$, we have the uniform estimate

$$(3.3) \quad \int_0^1 \left\| \sum_{\ell=1}^n r_\ell(t) f_\ell \right\|_{\mathcal{H}}^2 dt \asymp \sum_{\ell=1}^n \|f_\ell\|_{\mathcal{H}}^2,$$

where $\{r_\ell\}_{\ell \in \mathbb{N}}$ is the sequence of Rademacher functions on $[0, 1]$. In particular,

$$(3.4) \quad \int_0^1 \left\| \sum_{k=1}^N r_k(t) [\beta_k e_k] \right\|_{\mathcal{H}}^2 dt = \text{Avg}_{\epsilon_k=\pm 1} \left\| \sum_{k=1}^N \epsilon_k [\beta_k e_k] \right\|_{\mathcal{H}}^2.$$

We use (3.4) and (3.3), together with Lemma 3.1 applied to the quasi-greedy system $\{\beta_k e_k\}_{k \in \mathbb{N}}$, to obtain

$$\begin{aligned}
\left\| \sum_{k=1}^N \alpha_k e_k \right\|_{\mathcal{H}}^2 &\asymp \text{Avg}_{\epsilon_k = \pm 1} \left\| \sum_{k=1}^N \epsilon_k [\beta_k e_k] \right\|_{\mathcal{H}}^2 \\
&\asymp \sum_{k=1}^N \|\beta_k e_k\|_{\mathcal{H}}^2 \\
(3.5) \quad &\asymp N.
\end{aligned}$$

□

With Lemma 3.2 in hand we can now turn to our main result.

Theorem 3.3. *Let G be a countable abelian group G with an ordering given by $G = \{g_n\}_{n=1}^\infty$. Let $g_n \rightarrow T_{g_n}$ be a dual integrable unitary representation on a Hilbert space \mathcal{H} . Let $\psi \in \mathcal{H}$, and suppose that $\mathcal{Q} = \{T_{g_n}\psi : n \in \mathbb{N}\}$ is a quasi-greedy system in the cyclic space $\langle \psi \rangle$. Then \mathcal{Q} is in fact a Riesz basis for $\langle \psi \rangle$, i.e., there exist constants $0 < c \leq C < \infty$ such that*

$$(3.6) \quad c\|\{c_n\}\|_{\ell_2} \leq \left\| \sum_{n \in \mathbb{N}} c_n T_{g_n} \psi \right\|_{\mathcal{H}} \leq C\|\{c_n\}\|_{\ell_2},$$

for every finite sequence $\{c_n\}_{n \in \mathbb{N}}$.

Proof. The idea is to consider suitably translated Dirichlet kernels $D_N(\chi) = \sum_{n=1}^N \chi(g_n)$, $\chi \in \widehat{G}$, since they form “polynomials” with unimodular coefficients. In fact for fixed $\alpha \in \widehat{G}$,

$$\chi \rightarrow D_N(\alpha - \chi) = \sum_{n=1}^N \alpha(g_n) \overline{\chi(g_n)},$$

is a polynomial with unimodular coefficients since $|\alpha(g_k)| = 1$ for every $g_k \in G$. Now, according to by Proposition 3.2,

$$S_N := \sum_{n=1}^N \alpha(g_n) T_{g_n} \psi$$

satisfies the norm estimate $\|S_N\|_{\mathcal{H}}^2 \asymp N$ uniformly in N and $\alpha \in \widehat{G}$. From the linearity and sesqui-linearity of the bracket,

$$[S_N, S_N](\chi) = |D_N(\alpha - \chi)|^2 [\psi, \psi](\chi).$$

The system $\{T_{g_n}\psi : n \in \mathbb{N}\}$ is clearly complete in $\langle \psi \rangle$ since it forms a quasi-greedy Schauder basis, and one easily checks that $\Omega_\psi = \{\chi \in \widehat{G} : [\psi, \psi](\chi) > 0\} = \widehat{G}$. Hence, according to (2.5),

$$\|S_N\|_{\mathcal{H}} = \|S_\psi(S_N)\|_{L_2(\widehat{G}, [\psi, \psi] d\chi)},$$

which implies that,

$$N \asymp \|S_N\|^2 = \int_{\widehat{G}} |D_N(\alpha - \chi)|^2 [\psi, \psi](\chi) d\chi,$$

uniformly in N and $\alpha \in \widehat{G}$. Then clearly,

$$(3.7) \quad 1 \asymp \int_{\widehat{G}} \frac{|D_N(\alpha - \chi)|^2}{N} [\psi, \psi](\chi) d\chi, \quad N \in \mathbb{N}.$$

By the Fejér property (2.6), we have that for a.e. $\alpha \in \widehat{G}$,

$$(3.8) \quad \int_{\widehat{G}} \frac{|D_{N_j}(\alpha - \chi)|^2}{N_j} [\psi, \psi](\chi) d\chi \rightarrow [\psi, \psi](\alpha), \quad \text{as } j \rightarrow \infty,$$

for some subsequence $\{N_j\}_j \subset \mathbb{N}$. It follows at once from (3.7), and the pointwise convergence (3.8), that there exist $c, C > 0$ such that

$$(3.9) \quad c \leq [\psi, \psi](\alpha) \leq C, \quad \text{a.e. } \alpha \in \widehat{G}.$$

It is well-known that (3.9) implies that $\{T_{g_n}\psi : n \in \mathbb{N}\}$ satisfies (3.6), and the system therefore forms a Riesz basis for $\langle \psi \rangle$. \square

4. EXAMPLES: THE TRIGONOMETRIC SYSTEM

We first consider two examples related to the trigonometric system in $L^2(\mathbb{T}^d)$. The trigonometric system forms the dual group of continuous characters for the group of integers \mathbb{Z}^d .

The trigonometric system forms an unconditional basis for $L^2(\mathbb{T}^d)$, but when a weight is added the situation becomes more complicated. For weighted spaces, the specific ordering of the system may become important. We will focus on the following class of orderings of the trigonometric system.

For $\mathbf{a} \in \mathbb{N}^d$ we define the corresponding rectangle by

$$R_{\mathbf{a}} := \{-a_1, \dots, a_1\} \times \cdots \times \{-a_d, \dots, a_d\} \subset \mathbb{Z}^d.$$

We say that an ordering $\sigma : \mathbb{N} \rightarrow \mathbb{Z}^d$ is *adapted to rectangles* if there exist an increasing sequence $\{N_j\}_{j=1}^\infty \subset \mathbb{N}$, and associated rectangles $R_{\mathbf{a}_1} \subset R_{\mathbf{a}_2} \subset \cdots$, such that $\mathbb{Z}^d = \bigcup_j R_{\mathbf{a}_j}$ and

$$\sigma(\{1, 2, \dots, N_j\}) = R_{\mathbf{a}_j}, \quad j = 1, 2, \dots$$

It is not difficult to verify that a trigonometric system $\{e^{2\pi i k \cdot \xi}\}_{k \in \mathbb{Z}^d}$ with an ordering $\sigma : \mathbb{N} \rightarrow \mathbb{Z}^d$ adapted to rectangles satisfies the Fejér condition (2.6). In fact, the associated kernels are separable

$$F_{N_j}(x) = \frac{1}{N_j} \left| \sum_{j=1}^{N_j} e^{2\pi i \sigma(j) \cdot x} \right|^2 = \frac{1}{N_j} \prod_{k=1}^d \mathcal{F}_{a_{j,k}}(x_k),$$

with \mathcal{F}_n the usual univariate Fejér kernel. Specific examples of orderings $\sigma : \mathbb{N} \rightarrow \mathbb{Z}^d$ adapted to rectangles can be found in [11], see also [12].

The Muckenhoupt A_2 condition for rectangles plays a crucial role when it comes to stability of the trigonometric system in weighted L^2 -spaces. We say that a function $w : \mathbb{T}^d \rightarrow (0, \infty)$ is an $A_{2,R}(\mathbb{T}^d)$ weight provided that there exists $C < \infty$ such

that for any rectangle $R \subset \mathbb{T}^d$,

$$(4.1) \quad \frac{1}{|R|} \int_R w(x) dx \cdot \frac{1}{|R|} \int_R \frac{1}{w(x)} dx \leq C.$$

4.1. Integer translates. The map $\mathbb{Z}^d \ni k \longrightarrow T_k$ given by

$$(4.2) \quad T_k f(x) := f(x + k)$$

is a unitary representation of the integer group $(\mathbb{Z}^d, +)$ on $L^2(\mathbb{R}^d)$. The representation is dual integrable, which follows from the observation that

$$\begin{aligned} \langle \phi, T_k \psi \rangle_{L^2(\mathbb{R}^d)} &= \int_{\mathbb{R}^d} \phi(x) \overline{\psi(x+k)} dx \\ &= \int_{\mathbb{R}^d} \hat{\phi}(\xi) \overline{\hat{\psi}(\xi)} e^{2\pi i k \cdot \xi} d\xi \\ &= \int_{[0,1]^d} \left\{ \sum_{\ell \in \mathbb{Z}^d} \hat{\phi}(\xi + \ell) \overline{\hat{\psi}(\xi + \ell)} \right\} e^{2\pi i k \cdot \xi} d\xi, \end{aligned}$$

so one may choose

$$[\phi, \psi](\xi) := \sum_{\ell \in \mathbb{Z}^d} \hat{\phi}(\xi + \ell) \overline{\hat{\psi}(\xi + \ell)}, \quad \phi, \psi \in L^2(\mathbb{R}^d).$$

Now let $\psi \in L^2(\mathbb{R}^d)$. It is known that a sequence of partial sum operators for the trigonometric system $\mathcal{T} := \{e^{2\pi i k \cdot \xi}\}_{k \in \mathbb{Z}^d}$ adapted to rectangles is uniformly bounded on $L^2(\mathbb{T}^d, [\psi, \psi])$ if $[\psi, \psi]$ satisfies the product $A_{2,\mathcal{R}}$ -condition (4.1). With a suitable interpretation a converse statement also holds true, see [11, 12] for details. It follows that for a class of orderings of \mathbb{Z}^d adapted to rectangles, $[\psi, \psi] \in A_{2,\mathcal{R}}$ is sufficient for the trigonometric system to form a Schauder basis for

$$L^2(\mathbb{T}^d, [\psi, \psi]).$$

If in addition, \mathcal{T} forms a quasi-greedy basis for $L^2(\mathbb{T}^d, [\psi, \psi])$, or equivalently that $\mathcal{S} := \{\psi(\cdot - k)\}_{k \in \mathbb{Z}^d}$ is a quasi-greedy system in $\langle \psi \rangle$, then it follows from Theorem 3.3 that \mathcal{T} (resp. \mathcal{S}) in fact forms a Riesz basis in $L^2(\mathbb{T}^d, [\psi, \psi])$ (resp. in $\langle \psi \rangle$). This is exactly the main result obtained in [13].

4.2. Gabor systems. The Gabor representation of the LCA group $(\mathbb{Z}^d \times \mathbb{Z}^d, +)$ on $L^2(\mathbb{R}^d)$ is given by

$$\mathbb{Z}^d \times \mathbb{Z}^d \ni (m, n) \longrightarrow T_m M_n f(x) = f(x + m) e^{2\pi i n \cdot x},$$

with T_k given by (4.2) and M_ℓ is the modulation operator $M_\ell f(x) = f(x) e^{2\pi i \ell \cdot x}$. The Gabor representation is dual integrable through the Zak transform. The Zak transform is an isometry $Z : L^2(\mathbb{R}^d) \rightarrow L^2([0,1]^d \times [0,1]^d)$ given by

$$Zf(x, \xi) = \sum_{k \in \mathbb{Z}^d} f(x + k) e^{2\pi i k \cdot \xi}, \quad f \in L^2(\mathbb{R}^d).$$

One easily checks that

$$Z(M_\ell T_k \psi)(x, \xi) = e^{2\pi i l \cdot x} e^{2\pi i k \cdot \xi} Z\psi(x, \xi),$$

and it follows that for $\phi, \psi \in L^2(\mathbb{R}^d)$,

$$\langle \phi, M_\ell T_k \psi \rangle_{L^2(\mathbb{R}^d)} = \int_{[0,1]^d} \int_{[0,1]^d} Z\phi(x, \xi) \overline{Z\psi(x, \xi)} e^{-2\pi i l \cdot x} e^{-2\pi i k \cdot \xi} dx d\xi.$$

Consequently, we choose

$$[\phi, \psi](x, \xi) := Z\phi(x, \xi) \overline{Z\psi(x, \xi)}, \quad \phi, \psi \in L^2(\mathbb{R}^d).$$

Fix $\psi \in L^2(\mathbb{R}^d)$. It was demonstrated for $d = 1$ in [7], and for $d > 1$ in [11], that $|Z\psi|^2(x, \xi) \in A_{2,R}(\mathbb{T}^d \times \mathbb{T}^d)$ is sufficient for the associated Gabor system

$$\mathcal{G}(\psi) := \{M_\ell T_k \psi\}_{\ell, k \in \mathbb{Z}^d}$$

with a suitable ordering adapted to rectangles to form a Schauder basis for $L^2(\mathbb{R}^d)$. We can now apply Theorem 3.3 to obtain the following result.

Corollary 4.1. *Let $\psi \in L^2(\mathbb{R}^d)$. Suppose that the associated Gabor system $\mathcal{G}(\psi)$ forms a quasi-greedy basis for $L^2(\mathbb{R}^d)$ with an ordering adapted to rectangles. Then $\mathcal{G}(\psi)$ forms a Riesz basis for $L^2(\mathbb{R}^d)$.*

An immediate consequence of Corollary 4.1 is that whenever $\mathcal{G}(\psi)$ forms a quasi-greedy basis for $L^2(\mathbb{R}^d)$, with an ordering adapted to rectangles, then $\psi \in L^2(\mathbb{R}^d)$ must satisfy the Balian-Low theorem, namely

$$(4.3) \quad \int_{\mathbb{R}^d} |x|^2 |\psi(x)|^2 dx \cdot \int_{\mathbb{R}^d} |\xi|^2 |\hat{\psi}(\xi)|^2 d\xi = +\infty,$$

due to the fact that $\mathcal{G}(\psi)$ is a Riesz basis.

It is also interesting to note that there exists a conditional Gabor Schauder basis for $L^2(\mathbb{R}^d)$ for which the generator does not satisfy the strong form of the Balian-Low theorem (4.3), see [7]. According to Corollary 4.1, such systems cannot form a quasi-greedy basis for $L^2(\mathbb{R}^d)$.

5. EXAMPLE: VILENIN GROUPS

We now leave the trigonometric system setup and consider an example related to another well known system of group characters, namely the Walsh system.

Let $m = (m_k)_{k \in \mathbb{N}}$ be a sequence of positive integers, and let \mathbb{Z}_{m_k} be the m_k -th discrete cyclic group with Haar measure defined by assigning mass $1/m_k$ to each singleton. The associated Vilenkin group is the product group

$$\widehat{G}_m := \prod_{k=1}^{\infty} \mathbb{Z}_{m_k}.$$

For each $n \in \mathbb{N}$, and $x = (x_k)_{k \in \mathbb{N}} \in \widehat{G}_m$, we set

$$\rho_n(x) := \exp\left(\frac{2\pi i x_n}{m_n}\right).$$

Clearly, ρ_n is a character on \widehat{G}_m , and it can in fact be shown that any character of \widehat{G}_m can be expressed in terms of the ρ'_n 's. Put $M_0 := 1$ and for $k \in \mathbb{N}$ we define inductively $M_k := m_{k-1}M_{k-1}$. Then for $n \in \mathbb{N}$ we have a unique finite expansion

$$(5.1) \quad n = \sum_{k=0}^{\infty} n_k M_k,$$

where $0 \leq n_k < m_k$ for $k = 0, 1, 2, \dots$. Define $\psi_0 := 1$, and for $n \in \mathbb{N}$ given by (5.1), we let

$$(5.2) \quad \psi_n := \prod_{k=0}^{\infty} \rho_k^{n_k}.$$

The system $G_m := \{\psi_k\}_{k=0}^{\infty}$ is called a Vilenkin system and forms a complete orthonormal set of characters for \widehat{G}_m , see [14]. The well-known Walsh system corresponds to the special case $m_k := 2$, $k \in \mathbb{N}$.

The associated Dirichlet kernel are given by $D_n := \sum_{k=0}^{n-1} \psi_k$. The exact pointwise values of D_{M_n} are known, see [14, Appendix 0.7], and it can be deduced that G_m satisfies the Fejér property for the subsequence with indices $\{M_n\}_n$. The question about pointwise convergence a.e. relative to a Vilenkin system (i.e., not only for a subsequence) is more delicate, especially in the so-called unbounded case where $\sup_k m_k = \infty$, see [4].

The modular representation of $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ on $L^2(\widehat{G}_m, d\chi)$ is given by modulation operators

$$(5.3) \quad \mathbb{N}_0 \ni n \rightarrow M_n f := \psi_n f, \quad f \in L^2(\widehat{G}_m, d\chi).$$

The modular representation is dual integrable with $[\phi, \psi]_M := \phi \bar{\psi}$, since the product of any two $L^2(\widehat{G}_m, d\chi)$ -functions is in $L^1(\widehat{G}_m, d\chi)$. Moreover,

$$\langle \phi, M_n \psi \rangle = \int_{\widehat{G}} [\phi, \psi]_M \overline{\psi_n(\chi)} d\chi, \quad n \in \mathbb{N}_0.$$

We now apply Theorem 3.3 to obtain the following result, which concludes the paper.

Corollary 5.1. *Let $m = (m_k)_{k \in \mathbb{N}}$ be a sequence of positive integers, and let \widehat{G}_m be the associated Vilenkin group. Take $\psi \in L^2(\widehat{G}_m, d\chi)$, and suppose that the associated Vilenkin system G_m forms a quasi-greedy basis for $\langle \psi \rangle$ generated by the representation (5.3) with the ordering defined by (5.2). Then G_m forms a Riesz basis for $\langle \psi \rangle$.*

In particular, if G_m forms a quasi-greedy basis for $L^2(\widehat{G}_m, |\psi|^2 d\chi)$ then G_m is a Riesz basis for $L^2(\widehat{G}_m, |\psi|^2 d\chi)$ and there exists $c > 1$ such that

$$c^{-1} \leq |\psi(x)| \leq c, \quad a.e. x \in \widehat{G}_m.$$

REFERENCES

- [1] I. Daubechies. *Ten lectures on wavelets*, volume 61 of CBMS-NSF Regional Conference Series in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1992.

- [2] C. de Boor, R. A. DeVore, and A. Ron. Approximation orders of FSI spaces in $L_2(\mathbf{R}^d)$. *Constr. Approx.*, 14(4):631–652, 1998.
- [3] G. B. Folland. *A course in abstract harmonic analysis*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995.
- [4] G. Gát. Some convergence and divergence results with respect to summation of Fourier series on one and two-dimensional unbounded Vilenkin groups. *Ann. Univ. Sci. Budapest. Sect. Comput.*, 33:157–173, 2010.
- [5] K. Gröchenig. *Foundations of time-frequency analysis*. Applied and Numerical Harmonic Analysis. Birkhäuser Boston Inc., Boston, MA, 2001.
- [6] K. Gröchenig and S. Samarah. Nonlinear approximation with local Fourier bases. *Constr. Approx.*, 16(3):317–331, 2000.
- [7] C. Heil and A. M. Powell. Gabor Schauder bases and the Balian-Low theorem. *J. Math. Phys.*, 47(11):113506, 21, 2006.
- [8] E. Hernández, M. Nielsen, H. Šikić, and F. Soria. Democratic systems of translates. *J. Approx. Theory*, 171:105–127, 2013.
- [9] E. Hernández, H. Šikić, G. Weiss, and E. Wilson. Cyclic subspaces for unitary representations of LCA groups; generalized Zak transform. *Colloq. Math.*, 118(1):313–332, 2010.
- [10] S. V. Konyagin and V. N. Temlyakov. A remark on greedy approximation in Banach spaces. *East J. Approx.*, 5(3):365–379, 1999.
- [11] K. Moen. Multiparameter weights with connections to Schauder bases. *J. Math. Anal. Appl.*, 371(1):266–281, 2010.
- [12] M. Nielsen. On stability of finitely generated shift-invariant systems. *J. Fourier Anal. Appl.*, 16(6):901–920, 2010.
- [13] M. Nielsen and H. Šikić. Quasi-greedy systems of integer translates. *J. Approx. Theory*, 155(1):43–51, 2008.
- [14] F. Schipp, W. R. Wade, and P. Simon. *Walsh series*. Adam Hilger Ltd., Bristol, 1990. An introduction to dyadic harmonic analysis, With the collaboration of J. Pál.
- [15] P. Wojtaszczyk. *Banach spaces for analysts*, volume 25 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1991.
- [16] P. Wojtaszczyk. Greedy algorithm for general biorthogonal systems. *J. Approx. Theory*, 107(2):293–314, 2000.

DEPARTMENT OF MATHEMATICAL SCIENCES, AALBORG UNIVERSITY, FREDRIK BAJERS VEJ 7G,
DK - 9220 AALBORG EAST, DENMARK