

All P_3 -equipackable graphs

Randerath, Bert; Vestergaard, Preben D.

Publication date:
2008

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Randerath, B., & Vestergaard, P. D. (2008). *All P_3 -equipackable graphs*. Department of Mathematical Sciences, Aalborg University. Research Report Series No. R-2008-10

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

All P_3 -equipackable graphs

by

Bert Randerath and Preben Dahl Vestergaard

To appear in *Discrete Mathematics*

R-2008-10

October 2008

DEPARTMENT OF MATHEMATICAL SCIENCES
AALBORG UNIVERSITY

Fredrik Bajers Vej 7 G ■ DK-9220 Aalborg Øst ■ Denmark

Phone: +45 99 40 80 80 ■ Telefax: +45 98 15 81 29

URL: <http://www.math.aau.dk>



All P_3 -equipackable graphs

BERT RANDERATH

Institut für Informatik

Universität zu Köln

D-50969 Köln, Germany

rand Rath@informatik.uni-koeln.de

PREBEN DAHL VESTERGAARD

Department of Mathematical Sciences

Aalborg University

DK-9220 Aalborg Ø, Denmark

pdv@math.aau.dk

October 7, 2008

Accepted for publication in Discrete Mathematics

Abstract

A graph G is P_3 -equipackable if any sequence of successive removals of edge-disjoint copies of P_3 from G always terminates with a graph having at most one edge. All P_3 -equipackable graphs are characterised. They belong to a small number of families listed here.

Keywords: Packing, equipackable, randomly packable, covering, factor, decomposition, equiremovable

2000 Mathematics Subject Classification: 05C70, 05C35

1 Introduction

Let H be a subgraph of a graph G . An H -packing in G is a partition of the edges of G into disjoint sets, each of which is the edge set of a subgraph of G isomorphic to H , and possibly a remainder set. For short, $E(G)$ is partitioned into copies of H and maybe a remainder set. We list some references to an extensive literature at the back. A graph is called H -packable if G is the union of edge disjoint copies of H . An H -packing is *maximal* if the remainder set of edges is empty or contains no copy of H . An H -packing is *maximum* if $E(G)$ has been partitioned into a maximum number of sets isomorphic to H and a possible remainder set. A graph is called H -equipackable if any maximal H -packing is also a maximum H -packing, i.e., a graph G is H -equipackable if successive removals of copies of H from G can be done the same number of times regardless of the particular choices of edge sets for H in each step. If every maximal H -packing of a graph G uses all edges of G , then G is called *randomly H -packable*. Equivalently, G is randomly H -packable if each H -packing can be extended to an H -packing containing all edges of G , see e.g. [1, 2, 5, 6].

Zhang and Fan [9] have studied H -equipackable graphs for the case $H = 2K_2$. We shall consider path packing and in the following H will always be assumed to be the graph P_3 , the path of length two, and we may omit explicit reference to it. A graph G is then (P_3 -) equipackable if successive removals of two adjacent edges from G can be done the same number of times

regardless of the particular choices of edge pairs in each step. A component consisting of one vertex is called *trivial*, a *non-trivial* component contains an edge. A graph has *order* $|V(G)|$ and *size* $|E(G)|$. A graph of odd (even) size is called *odd* (even). A vertex of valency one is called a *leaf*. A star is called even if its size is even, and by $K_{1,2k}$ we denote the even star with $2k$ leaves.

Observation 1 *A graph is randomly H -packable if and only if it is H -packable and H -equipackable.*

S. Ruiz [7] characterised randomly P_3 -packable graphs.

Theorem 2 *A connected graph G is randomly packable if and only if $G \cong C_4$ or $G \cong K_{1,2k}$, $k \geq 1$.*

Y. Caro, J. Schönheim [3] and S. Ruiz [7] stated the following result.

Lemma 3 *A connected graph is packable if and only if it has even size.*

This immediately implies Corollary 4 below.

Corollary 4 *If a connected graph is equipackable, a maximal packing either contains all edges or all but one edge of the graph.*

From B.L. Hartnell, P.D. Vestergaard [4] and P.D. Vestergaard [8] we have the following observation.

Observation 5 *Let G be an equipackable graph. Then any sequence of P_3 -removals from G will produce an equipackable graph.*

From Corollary 4 and Observation 5 we obtain

Corollary 6 *Let G be a connected graph. If there is a sequence of P_3 -removals from G that creates more than one component of odd size, then G is not equipackable.*

We now state our main result, a characterisation of all equipackable graphs with at most one non-trivial component:

Theorem 7 *Let $G = (V, E)$ be a graph with at most one non-trivial component. Then G is equipackable if and only if its non-trivial component belongs to one of the thirteen families listed in Figure 1 or can be obtained by a sequence of P_3 -removals from such a graph.*

Clearly, we wish those thirteen families listed to be *maximal w.r.t. P_3 -removals*, i.e., no graph from one of the families can be obtained as a subgraph of a larger equipackable graph by removing a P_3 from it.

In the figures below we indicate by an arrow from which family of graphs we may obtain the given graph by a sequence of P_3 -deletions. The shaded vertex sets may vary in cardinality.




Figure 1: All connected, maximal with respect to P_3 -removal, P_3 -equipackable graphs

We will prove this characterisation in the following section.

2 Proof of Theorem 7

By Lemma 3 and Theorem 2 a graph with at most one non-trivial component, which has even size, is equipackable if and only if its non-trivial component is a 4-circuit or an even star (Figure 2). Thus it only remains to characterise equipackable graphs with at most one non-trivial component of odd size.




Figure 2: Connected P_3 -equipackable graphs of even size (Ruiz graphs)

In [8] P.D. Vestergaard examined equipackable graph with all degrees ≥ 2 and stated the following result.

Theorem 8 *A connected graph G with all degrees ≥ 2 is equipackable if and only if G is one of the graphs listed in Figure 3.*




Figure 3: All connected P_3 -equipackable graphs G without leaves

Observe that this solution contributes to our characterisation five graphs (F_6, F_3, F_4, F_5, F_9) maximal with respect to P_3 -removals. All other graphs of this solution are obtained by a sequence of P_3 -removals from graphs of the thirteen graph families of our characterisation. Thus it now remains to characterise equipackable graphs G which have only one non-trivial component, say H , where H has odd size and contains a leaf.

Since H has a leaf, it also has a bridge. Let $b = xy$ be a bridge of H . Throughout we shall denote the two components of $H - xy$ by H_1 and H_2 with $x \in V(H_1), y \in V(H_2)$. We shall first treat the case that G has a non-leaf bridge, then the case that all bridges are leaf bridges.

Case 1: Assume $b = xy$ is a non-leaf bridge of G , i.e., $\deg(x) \geq 2, \deg(y) \geq 2$.

Subcase 1.1: Assume further that H has a maximum P_3 -packing \mathcal{P} which does not contain b . Since \mathcal{P} by Corollary 4 contains all but one edge of G and $b \notin \mathcal{P}$, we have for $i = 1, 2$ that $\mathcal{P} \cap H_i$ is a P_3 -packing of H_i and therefore H_i has even size ≥ 2 .

Let $z \in N(x) \setminus \{y\}$. By P_3 -removal of zxy we obtain an equipackable graph which has an odd component contained in $H_1 - xz$, and $H - \{zx, xy\}$ also has the even component H_2 which is connected, randomly packable and hence, by Observation 1, is either a 4-circuit or an even star. By symmetry also H_1 is a 4-circuit or an even star. Therefore H belongs to one of the families of graphs depicted in Figure 4.




Figure 4: Connected, P_3 -equipackable graphs in Case 1.1

Note that only three new graph families (F_7, F_8, F_{10}) maximal with respect to P_3 -removals contribute in this case to our characterisation. All other graph families of this subcase are obtained by a sequence of P_3 -removals from graphs of the thirteen graph families of our characterisation.

Subcase 1.2: Assume now that each non-leaf bridge of H is contained in every maximum P_3 -packing.

With notation as above let $b = xy$ be a non-leaf bridge of H , the components of $H - xy$ are H_1, H_2 . Their sizes have the same parity since H has odd size. If H_1, H_2 both had even size they would be P_3 -packable and H would have a maximum P_3 -packing not containing b in contradiction to assumption. Therefore H_1, H_2 both have odd size.

CLAIM: At least one of H_1, H_2 is an odd star.

Proof. P_3 -removal from H of zxy , $z \in N(x) \setminus \{y\}$, creates an odd size component, namely H_2 . If H_2 is an odd star we are finished. Otherwise, we can isolate an odd component inside H_2 : If $\deg_{H_2}(y)$ is even we P_3 -remove all edges incident to y in pairs and if \deg_{H_2} is odd we P_3 -remove all but one edge incident to y in pairs and that remaining edge yw , $w \in N(y)$, together with wr , $r \in N(w) \setminus \{y\}$ (Since H_2 is not an odd star there has to exist at least one such vertex w). Then $H_1 \cup \{xy\}$ is even, connected, randomly packable and hence is either a 4-circuit or an even star. Since $H_1 \cup \{xy\}$ contains a leaf, it is an even star and hence H_1 is an odd star. That proves the claim. \square

Suppose H_1 and H_2 are both odd stars. Now assume that, say x , is not the center of H_1 and let v be the center of H_1 . Since vx is a non-leaf bridge and there obviously exists a maximum P_3 -packing \mathcal{P} which does not contain vx , we obtain a contradiction to the assumption of Subcase 1.2. Hence we find that H is obtained by adding an edge between the centers of H_1 and H_2 (see Figure 5). Consequently H can be obtained from one of the graphs of the family F_{12} in our characterisation by P_3 -deletions.

If, say, H_2 is an odd star and H_1 is not, then P_3 -removal of zxy from H , $z \in N(x) \setminus \{y\}$, gives that $H_1 - xz$ has even size.

Now assume that zx is a leaf bridge of H (and likewise of H_1), i.e., $\deg_H(z) = 1$.

Then P_3 -removal of zxy leaves the odd component H_2 and $H_1 - xz$ with one non-trivial even component. Thus the non-trivial even component of $H_1 - xz$ is either a 4-circuit or an even star. The former yields easily a non-equipackable graph, the latter gives that H_1 is an odd star, a contradiction to assumption on H_1 .

Suppose now that zx is a non-leaf bridge of H (and likewise for H_1).

The two components of $H_1 - xz$ have sizes of same parity. That cannot be odd since $G - zxy$ would then have three odd components in contradiction to Corollary 6. It cannot be even either because then we could easily construct a maximum P_3 -packing \mathcal{P} which does not contain the non-leaf bridge xz , a contradiction to the basic assumption of this subcase.

So we may for all $z \in N(x) \setminus \{y\}$ assume that xz is not a bridge of H (and H_1).

P_3 -removal of zxy for $z \in N(x) \setminus \{y\}$ produces the connected, even component $H_1 - xz$ which is then randomly P_3 -packable and hence is either an even star or a 4-circuit. If $H_1 - xz$ is a 4-circuit we are immediately led to H not being equipackable because, if a, b, c, d are the edges of this 4-circuit (in cyclic order) then the packing $\{xy, a\}, \{xz, c\}$ cannot be extended to a maximum packing of H . Observe that we have $N(x) \setminus \{y\} = \{z_1, z_2, \dots, z_p\}$ with $z_1 = z$ and $p \geq 2$. Thus for all $z_i \in N(x) \setminus \{y\}$ the connected graph $H_1 - xz_i$ is an even star. It follows that $p = 2$ and $H_1 - xz_i$ must always be isomorphic to a $P_3 = K_{1,2}$ with a center vertex z_{3-i} having neighbours x and z_i for $i = 1, 2$. Thus H_1 is a 3-circuit with vertices x, z_1, z_2 with x joined to y , and y has an odd number of leaves attached (see Figure 5).




Figure 5: Connected P_3 -equipackable graphs in Case 1.2

Observe that none of these equipackable graph families are new families maximal with respect to P_3 -removals for our characterisation. Both graph families of this subcase are obtained by a sequence of P_3 -removals from graphs of the graph families (F_{12}, F_{13}) of our characterisation. We may now assume that there exist no non-leaf bridge of H .

Case 2: All bridges of H are leaf bridges and there exists at least one bridge $b = xy$ of H , i.e. $H_2 = \{y\}$.

If all xz , $z \in N(x) \setminus \{y\}$, are bridges of H , then they are leaf bridges and H is an odd star, derivable from a member of our characterisation by a sequence of P_3 -removals. Thus we may assume that x is contained in at least one cycle of H_1 and there exist at least two edges incident to x , which are not bridges.

If x has an even number of neighbours in H_1 we can isolate xy by pairing up and P_3 -removing all xz , $z \in N(x) \setminus \{y\}$. If x has an odd number of neighbours in H_1 we isolate xy by P_3 -removing

all $xz, z \in N(x) \setminus \{y\}$, and one further edge $zw, w \in N(z) \setminus \{x\}$ (observe that such an edge has to exist). For simplicity, let E' be the set of edges of all P_3 's necessary to remove in order to isolate the bridge xy and $H' = H - E'$. Since xy is isolated in H' and H' is equipackable, we obtain by Lemmas 3, Observation 5 and Corollaries 4, 6 that all non-trivial components D not containing x and y are randomly packable and therefore of even size ≥ 2 . Thus every such non-trivial component D is either a 4-circuit or an even star.

Assume that one of these components is a 4-circuit C with vertices $\{c_0, c_1, c_2, c_3\}$ and edges $\{c_i c_{(i+1) \bmod 4} | 0 \leq i \leq 3\}$.

As all bridges of H are leaf bridges, with $E_C = \{xc | c \in N_{H_1}(x) \cap V(C)\}$ we have $|E_C| \geq 2$. It is easy to see that we can remove two (if $|E(C)| = 2$) or three P_3 's from the subgraph of H induced by $\{x\} \cup V(C)$ to produce two (if $|E(C)| = 3$) or three isolated edges (including xy) in contradiction with Corollary 6.

If $|E_C| = 2$ there exist $i, j, k, \ell = \{0, 1, 2, 3\}$ such that $xc_i, xc_j \in E_C$ and $xc_i c_k, xc_j c_\ell$ are P_3 's of H that isolate the two independent edges $c_i c_k, c_j c_\ell$ remaining in C . By Corollary 6 then H is not equipackable, a contradiction. If $|E_C| = 3$, without loss of generality we may assume that $E_C = \{xc_0, xc_1, xc_2\}$ and in that case $E' \cup \{c_0 c_3, c_1 c_2\}$ is an edge set of even size, which can be paired up in P_3 's whose removal isolate two edges $c_0 c_1$ and $c_2 c_3$ on C , by Corollary 6 that contradicts H being equipackable. If $|E_C| = 4$, again $E' \cup \{c_0 c_3, c_1 c_2\}$ has even size and can be paired up and P_3 -removed to leave two independent edges $c_0 c_1$ and $c_2 c_3$ on C , giving a contradiction to H being equipackable.

Hence every such non-trivial component D not containing x and y is an even star.

Now suppose there exist two different components R_1 and R_2 of this kind. Analogously to the previous argumentation let E_{R_i} be the subset of E' of edges incident to the vertices of R_i for $i = 1, 2$. Since H is connected, and all bridges of H are leaf bridges there has to exist for each $i = 1, 2$ at least two edges f'_i, f''_i of E_{R_i} adjacent to an edge of R_i . Pairing up f'_i with one edge of $E(R_i)$, say f_i , $i = 1, 2$, and P_3 -removing all remaining edges of E' (their number is even, recall that $f_i \notin E'$) will isolate two odd stars $E_{R_1} - f_1$ and $E_{R_2} - f_2$, a contradiction to Corollary 6. Thus there exists only one non-trivial component R of H' not containing x and y , and that is an even star.

We now distinguish between two cases depending on the parity of $\deg_{H_1}(x)$. Assume $\deg_{H_1}(x)$ is even. Then obviously H is, regardless of whether the centre r of R is adjacent to x or not, a member of the graph family F_{12} or can be obtained by a sequence of P_3 -removals from a member of F_{12} .

Now it remains to consider that $\deg_{H_1}(x)$ is odd, i. e. $d_H(x)$ is even. As already noted at the beginning of Case 2 the vertex x must be contained in at least one cycle of H_1 and there exist at least two edges incident to x , which are not bridges. Since R is an even star $K_{1,2l}$ with $l \geq 1$ it is not difficult to deduce that the cycle has length ≤ 5 . First let R be a star with at least four branches. Recall that E' is the set of edges of all P_3 's necessary to remove in order to isolate the bridge xy and let $H' = H - E'$. Moreover, since x has an odd number of neighbours in H_1 we isolate xy by P_3 -removing all $xz, z \in N(x) \setminus \{y\}$, and one further edge $zw, w \in N(z) \setminus \{x\}$. Regardless of the choice of this additional edge zw the remainder will be an even star with at least four edges. Concatenation of all ingredients builds up a member of F_{12} or a graph that can be obtained by a sequence of P_3 -removals from a member of F_{12} . Therefore we conclude that R is always an even star with two branches regardless of the choice of the additional edge zw . By

inspection we obtain that H is either the graph F_{11} or F_{13} depicted in Figure 6.




Figure 6: Connected P_3 -equipackable graphs in Case 2

This completes the proof of our main result. \square

The proof can also be done by induction on $|E(G)|$, but the arguments are not shorter.

Acknowledgement: The authors thank the anonymous referees for careful reading and valuable comments on this text.

References

- [1] L.W. Beineke, P. Hamburger, W.D. Goddard, *Random packings of graphs*, Discrete Math. 125(1994), 45-54.
- [2] J. Bosák, *Decompositions of graphs*, Kluwer Academic Publishers, Dordrecht, (1990).
- [3] Y. Caro, J. Schönheim, *Decompositions of trees into isomorphic subtrees*, Ars Combin. 9 (1980), 119-130.
- [4] B.L. Hartnell, P.D. Vestergaard, *Equipackable graphs*, Bull. Inst. Combin. Appl. 46 (2006), 35-48.
- [5] R. Molina, *On randomly P_k -decomposable graphs*, Congr. Numer. 148 (2001), 207-221.
- [6] R. Molina, M. McNally and K. Smith, *Characterising randomly P_k -decomposable graphs for $k \leq 9$* , Congr. Numer. 156 (2002), 211-221.
- [7] S. Ruiz, *Randomly decomposable graphs*, Discrete Math. 57 (1985), 123-128.
- [8] P.D. Vestergaard, *A short update on equipackable graphs*, Discrete Math. 308 (2008), 161-165.
- [9] Y. Zhang, Y. Fan, *M_2 -equipackable graphs*, Discrete Appl. Math. 154 (2006), 1766-1770.