



Stimulus-response evaluation of the antipruritic effect of homotopic, monophasic cold and TRP-agonist counter-stimulation on histamine-induced itch in healthy volunteers

Andersen, Hjalte Holm; Randers, Amalie; Simoni, Amalie H.; Jerwiarz, Anne; Melholt, Camilla; Pedersen, Jacob B.; Hilborg, Sigurd D.; Elberling, Jesper; Arendt-Nielsen, Lars

Published in:

Acta Dermato-Venereologica

DOI (link to publication from Publisher):

[10.2340/00015555-2211](https://doi.org/10.2340/00015555-2211)

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Andersen, H. H., Randers, A., Simoni, A. H., Jerwiarz, A., Melholt, C., Pedersen, J. B., Hilborg, S. D., Elberling, J., & Arendt-Nielsen, L. (2015). Stimulus-response evaluation of the antipruritic effect of homotopic, monophasic cold and TRP-agonist counter-stimulation on histamine-induced itch in healthy volunteers. *Acta Dermato-Venereologica*, 95(7), 902, No. PP3. <https://doi.org/10.2340/00015555-2211>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

POSTER ABSTRACTS (PP1–PP35)

BASIC RESEARCH TRACK –
ANIMAL MODEL

PP1

ITCH-RELATED BITING BEHAVIOR AND
NEURONAL RESPONSIVENESS INDUCED BY
INTRADERMAL INJECTION OF PRURITOGENS
IN HAIRLESS MICE

Yuhki Ueda¹, Kei Hashimoto¹, Takamichi Kitano¹, Kido Hiroko¹, Tatsumi Matsumoto¹, Ritsuko Ishii¹, Keiji Imoto², Hide-masa Furue²

¹Maruho Co., Ltd., Kyoto R&D Center, Kyoto, ²Department of Information Physiology, National Institute for Physiological Sciences, Okazaki, Japan

Hairless mice (HR-1) are useful to investigate the effects of topical drugs for various dermatological diseases such as atopic dermatitis and herpes simplex virus infection accompanied by itch and pain. We aimed to clarify the characteristics of itch-related behavioral and neuronal responses to pruritogens in HR-1 mice. Histamine (5–5,000 nmol), serotonin (5-HT, 10–300 nmol) and a PAR-2 agonist (SLIGRL-NH₂, 10–300 µg) were injected intradermally as a pruritogen into the hindpaw in HR-1 and ICR mice, and biting behavior and spinal neuronal response were measured for 30 minutes. Biting behaviors were dose-dependently observed after 5-HT and SLIGRL-NH₂ but not histamine injections in HR-1 mice. The 5-HT and SLIGRL-NH₂-induced biting behaviors were more prominent in HR-1 mice than ICR mice. An excitation of spinal dorsal horn neurons were evoked by the 5-HT and SLIGRL-NH₂ injections in HR-1 mice, and the frequency of action potentials were also dose-dependently increased. There was a positive correlation between the duration of biting behavior and the frequency of action potentials. These results indicate that the present recordings of itch-related behavioral and neuronal responses elicited in HR-1 mice enable us to study detailed mechanisms for topical antipruritic drug action on histamine-independent itch.

PP2

[LEU11]-HK-1-DERIVED PEPTIDES WITH D-TRP
PROLONG ANTIPRURICEPTIVE EFFECTS IN
RATS

Hideki Funahashi¹, Rumi Naono-Nakayama², Yu Miyahara¹, Yasushi Ishida¹, Toshikazu Nishimori¹

¹Dep. psychiatry Psychiatry, Fac. Medicine, Univ. Miyazaki, ²Div. Integrative Physiology, Fac. Medicine, Univ. Miyazaki, Miyazaki, Japan

Hemokinin-1 (HK-1) is a mammalian tachykinin peptide consisting of 11 amino acids. Recently, it was demonstrated that the previous treatment with [Leu11]-HK-1, in which Met at the C-terminal of HK-1 was replaced by Leu, reduced scratching induced by intrathecal injection of HK-1. Additionally, the pretreatment with [Leu11]-HK-1 attenuated the induction of scratching behavior by pruritogens, suggesting that [Leu11]-HK-1 may have an inhibitory effect on itch processing. Furthermore, it is believed that replacement of amino acids by D-tryptophan (D-Trp), prolongs the effective period. Therefore, to clarify the effective period of [Leu11]-HK-1-derived peptides, the effect

of previous treatment with [D-Trp7]-[Leu11]-HK-1, [D-Trp9]-[Leu11]-HK-1 and [D-Trp7,9]-[Leu11]-HK-1 on the induction of scratching behavior by the intrathecal injection of HK-1 and by the intradermal administration of chloroquine and histamine was evaluated. The induction of scratching by intrathecal injection of HK-1 was significantly suppressed until 24 hours after pretreatment with [D-Trp7]-[Leu11]-HK-1 and [D-Trp9]-[Leu11]-HK-1 and 4 hours after [D-Trp7,9]-[Leu11]-HK-1 treatment. On the other hand, intrathecal administration of [D-Trp7,9]-[Leu11]-HK-1 and [D-Trp9]-[Leu11]-HK-1 similarly inhibited the induction of scratching behavior by intradermal injection of a pruritogen, chloroquine and histamine. Taken together, these results suggest that the antipruriceptive effects of [Leu11]-HK-1-derived peptides replaced by D-Trp may be unrelated to the number of D-Trp.

BASIC RESEARCH TRACK –
RECEPTORS AND CHANNELS

PP3

STIMULUS-RESPONSE EVALUATION OF THE
ANTIPRURITIC EFFECT OF HOMOTOPIC,
MONOPHASIC COLD AND TRP-AGONIST
COUNTER-STIMULATION ON HISTAMINE-
INDUCED ITCH IN HEALTHY VOLUNTEERS

Hjalte H. Andersen¹, Amalie Randers¹, Amalie H. Simoni¹, Anne Jerwiarz¹, Camilla Melholt¹, Jacob B. Pedersen¹, Sigurd D. Hilborg¹, Jesper Elberling², Lars Arendt-Nielsen¹

¹Center for Sensory-Motor Interactions, Department of Health Science and Technology, Aalborg University, Aalborg, ²The Allergy Clinic, Copenhagen University Hospital, Gentofte, Copenhagen, Denmark

The alleviating effect of cold on itch is a common clinical observation. Contradictory results of the antipruritic effect of biphasic cold-modulation exist. We evaluated the counter-stimulatory efficacy of 3-min monophasic cold-stimulation and the effect of pretreatment with topical TRPA1- and TRPM8-agonists. In a two-session, randomized, single-blinded study in 13 volunteers (age: 22.8±3) superficial skin-puncture through a 1% histamine droplet was performed 12 times at their volar forearms. Thermal stimulation was performed 2 minutes after histamine application (3×3cm ATS-probe, Medoc). Applied temperatures were: 4, 12, 22, 28, 32 and 37°C. Chemical counter-stimulation was conducted with 40% l-menthol and 10% trans-cinnamaldehyde and compared to the effects of 5% topical doxepin. Cold detection and pain thresholds were assessed. Outcome measures were itch intensity (VAS 0–10), wheal-size and flare-response. Cold-stimulation alleviated histamine-induced itch in a stimulus-intensity-dependent manner with the temperatures; 4, 12 and 22°C having a significant inhibitory effect, $p < 0.05$, compared to 32°C. L-menthol, CA and doxepin reduced itch intensity but trans-cinnamaldehyde also gave rise to pain (VAS 1.6±0.47). Homotopic cold- or cold-like-stimulations in the lower innocuous or noxious range has an inhibitory effect on histaminergic itch. Selective and potent agonists of receptors conveying innocuous cold could be of potential therapeutic value as antipruritics.