



**AALBORG UNIVERSITY**  
DENMARK

**Aalborg Universitet**

## **Acid-induced experimental muscle pain and hyperalgesia with single and repeated infusion in human forearm**

Wang, Kelun; Luo, Yi; Asaki, Toshiyuki; Graven-Nielsen, Thomas; Cairns, Brian E.; Arendt-Nielsen, Thomas; Arendt-Nielsen, Lars

*Published in:*  
Scandinavian Journal of Pain

*DOI (link to publication from Publisher):*  
[10.1016/j.sjpain.2017.07.012](https://doi.org/10.1016/j.sjpain.2017.07.012)

*Creative Commons License*  
CC BY-NC-ND 4.0

*Publication date:*  
2017

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*  
Wang, K., Luo, Y., Asaki, T., Graven-Nielsen, T., Cairns, B. E., Arendt-Nielsen, T., & Arendt-Nielsen, L. (2017). Acid-induced experimental muscle pain and hyperalgesia with single and repeated infusion in human forearm. *Scandinavian Journal of Pain*, 17, 260-266. <https://doi.org/10.1016/j.sjpain.2017.07.012>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## ACID-INDUCED EXPERIMENTAL MUSCLE PAIN AND HYPERALGESIA WITH SINGLE AND REPEATED INFUSION IN HUMAN FOREARM

Kelun Wang<sup>\*1</sup>, Yi Luo<sup>1</sup>, Toshiyuki Asaki<sup>1</sup>, Thomas Graven-Nielsen<sup>2</sup>, Brian E Cairns<sup>1,2</sup>, Thomas Arendt-Nielsen<sup>1</sup>, Lars Arendt-Nielsen<sup>1,2</sup>

<sup>1</sup>SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Denmark

<sup>2</sup>Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, Aalborg University

**Original paper** for Scandinavian Journal of Pain

**Figures: 5**

**Table: 0**

**Correspondence:** Kelun Wang

SMI, Department of Health Science and Technology, The Faculty of Medicine, Aalborg University, Fredrik Bajers Vej 7D-3, 9220 Aalborg, Denmark.

Tel.: +45 9940 8745, Fax: +45 9815 4008

E-mail: kelun@hst.aau.dk

## ABSTRACT

**Background and purpose:** Acid has long been thought to play an important role in the pain process. Animal study showed that repeated acid stimulation induced central sensitization. The purpose of the study is to investigate muscle pain and hyperalgesia evoked by intramuscular infusion of saline at different pH levels, and to compare the effect of a single versus repeated acid infusions.

**Methods:** Twenty healthy subjects received infusions of buffered saline (pH 5.0, 6.0, and 7.4) into the brachioradialis muscle in a randomized order. Twelve of the subjects received repeated infusions. The subjects rated the pain intensity on visual analogue scale (VAS). Thermal pain sensitivity, and pressure pain threshold (PPT) were assessed in both arm before, during, immediately after, one hour after, and one day after the infusion. A McGill Pain Questionnaire and pain mapping were completed after each infusion.

**Results:** The pH 5 solution caused significantly higher pain and larger areas than pH 6.0 or 7.4. The local PPTs were significantly decreased (hyperalgesia) during and immediately after infusion of all three solutions. No significant differences were detected between the first and second infusion.

**Conclusions:** The intensity of acid-induced muscle pain is pH-dependent. All three solutions induced pressure hyperalgesia at the infusion site. Repeated infusions did not induce increased pain or prolonged hyperalgesia as compared with a single injection. Human intramuscular acidic saline infusion could not produce chronic pain model.

**Implications:** The acid-induced pain model may reflect the early stage responses to tissue injury of clinical conditions. Repeated intramuscular acidic saline injection model of prolonged hyperalgesia in rodents could not be translated into a human for modelling chronic musculoskeletal pain.

**Key words:** Acid-induced pain; Hyperalgesia; Muscle pain; Experimental pain; Gender

difference

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

## 1. Introduction

Tissue acidosis has been observed as a regular phenomenon following inflammation, ischemia, arthritis, cancer, hematomas, and exercise<sup>1,2,3</sup>. Local tissue pH has been found to drop to 5.4 in purulent exudates, 4.7 in fracture-related hematomas, 5.0-4.0 in bone cancer, and 6.0 in patients with occlusive arterial diseases in the leg<sup>3</sup>. Considering the painful nature of all the conditions above, the high proton concentration might be a significant contributor to the associated pain<sup>1,4</sup>. Previous studies also suggest that a strong reciprocal pain potentiating interaction exists between acidic pH and several inflammatory mediators and neurotransmitters, with low pH playing the dominant role<sup>5,6,7</sup>. The acid-sensing ion channels (ASICs) play an important role in the activation of nociceptors by low pH and thus may serve as potential targets for analgesic drug developing<sup>3,4,6,8,9,10</sup>.

Human and animal studies have shown that acid can induce both transient and sustained pain<sup>3,4,11</sup>. An acid-induced pain model in rats has been proven to be safe and without significant tissue damage by histological biopsy compared with other inflammation or tumour-induced pain models in rodents<sup>11</sup>. Using acid to produce pain in human skin and muscle has also proven safe<sup>12,13,14</sup>. Primary mechanical hyperalgesia was reported to be observed following the acid stimulation. In spite of the different methods adopted, the intensity of the acid-induced pain is pH-dependent<sup>3,7,13,14</sup>. A previous study including intramuscular acidic stimulation reported that women experienced higher referred pain and exhibited a stronger correlation between local and referred pain than men<sup>13</sup>.

Animal studies have reported that repeated intramuscular acidic stimulations induced spinal hyperexcitability with contralaterally spreading hyperalgesia<sup>8,11,15,16</sup>. Local anaesthetics applied to the muscle previously injected with acidic saline could not inhibit the acid-induced contralateral spreading of hyperalgesia<sup>11</sup> indicating a central origin of the phenomenon. However, spreading of pain has not been found after repeated acid injection

1 into the masseter muscle in neither human <sup>12</sup> nor animal studies <sup>17</sup>. Since the results from  
2 human research within this field are limited, further studies should be conducted to elucidate  
3 the possible central mechanism of acid-induced muscle pain.  
4  
5

6  
7 The aims of this human study were to investigate: 1) whether acid-induced muscle pain  
8 and 2) pressure hyperalgesia were pH dependent; 3) if spreading sensitization could be  
9 evoked by repeated versus single injection of acid stimulation; and 4) if there were gender  
10 differences in any of the parameters.  
11  
12  
13  
14  
15

## 16 17 18 19 **2. Methods**

### 20 21 22 **2.1 Subjects**

23  
24 Twenty healthy subjects (7 women, 24.3±3.1 years) participated in a three-session study with  
25 a single infusion in each session. Further, 12 of the 20 subjects (4 women, 24.1±2.8 years)  
26 participated in sessions with repeated infusions. None of the subjects had a history of pain or  
27 injuries or medical conditions that could interfere with normal somatosensory functioning.  
28  
29 Women in the menstrual period were avoided. The study was approved by the local ethics  
30 committee (N 2011-0081) and conducted in accordance with the Declaration of Helsinki. All  
31 subjects gave written informed consent.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 44 **2.2 Experimental Protocol**

45  
46 The subjects participated in three sessions; each with a single infusion of buffered saline with  
47 one of three different pH levels (pH 5.0, 6.0 and 7.4). The infusions were conducted in  
48 random order and with a one-week interval between sessions. Further, 12 subjects received  
49 repeated infusions of either pH 5.0 or pH 7.4 solution with a one-day interval between  
50 infusions. In each session, neutral phosphate buffered saline (10 ml) was infused into the  
51 brachioradialis muscle over 20 min using a computer-controlled infusion pump.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Cold pain threshold (CPT), heat pain threshold (HPT), mechanical pain sensitivity  
2 (MPS), and pressure pain threshold (PPT) were assessed before, during, after, one hour after,  
3 and one day after the infusion. The pain intensity was rated by means of an electronic visual  
4 analogue scale (VAS). A McGill Pain Questionnaire and a pain map were completed after  
5 each infusion.  
6  
7  
8  
9  
10

### 11 2.3 Acidic Infusions and Pain Assessment

12 The pH adjusted phosphate buffered saline (10 mL, Hospital Pharmacy of Aalborg University  
13 Hospital, Aalborg, Denmark) was randomly infused into the left brachioradialis muscle (2 cm  
14 from the superior border of cubital fossa) in a double-blinded manner with respect to the pH  
15 level. The infusion site was cleaned with alcohol and dried prior to the needle insertion. The  
16 needle (27 G, 19 mm, BD Microlance 3, Becton Dickinson, Ireland) was inserted into middle  
17 part of the brachioradialis muscle with a depth of 15 mm. The inserted needle was fixed to  
18 the skin using surgical tape and sterile cotton. A tube (200 cm, 1.5 ml, G30303M, Care  
19 Fusion, Switzerland) was connected to the needle from the syringe. The sterile buffers were  
20 infused at a constant rate of 30 ml/h for 20 minutes using a computer-controlled infusion  
21 pump (Asena CC MK-III, Alaris medical systems, USA). The needle and tube were removed  
22 immediately after completion of the infusion.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The subjects rated the induced pain intensity on an electronic VAS on which "0 cm"  
44 indicated "no pain" and "10 cm" represented "most pain imaginable". The VAS signal was  
45 sampled every 2 seconds from the beginning of the infusion until the pain intensity had  
46 returned to zero. The maximal pain (VAS peak) and the area under the curve (VAS area) were  
47 calculated. After the infusion, the subjects were asked to draw the pain areas on an arm  
48 drawing describe the quality of the pain on the McGill Pain Questionnaire (MPQ).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.4 Assessment Sites

Two sites in the infusion side and two sites in the contralateral side were assessed (Fig. 1). CPT, HPT, MPT, and PPT were assessed 1 cm from the infusion site (T2). All sensory assessments were also conducted at the infusion site (T1) at the same time points except during infusion. Further, the assessments were performed on the contralateral arm at the corresponding sites as controls (C2 and C1).

## 2.5 Cutaneous Thermal Pain Sensitivity

Cold pain threshold (CPT) and heat pain threshold (HPT) were measured (TSA 2001 II (CHEPS, Medoc, Israel) at T1 and C1 sites. The contact area of the thermode was 9 cm<sup>2</sup>. The baseline temperature was 32 C° (centre of neutral range). The method of limits was used by applying ramp stimuli at a velocity of 1 C°/s. The cut-off temperatures were 0 C° and 55 C°. The volunteers were asked to press a button when the respective thermal sensations were perceived. The mean threshold temperature of three consecutive measurements was calculated.

## 2.6 Cutaneous Mechanical Pain Sensitivity

The cutaneous sensitivity was assessed using weight-calibrated pins (128 mN, custom made Aalborg University) at all assessment sites. The subjects rated the cutaneous mechanical pain sensitivity (MPS) on a 0-5-10 Numeric Rating Scale (NRS) on which "0" represented "no sensation", "5" represented "pain threshold", and "10" presented "worst pain imaginable". The mean of the three measurements was used in the statistical analysis.

## 2.7 Pressure Pain Sensitivity

A hand-held pressure algometer (Somedic AB, Sweden) was used to assess the PPTs. The



1 pressure was applied to all assessment sites at a constant rate of 30 kPa/s through a 1 cm<sup>2</sup>  
2 probe. The subjects were instructed to push a button immediately when they felt the pressure  
3 turning into pain. The PPTs were measured twice at each site. The interval between the two  
4 PPT trials was at least 40 sec and the mean of the two measurements was used in the  
5 statistical analysis.  
6  
7  
8  
9  
10  
11  
12  
13

## 14 2.9 Statistics

15  
16 The normal distribution was checked for all data. The necessary logarithmic transformation  
17 was performed <sup>18</sup>. QST data were then analysed using a 3-way repeated measure analysis of  
18 variance (ANOVA) with gender as between-subject factor and testing site (T1, T2, C1, C2),  
19 pH levels (pH 5.0, 6.0, 7.4) and time (baseline, during, after, one hour after, and one day  
20 after) as within-subject factors. The VAS scores and pain areas of the different pH levels and  
21 of the single and repeated infusions were analysed by 2-way (pH level and gender) repeated  
22 measure of ANOVA. A Bonferroni test was employed for post-hoc comparisons in case of  
23 significant ANOVAs. All statistical calculations were performed using the Statistical Package  
24 for Social Sciences version 20 (SPSS, IBM). The significance level was set at P<0.05. The  
25 data are presented as mean and standard error of the mean (SEM).  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 3. Results

### 44 3.1 Acidic-evoked Pain

45  
46 The VAS profiles of the three infusions with different pH levels are shown in Fig. 2A.  
47 Significant differences were detected in VAS peak (ANOVA: P < 0.023, Fig. 2B) and VAS  
48 area (ANOVA: P < 0.012, Fig. 2C). The infusion of the pH 5.0 solution caused higher VAS  
49 scores (Peak 4.65) than the pH 6.0 solution (Peak 3.37) (P= 0.015) and the 7.4 solution (1.68)  
50 (P = 0.001). In addition, the pH 7.4 solution caused the lowest VAS scores compared with the  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 pH 6.0 solution (P= 0.016).

2  
3 The pain areas following the three different solutions are illustrated in Fig. 3. Infusion  
4 of the pH 5.0 solution evoked a larger area than infusion of pH 6.0 (P =0.036) and 7.4 (P  
5 =0.004), whereas the pH 7.4 solution evoked the smallest area compared with pH 6.0 and pH  
6  
7  
8  
9  
10 5.0 solution (P< 0.021).

11  
12 No gender differences were detected in VAS peak, VAS area, or drawing areas (P >  
13  
14 0.092).

### 15 16 17 18 19 3.2 Cutaneous Thermal and Mechanical Pain Sensitivity

20 The ANOVA of the CPT, HPT, or MPS among the different solutions at any of the tested sites  
21  
22 were not significant (ANOVA: P > 0.073). Likewise, no significant change was detected  
23  
24 before, during, immediately after, one hour after, and one day after the infusion (ANOVA: P  
25  
26  
27 > 0.103,) or between gender (ANOVA: P > 0.087, data not presented).

### 28 29 30 31 32 33 3.3 Pressure Pain Sensitivity

34 The ANOVA demonstrated that the PPTs were different over time (ANOVA: P < 0.017) but  
35  
36 not between solutions pH levels (P > 0.092). Compared with baseline, relative PPTs  
37  
38 decreased immediately after, 1 hour after, and 1 day after the infusion at T1 (p<0.017) and  
39  
40 during infusion at the T2 site (p<0.030) (Fig. 4AB). No significant difference was detected in  
41  
42 the contralateral sites (ANOVA: P > 0.105; Fig. 4CD) and no gender difference was found  
43  
44 (ANOVA: P > 0.087).

### 45 46 47 48 49 50 51 52 3.4 Repeated Infusions

53 Eight out of 12 subjects received repeated infusion of pH 5.0 solution, and 4 of them received  
54  
55 repeated infusion of pH 7.4 solution No significant difference in VAS scores was detected  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 between the two repeated sessions for neither pH 5.0 nor pH 7.4 ( $P > 0.195$ ) (Fig. 5A). No  
2 difference was detected in CPT, HPT, MPS or PPT between the two repeated sessions for  
3 either pH 5.0 or pH 7.4 (ANOVA:  $P > 0.106$ , data not presented). Normalized PPT values  
4 after the 1<sup>st</sup> and 2<sup>nd</sup> infusion of pH 5.0 is shown in Fig. 5B.  
5  
6  
7  
8  
9

#### 10 11 12 **4. Discussion**

13 The intensity of the acid-induced muscle pain was pH dependent whereas the deep tissue  
14 pressure hyperalgesia was not pH dependent. Repeated acid stimulation did not induce more  
15 pain or prolonged pressure hyperalgesia as compared with a single injection. No gender  
16 differences were found.  
17  
18  
19  
20  
21  
22  
23  
24  
25

##### 26 27 **4.1 Acid-induced Muscle Pain**

28 In the present study, the single infusions of the pH solution into the human forearm muscle  
29 produced significantly higher pain intensities than the neutral buffer (pH 7.4) and the pH 6.0  
30 buffer infusions, which is consistent with previous studies on acidic muscle pain models<sup>13,19</sup>.  
31 The study is the first to use infusion of buffered saline with different pH levels into the same  
32 group of human muscles. The results provide clear evidence that the acid-evoked pain was  
33 pH dependent. The thin myelinated Group III and unmyelinated Group IV nerve fibres in  
34 muscle are responsible for transmitting muscular nociceptive information and their endings  
35 are sensitive to inflammatory mediators including low pH stimulation<sup>20,21</sup>. The decrease in  
36 tissue pH following muscle ischemia is believed to activate ASICs in muscle nociceptors,  
37 thus contributing to e.g. ischaemic muscle pain<sup>8,19,22,23</sup>. Clinically it is known that local  
38 anaesthetics with pH levels as low as 5 occasionally produce transient pain upon injection<sup>24</sup>.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

66 Thus acidic infusion could be used as a muscle pain model in both animal and human  
67 studies.

#### 4.2 Factors affecting the pain evoked by acid infusions

Factors such as infusion volume and muscle size could affect acid-evoked pain intensity. In the present study, a total of 10 ml of buffered acid saline was infused into the brachioradialis muscle at a rate of 30ml/h. The average peak pain intensity of the evoked pain was 4.65 cm for the pH 5.0, 3.37 cm for the pH 6.0, and 1.68 pH 7.4 solution (Fig. 2). In previous animal and human studies, different volumes were used to induce pain. In studies examining male rats, acidic saline was injected in volumes of 20  $\mu$ L into the rat masseter<sup>26</sup> and rat gastrocnemius muscle<sup>8</sup>, which successfully evoked pain and hyperalgesia. However, in human studies with injection of 0.5 ml unbuffered acid saline, approximately 3% of the total volume of the human masseter muscle which was comparable to the relative injection volume used in the rats, no significant pain was induced when compared with neutral saline injections<sup>12</sup>. Infusion of five times the volume (2.5 mL) of unbuffered acidic saline into the masseter muscle of human subjects induced pain levels similar to the results of single injection (0.5 ml) in previous study<sup>27</sup>. Thus, the results of the previous human studies did not provide evidence that the injection/infusion volume had a major impact on the acid-induced pain intensity. It seems other factors, such as infusion rate, and using buffered saline may play more important role in the acid-evoked pain intensity. However, the injection or infusion volume should be considered according to the muscle volume when using an acid infusion as pain model.

The infusion rate may play an important role in the acid-evoked pain. A previous study on acid-induced human skin pain indicated that raising the infusion rate leads to increasing pain by lowering the local pH more effectively and by increasing the tissue volume in which the proton concentration exceeds the threshold to excite nociceptors<sup>28</sup>. Infusion of isotonic pH 5.2 phosphate buffer into the flexor carpi radialis muscle produced pain correlated with

1 the flow rate of the infusion <sup>19</sup>. In a recent human study, acid saline (pH 3.3) was infused into  
2 the masseter muscle with a slow infusion rate of 15ml/h. The infusion evoked only mild pain  
3 and no mechanical allodynia or increased release of algescic substances assessed by  
4 microdialysis were detected <sup>27</sup>. In our previous study, acid saline (pH 5.2) was infused into  
5 the anterior tibialis muscle and the pain level was higher and PPTs were lower following an  
6 infusion rate of 40 mL/h compared to the infusion rate of 20 ml/h <sup>13</sup>. In the present study, the  
7 infusion rate of 30 mL/h was selected since the expected pain intensity was evoked by both  
8 pH 5.0 and pH 6.0 buffered saline during the pilot experiments when the different infusion  
9 rates of 10ml, 20 ml, 30ml, and 40 ml were tested.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22 In addition, using buffered saline instead of unbuffered saline might be the necessary to  
23 evoke pain. Previous human studies have suggested that acid-induced muscle pain may be  
24 more effectively produced by infusion of low pH (~5) phosphate buffers <sup>13, 28</sup> than by  
25 injections of unbuffered acidic saline <sup>12, 27</sup>. Recent human studies did not evoke the expected  
26 pain by means of unbuffered acid saline with pH 3.3 <sup>12, 27</sup>. This difference is possibly due to  
27 the ability of the muscle tissue to rapidly buffer pH changes after injections of acidic  
28 solutions. Compared with a buffered saline solution, an unbuffered saline solution could  
29 physiologically regain pH level more quickly because of the buffering capacity of the muscle  
30 tissue. Since ASIC3 channels generate sustained currents as long as the pH is acidic <sup>29</sup>, the  
31 longer the pH in the muscle remains acidic and the longer the ASIC3 channels will be  
32 activated.

#### 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 4.3 Mechanical Hyperalgesia

52  
53 In the present healthy human study, PPT values at the infusion site (T1) and around the  
54 infused site (T2) were significantly decreased during the acidic infusion compared with  
55 baseline. However, no mechanical hyperalgesia was observed in the contralateral side  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 indicating that the acidic infusion caused local sensitisation without central mechanisms  
2 being involved. Further, no significant difference was detected in the three different pH  
3 solutions indicating that the local mechanical hyperalgesia was not pH-dependent but most  
4 likely a volume effect. Similar mechanical hyperalgesia was observed in the experimental  
5 muscle pain model conducted by injecting acidic buffer into the anterior tibialis muscle <sup>13</sup>. In  
6 contrast, in a recent human study acid-infusion into the masseter muscle did not evoke  
7 mechanical hyperalgesia in either the local or contralateral side <sup>27</sup>. It seems that only localised  
8 pain and short period local hyperalgesia were observed after infusion of acidic buffer in  
9 human studies.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 4.4 Gender Differences

25 No sex-related differences in pain intensity, pain areas, or induced local pressure hyperalgesia  
26 were observed among the three different infusions in the present study. A previous study of  
27 intramuscular acidic stimulation reported that women experienced higher referred pain and  
28 exhibited a stronger correlation between local and referred pain than men <sup>13</sup>. In rats,  
29 expression of ASIC<sub>3</sub> receptors is greater in masseter muscle sensory afferent fibers in females  
30 compared with males <sup>30</sup>. It is unclear if a similar difference in ASIC channel expression  
31 occurs in humans or if the expression of ASIC channels by sensory afferent fibers varies  
32 depending on the muscle assessed. The present study only included a relatively small study  
33 sample. The non-significant findings may have resulted from inadequate statistical power.  
34 Future studies in humans may help to address whether there indeed sex-related differences in  
35 acid induced muscle pain.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

#### 52 4.5 Effect of Repeated Acid Infusions

1 In animal studies, repeated intramuscular injections of acidic saline produced a prolonged  
2 bilateral mechanical hyperalgesia lasting up to 30 days <sup>11</sup> providing the first insight into the  
3 molecular mechanisms underlying the development of chronic muscle hyperalgesia <sup>8</sup>.  
4  
5 However, in the present human study, contralateral spreading of pain and hyperalgesia was  
6  
7 not observed following repeated infusions of acidic saline. In line, repeated infusion of  
8  
9 unbuffered acidic saline into human masseter muscle did not evoke any mechanical  
10  
11 hyperalgesia in either the local or the contralateral side <sup>12, 27</sup>. Repeated infusions into the  
12  
13 tibialis muscle induced short-lasting (20 minutes) local hyperalgesia without involving the  
14  
15 contralateral side <sup>13</sup>. It is not clear why the repeated acidic infusion in humans did not  
16  
17 reproduce any long-lasting and widespread hyperalgesia similar to those in animals. It should  
18  
19 be noted that conflicting results were also found in a previous animal study where the  
20  
21 mechanical allodynia could not be detected after two repeated injections of acidic saline into  
22  
23 the masseter muscle <sup>17</sup>. The modality differences between acidic saline, buffered or  
24  
25 unbuffered, flow rate, infusion volume, intervals between repeated infusions, different  
26  
27 muscles, trigeminal region vs spinal region, and evoked pain intensity are likely to contribute  
28  
29 to the controversial results. Another explanation might be the difference in the total amount  
30  
31 of acid stimulation between the animal and human studies as a larger part of the muscle was  
32  
33 actually stimulated in the animals; whereas only a small part of the muscle was affected in  
34  
35 humans. Again, the relatively smaller sample size may also contribute to the negative  
36  
37 result.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 **5. Conclusions**

51  
52 Infusions into the brachioradialis muscle induced pain that was pH-dependent and  
53  
54 mechanical hyperalgesia that was pH independent. However, repeated intramuscular acidic  
55  
56 saline injection model of prolonged hyperalgesia in rodents could not be translated into a  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 human for modelling chronic musculoskeletal pain.  
2  
3

## 4 **6. Implications**

5  
6  
7 The acid-induced pain model may reflect the early stage responses to tissue injury of clinical  
8 conditions. It was not possible to use this human intramuscular acidic saline infusion model  
9 to produce the type of prolonged local and widespread hyperalgesia that has previously been  
10 demonstrated to occur in animal models.  
11  
12  
13  
14  
15  
16

## 17 **Ethical issues**

18  
19 None to declare  
20  
21  
22  
23

## 24 **Conflict of Interest**

25  
26 None of the authors have potential conflicts of interest to be disclosed.  
27  
28  
29  
30

## 31 **Acknowledgements**

32  
33 This study was funded by The Danish Rheumatism Association, The Danish National  
34 Advanced Technology Foundation, Aase and Ejnar Danielsen's Foundation, Lions Club  
35 Denmark, The Shionogi Science Program, and The Danish Council for Technology and  
36 Innovation (09-052174). Center for Neuroplasticity and Pain (CNAP) is supported by the  
37 Danish National Research Foundation (DNRF121).  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2 **References:**  
3  
4

- 5 1. Garber K. Why it hurts: researchers seek mechanisms of cancer pain. *J Natl Cancer Inst*  
6 2003; 95: 770-772.  
7  
8  
9  
10 2. Hood VL, Chubert C, Keller U, Muller S. Effect of systemic pH on pHi and lactic acid  
11 generation in exhaustive forearm exercise. *Am J Physiol* 1988; 255: F479-F485.  
12  
13  
14 3. Jones NG, Slater R, Cadiou H, McNaughton P, McMahon SB. Acid-Induced Pain and Its  
15 Modulation in Humans. *J Neurosci* 2004; 24:10974 –10979.  
16  
17  
18  
19 4. McMahon SB, Jones NG. Plasticity of pain signalling: role of neurotrophic factors  
20 exemplified by acid-induced pain. *J Neurobiol* 2004; 61: 72-87.  
21  
22  
23  
24 5. Reeh PW, Steen KH. Tissue acidosis in nociception and pain *Prog Brain Res* 1996; 113:  
25 143-151.  
26  
27  
28  
29 6. Roche-Gonzalez HI, Herrejon-Abreu E, Lopez-Santillan FJ, Garcia-Lopez BE, Murbartian  
30 J, Granados-Soto V. Acid increase inflammatory pain in rats: effect of local peripheral ASICs  
31 inhibitors. *Eur J Pharmacol* 2009; 603: 56-61.  
32  
33  
34  
35  
36 7. Rukwied R, Chizh BA, Lorenz U, Obreja O, Margarit S, Schley M, Schmelz M.  
37 Potentiation of nociceptive responses to low pH injections in humans by prostaglandin E2. *J*  
38 *Pain* 2007; 8: 443-451.  
39  
40  
41  
42  
43 8. Sluka KA, Price MP, Breese NM, Stucky CL, Wemmie JA, Welsh MJ. Chronic  
44 hyperalgesia induced by repeated acid injections in muscle is abolished by the loss of ASIC3,  
45 but not ASIC1. *Pain* 2003;106: 229-239.  
46  
47  
48  
49  
50 9. Voilley N, de Weille J, Mamet J, Lazdunski M. Nonsteroidal anti-inflammatory drugs  
51 inhibit both the activity and the inflammation induced expression of acid-sensing ion  
52 channels in nociceptors. *J Neurosci* 2001; 21: 8026-8033.  
53  
54  
55  
56  
57 10. Ugawa S, Ueda T, Ishida T, Nishigaki M, Shibata Y, Shimada S. Amiloride-blockable  
58  
59  
60  
61  
62  
63  
64  
65

1 acid-sensing ion channels are leading acid sensors expressed in human nociceptors. *J Clin*  
2  
3 *Invest* 2002;110: 1185-1190.

4  
5 11. Sluka KA, Kalra A, Moore SA. Unilateral intramuscular injections of acidic saline  
6  
7 produce a bilateral, long-lasting hyperalgesia. *Muscle Nerve* 2001; 24: 37-46.

8  
9  
10 12. Castrillon EE, Cairns B, List T, Svensson P, Ernberg M. Acidic saline-induced pain as a  
11  
12 model for experimental masseter myalgia in healthy subjects. *Eur J Pain* 2013; 17:  
13  
14 1438-1746.

15  
16  
17 13. Frey Law LA, Sluka KA, McMullen T, Lee J, Arendt-Nielsen L, Graven-Nielsen T.  
18  
19 Acidic buffer induced muscle pain evokes referred pain and mechanical hyperalgesia in  
20  
21 humans. *Pain* 2008; 140: 254-264.

22  
23  
24 14. Steen KH, Reeh PW. Sustained graded pain and hyperalgesia from harmless experimental  
25  
26 tissue acidosis in human skin. *Neuroscience Letters* 1993; 154: 113-116.

27  
28  
29 15. Hoeger-Bement MK, Sluka KA. Phosphorylation of CREB and mechanical hyperalgesia  
30  
31 is reversed by blockade of the cAMP pathway in a time-dependent manner after repeated  
32  
33 intramuscular acid injections. *J Neurosci* 2003; 23:5437-5445.

34  
35  
36 16. Skyba DA, Lisi TL, Sluka KA. Excitatory amino acid concentrations increase in the  
37  
38 spinal cord dorsal horn after repeated intramuscular injection of acidic saline. *Pain* 2005;  
39  
40 119:142-149.

41  
42  
43 17. Ambalavanar R, Yallampalli C, Yallampalli U, Dessem D. Injection of adjuvant but not  
44  
45 acidic saline into craniofacial muscle evokes nociceptive behaviors and neuropeptide  
46  
47 expression. *Neuroscience* 2007; 149: 650–659.

48  
49  
50 18. Rolke R, Magerl W, Campbell KA, Schalber C, Caspari S, Birklein F, Treede RD.  
51  
52 Quantitative sensory testing: a comprehensive protocol for clinical trials. *Eur J Pain* 2006;  
53  
54 10:77-88.

55  
56  
57 19. Issberner U, Reeh PW, Steen KH. Pain due to acidosis: a mechanism for inflammatory  
58  
59

1 and ischemic myalgia? *Neurosci Lett* 1996; 208: 191-194.

2  
3 20. Hoheisel U, Reinöhl J, Unger T, Mense S. Acidic pH and capsaicin activate  
4 mechanosensitive group IV muscle receptors in the rat. *Pain* 2004;110:149-157.

5  
6  
7 21. Sluka KA, Gregory NS. The dichotomized role for acid sensing ion channels in  
8 musculoskeletal pain and inflammation. *Neuropharmacology* 2015; 94:58-63. Review

9  
10  
11 22. Fujii Y, Ozaki N, Taguchi T, Mizumura K, Furukawa K, Sugiura Y. TRP channels and  
12 ASICs mediate mechanical hyperalgesia in models of inflammatory muscle pain and delayed  
13 onset muscle soreness. *Pain* 2008; 140: 292–304.

14  
15  
16  
17 23. Sluka KA, Radhakrishnan R, Benson CJ, Eshcol JO, Price MP, Babinski K, Audette KM,  
18 Yeomans DC, Wilson SP. ASIC3 in muscle mediates mechanical, but not heat, hyperalgesia  
19 associated with muscle inflammation. *Pain* 2007; 129:102–112.

20  
21  
22  
23 24. Cepeda MS, Tzortzopoulou A, Thackrey M, Hudcova J, Arora Gandhi P, Schumann R.  
24 Adjusting the pH of lidocaine for reducing pain on injection. *Cochrane Database Syst Rev*  
25 2010; 12: 1-65.

26  
27  
28 25. Frank SG, Lalonde DH. How acidic is the lidocaine we are injecting, and how much  
29 bicarbonate should we add? *Can J Plast Surg* 2012; 20: 71-74.

30  
31  
32  
33 26. Lund JP, Sadeghi S, Athanassiadis T, Salas NC, Auclair F, Thivierge B, Arsenault I,  
34 Rompre P, Westberg K-G, Kolta A. Assessment of the potential role of muscle spindle  
35 mechanoreceptor afferents in chronic muscle pain in the rat masseter muscle. *PLoS ONE*  
36 2010; 5: 1–21.

37  
38  
39 27. Ernberg M, Castrillon EE, Ghafouri B, Larsson B, Gerdle B, List T, Svensson P.  
40 Experimental myalgia induced by repeated infusion of acidic saline into the human masseter  
41 muscle does not cause the release of algescic substances. *Eur J Pain* 2013; 17: 539–550.

42  
43  
44 28. Steen KH, Issberner U, Reeh PW. Pain due to experimental acidosis in human skin:  
45 evidence for non-adapting nociceptor excitation. *Neurosci Lett* 1995; 199: 29-32.

1 29. Deval E, Gasull X, Noël J, Salinas M, Baron A, Diochot S, Lingueglia E. Acid-sensing  
2 ion channels (ASICs): Pharmacology and implication in pain. *Pharmacol Ther* 2010; 128,  
3 549–558.  
4

5  
6  
7 30. Zhang E, Wang M, Dong X, Kumar U, Cairns B. Masseter ganglion neurons express acid  
8 sensing ion channels (ASIC). *J Dent Res* 2008; 87 (Spec Issue B): 3499.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 **Figure legends**

2  
3 **Fig. 1. Infusion Sites and Testing Sites**

4  
5 T1: infusion site on the left brachioradialis muscle; T2: local testing site, 1 cm from infusion  
6  
7 site. C1 and C2: testing sites on the contralateral brachioradialis muscle  
8  
9

10  
11  
12 **Fig. 2. Visual Analogue Scale (VAS) Profile and Pain Area Under the Curve**

13  
14 Mean VAS scores after pH 5.0 (blue), pH 6.0 (red), or pH 7.4 (green) infusion of acidic  
15  
16 buffered saline into the left brachioradialis muscle in healthy humans (Mean  $\pm$  Standard Error,  
17  
18 N=20). \* indicates significant difference ( $P < 0.05$ ).  
19  
20  
21  
22  
23

24 **Fig. 3. Pain Drawing Area**

25  
26 The pain distribution after pH 5.0, pH 6.0, or pH 7.4 infusion of acidic buffered saline into  
27  
28 the left brachioradialis muscle in healthy humans (N=20). Blue and red lines represent men  
29  
30 and women, respectively.  
31  
32  
33  
34  
35

36 **Fig. 4. Pressure Pain Threshold**

37  
38 Mean ( $\pm$  standard error of the mean, N=20) pressure pain thresholds relative (%) to baseline  
39  
40 measures on the infused site T1 (A), local site T2 (B) and contralateral side C1 (C), C2 (D)  
41  
42 by the infusion of pH 5.0 (blue), pH 6.0 (red), or pH 7.4 (green) buffered saline into the left  
43  
44 brachioradialis muscle in healthy humans. \* indicates significant difference ( $P < 0.05$ )  
45  
46 compared with baseline.  
47  
48  
49  
50  
51  
52

53 **Fig. 5. VAS scores and PPT Changes after Repeated Infusion**

54  
55 A: Mean VAS scores of the first (blue solid line) and second (blue dotted line) infusion of pH  
56  
57 5.0 acidic saline (N=8) and the first (green solid line) and second (green dotted line) infusion  
58  
59  
60  
61  
62  
63  
64  
65

1 of pH 7.4 neutral phosphate buffered saline (N=4) into the left brachioradialis muscle in  
2  
3 healthy humans (N=12). B: Means of % changes (relative changes to the baseline of the  
4  
5 respective days) of pressure pain threshold from the baseline at the T2 testing site after the  
6  
7 first (blue) and second (black) infusion of pH 5.0 acidic saline into the left brachioradialis  
8  
9 muscle in healthy humans. The relative changes of PPTs were significantly lower during  
10  
11 infusion of pH 5.0 acidic saline, but no significant difference was detected between two  
12  
13 repeated sessions.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure  
Fig. 1




Fig. 2






Fig. 3




Fig. 4









Fig. 5



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## ACID-INDUCED EXPERIMENTAL MUSCLE PAIN AND HYPERALGESIA WITH SINGLE AND REPEATED INFUSION IN HUMAN FOREARM

Kelun Wang<sup>\*1</sup>, Yi Luo<sup>1</sup>, Toshiyuki Asaki<sup>1</sup>, Thomas Graven-Nielsen<sup>2</sup>, Brian E Cairns<sup>1,2</sup>, Thomas Arendt-Nielsen<sup>1</sup>, Lars Arendt-Nielsen<sup>1,2</sup>

<sup>1</sup>SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Denmark

<sup>2</sup>Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, Aalborg University

**Original paper** for Scandinavian Journal of Pain

**Figures: 5**

**Table: 0**

**Correspondence:** Kelun Wang

SMI, Department of Health Science and Technology, The Faculty of Medicine, Aalborg University, Fredrik Bajers Vej 7D-3, 9220 Aalborg, Denmark.

Tel.: +45 9940 8745, Fax: +45 9815 4008

E-mail: kelun@hst.aau.dk

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## ABSTRACT

**Background and purpose:** Acid has long been thought to play an important role in the pain process. Animal study showed that repeated acid stimulation induced central sensitization. The purpose of the study is to investigate muscle pain and hyperalgesia evoked by intramuscular infusion of saline at different pH levels, and to compare the effect of a single versus repeated acid infusions.

**Methods:** Twenty healthy subjects received infusions of buffered saline (pH 5.0, 6.0, and 7.4) into the brachioradialis muscle in a randomized order. Twelve of the subjects received repeated infusions. The subjects rated the pain intensity on visual analogue scale (VAS). Thermal pain sensitivity, and pressure pain threshold (PPT) were assessed in both arm before, during, immediately after, one hour after, and one day after the infusion. A McGill Pain Questionnaire and pain mapping were completed after each infusion.

**Results:** The pH 5 solution caused significantly higher pain and larger areas than pH 6.0 or 7.4. The local PPTs were significantly decreased (hyperalgesia) during and immediately after infusion of all three solutions. No significant differences were detected between the first and second infusion.

**Conclusions:** The intensity of acid-induced muscle pain is pH-dependent. All three solutions induced pressure hyperalgesia at the infusion site. Repeated infusions did not induce increased pain or prolonged hyperalgesia as compared with a single injection. Human intramuscular acidic saline infusion could not produce chronic pain model.

**Implications:** The acid-induced pain model may reflect the early stage responses to tissue injury of clinical conditions. Repeated intramuscular acidic saline injection model of prolonged hyperalgesia in rodents could not be translated into a human for modelling chronic musculoskeletal pain.

**Key words:** Acid-induced pain; Hyperalgesia; Muscle pain; Experimental pain; Gender

difference

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

## 1. Introduction

Tissue acidosis has been observed as a regular phenomenon following inflammation, ischemia, arthritis, cancer, hematomas, and exercise<sup>1,2,3</sup>. Local tissue pH has been found to drop to 5.4 in purulent exudates, 4.7 in fracture-related hematomas, 5.0-4.0 in bone cancer, and 6.0 in patients with occlusive arterial diseases in the leg<sup>3</sup>. Considering the painful nature of all the conditions above, the high proton concentration might be a significant contributor to the associated pain<sup>1,4</sup>. Previous studies also suggest that a strong reciprocal pain potentiating interaction exists between acidic pH and several inflammatory mediators and neurotransmitters, with low pH playing the dominant role<sup>5,6,7</sup>. The acid-sensing ion channels (ASICs) play an important role in the activation of nociceptors by low pH and thus may serve as potential targets for analgesic drug developing<sup>3,4,6,8,9,10</sup>.

Human and animal studies have shown that acid can induce both transient and sustained pain<sup>3,4,11</sup>. An acid-induced pain model in rats has been proven to be safe and without significant tissue damage by histological biopsy compared with other inflammation or tumour-induced pain models in rodents<sup>11</sup>. Using acid to produce pain in human skin and muscle has also proven safe<sup>12,13,14</sup>. Primary mechanical hyperalgesia was reported to be observed following the acid stimulation. In spite of the different methods adopted, the intensity of the acid-induced pain is pH-dependent<sup>3,7,13,14</sup>. A previous study including intramuscular acidic stimulation reported that women experienced higher referred pain and exhibited a stronger correlation between local and referred pain than men<sup>13</sup>.

Animal studies have reported that repeated intramuscular acidic stimulations induced spinal hyperexcitability with contralaterally spreading hyperalgesia<sup>8,11,15,16</sup>. Local anaesthetics applied to the muscle previously injected with acidic saline could not inhibit the acid-induced contralateral spreading of hyperalgesia<sup>11</sup> indicating a central origin of the phenomenon. However, spreading of pain has not been found after repeated acid injection

1 into the masseter muscle in neither human <sup>12</sup> nor animal studies <sup>17</sup>. Since the results from  
2 human research within this field are limited, further studies should be conducted to elucidate  
3 the possible central mechanism of acid-induced muscle pain.  
4  
5

6  
7 The aims of this human study were to investigate: 1) whether acid-induced muscle pain  
8 and 2) pressure hyperalgesia were pH dependent; 3) if spreading sensitization could be  
9 evoked by repeated versus single injection of acid stimulation; and 4) if there were gender  
10 differences in any of the parameters.  
11  
12  
13  
14  
15

## 16 17 18 19 **2. Methods**

### 20 21 22 **2.1 Subjects**

23  
24 Twenty healthy subjects (7 women, 24.3±3.1 years) participated in a three-session study with  
25 a single infusion in each session. Further, 12 of the 20 subjects (4 women, 24.1±2.8 years)  
26 participated in sessions with repeated infusions. None of the subjects had a history of pain or  
27 injuries or medical conditions that could interfere with normal somatosensory functioning.  
28  
29 Women in the menstrual period were avoided. The study was approved by the local ethics  
30 committee (N 2011-0081) and conducted in accordance with the Declaration of Helsinki. All  
31 subjects gave written informed consent.  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 44 **2.2 Experimental Protocol**

45  
46 The subjects participated in three sessions; each with a single infusion of buffered saline with  
47 one of three different pH levels (pH 5.0, 6.0 and 7.4). The infusions were conducted in  
48 random order and with a one-week interval between sessions. Further, 12 subjects received  
49 repeated infusions of either pH 5.0 or pH 7.4 solution with a one-day interval between  
50 infusions. In each session, neutral phosphate buffered saline (10 ml) was infused into the  
51 brachioradialis muscle over 20 min using a computer-controlled infusion pump.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 Cold pain threshold (CPT), heat pain threshold (HPT), mechanical pain sensitivity  
2 (MPS), and pressure pain threshold (PPT) were assessed before, during, after, one hour after,  
3 and one day after the infusion. The pain intensity was rated by means of an electronic visual  
4 analogue scale (VAS). A McGill Pain Questionnaire and a pain map were completed after  
5 each infusion.  
6  
7  
8  
9  
10

### 11 2.3 Acidic Infusions and Pain Assessment 12

13 The pH adjusted phosphate buffered saline (10 mL, Hospital Pharmacy of Aalborg University  
14 Hospital, Aalborg, Denmark) was randomly infused into the left brachioradialis muscle (2 cm  
15 from the superior border of cubital fossa) in a double-blinded manner with respect to the pH  
16 level. The infusion site was cleaned with alcohol and dried prior to the needle insertion. The  
17 needle (27 G, 19 mm, BD Microlance 3, Becton Dickinson, Ireland) was inserted into middle  
18 part of the brachioradialis muscle with a depth of 15 mm. The inserted needle was fixed to  
19 the skin using surgical tape and sterile cotton. A tube (200 cm, 1.5 ml, G30303M, Care  
20 Fusion, Switzerland) was connected to the needle from the syringe. The sterile buffers were  
21 infused at a constant rate of 30 ml/h for 20 minutes using a computer-controlled infusion  
22 pump (Asena CC MK-III, Alaris medical systems, USA). The needle and tube were removed  
23 immediately after completion of the infusion.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 The subjects rated the induced pain intensity on an electronic VAS on which "0 cm"  
44 indicated "no pain" and "10 cm" represented "most pain imaginable". The VAS signal was  
45 sampled every 2 seconds from the beginning of the infusion until the pain intensity had  
46 returned to zero. The maximal pain (VAS peak) and the area under the curve (VAS area) were  
47 calculated. After the infusion, the subjects were asked to draw the pain areas on an arm  
48 drawing describe the quality of the pain on the McGill Pain Questionnaire (MPQ).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 2.4 Assessment Sites

Two sites in the infusion side and two sites in the contralateral side were assessed (Fig. 1). CPT, HPT, MPT, and PPT were assessed 1 cm from the infusion site (T2). All sensory assessments were also conducted at the infusion site (T1) at the same time points except during infusion. Further, the assessments were performed on the contralateral arm at the corresponding sites as controls (C2 and C1).

## 2.5 Cutaneous Thermal Pain Sensitivity

Cold pain threshold (CPT) and heat pain threshold (HPT) were measured (TSA 2001 II (CHEPS, Medoc, Israel) at T1 and C1 sites. The contact area of the thermode was 9 cm<sup>2</sup>. The baseline temperature was 32 C° (centre of neutral range). The method of limits was used by applying ramp stimuli at a velocity of 1 C°/s. The cut-off temperatures were 0 C° and 55 C°. The volunteers were asked to press a button when the respective thermal sensations were perceived. The mean threshold temperature of three consecutive measurements was calculated.

## 2.6 Cutaneous Mechanical Pain Sensitivity

The cutaneous sensitivity was assessed using weight-calibrated pins (128 mN, custom made Aalborg University) at all assessment sites. The subjects rated the cutaneous mechanical pain sensitivity (MPS) on a 0-5-10 Numeric Rating Scale (NRS) on which "0" represented "no sensation", "5" represented "pain threshold", and "10" presented "worst pain imaginable". The mean of the three measurements was used in the statistical analysis.

## 2.7 Pressure Pain Sensitivity

A hand-held pressure algometer (Somedic AB, Sweden) was used to assess the PPTs. The

1 pressure was applied to all assessment sites at a constant rate of 30 kPa/s through a 1 cm<sup>2</sup>  
2 probe. The subjects were instructed to push a button immediately when they felt the pressure  
3 turning into pain. The PPTs were measured twice at each site. The interval between the two  
4 PPT trials was at least 40 sec and the mean of the two measurements was used in the  
5 statistical analysis.  
6  
7  
8  
9  
10  
11  
12  
13

## 14 2.9 Statistics

15 The normal distribution was checked for all data. The necessary logarithmic transformation  
16 was performed <sup>18</sup>. QST data were then analysed using a 3-way repeated measure analysis of  
17 variance (ANOVA) with gender as between-subject factor and testing site (T1, T2, C1, C2),  
18 pH levels (pH 5.0, 6.0, 7.4) and time (baseline, during, after, one hour after, and one day  
19 after) as within-subject factors. The VAS scores and pain areas of the different pH levels and  
20 of the single and repeated infusions were analysed by 2-way (pH level and gender) repeated  
21 measure of ANOVA. A Bonferroni test was employed for post-hoc comparisons in case of  
22 significant ANOVAs. All statistical calculations were performed using the Statistical Package  
23 for Social Sciences version 20 (SPSS, IBM). The significance level was set at P<0.05. The  
24 data are presented as mean and standard error of the mean (SEM).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## 43 3. Results

### 44 3.1 Acidic-evoked Pain

45 The VAS profiles of the three infusions with different pH levels are shown in Fig. 2A.  
46 Significant differences were detected in VAS peak (ANOVA: P < 0.023, Fig. 2B) and VAS  
47 area (ANOVA: P < 0.012, Fig. 2C). The infusion of the pH 5.0 solution caused higher VAS  
48 scores (Peak 4.65) than the pH 6.0 solution (Peak 3.37) (P= 0.015) and the 7.4 solution (1.68)  
49 (P = 0.001). In addition, the pH 7.4 solution caused the lowest VAS scores compared with the  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 pH 6.0 solution (P= 0.016).

2  
3 The pain areas following the three different solutions are illustrated in Fig. 3. Infusion  
4 of the pH 5.0 solution evoked a larger area than infusion of pH 6.0 (P =0.036) and 7.4 (P  
5 =0.004), whereas the pH 7.4 solution evoked the smallest area compared with pH 6.0 and pH  
6  
7  
8  
9  
10 5.0 solution (P< 0.021).

11  
12 No gender differences were detected in VAS peak, VAS area, or drawing areas (P >  
13  
14 0.092).

### 15 16 17 18 19 3.2 Cutaneous Thermal and Mechanical Pain Sensitivity

20 The ANOVA of the CPT, HPT, or MPS among the different solutions at any of the tested sites  
21 were not significant (ANOVA: P > 0.073). Likewise, no significant change was detected  
22 before, during, immediately after, one hour after, and one day after the infusion (ANOVA: P  
23  
24 > 0.103,) or between gender (ANOVA: P > 0.087, data not presented).

### 25 26 27 28 29 30 31 32 33 3.3 Pressure Pain Sensitivity

34 The ANOVA demonstrated that the PPTs were different over time (ANOVA: P < 0.017) but  
35  
36 not between solutions pH levels (P > 0.092). Compared with baseline, relative PPTs  
37 decreased immediately after, 1 hour after, and 1 day after the infusion at T1 (p<0.017) and  
38 during infusion at the T2 site (p<0.030) (Fig. 4AB). No significant difference was detected in  
39 the contralateral sites (ANOVA: P > 0.105; Fig. 4CD) and no gender difference was found  
40  
41 (ANOVA: P > 0.087).

### 42 43 44 45 46 47 48 49 50 51 52 3.4 Repeated Infusions

53 Eight out of 12 subjects received repeated infusion of pH 5.0 solution, and 4 of them received  
54  
55 repeated infusion of pH 7.4 solution No significant difference in VAS scores was detected  
56  
57  
58  
59  
60

1 between the two repeated sessions for neither pH 5.0 nor pH 7.4 ( $P > 0.195$ ) (Fig. 5A). No  
2 difference was detected in CPT, HPT, MPS or PPT between the two repeated sessions for  
3 either pH 5.0 or pH 7.4 (ANOVA:  $P > 0.106$ , data not presented). Normalized PPT values  
4 after the 1<sup>st</sup> and 2<sup>nd</sup> infusion of pH 5.0 is shown in Fig. 5B.  
5  
6  
7  
8  
9

#### 10 11 12 **4. Discussion**

13 The intensity of the acid-induced muscle pain was pH dependent whereas the deep tissue  
14 pressure hyperalgesia was not pH dependent. Repeated acid stimulation did not induce more  
15 pain or prolonged pressure hyperalgesia as compared with a single injection. No gender  
16 differences were found.  
17  
18  
19  
20  
21  
22  
23  
24  
25

##### 26 27 **4.1 Acid-induced Muscle Pain**

28 In the present study, the single infusions of the pH solution into the human forearm muscle  
29 produced significantly higher pain intensities than the neutral buffer (pH 7.4) and the pH 6.0  
30 buffer infusions, which is consistent with previous studies on acidic muscle pain models<sup>13,19</sup>.  
31 The study is the first to use infusion of buffered saline with different pH levels into the same  
32 group of human muscles. The results provide clear evidence that the acid-evoked pain was  
33 pH dependent. The thin myelinated Group III and unmyelinated Group IV nerve fibres in  
34 muscle are responsible for transmitting muscular nociceptive information and their endings  
35 are sensitive to inflammatory mediators including low pH stimulation<sup>20,21</sup>. The decrease in  
36 tissue pH following muscle ischemia is believed to activate ASICs in muscle nociceptors,  
37 thus contributing to e.g. ischaemic muscle pain<sup>8,19,22,23</sup>. Clinically it is known that local  
38 anaesthetics with pH levels as low as 5 occasionally produce transient pain upon injection<sup>24</sup>.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

#### 4.2 Factors affecting the pain evoked by acid infusions

Factors such as infusion volume and muscle size could affect acid-evoked pain intensity. In the present study, a total of 10 ml of buffered acid saline was infused into the brachioradialis muscle at a rate of 30ml/h. The average peak pain intensity of the evoked pain was 4.65 cm for the pH 5.0, 3.37 cm for the pH 6.0, and 1.68 pH 7.4 solution (Fig. 2). In previous animal and human studies, different volumes were used to induce pain. In studies examining male rats, acidic saline was injected in volumes of 20  $\mu$ L into the rat masseter<sup>26</sup> and rat gastrocnemius muscle<sup>8</sup>, which successfully evoked pain and hyperalgesia. However, in human studies with injection of 0.5 ml unbuffered acid saline, approximately 3% of the total volume of the human masseter muscle which was comparable to the relative injection volume used in the rats, no significant pain was induced when compared with neutral saline injections<sup>12</sup>. Infusion of five times the volume (2.5 mL) of unbuffered acidic saline into the masseter muscle of human subjects induced pain levels similar to the results of single injection (0.5 ml) in previous study<sup>27</sup>. Thus, the results of the previous human studies did not provide evidence that the injection/infusion volume had a major impact on the acid-induced pain intensity. It seems other factors, such as infusion rate, and using buffered saline may play more important role in the acid-evoked pain intensity. However, the injection or infusion volume should be considered according to the muscle volume when using an acid infusion as pain model.

The infusion rate may play an important role in the acid-evoked pain. A previous study on acid-induced human skin pain indicated that raising the infusion rate leads to increasing pain by lowering the local pH more effectively and by increasing the tissue volume in which the proton concentration exceeds the threshold to excite nociceptors<sup>28</sup>. Infusion of isotonic pH 5.2 phosphate buffer into the flexor carpi radialis muscle produced pain correlated with

1 the flow rate of the infusion <sup>19</sup>. In a recent human study, acid saline (pH 3.3) was infused into  
2 the masseter muscle with a slow infusion rate of 15ml/h. The infusion evoked only mild pain  
3 and no mechanical allodynia or increased release of algescic substances assessed by  
4 microdialysis were detected <sup>27</sup>. In our previous study, acid saline (pH 5.2) was infused into  
5 the anterior tibialis muscle and the pain level was higher and PPTs were lower following an  
6 infusion rate of 40 mL/h compared to the infusion rate of 20 ml/h <sup>13</sup>. In the present study, the  
7 infusion rate of 30 mL/h was selected since the expected pain intensity was evoked by both  
8 pH 5.0 and pH 6.0 buffered saline during the pilot experiments when the different infusion  
9 rates of 10ml, 20 ml, 30ml, and 40 ml were tested.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22 In addition, using buffered saline instead of unbuffered saline might be the necessary to  
23 evoke pain. Previous human studies have suggested that acid-induced muscle pain may be  
24 more effectively produced by infusion of low pH (~5) phosphate buffers <sup>13, 28</sup> than by  
25 injections of unbuffered acidic saline <sup>12, 27</sup>. Recent human studies did not evoke the expected  
26 pain by means of unbuffered acid saline with pH 3.3 <sup>12, 27</sup>. This difference is possibly due to  
27 the ability of the muscle tissue to rapidly buffer pH changes after injections of acidic  
28 solutions. Compared with a buffered saline solution, an unbuffered saline solution could  
29 physiologically regain pH level more quickly because of the buffering capacity of the muscle  
30 tissue. Since ASIC3 channels generate sustained currents as long as the pH is acidic <sup>29</sup>, the  
31 longer the pH in the muscle remains acidic and the longer the ASIC3 channels will be  
32 activated.

#### 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 4.3 Mechanical Hyperalgesia

52  
53 In the present healthy human study, PPT values at the infusion site (T1) and around the  
54 infused site (T2) were significantly decreased during the acidic infusion compared with  
55 baseline. However, no mechanical hyperalgesia was observed in the contralateral side  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 indicating that the acidic infusion caused local sensitisation without central mechanisms  
2 being involved. Further, no significant difference was detected in the three different pH  
3 solutions indicating that the local mechanical hyperalgesia was not pH-dependent but most  
4 likely a volume effect. Similar mechanical hyperalgesia was observed in the experimental  
5 muscle pain model conducted by injecting acidic buffer into the anterior tibialis muscle <sup>13</sup>. In  
6 contrast, in a recent human study acid-infusion into the masseter muscle did not evoke  
7 mechanical hyperalgesia in either the local or contralateral side <sup>27</sup>. It seems that only localised  
8 pain and short period local hyperalgesia were observed after infusion of acidic buffer in  
9 human studies.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 4.4 Gender Differences

25 No sex-related differences in pain intensity, pain areas, or induced local pressure hyperalgesia  
26 were observed among the three different infusions in the present study. A previous study of  
27 intramuscular acidic stimulation reported that women experienced higher referred pain and  
28 exhibited a stronger correlation between local and referred pain than men <sup>13</sup>. In rats,  
29 expression of ASIC<sub>3</sub> receptors is greater in masseter muscle sensory afferent fibers in females  
30 compared with males <sup>30</sup>. It is unclear if a similar difference in ASIC channel expression  
31 occurs in humans or if the expression of ASIC channels by sensory afferent fibers varies  
32 depending on the muscle assessed. The present study only included a relatively small study  
33 sample. The non-significant findings may have resulted from inadequate statistical power.  
34 Future studies in humans may help to address whether there indeed sex-related differences in  
35 acid induced muscle pain.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

#### 52 4.5 Effect of Repeated Acid Infusions



1 In animal studies, repeated intramuscular injections of acidic saline produced a prolonged  
2 bilateral mechanical hyperalgesia lasting up to 30 days <sup>11</sup> providing the first insight into the  
3 molecular mechanisms underlying the development of chronic muscle hyperalgesia <sup>8</sup>.  
4  
5 However, in the present human study, contralateral spreading of pain and hyperalgesia was  
6  
7 not observed following repeated infusions of acidic saline. In line, repeated infusion of  
8  
9 unbuffered acidic saline into human masseter muscle did not evoke any mechanical  
10  
11 hyperalgesia in either the local or the contralateral side <sup>12, 27</sup>. Repeated infusions into the  
12  
13 tibialis muscle induced short-lasting (20 minutes) local hyperalgesia without involving the  
14  
15 contralateral side <sup>13</sup>. It is not clear why the repeated acidic infusion in humans did not  
16  
17 reproduce any long-lasting and widespread hyperalgesia similar to those in animals. It should  
18  
19 be noted that conflicting results were also found in a previous animal study where the  
20  
21 mechanical allodynia could not be detected after two repeated injections of acidic saline into  
22  
23 the masseter muscle <sup>17</sup>. The modality differences between acidic saline, buffered or  
24  
25 unbuffered, flow rate, infusion volume, intervals between repeated infusions, different  
26  
27 muscles, trigeminal region vs spinal region, and evoked pain intensity are likely to contribute  
28  
29 to the controversial results. Another explanation might be the difference in the total amount  
30  
31 of acid stimulation between the animal and human studies as a larger part of the muscle was  
32  
33 actually stimulated in the animals; whereas only a small part of the muscle was affected in  
34  
35 humans. Again, the relatively smaller sample size may also contribute to the negative  
36  
37 result.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 **5. Conclusions**

51  
52 Infusions into the brachioradialis muscle induced pain that was pH-dependent and  
53  
54 mechanical hyperalgesia that was pH independent. However, repeated intramuscular acidic  
55  
56 saline injection model of prolonged hyperalgesia in rodents could not be translated into a  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 human for modelling chronic musculoskeletal pain.  
2  
3  
4

## 5 **6. Implications**

7 The acid-induced pain model may reflect the early stage responses to tissue injury of clinical  
8 conditions. It was not possible to use this human intramuscular acidic saline infusion model  
9 to produce the type of prolonged local and widespread hyperalgesia that has previously been  
10 demonstrated to occur in animal models.  
11  
12  
13  
14  
15  
16  
17

## 18 **Ethical issues**

19 None to declare  
20  
21  
22  
23

## 24 **Conflict of Interest**

25 None of the authors have potential conflicts of interest to be disclosed.  
26  
27  
28  
29  
30

## 31 **Acknowledgements**

32 This study was funded by The Danish Rheumatism Association, The Danish National  
33 Advanced Technology Foundation, Aase and Ejnar Danielsen's Foundation, Lions Club  
34 Denmark, The Shionogi Science Program, and The Danish Council for Technology and  
35 Innovation (09-052174). Center for Neuroplasticity and Pain (CNAP) is supported by the  
36 Danish National Research Foundation (DNRF121).  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 **References:**  
3  
4

- 5 1. Garber K. Why it hurts: researchers seek mechanisms of cancer pain. *J Natl Cancer Inst*  
6 2003; 95: 770-772.  
7  
8  
9  
10 2. Hood VL, Chubert C, Keller U, Muller S. Effect of systemic pH on pHi and lactic acid  
11 generation in exhaustive forearm exercise. *Am J Physiol* 1988; 255: F479-F485.  
12  
13  
14 3. Jones NG, Slater R, Cadiou H, McNaughton P, McMahon SB. Acid-Induced Pain and Its  
15 Modulation in Humans. *J Neurosci* 2004; 24:10974 –10979.  
16  
17  
18  
19 4. McMahon SB, Jones NG. Plasticity of pain signalling: role of neurotrophic factors  
20 exemplified by acid-induced pain. *J Neurobiol* 2004; 61: 72-87.  
21  
22  
23  
24 5. Reeh PW, Steen KH. Tissue acidosis in nociception and pain *Prog Brain Res* 1996; 113:  
25 143-151.  
26  
27  
28  
29 6. Roche-Gonzalez HI, Herrejon-Abreu E, Lopez-Santillan FJ, Garcia-Lopez BE, Murbartian  
30 J, Granados-Soto V. Acid increase inflammatory pain in rats: effect of local peripheral ASICs  
31 inhibitors. *Eur J Pharmacol* 2009; 603: 56-61.  
32  
33  
34  
35  
36 7. Rukwied R, Chizh BA, Lorenz U, Obreja O, Margarit S, Schley M, Schmelz M.  
37 Potentiation of nociceptive responses to low pH injections in humans by prostaglandin E2. *J*  
38 *Pain* 2007; 8: 443-451.  
39  
40  
41  
42  
43 8. Sluka KA, Price MP, Breese NM, Stucky CL, Wemmie JA, Welsh MJ. Chronic  
44 hyperalgesia induced by repeated acid injections in muscle is abolished by the loss of ASIC3,  
45 but not ASIC1. *Pain* 2003;106: 229-239.  
46  
47  
48  
49  
50 9. Voilley N, de Weille J, Mamet J, Lazdunski M. Nonsteroidal anti-inflammatory drugs  
51 inhibit both the activity and the inflammation induced expression of acid-sensing ion  
52 channels in nociceptors. *J Neurosci* 2001; 21: 8026-8033.  
53  
54  
55  
56  
57 10. Ugawa S, Ueda T, Ishida T, Nishigaki M, Shibata Y, Shimada S. Amiloride-blockable  
58  
59  
60  
61  
62  
63  
64  
65

1 acid-sensing ion channels are leading acid sensors expressed in human nociceptors. *J Clin*  
2  
3 *Invest* 2002;110: 1185-1190.

4  
5 11. Sluka KA, Kalra A, Moore SA. Unilateral intramuscular injections of acidic saline  
6  
7 produce a bilateral, long-lasting hyperalgesia. *Muscle Nerve* 2001; 24: 37-46.

8  
9  
10 12. Castrillon EE, Cairns B, List T, Svensson P, Ernberg M. Acidic saline-induced pain as a  
11  
12 model for experimental masseter myalgia in healthy subjects. *Eur J Pain* 2013; 17:  
13  
14 1438-1746.

15  
16  
17 13. Frey Law LA, Sluka KA, McMullen T, Lee J, Arendt-Nielsen L, Graven-Nielsen T.  
18  
19 Acidic buffer induced muscle pain evokes referred pain and mechanical hyperalgesia in  
20  
21 humans. *Pain* 2008; 140: 254-264.

22  
23  
24 14. Steen KH, Reeh PW. Sustained graded pain and hyperalgesia from harmless experimental  
25  
26 tissue acidosis in human skin. *Neuroscience Letters* 1993; 154: 113-116.

27  
28  
29 15. Hoeger-Bement MK, Sluka KA. Phosphorylation of CREB and mechanical hyperalgesia  
30  
31 is reversed by blockade of the cAMP pathway in a time-dependent manner after repeated  
32  
33 intramuscular acid injections. *J Neurosci* 2003; 23:5437-5445.

34  
35  
36 16. Skyba DA, Lisi TL, Sluka KA. Excitatory amino acid concentrations increase in the  
37  
38 spinal cord dorsal horn after repeated intramuscular injection of acidic saline. *Pain* 2005;  
39  
40 119:142-149.

41  
42  
43 17. Ambalavanar R, Yallampalli C, Yallampalli U, Dessem D. Injection of adjuvant but not  
44  
45 acidic saline into craniofacial muscle evokes nociceptive behaviors and neuropeptide  
46  
47 expression. *Neuroscience* 2007; 149: 650–659.

48  
49  
50 18. Rolke R, Magerl W, Campbell KA, Schalber C, Caspari S, Bircklein F, Treede RD.  
51  
52 Quantitative sensory testing: a comprehensive protocol for clinical trials. *Eur J Pain* 2006;  
53  
54 10:77-88.

55  
56  
57 19. Issberner U, Reeh PW, Steen KH. Pain due to acidosis: a mechanism for inflammatory  
58  
59

1 and ischemic myalgia? *Neurosci Lett* 1996; 208: 191-194.

2  
3 20. Hoheisel U, Reinöhl J, Unger T, Mense S. Acidic pH and capsaicin activate  
4 mechanosensitive group IV muscle receptors in the rat. *Pain* 2004;110:149-157.

5  
6  
7 21. Sluka KA, Gregory NS. The dichotomized role for acid sensing ion channels in  
8 musculoskeletal pain and inflammation. *Neuropharmacology* 2015; 94:58-63. Review

9  
10  
11 22. Fujii Y, Ozaki N, Taguchi T, Mizumura K, Furukawa K, Sugiura Y. TRP channels and  
12 ASICs mediate mechanical hyperalgesia in models of inflammatory muscle pain and delayed  
13 onset muscle soreness. *Pain* 2008; 140: 292–304.

14  
15  
16  
17 23. Sluka KA, Radhakrishnan R, Benson CJ, Eshcol JO, Price MP, Babinski K, Audette KM,  
18 Yeomans DC, Wilson SP. ASIC3 in muscle mediates mechanical, but not heat, hyperalgesia  
19 associated with muscle inflammation. *Pain* 2007; 129:102–112.

20  
21  
22  
23 24. Cepeda MS, Tzortzopoulou A, Thackrey M, Hudcova J, Arora Gandhi P, Schumann R.  
24 Adjusting the pH of lidocaine for reducing pain on injection. *Cochrane Database Syst Rev*  
25 2010; 12: 1-65.

26  
27  
28 25. Frank SG, Lalonde DH. How acidic is the lidocaine we are injecting, and how much  
29 bicarbonate should we add? *Can J Plast Surg* 2012; 20: 71-74.

30  
31  
32  
33 26. Lund JP, Sadeghi S, Athanassiadis T, Salas NC, Auclair F, Thivierge B, Arsenault I,  
34 Rompre P, Westberg K-G, Kolta A. Assessment of the potential role of muscle spindle  
35 mechanoreceptor afferents in chronic muscle pain in the rat masseter muscle. *PLoS ONE*  
36 2010; 5: 1–21.

37  
38  
39 27. Ernberg M, Castrillon EE, Ghafouri B, Larsson B, Gerdle B, List T, Svensson P.  
40 Experimental myalgia induced by repeated infusion of acidic saline into the human masseter  
41 muscle does not cause the release of algescic substances. *Eur J Pain* 2013; 17: 539–550.

42  
43  
44 28. Steen KH, Issberner U, Reeh PW. Pain due to experimental acidosis in human skin:  
45 evidence for non-adapting nociceptor excitation. *Neurosci Lett* 1995; 199: 29-32.

1 29. Deval E, Gasull X, Noël J, Salinas M, Baron A, Diochot S, Lingueglia E. Acid-sensing  
2 ion channels (ASICs): Pharmacology and implication in pain. *Pharmacol Ther* 2010; 128,  
3 549–558.  
4

5  
6  
7 30. Zhang E, Wang M, Dong X, Kumar U, Cairns B. Masseter ganglion neurons express acid  
8 sensing ion channels (ASIC). *J Dent Res* 2008; 87 (Spec Issue B): 3499.  
9

1 **Figure legends**

2  
3 **Fig. 1. Infusion Sites and Testing Sites**

4  
5 T1: infusion site on the left brachioradialis muscle; T2: local testing site, 1 cm from infusion  
6  
7 site. C1 and C2: testing sites on the contralateral brachioradialis muscle  
8  
9

10  
11  
12 **Fig. 2. Visual Analogue Scale (VAS) Profile and Pain Area Under the Curve**

13  
14 Mean VAS scores after pH 5.0 (blue), pH 6.0 (red), or pH 7.4 (green) infusion of acidic  
15 buffered saline into the left brachioradialis muscle in healthy humans (Mean  $\pm$  Standard Error,  
16  
17 N=20). \* indicates significant difference ( $P < 0.05$ ).  
18  
19  
20  
21  
22  
23

24 **Fig. 3. Pain Drawing Area**

25  
26 The pain distribution after pH 5.0, pH 6.0, or pH 7.4 infusion of acidic buffered saline into  
27 the left brachioradialis muscle in healthy humans (N=20). Blue and red lines represent men  
28  
29 and women, respectively.  
30  
31  
32  
33  
34  
35

36 **Fig. 4. Pressure Pain Threshold**

37  
38 Mean ( $\pm$  standard error of the mean, N=20) pressure pain thresholds relative (%) to baseline  
39 measures on the infused site T1 (A), local site T2 (B) and contralateral side C1 (C), C2 (D)  
40  
41 by the infusion of pH 5.0 (blue), pH 6.0 (red), or pH 7.4 (green) buffered saline into the left  
42  
43 brachioradialis muscle in healthy humans. \* indicates significant difference ( $P < 0.05$ )  
44  
45 compared with baseline.  
46  
47  
48  
49  
50  
51  
52

53 **Fig. 5. VAS scores and PPT Changes after Repeated Infusion**

54  
55 A: Mean VAS scores of the first (blue solid line) and second (blue dotted line) infusion of pH  
56  
57 5.0 acidic saline (N=8) and the first (green solid line) and second (green dotted line) infusion  
58  
59  
60  
61  
62  
63  
64  
65

1 of pH 7.4 neutral phosphate buffered saline (N=4) into the left brachioradialis muscle in  
2  
3 healthy humans (N=12). B: Means of % changes (relative changes to the baseline of the  
4  
5 respective days) of pressure pain threshold from the baseline at the T2 testing site after the  
6  
7 first (blue) and second (black) infusion of pH 5.0 acidic saline into the left brachioradialis  
8  
9 muscle in healthy humans. The relative changes of PPTs were significantly lower during  
10  
11 infusion of pH 5.0 acidic saline, but no significant difference was detected between two  
12  
13 repeated sessions.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65