



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes

Imamura, Minako; Takahashi, Atsushi; Yamauchi, Toshimasa; Hara, Kazuo; Yasuda, Kazuki; Grarup, Niels; Zhao, Wei; Wang, Xu; Huerta-Chagoya, Alicia; Hu, Cheng; Moon, Sanghoon; Long, Jirong; Kwak, Soo Heon; Rasheed, Asif; Saxena, Richa; Ma, Ronald C W; Okada, Yukinori; Iwata, Minoru; Hosoe, Jun; Shojima, Nobuhiro; Iwasaki, Minaka; Fujita, Hayato; Suzuki, Ken; Danesh, John; Jørgensen, Torben; Jørgensen, Marit E; Witte, Daniel R; Brandslund, Ivan; Christensen, Cramer; Hansen, Torben; Mercader, Josep M; Flannick, Jason; Moreno-Macías, Hortensia; Burtt, Noël P; Zhang, Rong; Kim, Young Jin; Zheng, Wei; Singh, Jai Rup; Tam, Claudia H T; Hirose, Hiroshi; Maegawa, Hiroshi; Ito, Chikako; Kaku, Kohei; Watada, Hirotaka; Tanaka, Yasushi; Tobe, Kazuyuki; Kawamori, Ryuzo; Kubo, Michiaki; Cho, Yoon Shin; Chan, Juliana C N; Sanghera, Dharambir; Frossard, Philippe; Park, Kyong Soo; Shu, Xiao-Ou; Kim, Bong-Jo; Florez, Jose C; Tusié-Luna, Teresa; Jia, Weiping; Tai, E Shyong; Pedersen, Oluf; Saleheen, Danish; Maeda, Shiro; Kadowaki, Takashi

Published in:
Nature Communications

DOI (link to publication from Publisher):
[10.1038/ncomms10531](https://doi.org/10.1038/ncomms10531)

Creative Commons License
CC BY 4.0

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

ARTICLE

Received 26 Jun 2015 | Accepted 22 Dec 2015 | Published 28 Jan 2016

DOI: 10.1038/ncomms10531

OPEN

Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes

Minako Imamura *et al.*[#]

Genome-wide association studies (GWAS) have identified more than 80 susceptibility loci for type 2 diabetes (T2D), but most of its heritability still remains to be elucidated. In this study, we conducted a meta-analysis of GWAS for T2D in the Japanese population. Combined data from discovery and subsequent validation analyses (23,399 T2D cases and 31,722 controls) identify 7 new loci with genome-wide significance ($P < 5 \times 10^{-8}$), rs1116357 near *CCDC85A*, rs147538848 in *FAM60A*, rs1575972 near *DMRTA1*, rs9309245 near *ASB3*, rs67156297 near *ATP8B2*, rs7107784 near *MIR4686* and rs67839313 near *INAFM2*. Of these, the association of 4 loci with T2D is replicated in multi-ethnic populations other than Japanese (up to 65,936 T2Ds and 158,030 controls, $P < 0.007$). These results indicate that expansion of single ethnic GWAS is still useful to identify novel susceptibility loci to complex traits not only for ethnicity-specific loci but also for common loci across different ethnicities.

Correspondence and requests for materials should be addressed to S.M. (email: smaeda@med.u-ryukyu.ac.jp) or to T.K. (email: kadowaki-3im@h.u-tokyo.ac.jp).
[#]A full list of authors and their affiliations appears at the end of the paper.

To date, more than 80 susceptibility loci for type 2 diabetes (T2D) have been identified through genome-wide association studies (GWAS)^{1–18}. However, the joint effects of these variants account for <10% of the heritability for T2D^{10,19}. GWAS for T2D have been extensively conducted in populations of European descent and, accordingly, the majority of established T2D susceptibility genetic loci were originally identified by European GWAS^{1,2,8–11,16}. Cumulative evidence suggests that Asian populations may be more genetically susceptible to T2D than populations with European ancestry²⁰. In addition, there are significant interethnic differences in the risk allele frequency or in effect sizes at several loci, which may affect the power to detect associations in these populations². On the other hand, both overlap in T2D susceptibility loci among different ancestry groups and coincident risk alleles at lead single-nucleotide polymorphisms (SNPs) across diverse populations have been reported, suggesting that causal variants at many of these loci are shared across different ancestry groups¹². Moreover, a recently published transethnic GWAS has successfully identified seven novel T2D susceptibility loci by combining the association data from European, South Asian, East Asian and Mexican/Latinos GWAS¹². Therefore, it is valuable to perform GWAS for T2D using non-European and European populations, to facilitate identification of both ethnicity-specific and common-susceptibility loci among different ethnic groups.

Four T2D GWAS loci discovered in a Japanese population earlier have been shown to be significantly associated with T2D in the largest European GWAS meta-analysis¹⁰: *KCNQ1* (refs 3,4), *UBE2E2* (ref. 5), *C2CD4A–C2CD4B*⁵ and *ANK1* (ref. 6), highlighting that there are common loci conferring susceptibility to T2D among the different ethnic groups studied. Three additional loci (*MIR129-LEP*, *GPSM1* and *SLC16A11-SLC16A13*) have been identified by a large-scale Japanese GWAS ($n = \sim 25,000$) based on the imputation of genotypes using the 1000 Genomes Project data as a reference⁷. One of the findings, the association in the *SLC16A11-SLC16A13* was also confirmed in the Mexican GWAS study¹⁵.

To identify novel loci for susceptibility to T2D, we have expanded the Japanese GWAS data set by incorporating new Japanese GWAS data (9,817 T2D cases and 6,763 controls) with GWAS data in previously reported case-control individuals (5,646 T2D cases and 19,420 controls)⁷ followed by a validation study using independent Japanese case-control individuals (7,936 T2D and 5,539 controls) and multi-ethnic replication studies (East Asians: 12,554 T2D and 17,383 controls; Europeans: 38,947 T2D and 121,903 controls; South Asians: 10,587 T2D and 14,378 controls; and Mexican/Latinos: 3,848 T2D and 4,366 controls). As a result, we identify seven novel loci for T2D and the result indicates that expansion of single ethnic GWAS is still useful to identify novel susceptibility loci to complex traits.

Results

GWAS meta-analysis and validation in the Japanese population.

Imputed genotype dosage data for 9,817 T2D cases and 6,763 controls for 7,521,072 autosomal SNPs (Stage-1, set-1) were obtained and combined with an independent GWAS data of previously reported case-control individuals⁷ (Stage-1, set-2: 5,646 T2D cases and 19,420 controls; 7,521,072 autosomal SNPs), as shown in Fig. 1a. There was no obvious inflation in the quantile-quantile plots for each study (Stage-1: $\lambda_{GC} = 1.13$ and λ_{GC} adjusted for 1,000 cases and controls ($\lambda_{GC-1000}$)²¹ = 1.012; Stage-2: $\lambda_{GC} = 1.082$ and $\lambda_{GC-1000} = 1.009$), as shown in Supplementary Fig. 1A,B. SNPs with a low imputation quality ($r^2 < 0.7$ either in set-1 or set-2) or with an inconsistent direction of effect between the studies were excluded from the analysis. We obtained 42 loci exhibiting a suggestive

association with T2D ($P < 1 \times 10^{-6}$). The most significant association in this meta-analysis was rs2237896 located at intron 15 of *KCNQ1* ($P = 2.81 \times 10^{-70}$), which was previously identified in Japanese GWAS^{3,4} (Supplementary Fig. 1C). Out of the 42 loci, 25 were previously established T2D susceptibility loci (Supplementary Table 1) and the remaining 17 were further evaluated using an independent Japanese case-control study (Stage-2: 7,936 T2D cases and 5,539 controls, multi-centre) and *de novo* genotyping (Supplementary Tables 2 and 3).

To explore the novel T2D susceptibility loci, we combined the association data for the candidate SNPs from all the Japanese case-control samples (Stage-1 set-1, Stage-1 set-2 and Stage-2) using a meta-analysis (Fig. 1b and Supplementary Table 3) and identified seven T2D susceptibility loci with a significant association ($P < 5 \times 10^{-8}$; Table 1, Fig. 2 and Supplementary Table 3): rs1116357 near *CCDC85A* ($P = 6.97 \times 10^{-10}$, odds ratio (OR) = 1.09, 95% confidence interval (CI) = 1.06–1.12), rs147538848 in *FAM60A* ($P = 7.83 \times 10^{-10}$, OR = 1.11, 95% CI = 1.07–1.15), rs1575972 near *DMRTA1* ($P = 1.50 \times 10^{-9}$, OR = 1.19, 95% CI = 1.13–1.26), rs9309245 near *ASB3* ($P = 1.25 \times 10^{-8}$, OR = 1.10, 95% CI = 1.07–1.14), rs67156297 near *ATP8B2* ($P = 1.95 \times 10^{-8}$, OR = 1.14, 95% CI = 1.09–1.19), rs7107784 near *MIR4686* ($P = 2.07 \times 10^{-8}$, OR = 1.14, 95% CI = 1.09–1.20) and rs67839313 near *INAFM2* ($P = 2.42 \times 10^{-8}$, OR = 1.09, 95% CI = 1.06–1.12). The effect sizes (OR) of these seven SNPs were similar before and after adjusting for age, sex and body mass index (BMI) (Supplementary Table 4). The rs1575972 locus in *DMRTA1* was located 170 kbp away from the T2D locus *CDKN2A/B* at 9p21 (refs 10,22–24) and the rs7107784 locus near *MIR4686* was located ~ 620 kbp upstream of *KCNQ1* (refs 3,4). The linkage disequilibrium (LD) between rs1575972 and rs10811661, which was a lead SNP within the *CDKN2A/B* locus¹⁰, was weak (JPT: $r^2 = 0.01$, CEU: $r^2 = 0.02$). The association of rs1575972 was significant even after conditioning on rs10811661 (P for meta-analysis = 2.45×10^{-9} , OR = 1.19, 95% CI = 1.12–1.26; Supplementary Table 5); therefore, we considered the rs1575972 locus as a novel T2D susceptibility locus, independent of the *CDKN2A/B* locus. We also observed that the rs7107784 locus near *MIR4686* was not in LD with rs2237897 (JPT: $r^2 < 0.01$, CEU: $r^2 < 0.01$), which was a lead SNP within the *KCNQ1* locus³, and the association was similar after conditioning on rs2237897 (P for meta-analysis = 2.75×10^{-8} , OR = 1.16, 95% CI = 1.10–1.22; Supplementary Table 6).

We also examined the association of these seven SNPs with glycaemic traits in Stage-2 control individuals, including fasting plasma glucose, homeostasis model assessment (HOMA) of β -cell function (HOMA- β) and HOMA of insulin resistance (IR). However, we did not detect any significant associations between the T2D risk alleles and these glycaemic traits ($P \geq 0.0024$ Supplementary Table 7). We also searched the publicly available European GWAS data^{11,25,26} (MAGIC, <http://www.magicinvestigators.org>) and found that the T2D risk allele at the *DMRTA1* locus (rs11791293-C; proxy for rs1575972-T, CEU $r^2 = 1$) and at the *MIR4686* locus (rs7111341-T; proxy for rs7107784-G, CEU $r^2 = 0.95$) were associated with a decrease in fasting plasma insulin (FPI) ($P = 0.0039$; Supplementary Table 8) and with an increase in FPI ($P = 0.0066$; Supplementary Table 8), respectively, although these associations were not statistically significant ($P \geq 0.0008 = 0.05/63$ (7 SNPs \times 9 traits)).

Examination of seven novel loci in diverse ethnic groups. We analysed the association of these seven variants with disease susceptibility in populations other than Japanese. We obtained association data for four ethnic groups using *de novo* genotyping, *in silico* replication and by examining publicly available GWAS data¹⁰ including East Asian ($n =$ up to 29,937: 12,554 T2Ds and




Figure 1 | Outline of the present study. (a) GWAS meta-analysis and subsequent validation analysis. (b) Follow-up analyses for seven novel susceptibility loci for T2D in populations other than Japanese. SNP, single nucleotide polymorphism; MAF, minor allele frequency; HWE P, Hardy-Weinberg Equilibrium test P; RSQ, r square.

17,383 controls), South Asian ($n = \text{up to } 24,965$: 10,587 T2Ds and 14,378 controls), European ($n = \text{up to } 160,850$: 38,947 T2Ds and 121,903 controls) and Mexican ($n = \text{up to } 8,214$: 3,848 T2Ds and 4,366 controls) populations (Supplementary Table 9). Meta-analyses of the combined data from the four non-Japanese ethnicities indicated that four SNP loci, namely rs147538848 in *FAM60A*, rs1575972 near *DMRTA1*, rs7107784 near *MIR4686* and rs67839313 near *INAFM2* were associated with the disease after Bonferroni's correction ($P < 0.00714 = 0.05/7$; Table 2). The disease association of these four SNPs was further corroborated by combining the Japanese data with the multi-ethnic replication data sets (Supplementary Table 10). The rs67156297 locus in *ATP8B2* was nominally associated with T2D in the combined meta-analysis for multi-ethnic groups other than the Japanese populations. We did not detect any disease association for the remaining two SNP loci other than in the Japanese population; however, the effect direction for each of the seven loci was consistent with that in the Japanese population.

Sex- and BMI-stratified analyses in the Japanese population. We performed BMI-stratified (BMI < 25 or ≥ 25) and sex-stratified analyses in the novel and established GWAS loci,

to determine whether significant heterogeneity in allelic effects existed between non-obese and obese individuals or males and females in the Japanese population. BMI-stratified analysis for 83 previously established loci revealed evidence of significant heterogeneity in the effect size between non-obese and obese individuals at *KCNQ1* (P for heterogeneity = 8.89×10^{-5} ; Supplementary Table 11). The effect size of *KCNQ1* was greater in the non-obese group than in the obese group (Supplementary Table 11). In sex-stratified analyses, individual established loci did not show significant heterogeneity in effect sizes between men and women ($P > 6 \times 10^{-4}$; Supplementary Table 12).

For the seven novel T2D-associated loci identified in this study, no significant heterogeneity was detected in BMI-stratified or sex-stratified analyses (Supplementary Tables 13 and 14).

Fine mapping analyses for established T2D loci. We examined the association data of 83 previously identified T2D susceptibility loci in the Japanese GWAS meta-analysis data (Supplementary Data 1 and Supplementary Fig. 2). Variants at 19 loci were found associated with T2D at a genome-wide level of significance and additional 30 loci were determined to be significantly associated with T2D ($P < 6.02 \times 10^{-4} = 0.05/83$). Of the above




Figure 2 | Regional association plots of the two-stage GWAS meta-analysis for seven novel T2D susceptibility loci in the Japanese. Each plot shows $-\log_{10} P$ -values against the chromosomal positions of SNPs in the specific region. The SNP with the strongest association signal (lead SNP) in each locus is represented as a purple diamond; the other SNPs are coloured according to the extent of LD with the lead SNP. Estimated recombination rates from the hg19/1000 Genomes Project March 2012 East Asian references are shown as light-blue lines. (a) *CCDC85A*, (b) *FAM60A*, (c) *DMRTA1*, (d) *ASB3*, (e) *ATP8B2*, (f) *MIR4686* and (g) *INAFM2*. *Gene position for *INAFM2* was added manually.

49 significant associations, *ADCY5*, *HNF1A* and *PRC1* were not previously evaluated in the Japanese population, because the lead SNPs within these loci in the European GWAS (rs11708067 and rs11717195 at the *ADCY5* locus^{9,10}, rs12427353 and rs7957197 at the *HNF1A* locus^{8,10} and rs8042680 at the *PRC1* locus⁸) were monoallelic in the Japanese population. In this study, rs79223353 at the *ADCY5* locus, rs55783344 at the *HNF1A* locus and rs79548680 at the *PRC1* locus were determined to be significantly associated with T2D ($P < 6.02 \times 10^{-4}$; Supplementary Data 1 and Supplementary Fig. 3). Meta-analysis combining the GWAS data with *de novo* genotyping data for Stage-2 individuals revealed that the association of rs79223353 within the *ADCY5* locus and rs79548680 within the *PRC1* locus reached genome-wide significance in the Japanese population (Supplementary Table 15). We did not detect any disease-associated SNPs within the 16 loci (9 derived from European GWAS, 3 from East Asian, 2 from trans-ethnic, 1 from South Asian and 1 from African

American, $P \geq 0.05$) using fine mapping analyses (Supplementary Data 1 and Supplementary Fig. 2). We also identified a secondary association signal located at *EXOC6* near the *IDE-HHEX* locus¹⁰. The associations of rs78627331 and rs34773007 within the *EXOC6* locus were significant after conditioning on rs1111875 ($r^2 = 0.01$ for rs78627331 and $r^2 = 0.04$ for rs34773007 in *JPT*), which was a previously reported lead SNP within the *IDE-HHEX* locus ($P = 1.49 \times 10^{-8}$ for rs78627331, $P = 2.20 \times 10^{-8}$ for rs34773007; Supplementary Table 16).

Drug targets search by a bioinformatics approach. We applied the genetic information from previously reported and the present GWAS, to investigate potential drug targets for the treatment of T2D. First, we defined 286 T2D potential risk genes located in any of the 90 T2D risk loci (7 novel T2D loci that were identified in the present study and 83 previously identified T2D loci; see

Table 1 | Newly identified T2D susceptibility loci achieving genome-wide significance (combined meta-analysis $P < 5 \times 10^{-8}$) in Japanese populations.

SNP ID	Chr.	Position (Build 37)	Alleles		Study	RAF		OR (95% CI)	P-value	Phet
			Risk	Other		Case	Control			
rs1116357			G	A	Stage-1, set-1	0.292	0.269	1.12 (1.07-1.17)	6.55×10^{-6}	
2					Stage-1, set-2	0.299	0.279	1.10 (1.05-1.15)	2.61×10^{-5}	
57287411					Stage-2	0.294	0.284	1.05 (0.99-1.11)	8.65×10^{-2}	
CCDC85A					Combined			1.09 (1.06-1.12)	6.97×10^{-10}	0.21
rs147538848			A	G	Stage-1, set-1	0.205	0.190	1.11 (1.05-1.18)	5.81×10^{-4}	
12					Stage-1, set-2	0.199	0.184	1.12 (1.06-1.18)	1.15×10^{-4}	
31466613					Stage-2	0.210	0.193	1.11 (1.04-1.18)	1.10×10^{-3}	
FAM60A					Combined			1.11 (1.07-1.15)	7.83×10^{-10}	0.99
rs1575972			T	A	Stage-1, set-1	0.947	0.935	1.24 (1.13-1.36)	8.64×10^{-6}	
9					Stage-1, set-2	0.943	0.935	1.17 (1.07-1.29)	1.13×10^{-3}	
22301092					Stage-2	0.950	0.942	1.16 (1.04-1.29)	7.60×10^{-3}	
DMRTA1					Combined			1.19 (1.13-1.26)	1.50×10^{-9}	0.62
rs9309245			G	C	Stage-1, set-1	0.178	0.164	1.10 (1.04-1.17)	1.57×10^{-3}	
2					Stage-1, set-2	0.186	0.168	1.14 (1.08-1.20)	3.43×10^{-6}	
53397048					Stage-2	0.181	0.172	1.06 (0.998-1.14)	5.95×10^{-2}	
ASB3					Combined			1.10 (1.07-1.14)	1.25×10^{-8}	0.30
rs67156297			A	G	Stage-1, set-1	0.102	0.086	1.21 (1.12-1.30)	1.28×10^{-6}	
1					Stage-1, set-2	0.097	0.087	1.13 (1.05-1.21)	1.34×10^{-3}	
154336716					Stage-2	0.092	0.087	1.07 (0.98-1.16)	1.33×10^{-1}	
ATP8B2					Combined			1.14 (1.09-1.19)	1.95×10^{-8}	0.10
rs7107784			G	A	Stage-1, set-1	0.099	0.089	1.16 (1.07-1.26)	4.91×10^{-4}	
11					Stage-1, set-2	0.101	0.090	1.18 (1.09-1.27)	3.88×10^{-5}	
2215089					Stage-2	0.093	0.086	1.09 (0.998-1.19)	5.58×10^{-2}	
MIR4686					Combined			1.14 (1.09-1.20)	2.07×10^{-8}	0.42
rs67839313			C	T	Stage-1, set-1	0.280	0.261	1.10 (1.05-1.16)	7.87×10^{-5}	
15					Stage-1, set-2	0.278	0.264	1.08 (1.03-1.14)	1.75×10^{-3}	
40619724					Stage-2	0.281	0.267	1.07 (1.01-1.13)	1.36×10^{-2}	
INAFM2					Combined			1.09 (1.06-1.12)	2.42×10^{-8}	0.73

CI, confidence interval; Chr., chromosome; OR, odds ratio; Phet, P-value for Cochran's Q-test for heterogeneity; RAF, risk allele frequency; SNP, single-nucleotide polymorphism. Alleles are aligned to the forward strand of NCBI Build 37.

Supplementary Note). Among the 286 genes, by using a previously described scoring system²⁷, we selected 40 genes with a score of 2 or higher (Supplementary Note, Supplementary Tables 17–20, Supplementary Data 2 and 3, and Supplementary Figs 4 and 5) as 'biological T2D risk genes' (Fig. 3 and Supplementary Data 4). In brief, we scored each of the 286 biological candidate genes by adopting the following six selection criteria and calculating the number of satisfied criteria as follows: (1) genes for which T2D risk SNPs or any of the SNPs in LD ($r^2 \geq 0.80$) with them were annotated as missense variants; (2) genes for which *cis*-eQTL genes of any of lymphoblastoid cell lines, adipose tissue or liver tissues were observed for T2D risk SNPs ($P < 0.05$ for lymphoblastoid cell lines and adipose tissues, and $P < 0.004$ for liver tissues); (3) monogenic diabetes genes; (4) genes for which at least three out of six associated phenotype labels (homeostasis/metabolism, liver/biliary system, endocrine/exocrine gland, growth/size/body, mortality/ageing and embryogenesis; $P < 9.2 \times 10^{-5}$) were observed in knockout mouse²⁸; (5) genes prioritized by PubMed text mining genes using GRAIL²⁹ with gene-based $P < 0.05$; and (6) genes prioritized by protein–protein interaction (PPI) network using DAPPLE³⁰ with gene based $P < 0.05$.

As these criteria exhibited weak correlations with each other ($r^2 < 0.34$; Supplementary Fig. 5), each gene was given a score based on the number of criteria that were met (scores ranged from 0 to 6). Genes with a score of 2 or higher were defined as biological T2D risk genes.

We searched for overlapping genes between the 871 drug target genes corresponding to approved, in clinical trials or experimental drugs for various human diseases described in the previous report²⁷, and the 40 biological T2D risk genes plus 712 genes that are known to have products that have direct PPI³⁰ with the

biological T2D risk gene products. We identified a total of 83 overlapping genes (Supplementary Fig. 6 and Supplementary Table 21). Fourteen drug target genes with approved T2D treatments demonstrated significant overlap with the 40 biological T2D risk genes and 712 genes with direct PPI (4 genes overlapped with 5.6-fold enrichment as determined using permutation analysis, $P = 0.0042$; Supplementary Table 22 and Supplementary Fig. 6). The 871 drug target genes had overlap with the identified 83 genes, which is 1.8-fold more enrichment than would be expected by chance, but this is 3.1-fold less enrichment compared with overlap of the targets of T2D drugs (Supplementary Fig. 6).

Of the 83 overlapping genes, 5 were biological T2D risk genes (*PPARG*, *KCNJ11*, *ABCC8*, *GSK3B* and *KIF11*; Fig. 4). Three of these are targets of approved T2D drug treatments: *PPARG*, thiazolidinediones; *KCNJ11*, sulfonylurea; *ABCC8*, sulfonylureas and glinide. *GSK3B* is a target gene of a GSK3B activator that was in clinical trials as of August 2014 (Supplementary Table 23). Of the remaining 78 genes, 2 genes exhibit PPI with 3 biological T2D risk gene products. *GSK3B* interacts with *NOTCH1*, *NOTCH2* and *CCND2*, whereas *JUN* interacts with *FBXW7*, *HHEX* and *CCND2*. Eight genes interact with 2 biological T2D risk gene products and 68 genes interact with a single biological T2D risk gene product (Supplementary Table 21). A list of therapeutic drugs that are currently under clinical trials targeting *GSK3B*, *KIF11*, *GSK3B* and *JUN* is shown in Supplementary Table 23.

Discussion

In this study, we performed a GWAS meta-analysis in the Japanese population followed by validation using an independent Japanese sample. Integration of the results for ~55,000 Japanese individuals identified 7 novel loci associated with T2D that

Table 2 | Association of novel seven SNP loci with type 2 diabetes risk in the population other than Japanese.

SNP ID	Alleles		Population	OR (95% CI)	P-value	Phet	RAF*
	Risk	Other					
rs1116357 <i>CCDC85A</i>	G	A	All replication set [†]	1.01 (0.99-1.03)	1.99×10^{-1}	0.96	
			East Asian	1.03 (0.99-1.07)	2.06×10^{-1}	0.98	0.28
			South Asian	1.02 (0.98-1.05)	4.18×10^{-1}	0.34	0.57
			European	1.01 (0.98-1.04)	5.45×10^{-1}	0.98	0.53
			Mexican/Latino	0.99 (0.93-1.05)	5.96×10^{-1}	NA	0.48
		All Japanese	1.09 (1.06-1.12)	6.97×10^{-10}	0.21	0.28	
rs147538848 <i>FAM60A</i>	A	G	All replication set	1.10 (1.05-1.16)	2.25×10^{-4}	0.82	
			East Asian	1.13 (1.05-1.20)	3.58×10^{-4}	0.88	0.19
			South Asian	1.07 (0.97-1.17)	1.59×10^{-1}	0.35	0.06
			European	NA	NA	NA	0.01
			Mexican/Latino	1.01 (0.74-1.37)	8.98×10^{-1}	NA	0.01
		All Japanese	1.11 (1.07-1.15)	7.83×10^{-10}	0.99	0.23	
rs1575972 <i>DMRTA1</i>	T	A	All replication set	1.13 (1.08-1.18)	2.26×10^{-7}	0.13	
			East Asian	1.14 (1.02-1.26)	1.62×10^{-2}	0.03	0.96
			South Asian	1.13 (1.03-1.24)	1.21×10^{-2}	0.45	0.94
			European	1.14 (1.07-1.21)	5.58×10^{-5}	0.56	0.97
			Mexican/Latino	1.004 (0.83-1.21)	9.92×10^{-1}	NA	0.98
		All Japanese	1.19 (1.13-1.26)	1.50×10^{-9}	0.62	0.93	
rs9309245 <i>ASB3</i>	G	C	All replication set	1.01 (0.99-1.03)	5.50×10^{-1}	0.58	
			East Asian	1.04 (0.99-1.09)	1.64×10^{-1}	0.40	0.20
			South Asian	0.996 (0.96-1.04)	8.21×10^{-1}	0.63	0.31
			European	1.01 (0.98-1.04)	4.74×10^{-1}	0.80	0.35
			Mexican/Latino	0.96 (0.90-1.03)	2.31×10^{-1}	NA	0.26
		All Japanese	1.10 (1.07-1.14)	1.25×10^{-8}	0.30	0.17	
rs67156297 <i>ATP8B2</i>	A	G	All replication set	1.03 (1.001-1.05)	4.08×10^{-2}	0.38	
			East Asian	1.05 (0.99-1.12)	1.22×10^{-1}	0.26	0.09
			South Asian	0.98 (0.94-1.03)	4.79×10^{-1}	0.99	0.20
			European	1.04 (0.999-1.08)	5.90×10^{-2}	0.88	0.25
			Mexican/Latino	1.08 (0.995-1.18)	8.19×10^{-2}	NA	0.16
		All Japanese	1.14 (1.09-1.19)	1.95×10^{-8}	0.10	0.10	
rs7107784 <i>MIR4686</i>	G	A	All replication set	1.05 (1.03-1.08)	7.05×10^{-7}	0.0079	
			East Asian	1.001 (0.91-1.10)	9.83×10^{-1}	0.010	0.12
			South Asian	1.09 (1.03-1.15)	1.60×10^{-3}	0.20	0.25
			European	1.04 (1.02-1.07)	4.39×10^{-4}	0.43	0.28
			Mexican/Latino	1.15 (1.05-1.25)	6.00×10^{-4}	NA	0.16
		All Japanese	1.14 (1.09-1.20)	2.07×10^{-8}	0.42	0.08	
rs67839313 <i>INAFM2</i>	C	T	All replication set	1.05 (1.02-1.07)	2.91×10^{-4}	0.41	
			East Asian	1.09 (1.04-1.15)	1.93×10^{-4}	0.50	0.25
			South Asian	1.03 (0.99-1.08)	1.68×10^{-1}	0.77	0.32
			European	1.02 (0.98-1.06)	3.61×10^{-1}	0.18	0.11
			Mexican/Latino	1.06 (0.97-1.14)	1.16×10^{-1}	NA	0.19
		All Japanese	1.09 (1.06-1.12)	2.42×10^{-8}	0.73	0.27	

CI, confidence interval; OR, odds ratio; Phet, P-value for Cochran's Q-test for heterogeneity; RAF, risk allele frequency; SNP, single-nucleotide polymorphism.

Alleles are aligned to the forward strand of NCBI Build 37.

*Risk allele frequency in each population. Data are from 1000 Genomes Project Phase 3 allele frequencies (ASN, SAS, EUR, MXL and JPT).

[†]Combined association data from 21 studies (11 East Asian other than Japanese, 6 South Asian, 3 European and one Mexican/Latino).

reached genome-wide significance. In a subsequent transethnic meta-analysis, four loci were confirmed and one locus was suggested as common susceptibility loci for T2D in populations other than the Japanese population.

GWAS have been extensively performed in diverse ethnic groups, including populations of European, East Asian, South Asian and Mexican decent¹⁻¹⁸. To this point, the sample size of GWAS for European populations has grown to over 100,000 (ref. 10) and these studies have identified nearly 50 loci associated with T2D. GWAS on populations of non-European origin and transethnic GWAS meta-analysis have identified more than 30 loci associated with T2D, which were not detected in earlier European GWAS^{3-7,12-15,18}. Among these, several loci have been shown to be associated with T2D in larger European populations¹⁰, which suggests that further expansion of GWAS

for non-European populations could prove useful in identifying additional susceptibility loci associated with T2D.

Among the seven novel loci identified in this study, rs147538848 in *FAM60A*, rs1575972 near *DMRTA1*, rs7107784 near *MIR4686* and rs67839313 near *INAFM2* were shown to be common susceptibility loci for T2D across different ethnicities, although the significance of the association differed among individual ethnic groups for several loci.

rs147538848 is located in the intron of *FAM60A*, which encodes a subunit of the Sin3 deacetylase complex (Sin3/HDAC1) that has been shown to be important for the repression of genes encoding components of the transforming growth factor- β signalling pathway³¹. Studies using a rat intrauterine growth retardation model have suggested that the Sin3/HDAC1 complex may negatively regulate the expression level of pancreatic and

SNP	Chr	Gene	Score	Biological gene criteria									
				T2D risk missense variant	Monogenic diabetes	PPI	Knockout mouse phenotype	Pubmed text mining	eQTL	Nearest gene	Drug target gene	T2D drug target gene	PPI with T2D drug target gene
rs10923931	1	NOTCH2	2										
rs780094	2	GCKR	3										
rs780094	2	NRBP1	2										
rs780094	2	PPM1G	2										
rs10203174	2	THADA	2										
rs1801282	3	PPARG	5										
rs1801282	3	SYN2	2										
rs4458523	4	WFS1	5										
rs6813195	4	FBXW7	3										
rs702634	5	ARL15	2										
rs35658696	5	PAM	3										
rs9505118	6	SSR1	2										
rs115164593	6	POU5F1	2										
rs115164593	6	TCF19	2										
rs10278336	7	GCK	3										
rs10229583	7	PAX4	3										
rs4731420	7	LEP	3										
rs12549902	8	ANK1	2										
rs7845219	8	TP53INP1	2										
rs7845219	8	CCNE2	2										
rs3802177	8	SLC30A8	2										
rs7041847	9	GLIS3	4										
rs28642252	9	GPSM1	2										
rs28642252	9	INPP5E	2										
rs28642252	9	NOTCH1	2										
rs1802295	10	VPS26A	2										
rs1802295	10	SUPV3L1	2										
rs1111875	10	KIF11	2										
rs1111875	10	HHEX	3										
rs7903146	10	TCF7L2	2										
rs5215	11	ABCC8	3										
rs5215	11	KCNJ11	4										
rs11063069	12	CCND2	2										
rs12427353	12	HNF1A	4										
rs4275659	12	SETD8	2										
rs11634397	15	FAH	2										
rs11651052	17	HNF1B	5										
rs10401969	19	GATAD2A	2										
rs10401969	19	TM6SF2	2										
rs4812829	20	HNF4A	3										

Figure 3 | Prioritized biological T2D risk genes. The gene score for each gene was calculated by summing up the number of criteria satisfied (filled red box indicates criterion satisfied; 40 genes with a score ≥ 2 out of 286 genes included in the T2D risk loci were defined as 'biological candidate genes'; see Supplementary Data 4). Filled blue boxes indicate the nearest gene to the T2D risk SNP. Filled green boxes indicate overlap with drug target genes.

duodenal homeobox 1 (*PDX1*), which is known as an important transcription factor for the development of pancreas and β -cell maturation³² via histone modification of its proximal promoter³³. A T2D risk allele at the *FAM60A* locus might contribute to disease susceptibility by impairing the transcriptional regulation of genes that are important for glucose metabolism.

INAFM2 encodes InaF-motif containing 2 and has previously been known as Osteogenesis upregulated transcript 1 (*OGU1*) or long intergenic non-protein coding RNA 984 (*LINC00984*), which is a putative long non-coding RNA. Although the expression of *OGU1* has been shown to be upregulated during osteogenesis³⁴, the function of *INAFM2* encoding protein is still unknown. Around rs67839313, there are two plausible genes for

susceptibility to T2D: *PLCB2* and *DISP2*. *PLCB2* encodes phospholipase C isoform β -2 and phospholipase C is a known regulator of insulin secretion through hydrolysis of islet phosphoinositide pools³⁵. Therefore, it is feasible that this locus is associated with impaired glucose-stimulated insulin secretion machinery. *DISP2* encodes dispatched homologue 2, which is a cell surface marker on insulin-positive cells³⁶. Although the functional role of this molecule in glucose homeostasis is not well understood, it is potentially involved in the maturation of pancreatic β -cells or it might have a role in already matured pancreatic β -cells.

The effect size for the T2D association of rs1575972 near *DMRTA1* was similar among all populations in this study, except




Figure 4 | Connection of biological T2D risk genes to drug targets. Representative connections between T2D risk SNPs (blue), T2D biological genes (green), drug target genes (purple) and targeted drugs are shown (see Supplementary Table 21). *Compounds under clinical trial; **see Supplementary Table 23.

for the Mexican/Latino population (Table 2 and Supplementary Table 10). The risk allele of rs1575972 in the *DMRTA1* locus was nominally correlated with a decrease in FPI (Supplementary Table 8), which suggests that this locus might contribute to T2D susceptibility through affecting insulin secretion in pancreatic β -cells. The *DMRTA1* encodes doublesex and mab-3-related transcription factor-like family A1, which has been recently reported to be involved in neuronal development by regulating the Pax6-Neurog2 transcriptional cascade³⁷. Although the relevance of *DMRTA1* to pancreatic development has not been established, *DMRTA1* might play a role in β -cell development, because Pax6 and Neurog3, other member of the neurogenin subfamily, are key transcriptional regulators of pancreatic endocrine cell differentiation³⁸.

INS, *IGF-2* and *TH* are located at approximately rs7107784 near the *MIR4686* locus. *IGF-2* plays a key role in embryonic growth and may also influence body weight in adulthood³⁹, and *TH* (tyrosine hydroxylase) has been shown to play a role in β -cell development⁴⁰. This locus is known to be associated with risk of type-1 diabetes (rs1004446-C, 45 kbp from rs7107784; $r^2 = 0.003$, $D' = 0.275$)⁴¹. The risk allele of rs7107784-G is nominally associated with the increase of FPI levels in the European population (MAGIC data; Supplementary Table 8) and an increase of HOMA-IR in our Japanese data set (Supplementary Table 7). This suggests that the effects of rs7107784-G are probably not mediated by an impairment of insulin production or secretion, but rather by an impairment of insulin sensitivity.

rs67156297 near *ATP8B2* was nominally associated with T2D in the transethnic replication meta-analysis (Table 2 and Supplementary Table 10). *ATP8B2* encodes a member of the P4 family of ATPases (type 4P-type ATPase), which are multispan transmembrane proteins that have been implicated in phospholipid translocation from the exoplasmic to the cytoplasmic membrane leaflet⁴². The role of *ATP8B2* in the pathogenesis of T2D has not been established. However, another member of the P4 ATPase family, *atp10a*, has been shown to be important for the biogenesis and/or membrane-directed trafficking of Glut4 receptors, and loss-of-function of *atp10a* induces IR and obesity in mice⁴².

The remaining two loci, rs1116357 near *CCDC85A* and rs9309245 near *ASB3*, were not associated with T2D ($P > 0.05$) in the replication meta-analysis for non-Japanese populations, which suggests that the effect of these loci might be specific to the Japanese population. As heterogeneity in effect sizes was observed for rs1116357 or rs9309245 between Japanese and other ethnic groups, including European, South Asian and Mexican (Supplementary Tables 25 and 27), two possibilities might exist for the two SNP loci: (1) the LD between the causal alleles and the Japanese lead SNPs are consistent across the populations, but the risk alleles have effects only in the Japanese, and (2) the causal alleles are in LD with these SNPs only in the Japanese. By a systematic evaluation for effect sizes and LDs within these loci, we did not identify any SNPs associated with T2D in European populations, which are in LD with our lead SNPs in the Japanese, whereas not in LD in European populations (Supplementary Fig. 8 and Supplementary Tables 26 and 27). Therefore, the causal allele in the two loci might have an effect only in Japanese populations; however, further evaluation is required to elucidate the precise mechanism how these loci contribute to T2D susceptibility in the Japanese.

While searching for potential drug targets for T2D using a systematic bioinformatics approach, 83 overlapping genes were identified from 752 genes (40 biological T2D GWAS genes and 712 genes that encode products in direct PPI with 40 biological T2D GWAS genes) and 871 drug target genes for various human diseases²⁷. Of these, 5 were T2D GWAS genes: *PPARG*, *KCNJ11*, *ABCC8*, *GCK* and *KIF11*. *PPARG*, *KCNJ11* and *ABCC8* have approved T2D treatment options. In addition, a GCK activator is currently undergoing clinical trials for the treatment of T2D. *KIF11*, which encodes kinesin family member 11 (also known as EG5), has been shown to be involved in regulating cell mitosis and inhibitors targeting this gene product have been developed as chemotherapeutic agents in the treatment of cancer⁴³. Although the role of *KIF11* in the regulation of glucose metabolism has not been well established, a recent study reported that knockdown of *KIF11* using small interfering RNA resulted in increased glycogenesis in human primary hepatocytes⁴⁴. Thus, a *KIF11* inhibitor might ameliorate glucose homeostasis by suppressing gluconeogenesis from the liver.

We identified two genes, *GSK3B* and *JUN*, which directly interact with multiple biological T2D susceptibility genes. *GSK3B* encodes glycogen synthase kinase 3 β , which is a constitutively active multifunctional serine/threonine kinase and is involved in diverse physiological pathways, including metabolism, cell cycle regulation, gene expression, development, oncogenesis and neuroprotection⁴⁵. Several studies using *Gsk3b*-modified mouse models have suggested that inhibition of *GSK3B* function may have beneficial effects on glucose metabolism through pancreatic β -cell preservation or enhancement of insulin-stimulated glycogen synthase regulation and glycogen deposition^{45–47}. Currently, *GSK3B* inhibitors are under clinical trial for the treatment of cancers (Supplementary Table 23), but these compounds could also be potential treatments for T2D.

JUN encodes the proto-oncogene c-Jun and the role of c-Jun in the pathogenesis of T2D is not well understood. However, c-Jun has been shown to decrease the expression of the human insulin gene by repressing insulin promoter activity⁴⁸. c-Jun is a transactivation component of the heterodimeric transcription factor AP-1 and activated through phosphorylation of serines 63 and 73 by Jun N-terminal kinase 2 (ref. 49), and inhibition of JNK has been shown to ameliorate glucose intolerance in a mouse model for T2D⁵⁰. Currently, *AP-1* inhibitor is under clinical trial for the treatment of rheumatoid arthritis (Supplementary Table 23) and might also be potential treatments for T2D.

Although these results suggest that these loci are potential therapeutic targets for treating T2D, the pipeline used to identify these genes has some limitations. As eQTL effects have often been observed for genes far from each locus, it is possible that some biological genes located outside of LD block in each locus were overlooked. In addition, the selection criteria for PubMed text-mining or knockout mouse studies were based on the known functions; therefore, T2D-associated genes whose functions have not been established may have been missed. The number of criteria that were met for individual genes were simply summed for scoring, although the relative impact of the six criteria used here on biological significance may not be equal. We used the previously described scoring method²⁷, to prioritize genes in an objective manner; however, it would be worthwhile to refine the pipeline by modifying the selection criteria for genes in future studies. Finally, the potential therapeutic targets or treatments identified through the *in silico* pipeline have not yet been validated through an experimental approach. Furthermore, *in vivo* evaluation is essential to clarify the therapeutic effect of these potential T2D treatments.

In conclusion, we have identified seven novel T2D susceptibility loci using a large-scale Japanese GWAS meta-analysis. The T2D association for four of these was also observed in non-Japanese populations. In addition, we have proposed several new potential pharmacological targets for T2D treatment using a systematic bioinformatics approach. These results indicate that expansion of single ethnic GWAS is still useful to identify novel susceptibility loci to complex traits not only for ethnicity-specific but also for common loci across different ethnicities. Moreover, systematic approaches for integrating the findings of genetic, biological and pharmacological studies could be useful for developing new T2D treatments, although additional pipeline refinement would be required.

Methods

Subjects. *Discovery stage (Stage-1).* We selected T2D cases from individuals registered in BioBank Japan as having T2D (set-1 cases, $n = 9,817$). Control groups consisted of individuals registered in BioBank Japan as not having T2D but with diseases other than T2D (cerebral aneurysm, oesophageal cancer, endometrial cancer, chronic pulmonary emphysema or glaucoma) or volunteers from the Osaka-Midosuji Rotary Club and Pharma SNP consortium (set-1 controls, $n = 6,763$; Supplementary Table 24). We also used case and control individuals registered in the BioBank Japan that were previously analysed and reported (set-2 cases, $n = 5,646$ and set-2 controls, $n = 19,420$)⁷. There was no overlap in individuals in set-1 and set-2.

Validation analysis (Stage-2). We examined 7,936 T2D cases from the BioBank Japan that were not included in the discovery stage and from subjects with T2D, who visited outpatient clinics at The University of Tokyo, Juntendo University, National Center for Global Health and Medicine, Hiranuma Clinic, St Marianna University School of Medicine, The Hiroshima Atomic Bomb Casualty Council Health Management Center, Kawasaki Medical School, Toyama University Hospital or Shiga University of Medical Science. We also examined 5,539 controls from individuals that enrolled during an annual health check-up at six institutions: The Hiroshima Atomic Bomb Casualty Council Health Management Center, The National Center for Global Health and Medicine, Keio University, Hiranuma Clinic, St Marianna University School of Medicine and Toyama University Hospital. T2D was diagnosed according to World Health Organization criteria⁵¹. We excluded individuals who were positive for antibodies against glutamic acid decarboxylase and those with diabetes due to liver dysfunction, steroids and other drugs that might raise glucose levels, malignancy or a monogenic disorder known to cause diabetes.

Clinical characteristics of Stage-1 and Stage-2 participants are shown in Supplementary Table 24. Genomic DNA was extracted from peripheral leukocytes using the standard procedure. All individuals provided written informed consent to participate in this study. The protocol of this study conformed to the provisions of the Declaration of Helsinki and was approved by the ethical committees at the RIKEN Yokohama Institute and all other institutions.

Genotyping and quality control in the discovery stage. Set-1 samples were genotyped using the Human Omni Express Exome Bead Chip. There were 535,686 autosomal SNPs that passed quality control, with a call rate ≥ 0.99 , for Hardy–Weinberg equilibrium test $P \geq 1 \times 10^{-6}$ in controls and minor allele frequency (MAF) ≥ 0.01 . Set-2 samples were genotyped using the Illumina Human 610K SNP array. There were 480,426 autosomal SNPs that passed quality control and were used for further analysis. For sample quality control, we evaluated cryptic relatedness for each sample using an identity-by-state method and removed samples that exhibited second-degree or closer relatedness. We further performed principal component analysis to select individuals within the major Japanese (Hondo) cluster as reported previously^{5–7,52}, and data for 16,580 individuals (9,817 T2D cases and 6,763 controls) in set-1 and 25,066 individuals (5,646 T2D cases and 19,420 controls) in set-2 were used in subsequent analyses. To evaluate the potential effect of population stratification, we used a quantile–quantile plot of the observed P -values (Supplementary Fig. 1A,B).

Imputation. We performed genotype imputation using MACH and Minimac^{53,54} with individuals from the 1000 Genomes Project (phased JPT, CHB and Han Chinese South data $n = 275$, March 2012) as reference populations⁵⁵. We selected SNPs with MAF ≥ 0.01 and a Minimac software quality score (r^2) ≥ 0.7 . Individual genotype dosage data were used for association studies using mach2dat^{53,54}.

Genotyping and quality control in the Stage-2 analysis. We genotyped 7,936 individuals with T2D and 5,539 controls using a multiplex PCR–Invader assay, as described previously^{3,5–7}. Genotyping success rates $< 95\%$ or concordance rates $< 99.9\%$ were excluded from further evaluation.

Follow-up analyses. We obtained follow-up analysis data ($n =$ up to 223,966: 65,936 T2Ds and 158,030 controls) from multiple cohorts or a publicly available database, as described below.

East Asian populations. We obtained genotype data for up to 29,937 individuals (12,554 T2Ds and 17,383 controls), *de novo* genotyping from 2 cohorts and *in silico* replication data from 9 cohorts (Supplementary Table 9).

South Asian populations. We obtained *in silico* genotype data for a total of up to 24,965 individuals (10,587 T2Ds and 14,378 controls) from 6 cohorts (Supplementary Table 9).

European populations. We obtained genotype data for up to 160,850 individuals (38,947 T2Ds and 121,903 controls), *de novo* genotyping data from the Danish case–control study and from a publicly available database (DIAGRAM3 <http://diagram-consortium.org/downloads.html>)¹⁰. The Danish case–control study consisted of individuals from the Inter99 cohort⁵⁶, Health2006 cohort⁵⁷, Vejle Biobank⁵⁸, T2D cases from the Danish ADDITION screening cohort⁵⁹ and a T2D case–control study obtained at Steno Diabetes Center (SDC). Two SNPs were genotyped by Illumina MetaboChip in 8,781 individuals from Inter99, Health2006 and SDC, whereas four SNPs were genotyped by LGC Genomics, UK, in individuals from Inter99, Vejle Biobank, ADDITION and SDC samples (Supplementary Table 9).

Mexican/Latino population. We obtained *in silico* genotype data for up to 8,214 individuals (3,848 T2Ds and 4,366 controls) from the SIGMA Type 2 Diabetes Consortium (Supplementary Table 9).

Ethnicity was self-reported by the enrolled individuals. For each study, approval was obtained from the institutional review boards of the participating institutions and written informed consent was obtained from all participants. We excluded association data obtained by imputed genotyped data with a low quality of imputation ($r^2 < 0.7$ or info < 0.7). Details of the study samples are described in Supplementary Table 9.

Statistical analysis. The association between each SNP and T2D was assessed using the logistic regression test with an additive model with or without adjusting for age, sex and log-transformed BMI. We combined data from the each GWAS and our validation analyses using an inverse variance method and examined heterogeneity with a Cochran's *Q* test using METAL⁶⁰. Regional association plots were generated using LocusZoom⁶¹.

We also performed quantitative traits analysis for fasting plasma glucose, HOMA- β and HOMA-IR using multiple linear regression analysis in an additive association model with or without adjusting for age, sex and log-transformed BMI. The Japanese samples studied here show skewed distribution values for BMI, HOMA-IR and HOMA- β ; therefore, we have analysed the quantitative traits using log-transformed BMI, HOMA-IR and HOMA- β .

Drug discovery. We performed a search for potential drug targets using genetic information of confirmed T2D susceptibility loci and publicly available bioinformatics tools^{29,30} and databases^{28,62–65} using a method that has been previously described by Okada *et al.*²⁷ (Supplementary Note).

References

- McCarthy, M. I. Genomics, type 2 diabetes, and obesity. *N. Engl. J. Med.* **363**, 2339–2350 (2010).
- Imamura, M. *et al.* Genetics of type 2 diabetes: the GWAS era and future perspectives. *Endocr. J.* **58**, 723–739 (2011).
- Unoki, H. *et al.* SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nat. Genet.* **40**, 1098–1102 (2008).
- Yasuda, K. *et al.* Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nat. Genet.* **40**, 1092–1097 (2008).
- Yamauchi, T. *et al.* A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at *UBE2E2* and *C2CD4A-C2CD4B*. *Nat. Genet.* **42**, 864–868 (2010).
- Imamura, M. *et al.* A single-nucleotide polymorphism in *ANK1* is associated with susceptibility to type 2 diabetes in Japanese populations. *Hum. Mol. Genet.* **21**, 3042–3049 (2012).
- Hara, K. *et al.* Genome-wide association study identifies three novel loci for type 2 diabetes. *Hum. Mol. Genet.* **23**, 239–446 (2014).
- Voight, B. F. *et al.* Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nat. Genet.* **42**, 579–589 (2010).
- Dupuis, J. *et al.* New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat. Genet.* **42**, 105–116 (2010).
- Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat. Genet.* **44**, 981–990 (2012).
- Scott, R. A. *et al.* Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. *Nat. Genet.* **44**, 991–1005 (2012).
- Mahajan, A. *et al.* Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. *Nat. Genet.* **46**, 234–244 (2014).
- Kooner, J. S. Genome-wide association study in individuals of South Asian ancestry identifies six new type 2 diabetes susceptibility loci. *Nat. Genet.* **43**, 984–989 (2011).
- Cho, Y. S. Meta-analysis of genome-wide association studies identifies eight new loci for type 2 diabetes in East Asians. *Nat. Genet.* **44**, 67–72 (2011).
- The SIGMA Type 2 Diabetes Consortium. Sequence variants in *SLC16A11* are a common risk factor for type 2 diabetes in Mexico. *Nature* **506**, 97–101 (2014).
- Steinthorsdottir, V. *et al.* Identification of low-frequency and rare sequence variants associated with elevated or reduced risk of type 2 diabetes. *Nat. Genet.* **46**, 294–298 (2014).
- Moltke, I. *et al.* A common Greenlandic TBC1D4 variant confers muscle insulin resistance and type 2 diabetes. *Nature* **512**, 190–193 (2014).
- Saxena, R. *et al.* Genome-wide association study identifies a novel locus contributing to type 2 diabetes susceptibility in Sikhs of Punjabi origin from India. *Diabetes* **62**, 1746–1755 (2013).
- Stahl, E. A. *et al.* Bayesian inference analyses of the polygenic architecture of rheumatoid arthritis. *Nat. Genet.* **44**, 483–489 (2012).
- Chan, J. C. N. *et al.* Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA* **301**, 2129–2140 (2009).
- Freedman, M. L. *et al.* Assessing the impact of population stratification on genetic association studies. *Nat. Genet.* **36**, 388–393 (2004).
- Saxena, R. *et al.* Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* **316**, 1331–1336 (2007).
- Zeggini, E. *et al.* Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science* **316**, 1336–1341 (2007).
- Scott, L. J. *et al.* A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. *Science* **316**, 1341–1345 (2007).
- Manning, A. K. *et al.* A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nat. Genet.* **44**, 659–669 (2012).
- Soranzo, N. *et al.* Common variants at 10 genomic loci influence hemoglobin A_{1c} levels via glycemic and nonglycemic pathways. *Diabetes* **59**, 3229–3239 (2010).
- Okada, Y. *et al.* Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* **20**, 376–381 (2014).
- Eppig, J. T. *et al.* The Mouse Genome Database (MGD): comprehensive resource for genetics and genomics of the laboratory mouse. *Nucleic Acids Res.* **40**, D881–D886 (2012).
- Raychaudhuri, S. *et al.* Identifying relationships among genomic disease regions: predicting genes at pathogenic SNP associations and rare deletions. *PLoS Genet.* **5**, e1000534 (2009).
- Rossin, E. J. *et al.* Proteins encoded in genomic regions associated with immune-mediated disease physically interact and suggest underlying biology. *PLoS Genet.* **7**, e1001273 (2011).
- Rouse, R. *et al.* Complex SIN3-HDAC1 histone deacetylase cycle-fluctuating regulator of the protein is a cell family with sequence similarity 60A gene regulation. *J. Biol. Chem.* **287**, 32346–32353 (2012).
- Holland, A. M. *et al.* Experimental control of pancreatic development and maintenance. *Proc. Natl Acad. Sci. USA* **99**, 12236–12241 (2011).
- Park, J. H. Development of type 2 diabetes following intrauterine growth retardation in rats is associated with progressive epigenetic silencing of Pdx1. *J. Clin. Invest.* **118**, 2316–2324 (2008).
- Kikuchi, K. *et al.* Transcripts of unknown function in multiple-signaling pathways involved in human stem cell differentiation. *Nucleic Acids Res.* **37**, 4987–5000 (2009).
- Walstar, S. *et al.* Regulation of insulin secretion by phospholipase C. *Am. J. Physiol.* **271**, E409–E416 (1996).
- Hald, J. *et al.* Pancreatic islet and progenitor cell surface markers with cell sorting potential. *Diabetologia* **55**, 154–165 (2012).
- Kikkawa, T. *et al.* Dmrt1 regulates proneural gene expression downstream of Pax6 in the mammalian telencephalon. *Genes Cells* **18**, 636–649 (2013).
- Wilson, M. E. *et al.* Gene expression cascades in pancreatic development. *Mech. Dev.* **120**, 65–80 (2003).
- O'Dell, S. D. *et al.* Molecules in focus Insulin-like growth factor II (IGF-II). *Int. J. Biochem. Cell Biol.* **30**, 767–771 (1998).
- Vázquez, P. *et al.* Non-neural tyrosine hydroxylase, via modulation of endocrine pancreatic precursors, is required for normal development of beta cells in the mouse pancreas. *Diabetologia* **57**, 2339–2347 (2014).
- Hakonarson, H. *et al.* A genome-wide association study identifies KIAA0350 as a type 1 diabetes gene. *Nature* **448**, 591–594 (2007).
- Flomer, D. E. P4 ATPases—Lipid flippases and their role in disease. *Biochim. Biophys. Acta* **1791**, 628–635 (2009).
- Salmela, A. L. *et al.* Mitosis as an anti-cancer drug target. *Chromosoma* **122**, 431–449 (2013).
- Haney, S. *et al.* RNAi screening in primary human hepatocytes of genes implicated in genome-wide association studies for roles in type 2 diabetes identifies roles for CAMK1D and CDKAL1, among others, in hepatic glucose regulation. *PLoS ONE* **8**, e64946 (2013).
- Rayasam, G. V. *et al.* Glycogen synthase kinase 3: more than a namesake. *Br. J. Pharmacol.* **156**, 885–898 (2009).
- Liu, Y. *et al.* Conditional ablation of Gsk-3 β in islet beta cells results in expanded mass and resistance to fat feeding-induced diabetes in mice. *Diabetologia* **53**, 2600–2610 (2010).
- Patel, S. *et al.* Tissue-specific role of glycogen synthase kinase 3beta in glucose homeostasis and insulin action. *Mol. Cell Biol.* **28**, 6314–6328 (2008).
- Inagaki, N. *et al.* c-Jun represses the human insulin promoter activity that depends on multiple cAMP response elements. *Proc. Natl Acad. Sci. USA* **89**, 1045–1049 (1992).
- Meng, Q. H. *et al.* c-JUN, at the crossroad of the signaling network. *Protein Cell* **2**, 889–898 (2011).
- Bennett, B. L. *et al.* JNK: a new therapeutic target for diabetes. *Curr. Opin. Pharmacol.* **3**, 420–425 (2003).
- Alberti, K. G. *et al.* Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet. Med.* **15**, 539–553 (1998).
- Yamaguchi-Kabata, Y. *et al.* Japanese population structure, based on SNP genotypes from 7003 individuals compared to other ethnic groups: effects on population-based association studies. *Am. J. Hum. Genet.* **83**, 445–456 (2008).
- Li, Y. *et al.* MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genet. Epidemiol.* **34**, 816–834 (2010).
- Howe, B. *et al.* Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat. Genet.* **44**, 955–959 (2012).
- 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature* **467**, 1061–1073 (2010).
- Jørgensen, T. *et al.* A randomized non-pharmacological intervention study for prevention of ischaemic heart disease: baseline results Inter99(1). *Eur. J. Cardiovasc. Prev. Rehab.* **10**, 377–386 (2003).

57. Thuesen, B. H. *et al.* Cohort Profile: the Health2006 cohort, Research Centre for Prevention and Health. *Int. J. Epidemiol.* **43**, 568–575 (2014).
58. Albrechtsen, A. *et al.* Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. *Diabetologia* **56**, 298–310 (2013).
59. Lauritzen, T. *et al.* The ADDITION study: proposed trial of the cost-effectiveness of an intensive multifactorial intervention on morbidity and mortality among people with Type 2 diabetes detected by screening. *Int. J. Obes. Relat. Metab. Disord.* **24**, S6–S11 (2000).
60. Cristen, J. W. *et al.* METAL: fast and efficient meta-analysis of genome wide association scan. *Bioinformatics* **26**, 2190–2191 (2010).
61. Pruim, R. J. *et al.* LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336–2337 (2010).
62. Grundberg, E. *et al.* Mapping *cis*- and *trans*-regulatory effects across multiple tissues in twins. *Nat. Genet.* **44**, 1084–1089 (2012).
63. Schadt, E. E. *et al.* Mapping the genetic architecture of gene expression in human liver. *PLoS Biol.* **6**, e107 (2008).
64. Wishart, D. S. *et al.* DrugBank: a comprehensive resource for in silico drug discovery and exploration. *Nucleic Acids Res.* **34**, D668–D672 (2006).
65. Zhu, F. *et al.* Therapeutic target database update 2012: a resource for facilitating target-oriented drug discovery. *Nucleic Acids Res.* **40**, D1128–D1136 (2012).

Acknowledgements

The Mexican/Latino association data were provided by SIGMA T2D Consortium (see Supplementary Note for the contributors list). Data on glycaemic traits in European populations have been contributed by MAGIC investigators and have been downloaded from www.magicinvestigators.org/. Data on GWAS meta-analysis for T2D in European populations have been contributed by DIAGRAM consortium and have been downloaded from <http://diagram-consortium.org/>. This work was partly supported by a grant from the Leading Project of Ministry of Education, Culture, Sports, Science and Technology-Japan. The work of the Shanghai Jiao Tong University was supported from grants from the National 973 Program (2011CB504001), 863 Program (2012AA02A509) and National Science Foundation of China (81322010). R.C.W.M. and J.C.N.C. acknowledge support from the Hong Kong Foundation for Research and Development in Diabetes, established under the auspices of the Chinese University of Hong Kong, the Innovation and Technology Fund (ITS/088/08 and ITS/487/09FP), and the Research Grants Council Theme-based Research Scheme (T12-402/13-N). The work by the Shanghai Diabetes Genetic Study (SDGS) was supported in part by the US National Institutes of Health grants R37CA070867, R01CA124558, R01CA64277 and UL1 RR024975, the Department of Defense Idea Award BC050791, Vanderbilt Ingram professorship funds and the Allen Foundation Fund. We thank the dedicated investigators and staff members from research teams at Vanderbilt University, Shanghai Cancer Institute and the Shanghai Institute of Preventive Medicine, and especially the study participants for their contributions in the studies. This study was provided with data from the Korean Genome Analysis Project (4845-301), the Korean Genome and Epidemiology Study (4851-302) and Korea Biobank Project (4851-307, KBP-2013-11 and KBP-2014-68) that were supported by the Korea Center for Disease Control and Prevention, Republic of Korea. This research was supported by an intramural grant from the Korea National Institute of Health (2014-NI73001-00), Republic of Korea. This study was supported by a grant of the Korea Health Technology R&D Project, Ministry of Health and Welfare, Republic of Korea (HI14C0060). The Novo Nordisk Foundation Center for Basic Metabolic Research is an independent Research Center at the University of Copenhagen partially funded by an unrestricted donation from the Novo Nordisk Foundation (www.metabol.ku.dk). The Danish studies, Inter99 and Health2006, were partly funded by the Lundbeck Foundation and produced by The Lundbeck Foundation Centre for Applied Medical Genomics in Personalised Disease Prediction, Prevention and Care (LuCamp, www.lucamp.org). The Asian Indian Diabetic Heart Study/Sikh Diabetes Study (AIDHS/SDS) was supported by the National Institute of Health grants KO1TW006087 funded by the Fogarty International Center, R01DK082766 funded by National Institute of Diabetes and

Digestive and Kidney Diseases, and a seed grant from University of Oklahoma Health Sciences Center, Oklahoma City, USA. We thank the research participants for their contribution and support for making this study possible. A.H.C. was supported by a fellowship from CONACyT-Mexico. J.M.M. was supported by Sara Borrell Fellowship from the Instituto Carlos III, grant SEV-2011-00067 of Severo Ochoa Program and EMBO short-term fellowship, EFSO/Lilly research fellowship and Beatriu de Pinós fellowship from the Agency for Management of University and Research Grants (AGAUR). SIGMA study was supported by the Slim Foundation. Y.S.C. acknowledges support from the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (2012R1A2A1A03006155). Field-work, genotyping and standard clinical chemistry assays in PROMIS were principally supported by grants awarded to the University of Cambridge from the British Heart Foundation, UK Medical Research Council, Wellcome Trust, EU Framework 6-funded Bloodomics Integrated Project, Pfizer, Novartis and Merck. J.D. acknowledges that this work was funded by the UK Medical Research Council (G0800270), British Heart Foundation (SP/09/002), UK National Institute for Health Research Cambridge Biomedical Research Centre, European Research Council (268834) and European Commission Framework Programme 7 (HEALTH-F2-2012-279233).

Author contributions

S. Maeda and T.K. supervised the study. M. Imamura, T.Y., K.H., S. Maeda and T.K. conceived and designed the experiments. M. Imamura, T.Y., K.H., S. Maeda and T.K. wrote the manuscript. K.Y., M. Iwata, H.H., H.M., C.I., K.K., H.W., Y.T., K.T., R.K. and M.K. collected Japanese samples and generated genotype data. M.Imamura, A.T., J.H., N.S., M. Iwasaki, H.F. and K.S. performed statistical analyses for the Japanese GWAS meta-analysis. C.H., R.Z., W.J., J.L., X.-O.S., W. Zheng, R.C.W.M., J.C.N.C. and C.H.T.T. contributed in design, sample collection, GWAS assays or statistical analyses for the Chinese GWA studies. S. Moon, Y.J.K., B.-J.K., S.H.K., K.S.P. and Y.S.C. performed statistical analyses for the Korean GWAS data. N.G., T.J., M.E.J., D.R.W., I.B., C.C., T.H. and O.P. performed statistical analyses for the Danish study. E.S.T. and X.W. performed statistical analyses for the Singapore GWAS data. A.H.C., J.M.M. and J.F. performed statistical analyses for the Mexican/Latin GWAS data. W. Zhao, J.D., D. Saleheen, A.R., P.F., J.R.S., R.S. and D. Sanghera contributed study data sets and performed statistical analyses for the South Asian GWAS data. M. Imamura and Y.O. performed *in silico* analyses for drug discovery.

Additional information

Data availability: Summary statistics of 2 Japanese GWAS (study 1: 9,817 cases, 6,763 controls; study2: 5,646 cases, 19,420 controls) for directly genotyped data are available through a NBDC Human Database website (<http://humandbs.biosciencedbc.jp/en/>).

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Imamura, M. *et al.* Genome-wide association studies in the Japanese population identifies seven novel loci for type 2 diabetes. *Nat. Commun.* **7**:10531 doi: 10.1038/ncomms10531 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Minako Imamura^{1,*}, Atsushi Takahashi^{2,3,*}, Toshimasa Yamauchi^{4,*}, Kazuo Hara^{4,5,*}, Kazuki Yasuda⁶, Niels Grarup⁷, Wei Zhao⁸, Xu Wang⁹, Alicia Huerta-Chagoya¹⁰, Cheng Hu¹¹, Sanghoon Moon¹², Jirong Long¹³, Soo Heon Kwak¹⁴, Asif Rasheed¹⁵, Richa Saxena¹⁶, Ronald C.W. Ma^{17,18,19}, Yukinori Okada^{2,20}, Minoru Iwata^{21,22}, Jun Hosoe⁴, Nobuhiro Shojima⁴, Minaka Iwasaki⁴, Hayato Fujita⁴, Ken Suzuki⁴, John Danesh^{23,24,25}, Torben Jørgensen^{26,27,28}, Marit E. Jørgensen²⁹, Daniel R. Witte^{30,31}, Ivan Brandslund^{32,33}, Cramer Christensen³⁴, Torben Hansen⁷, Josep M. Mercader^{35,36,37}, Jason Flannick^{35,38}, Hortensia Moreno-Macías³⁹, Noël P. Burt³⁵, Rong Zhang¹¹, Young Jin Kim¹², Wei Zheng¹³, Jai Rup Singh⁴⁰, Claudia H.T. Tam¹⁷, Hiroshi Hirose⁴¹,

Hiroshi Maegawa⁴², Chikako Ito⁴³, Kohei Kaku⁴⁴, Hirotaka Watada^{45,46}, Yasushi Tanaka⁴⁷, Kazuyuki Tobe²¹, Ryuzo Kawamori⁴⁶, Michiaki Kubo⁴⁸, Yoon Shin Cho⁴⁹, Juliana C.N. Chan^{17,18,19}, Dharambir Sanghera^{50,51}, Philippe Frossard¹⁵, Kyong Soo Park^{14,52,53}, Xiao-Ou Shu¹³, Bong-Jo Kim¹², Jose C. Florez^{35,37,54}, Teresa Tusié-Luna¹⁰, Weiping Jia¹¹, E. Shyong Tai^{9,55,56}, Oluf Pedersen⁷, Danish Saleheen^{8,15}, Shiro Maeda^{1,57,58} & Takashi Kadowaki⁴

¹Laboratory for Endocrinology, Metabolism and Kidney Diseases, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan.

²Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan. ³Laboratory for Omics Informatics, Omics Research Center, National Cerebral And Cardiovascular Center, Suita 565-8565, Japan. ⁴Department of Diabetes and Metabolic Diseases, Graduate School of Medicine, The University of Tokyo, Tokyo 113-0033, Japan. ⁵Department of Diabetes Endocrinology, Metabolism and Rheumatology, Tokyo Medical University, Tokyo 160-0023, Japan. ⁶Department of Metabolic Disorder, Diabetes Research Center, Research Institute, National Center for Global Health and Medicine, Tokyo 162-8655, Japan. ⁷The Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen DK-2200, Denmark. ⁸Department of Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6021, USA. ⁹Saw Swee Hock School of Public Health, National University of Singapore, Singapore 138672, Singapore. ¹⁰Unidad de Biología Molecular y Medicina Genómica, Instituto de Investigaciones Biomédicas UNAM/Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City C.P.14000, Mexico. ¹¹Shanghai Diabetes Institute, Department of Endocrinology and Metabolism, Shanghai Key Laboratory of Diabetes Mellitus, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200233, China. ¹²Division of Structural and Functional Genomics, Center for Genome Science, National Institute of Health, Chungcheongbuk-do 28159, Korea. ¹³Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, Tennessee 37203-1738, USA. ¹⁴Department of Internal Medicine, Seoul National University Hospital, Seoul 03080, Korea. ¹⁵Center for Non-Communicable Diseases, Karachi, Pakistan. ¹⁶Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. ¹⁷Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China. ¹⁸Hong Kong Institute of Diabetes and Obesity, The Chinese University of Hong Kong, Hong Kong, China. ¹⁹Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China. ²⁰Department of Human Genetics and Disease Diversity, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo 113-8510, Japan. ²¹First Department of Internal Medicine, University of Toyama, Toyama 930-0194, Japan. ²²Health Administration Center, University of Toyama, Toyama 930-0194, Japan. ²³Department of Public Health and Primary Care, University of Cambridge CB1 8RN, UK. ²⁴Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus Hinxton, Cambridge CB10 1RQ, UK. ²⁵NIHR Blood and Transplant Research Unit in Donor Health and Genomics, Department of Public Health and Primary Care, University of Cambridge, Cambridge CB1 8RN, UK. ²⁶Research Centre for Prevention and Health, Capital Region of Denmark, Glostrup DK-2600, Denmark. ²⁷Department of Public Health, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen 2200, Denmark. ²⁸Faculty of Medicine, University of Aalborg, Aalborg 9220, Denmark. ²⁹Steno Diabetes Center, Gentofte 2820, Denmark. ³⁰Department of Public Health, Aarhus University, Aarhus 8000, Denmark. ³¹Danish Diabetes Academy, Odense 5000, Denmark. ³²Department of Clinical Biochemistry, Lillebaelt Hospital, Vejle 7100, Denmark. ³³Institute of Regional Health Research, University of Southern Denmark, Odense 5230, Denmark. ³⁴Medical Department, Lillebaelt Hospital, Vejle 7100, Denmark. ³⁵Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge, Massachusetts 02142, USA. ³⁶Joint BSC-CRG-IRB Research Program in Computational Biology, Barcelona Supercomputing Center, Barcelona 08034, Spain. ³⁷Center for Human Genetic Research and Diabetes Research Center (Diabetes Unit), Massachusetts General Hospital, Boston, Massachusetts 02114, USA. ³⁸Department of Molecular Biology, Harvard Medical School, Boston, Massachusetts 02115, USA. ³⁹Universidad Autónoma Metropolitana, Mexico City 09340, Mexico. ⁴⁰Central University of Punjab, Bathinda, Punjab 151001, India. ⁴¹Health Center, Keio University, Tokyo 160-8582, Japan. ⁴²Department of Medicine, Shiga University of Medical Science, Otsu 520-2192, Japan. ⁴³Grand Tower Medical Court Life Care Clinic, Hiroshima 730-0012, Japan. ⁴⁴Department of Internal Medicine, Kawasaki Medical School, Kurashiki 701-0192, Japan. ⁴⁵Department of Metabolism and Endocrinology, Juntendo University Graduate School of Medicine, Tokyo 113-8421, Japan. ⁴⁶Sportology Center, Juntendo University Graduate School of Medicine, Tokyo 113-0034, Japan. ⁴⁷Division of Metabolism and Endocrinology, Department of Internal Medicine, St Marianna University School of Medicine, Kawasaki 216-8511, Japan. ⁴⁸RIKEN Center for Integrative Medical Sciences, Yokohama 230-0045, Japan. ⁴⁹Department of Biomedical Science, Hallym University, Chunchon, Gangwon-do 24252, Republic of Korea. ⁵⁰Department of Pediatrics, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma 73104, USA. ⁵¹Department of Pharmaceutical Sciences, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma 73104, USA. ⁵²Department of Internal Medicine, Seoul National University College of Medicine, Seoul 03080, Korea. ⁵³Department of Molecular Medicine and Biopharmaceutical Sciences, Graduate School of Convergence Science and Technology, Seoul National University, Seoul 03080, Korea. ⁵⁴Department of Medicine, Harvard Medical School, Boston, Massachusetts 02115, USA. ⁵⁵Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119228, Singapore. ⁵⁶Duke-National University of Singapore Graduate School, Singapore 169857, Singapore. ⁵⁷Department of Advanced Genomic and Laboratory Medicine, Graduate School of Medicine, University of the Ryukyus, Nishihara 903-0215, Japan. ⁵⁸Division of Clinical Laboratory and Blood Transfusion, University of the Ryukyus Hospital, Nishihara 903-0215, Japan.

* These authors contributed equally to this work.