

Rivaroxaban for stroke prevention in people with atrial fibrillation and diabetes mellitus

Li, Y-G; Pastori, Daniele; Lip, G Y H

Published in:
Diabetic Medicine Online

DOI (link to publication from Publisher):
[10.1111/dme.13685](https://doi.org/10.1111/dme.13685)

Publication date:
2018

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Li, Y.-G., Pastori, D., & Lip, G. Y. H. (2018). Rivaroxaban for stroke prevention in people with atrial fibrillation and diabetes mellitus. *Diabetic Medicine Online*, 35(8), 1134–1135. <https://doi.org/10.1111/dme.13685>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Article type : Letter

Title: Diabetic Medicine
Created by: Maria Davie
Email proofs to: g.y.h.lip@bham.ac.uk
Article no.: DME-2018-00186
Article type: Letter
Figures:0; Tables:0; Equations:0; References: 9

Rivaroxaban for stroke prevention in people with atrial fibrillation and diabetes mellitus

Atrial fibrillation (AF) is associated with various cardiovascular comorbidities, such as hypertension, heart failure and diabetes mellitus, which increases not only the risk of ischaemic stroke but also of myocardial infarction and mortality. Oral anticoagulation for stroke prevention is the cornerstone of management for people with AF, both with vitamin K antagonists, or with non-vitamin K antagonist oral anticoagulants (NOACs), which show efficacy, safety and convenience compared with vitamin K antagonists [1].

In a recent issue of *Diabetic Medicine*, Coleman *et al.* [2] reported on their 'real-world' retrospective study which used US national MarketScan claims data. Their study included a large sample size of 11 034 people with AF, with a median 1.5 years of follow-up, and compared the performance of rivaroxaban with that of propensity-score matched warfarin ($n=5517$ in each arm). The authors found

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/dme.13685

This article is protected by copyright. All rights reserved.

that standard-dose rivaroxaban (20 mg) resulted in a non-significant difference in the efficacy endpoint (stroke or systemic embolism and ischaemic stroke alone), but reduced-dose rivaroxaban (15 mg, used in ~20% of participants) was associated with a 67% lower risk of stroke or systemic embolism (hazard ratio 0.33, 95% CI 0.13--0.79) and an 80.0% lower risk of ischaemic stroke (hazard ratio 0.20, 95% CI 0.07--0.62). Also, both standard and reduced doses of rivaroxaban were associated with similar major bleeding and intracranial haemorrhage compared with warfarin. Based on the results of this study, rivaroxaban was similar to warfarin with regard to effectiveness and safety in people with non-valvular AF and diabetes who were seen in routine clinical practice.

The effectiveness results were generally consistent with the overall results from the phase III randomized ROCKET-AF (Rivaroxaban Once Daily Oral Direct Factor Xa Inhibition Compared with Vitamin K Antagonism for Prevention of Stroke and Embolism Trial in Atrial Fibrillation) trial, which showed that rivaroxaban was non-inferior to warfarin in reducing ischaemic stroke or systemic embolism in participants [3]. In the ROCKET-AF trial, 40.0% participants had diabetes, and a subgroup analysis showed a twofold increase in the risk of ischaemic stroke amongst people with diabetes compared with those without diabetes; nevertheless, rivaroxaban showed similar results in terms of efficacy and safety in people with AF with and without diabetes [4].

Interestingly, Coleman *et al.* [2] found that rates of ischaemic stroke/systemic embolism in people with diabetes were lower than those seen in the participants with diabetes in the ROCKET-AF trial (0.87 vs 1.74 per 100 patient-years, respectively) [4]. In the international XANTUS (Xarelto on prevention of stroke and non-central nervous system systemic embolism in people with non-valvular atrial fibrillation) registry, which included 6784 participants treated with rivaroxaban from 311 centres in Europe, Israel and Canada, the overall rate of ischaemic stroke/systemic embolism was 0.8 per 100 patient-years [5]. Lower incidence rates of major bleeding on rivaroxaban were also reported by Coleman *et al.* [2] (2.7 per 100 patient-years) than in people with diabetes in the ROCKET-AF trial (3.79 per 100 patient-years) [4]. Major bleeding rates with rivaroxaban treatment

were also low, at 2.1 events per 100 patient-years in the XANTUS study [5] and 3.0 per 100 patient-years in the Dresden NOAC registry [6].

These inter-study rates should be interpreted with caution, however, given the differences in study designs, settings and patient profiles. Also, outcomes in observational registries are not necessarily independently adjudicated as they are in randomized trials, and the reimbursement for sponsored registries may introduce some channelling biases; that is, negative results are less likely to be published. Indeed, evidence from real-world studies is confounded by methodology issues including patient selection, duration of follow-up and definition of outcomes. Despite reflecting more accurately the clinical characteristics of patients encountered in clinical practice, observational data can only establish an association between the studied variables rather than a cause--effect relationship.

Overall, these data would suggest that in the real-world of people with AF treated with rivaroxaban, the rates of outcomes (both ischaemic and bleeding) are lower than those observed in clinical trials [7]. We also need to see results on cardiovascular outcomes from the cohort studied by Coleman *et al.* [2], as people with diabetes are at elevated risk of myocardial infarction and vascular death. In the subgroup of people with diabetes in the ROCKET-AF trial, for example, the rate of myocardial infarction was 1.35 vs 0.75 ($P<0.0001$) per 100 patient-years in people with vs without diabetes, respectively [4]. Explanations for the difference in stroke outcomes between standard- and reduced-dose rivaroxaban regimens also require further study. Treatment adherence and patient values and preferences should also be addressed, given the importance of these aspects in AF management in the era of NOAC [8,9].

In conclusion, the study by Coleman *et al.* [2] gives us some confidence in prescribing rivaroxaban in people with AF and diabetes. Indeed, evidence from observational studies will continue playing an important role in orientating NOAC prescription in everyday clinical practice.

Competing interests

The authors have no competing interests directly related to this paper. G.Y.H.L. has been a consultant for Bayer/Janssen, BMS/Pfizer, Medtronic, Boehringer Ingelheim, Novartis, Verseon and Daiichi-Sankyo and a speaker for Bayer, BMS/Pfizer, Medtronic, Boehringer Ingelheim, and Daiichi-Sankyo. No fees were directly received personally.

Y.-g. Li^{1,2}, Daniele Pastori^{1,3} and G. Y. H. Lip^{1,3,4}

¹Institute of Cardiovascular Sciences, University of Birmingham, Birmingham, UK, ²Department of Cardiology, Chinese PLA General Hospital, Chinese PLA Medical School, Beijing, China, ³I Clinica Medica, Atherothrombosis Centre, Department of Internal Medicine and Medical Specialties, Sapienza University, Rome, Italy and ⁴Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

References

1. Lip G, Freedman B, De Caterina R, Potpara TS. Stroke prevention in atrial fibrillation: Past, present and future. Comparing the guidelines and practical decision-making. *Thromb Haemost* 2017; **117**: 1230--1239.
2. Coleman CI, Bunz TJ, Eriksson D, Meinel AK, Soode NA. Effectiveness and safety of rivaroxaban vs warfarin in people with non-valvular atrial fibrillation and diabetes: an administrative claims database analysis. *Diabet Med* 2018; doi: 10.1111/dme.13648. [Epub ahead of print].
3. Patel MR, Mahaffey KW, Garg J, Pan G, Singer DE, Hacke W *et al.* Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med* 2011; **365**: 883--891.
4. Bansilal S, Bloomgarden Z, Halperin JL, Hellkamp AS, Lokhnygina Y, Patel MR *et al.* Efficacy and safety of rivaroxaban in patients with diabetes and nonvalvular atrial fibrillation: the

Rivaroxaban Once-daily, Oral, Direct Factor Xa Inhibition Compared with Vitamin K Antagonism for Prevention of Stroke and Embolism Trial in Atrial Fibrillation (ROCKET AF Trial). *Am Heart J* 2015; **170**: 675--682 e678.

5. Camm AJ, Amarenco P, Haas S, Hess S, Kirchhof P, Kuhls S *et al*. XANTUS: a real-world, prospective, observational study of patients treated with rivaroxaban for stroke prevention in atrial fibrillation. *Eur Heart J* 2016; **37**: 1145--1153.

6. Hecker J, Marten S, Keller L, Helmert S, Michalski F, Werth S *et al*. Effectiveness and safety of rivaroxaban therapy in daily-care patients with atrial fibrillation. Results from the Dresden NOAC Registry. *Thromb Haemost* 2016; **115**: 939--949.

7. Beyer-Westendorf J, Camm AJ, Coleman CI, Tamayo S. Rivaroxaban real-world evidence: Validating safety and effectiveness in clinical practice. *Thromb Haemost* 2016; **116**: S13--S23.

8. Martinez C, Katholing A, Wallenhorst C, Freedman SB. Therapy persistence in newly diagnosed non-valvular atrial fibrillation treated with warfarin or NOAC. A cohort study. *Thromb Haemost* 2016; **115**: 31--39.

9. Raparelli V, Proietti M, Cangemi R, Lip GY, Lane DA, Basili S. Adherence to oral anticoagulant therapy in patients with atrial fibrillation. Focus on non-vitamin K antagonist oral anticoagulants. *Thromb Haemost* 2017; **117**: 209--218.