

There it is! *Fusarium pseudograminearum* did not lose the fusaristatin gene cluster after all

Wollenberg, Rasmus Dam; Sondergaard, Teis Esben; Nielsen, Mikkel Rank; Knutsson, Simon; Pedersen, Tobias Bruun; Westphal, Klaus Ringsborg; Wimmer, Reinhard; Gardiner, Donald Max; Sørensen, Jens Laurids

Published in:
Fungal Biology

DOI (link to publication from Publisher):
[10.1016/j.funbio.2018.10.004](https://doi.org/10.1016/j.funbio.2018.10.004)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2019

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Wollenberg, R. D., Sondergaard, T. E., Nielsen, M. R., Knutsson, S., Pedersen, T. B., Westphal, K. R., Wimmer, R., Gardiner, D. M., & Sørensen, J. L. (2019). There it is! *Fusarium pseudograminearum* did not lose the fusaristatin gene cluster after all. *Fungal Biology*, 123(1), 10-17. <https://doi.org/10.1016/j.funbio.2018.10.004>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: June 18, 2025

Accepted Manuscript

There it is! *Fusarium pseudograminearum* did not lose the fusaristatin gene cluster after all

Rasmus Dam Wollenberg, Teis Esben Sondergaard, Mikkel Rank Nielsen, Simon Knutsson, Tobias Bruun Pedersen, Klaus Westphal, Reinhard Wimmer, Donald Max Gardiner, Jens Laurids Sørensen

PII: S1878-6146(18)30350-7

DOI: <https://doi.org/10.1016/j.funbio.2018.10.004>

Reference: FUNBIO 965

To appear in: *Fungal Biology*

Received Date: 8 February 2018




Revised Date: 27 August 2018

Accepted Date: 17 October 2018

Please cite this article as: Wollenberg, R.D., Sondergaard, T.E., Nielsen, M.R., Knutsson, S., Pedersen, T.B., Westphal, K., Wimmer, R., Gardiner, D.M., Sørensen, J.L., There it is! *Fusarium pseudograminearum* did not lose the fusaristatin gene cluster after all, *Fungal Biology*, <https://doi.org/10.1016/j.funbio.2018.10.004>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.





There it is! *Fusarium pseudograminearum* did not lose the fusaristatin gene cluster after all.

Rasmus Dam Wollenberg¹, Teis Esben Sondergaard¹, Mikkil Rank Nielsen², Simon Knutsson², Tobias Bruun Pedersen², Klaus Westphal¹, Reinhard Wimmer¹, Donald Max Gardiner³, Jens Laurids Sørensen^{2,*}

¹ Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark.

² Department of Chemistry and Bioscience, Aalborg University, Esbjerg, Denmark.

³ Commonwealth Scientific and Industrial Research Organization (CSIRO) Agriculture and Food, Queensland Bioscience Precinct, Brisbane, Australia.

* Corresponding author: Jens Laurids Sørensen e-mail: jls@bio.aau.dk. Aalborg University Esbjerg, Department of Chemistry and Bioscience, Niels Bohrs Vej 8, 6700 Esbjerg, Denmark

Summary

Fusarium pseudograminearum is a significant pathogen of cereals in arid regions worldwide and has the ability to produce numerous bioactive secondary metabolites. The genome sequences of seven *F. pseudograminearum* strains have been published and in one of these strains, C5834, we identified an intact gene cluster responsible for biosynthesis of the cyclic lipopeptide fusaristatin A. The high level of sequence identity of the fusaristatin cluster remnant in strains that do not produce fusaristatin suggests that the absence of the cluster evolved once, and subsequently the resulting locus with the cluster fragments became widely dispersed among strains of *F. pseudograminearum* in Australia. We examined a selection of 99 Australian *F. pseudograminearum* isolates to determine how widespread the ability to produce fusaristatin A is in *F. pseudograminearum*. We identified 15 fusaristatin producing strains, all originating from Western Australia. Phylogenetic analyses could not support a division of *F. pseudograminearum* into fusaristatin producing and nonproducing populations, which could indicate the loss has occurred relatively recent.

Keyword: Secondary metabolites; polyketides; non-ribosomal peptides; Fusarium Crown Rot; evolution

Introduction

Fusarium pseudograminearum is the primary cause of Fusarium crown rot (FCR) of wheat and barley in the arid cereal growing regions of the world including Australia (Burgess et al. 2001), Southern Europe (Balmas 1994), Northern Africa (Gargouri et al. 2011), South Africa (Lamprecht et al. 2006), China (Ji et al. 2016; Li et al. 2012; Xu et al. 2017) and the United States of America (Smiley et al. 2005). The disease is one of the most severe in cereals in Australia with yearly economic losses of approximately 100 million Australian dollars (Murray and Brennan 2009, 2010). *F. pseudograminearum* is heterothallic (Aoki and O'Donnell 1999b; Summerell et al. 2001) and was initially recognized as a population within the *F. graminearum* species group (Group 1) based on cultivation and its inability to form homothallic perithecia (Burgess et al. 1975; Francis and Burgess 1977). Later, the two species were formally segregated by molecular analyses (Aoki and O'Donnell 1999a) and further sequence analyses suggested that *F. pseudograminearum* is a single globally occurring species (Scott and Chakraborty 2006), while *F. graminearum* can be divided into more than 16 phylogenetically distinct species (Aoki et al. 2012; O'Donnell et al. 2000). *F. graminearum* is involved in Fusarium head blight (FHB) in cereals, a disease which *F. pseudograminearum* has only been observed to cause in Australia (Backhouse et al. 2004) and China (Ji et al. 2016). Both species are known producers of the trichothecene mycotoxin deoxynivalenol (and derivatives) and of the mycoestrogen zearalenone (Sydenham et al. 1991).

Comparative analyses of the first genome sequenced strains of *F. graminearum* (NRRL 31084) and *F. pseudograminearum* (CS3096) revealed only minor differences in the composition of polyketide synthases (PKSs) and non-ribosomal peptide synthetases (NRPSs) (Hansen et al. 2015). The two strains differ, however, in their ability to produce polyketide lipopeptides: in their ability to produce two polyketide lipopeptides: *F. graminearum* NRRL 31084 produces fusaristatin A but not W493,

while *F. pseudograminearum* CS3096 produces W493 but not fusaristatin A (**Figure 1**; (Sørensen et al. 2014a)).

Biosynthesis of W493 and fusaristatin A are suggested to follow similar routes starting with production of a partially reduced polyketide which serves as a substrate for a NRPS that catalyzes the condensation of the polyketide and amino acids before the compounds are released by cyclization (Sørensen et al. 2014a). The key enzymes involved in biosynthesis of W493 are PKS32, which produces a reduced polyketide (C₁₄) chain and NRPS40, which catalyzes condensation of six amino acids (threonine, alanine, alanine, glutamine, tyrosine and valine/isoleucine (W493-A/W493-B)). Fusaristatin biosynthesis is initiated by production of a reduced polyketide (C₂₄) by PKS6 prior to incorporation of three amino acids (dehydroalanine, β -aminoisobutyric acid and glutamine) by NRPS7.

The fusaristatin gene cluster has also been identified in the more distantly related *Botrytis fuckeliana*, *Cochliobolus heterostrophus* and *Pyrenophora teres* (Sieber et al. 2014). Following the first genome release of a *F. pseudograminearum* strain, six additional strains were published (Gardiner et al. 2017; Moolhuijzen et al. 2013). In one of these strains, CS5834, we identified the intact fusaristatin gene cluster and the aim of the current study was to determine how common this cluster is in *F. pseudograminearum* and whether its presence or absence arose from a gain or loss event.

79 **Materials and methods**

80 *Fungal strains*

81 Ninety-nine strains of *F. pseudograminearum* were obtained from the CSIRO collection in Brisbane
82 Australia. These strains were isolated from four different Australian states; New South Wales (42
83 strains), Queensland (18 strains), South Australia (4) and Western Australia (35).

84

85 *Fusaristatin gene cluster analyses*

86 The available genome sequences of seven *F. pseudograminearum* strains (CS3096, CS3220,
87 CS3270, CS3427, CS3487, CS5834 and RBG5266) were screened for presence of the fusaristatin
88 gene cluster using the published gene cluster from *F. graminearum* (Sørensen et al. 2014a).
89 Remnant fragments of the fusaristatin gene cluster were identified though BlastN analyses (Altschul
90 et al. 1990) using the fusaristatin gene cluster from *F. pseudograminearum* CS5834 against the
91 whole-genome sequence (WGS) database of the six other *F. pseudograminearum* strains.

92

93 *Analyses of W493-B and fusaristatin A production*

94 For secondary metabolite analyses the 99 *F. pseudograminearum* strains were cultivated on solid
95 yeast extract sucrose (YES) agar medium (Sørensen and Sondergaard 2014) and corn meal agar
96 (CM; corn meal 60 g/L, ZnSO₄ x 7 H₂O 10 mg/L, CuSO₄ x 5 H₂O 5 mg/L, agar 20 g/L) medium for
97 two weeks in the dark at 25 °C. The extraction of secondary metabolites were performed as
98 previously described (Smedsgaard 1997). The resulting extracts were analyzed on a Hitachi Elite
99 LaChrom HPLC system equipped with a 150 x 4.6 mm Ascentis Xpress 2.7 µm phenyl-hexyl
100 column (Sigma-Aldrich, USA) and coupled to a high resolution mass spectrometer (compact qTOF,
101 Bruker, Germany) with an electrospray source using a 3:97 flowsplitter. 40 µL extract was

separated using a flow of 1 mL/min with a linear water–acetonitrile gradient, with both eluents buffered with 0.1% formic acid. The gradient started at 10% acetonitrile and reached 100% in 20 min, which was held for 5 min.

Determination of presence or absence of the fusaristatin gene cluster

The fungal strains were cultivated in 30 mL liquid Czapek dox (Sigma-Aldrich) medium prior to DNA extraction. The cultivated fungi were filtered through sterile MiraCloth (Calbiochem®) and ground in liquid nitrogen before genomic DNA was extracted with the DNeasy® Plant Mini Kit (Qiagen, Hilden, Germany) (Droce et al. 2013). The isolated genomic DNA served as template in a polymerase chain reaction (PCR) targeting *PKS6* with primers *PKS6conFw* (5'-3': CTG TTG TTG GCA TGA GTT GC) and *PKS6conRv* (5'-3': TGG CCC ATG CGA GGA TAC TG), which amplify a 1751 bp product in strains with intact *PKS6* and 1564 bp product in strains with *PKS6* remnants. The PCR reactions were performed in 50 µL volume using the Phusion Hot Start II DNA Polymerase (Thermo Fisher Scientific) according to manufactures protocol. The resulting PCR products were run on 1% agarose gels with 1 kbp plus DNA ladder (Thermo Fisher Scientific).

*Phylogenetic analyses of *F. pseudograminearum* strains*

For phylogenetic analyses the primers *PHO1* (5'-3': ATC TTC TGG CGT GTT ATC ATG) and *PHO6* (5'-3': GAT GTG GTT GTA AGC AAA GCC C) were used to amplify a fragment of the Phosphate permease gene (FPSE_11047 in *F. pseudograminearum* CS3096) (Scott and Chakraborty 2006) by PCR. The PCR products were purified with the QIAquick PCR purification kit (Qiagen, Hilden, Germany) and sequenced at Eurofins Genomics (Ebersberg, Germany) using the forward primer *PHO1*. The sequences were aligned with by multiple alignment using fast

fourier transform (MAFFT) at the T-REX web server (Boc et al. 2012). The alignments were analysed with CLC main workbench (CLC Bio, Qiagen, Germany) using maximum likelihood with 1000 bootstraps and visualized with EvolView (<http://evolgenius.info/evolview>) (Zhang et al. 2012).

Whole-Genome Sequencing

With minor modifications, genomic DNA was extracted from strains CS3894, CS3900, CS5541, CS7093, CS7108, CS7081, CS7088, CS7065 and CS7060 using the FastDNATM SPIN kit for Soil (MP Biomedicals, USA). Following clean-up with Agencourt AMPure XP beads (Beckman Coulter, USA), 2 µg DNA was used as input for the SQK-LSK8 ligation sequencing kit protocol (NBE_9006_v103_revQ_21Dec2016). The protocol was modified to allow for barcoding with the Native Barcoding Kit (EXP-NBD103, Oxford Nanopore Technologies, UK) directly following the end-prep step and for downstream compatibility with sequencing on the PromethION alfa/beta sequencer (Oxford Nanopore Technologies, UK). Briefly, 10 µL Native barcode (NB01-NB9) was mixed with 30 µL end-prepped DNA mix (2 µg DNA), 10 µL nuclease-free water, 40 µL Ultra II ligation master mix (New-England Biolabs, USA), 1 µL ligation enhancer (New-England Biolabs, USA) and incubated at room temperature for 10 minutes before being further processed according to the PromethION SQK-LSK9 protocol (GDLE_9056_v109_revE_02Feb2018). Approximately 600 ng of pooled DNA was loaded onto a primed FLO-PRO001 flow-cell (Oxford Nanopore Technologies, UK) and sequenced on the PromethION alfa/beta sequencer with live base-calling enabled. Approximately 60 Gbp reads were demultiplexed and trimmed in Porechop version 0.2.3 and subsequently mapped to the reference genome of *F. pseudograminearum* CS3096 (Gardiner et al. 2017) in CLC Genomics Workbench version 9.5.5 (CLC Bio, Qiagen, Germany). Consensus sequences from the complete genes of beta-tubulin (FPSE_03337), translation elongation factor 1-

alfa (FPSE_11980), trichothecene 3-O-acetyltransferase (FPSE_11049), ammonia-ligase (FPSE_11050) and phosphate permease (FPSE_11047) were finally extracted for phylogenetic analysis (O'Donnell et al. 2000). The alignment was executed with MUSCLE (Edgar 2004). A few nucleotides (1-3 pr. sequence) resulting in non-sense mutation were excluded from the final alignments to eliminate Nanopore sequencing-biases (in some homopolymeric nucleotide-region). The alignments were fused and analysed using the same approach as for the phosphate permease gene. CANU version 1.7 was used to assemble the genome of CS3894 with default settings (genome size set at 36 gbp) (Koren et al. 2017).

Results and discussion

The fusaristatin cluster is conserved in F. pseudograminearum CS5834

The predicted fusaristatin cluster in *F. pseudograminearum* CS5834 was initially compared to the published clusters in *F. graminearum* and *F. avenaceum* (Sørensen et al. 2014a; Sørensen et al. 2014b). The comparison showed that the hypothetical proteins are of comparable length and identity (**Table 1**) suggesting that the gene cluster is also functional in *F. pseudograminearum* CS5834. Based on their phylogenetic relationship (Kristensen et al. 2005; O'Donnell et al. 2013) it was not surprising that a higher identity was observed to *F. graminearum* (94-98 %) than to *F. avenaceum* (73-86 %).

Further analyses of the available *Fusarium* genome sequences revealed that the fusaristatin gene cluster is present with conserved synteny in *F. pseudograminearum* CS5834, *F. graminearum*, *F. culmorum*, *F. meridionale*, *F. asiaticum*, *F. langsethiae*, *F. acuminatum* and *F. avenaceum* (**Figure 2**). The flanking genes were, however, different in *F. avenaceum* and *F. acuminatum* compared to the other *Fusarium* species, indicating that the cluster is present in a different genomic location

these two species. The identical location of the fusaristatin cluster in *F. pseudograminearum* CS5834 and the majority of other *Fusarium* species suggests that CS5834 did not acquire the cluster through horizontal gene transfer. This in turn suggests that the fusaristatin cluster was present in *F. pseudograminearum* after it diverged from other fusaria but was subsequently lost. To further investigate the nature of the loss, we examined the genomic region between the flanking genes of the fusaristatin gene cluster by which five conserved remnant fragments (88-95 % sequence identity) of the cluster could be found in all six *F. pseudograminearum* strains (**Figure 2A**). One of the fragments (R1; 897 bp) originates from a predicted aminotransferase gene (BN849_0052070), three other fragments (R2-R4; 120, 446 and 273 bp, respectively) originate from *PKS6* (BN849_0052040) while a fifth fragment (R5; 407 bp) originates from *NRPS7* (BN849_0052030). To illustrate that the fragments originate from *PKS6* the three remnant fragments of *PKS6* in *F. pseudograminearum* CS3096 were translated into amino acid sequences and aligned against the functional *PKS6* of *F. pseudograminearum* CS5834 (**Figure 2B**). In these alignments, a high sequence identity was observed for the three fragments as R2 had 90% (60 amino acids), R3 had 89% (148 amino acids) and R4 had 82% identity (91 amino acids). The presence of conserved remnant fragments suggests that the missing fusaristatin gene cluster is a result of a deletion event in a common ancestor.

Fusaristatin-producing F. pseudograminearum strains are geographically co-localized

The distribution of the fusaristatin-producing ability in Australian *F. pseudograminearum* strains was further investigated through chemical analyses of the 99 strains, which originated from five different states (New South Wales, Queensland, South Australia and Western Australia). The analyses showed that while nearly all strains (except CS3002 and CS5897) were able to produce W493-B only 15 strains produced fusaristatin when cultivated on solid YES or CM medium (**Table**

2). The ability to produce fusaristatin seemed to be geographically confined, because all 15 fusaristatin A-producing strains were isolated from Western Australia. Although a slight decrease in W493-B levels was observed in the fusaristatin A producers, this difference was not significant ($P>0.05$; Supplementary Figure 1).

Fusaristatin producing isolates do not form a unique lineage

The phosphate permease gene was partially sequenced (807 of 1851 bp) to investigate whether fusaristatin-producing and nonproducing strains constitute phylogenetically distinct lineages of *F. pseudograminearum*. Assumedly, this locus is inherited independently of the fusaristatin gene cluster, as they are located on two different chromosomes. The phosphate permease gene is located near the middle of chromosome IV, while the fusaristatin gene cluster is located near and end of chromosome II. Phylogenetic analyses of the resulting sequences resulted in a tree with two major clades (**Figure 3A**), separated by 26 variable sites (3%). The first clade contained the majority of the strains isolated from New South Wales (40/42) and Queensland (16/18). Three nonproducers of fusaristatin A from Western Australia were also present in clade I, while the remaining thirty-two strains were located in the second clade. This second clade consisted of two different sequence types, sharing 805 of 807 nucleotides and contained both fusaristatin producers and nonproducers without any signs of segregation.

In a further attempt to achieve a phylogenetic separation of fusaristatin producers and nonproducers, we performed a multiplexed genome sequencing of four producers (CS5541, CS7108, CS7081, and CS7060) and five nonproducers (CS3894, CS3900, CS7065, CS7088, CS7093). In addition to the phosphate permease gene, sequences of five genes were extracted (β -tubulin, translation elongation factor 1 α , trichothecene 3-O-acetyltransferase and ammonia-ligase) and used to generate an additional phylogenetic tree. The resulting tree failed to separate fusaristatin producers and

nonproducers, although this combination of genes has previously been used to separate *F. graminearum* into different phylogenetic species (**Figure 3C**). Due to the inadequacy of this multigene approach, future studies could focus on full genome analyses in order to determine whether producers and nonproducers of fusaristatin can be separated into two groups.

The lack of fusaristatin production in a strain does not necessarily mean that the strain does not have a functional fusaristatin gene cluster, because lack of production can also be caused by too low production levels or repression under the tested conditions. A PCR based strategy was used to determine the presence or absence of a functional *PKS6* yielding predicted products of 1751 bp in strains with an intact *PKS6* and 1564 bp in strains with *PKS6* remnant fragments. Thus, the two fragments are markers for the two alternative alleles of the locus (i.e., an intact and a deleted gene cluster) based on available genome sequence data. The results showed that the PCR of the 15 fusaristatin producing strains resulted in amplified fragments of the expected size for the intact and functional *PKS6* (**Figure 3B**). The PCR fragments for all the nonproducing strains, except CS3894, were smaller, which corresponds to the presence of the *PKS6* remnant region. The slightly larger PCR fragment in CS3894 was investigated further using the full genome sequence of CS3894, which showed that overall the sequence was very similar to the nonproducing CS3096 remnant region with the exception of an additional 100 bp (**Supplementary Figure 2**) which accounts for the intermediate size of the band observed for this isolate (**Figure 3B**).

Together the molecular analyses suggests that the presence of the fusaristatin gene cluster is reflected to some extent in the phylogenetic analyses of genes used in the present study. However, the genes do not contain sufficient variation to segregate the strains into clades reflecting the ability to produce fusaristatin A. A phylogenetic analysis of *F. pseudograminearum* based on the phosphate permease, reductase, translation elongation factor-1 α and β -tubulin genes concluded that *F. pseudograminearum* is a single monophyletic species (Scott and Chakraborty 2006). The high

sequence conservation within *F. pseudograminearum* is also reflected in the RNA polymerase II largest (RPB1) and second largest subunit (RPB2) genes, which have been successfully used for separating closely related *Fusarium* species (O'Donnell et al. 2013). In these genes CS3096 and CS5834 share high sequence identity (1604/1606 and 901/902).

The loss of the fusaristatin gene cluster in *F. pseudograminearum* could represent an evolutionary development where the compound is not needed for spread and survival. Biosynthesis of huge proteins, like PKS6 and NRPS7, represent a significant energy cost for the fungus; thus, losing the redundant gene cluster can result in an improved fitness.

One of the reasons for losing the fusaristatin gene cluster could be due to an overlapping mode of action for W493 and fusaristatin A, which is not an unlikely scenario given their similar biosynthetic background and structural similarities. The high level of identity of the sequence of the remnant fusaristatin cluster in strains CS3096, CS3220, CS3270, CS3427, CS3487 and RBG5266 suggests that presence a deletion event occurred in one strain or lineage of the fungus rather than multiple times in multiple strains or lineages. However, the presence of the additional region in CS3894 suggests that some modifications has occurred locus where the fusaristatin gene cluster was lost. Understanding when this loss event occurred may provide some indication of the evolutionary reason for the absence of the cluster in most strains. The climatic conditions (and native grass populations) in WA can be drastically different to the eastern states of Australia. The restricted geographic location of isolates containing the fusaristatin cluster may suggest different evolutionary pressures exist in WA but the widespread (and overlapping) presence of isolates carrying the cluster loss in the same location and the absence of obvious lineages are contrary to this scenario.

Although *Fusarium* crown rot has likely been present in WA for a long time, it has only recently emerged as a significant economic impediment to wheat production in this area (Murray and Brennan 2009). Further complicating our understanding of the evolutionary pressures that have

shaped the *F. pseudograminearum* genome is the likelihood that *F. pseudograminearum*, like *F. graminearum*, has not co-evolved with wheat (Lofgren et al. 2018) and can be considered an opportunistic pathogen of wheat. Thus, it will be extremely challenging to pinpoint the reason for loss of the cluster or even whether maintaining the clusters provides some advantage in the WA environment.

Acknowledgements

The study was funded by the Danish Research Council, Technology and Production (grant no. 4005-00204B) and the Novo Nordisk Foundation (NNF15OC0016028 and NNF15OC0016186). The collection of isolates used in this study was funded in various projects by the Australian Grain Research and Development and Research Corporation and CSIRO. Isolate RBG5266 was a kind gift from the Royal Botanic Gardens of Sydney.

The authors wish to thank laboratory trainees Belinda Sandvang Jensen and Karoline Andersen for extracting DNA and Camilla Juel Høj for extracting secondary metabolites from *F. pseudograminearum*. The NMR laboratory at Aalborg University is supported by the Obel, SparNord and Carlsberg Foundations.

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ, 1990. Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403-410.
- Aoki T, O'Donnell K, 1999a. Morphological and molecular characterization of *Fusarium pseudograminearum* sp nov., formerly recognized as the Group 1 population of *F. graminearum*. *Mycologia* **91**, 597-609.
- Aoki T, O'Donnell K, 1999b. Morphological characterization of *Gibberella coronicola* sp. nov., obtained through mating experiments of *Fusarium pseudograminearum*. *Mycoscience* **40**, 443-453.

- Aoki T, Ward TJ, Kistler HC, O'Donnell K, 2012. Systematics, phylogeny and trichothecene mycotoxin potential of *Fusarium* head blight cereal pathogens. *JSM Mycotoxins* **62**, 91-102.
- Backhouse D, Abubakar AA, Burgess LW, Dennisc JI, Hollaway GJ, Wildermuth GB, Wallwork H, Henry FJ, 2004. Survey of *Fusarium* species associated with crown rot of wheat and barley in eastern Australia. *Australasian Plant Pathology* **33**, 255-261.
- Balmas V, 1994. Root rot of wheat in Italy caused by *Fusarium graminearum* group I. *Plant Disease* **78**, 317.
- Boc A, Diallo AB, Makarenkov V, 2012. T-REX: a web server for inferring, validating and visualizing phylogenetic trees and networks. *Nucleic Acids Research* **40**, W573-W579.
- Burgess LW, Backhouse D, Summerell BA, Swan LJ, 2001. Crown rot of wheat, in: Summerell BA, Leslie JF, Backhouse D, Bryden WL, Burgess LW (eds), *Fusarium*. APS Press, St Paul, MN, USA, pp. 271-294.
- Burgess LW, Wearing AH, Toussoun TA, 1975. Surveys of fusaria associated with crown rot of wheat in eastern Australia. *Australian Journal of Agricultural Research* **26**, 791-799.
- Droce A, Sørensen JL, Giese H, Sondergaard TE, 2013. Glass bead cultivation of fungi: Combining the best of liquid and agar media. *Journal of Microbiological Methods* **94**, 4.
- Edgar RC, 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**, 1792-1797.
- Francis RG, Burgess LW, 1977. Characteristics of two populations of *Fusarium roseum* 'Graminearum' in Eastern Australia. *Transactions of the British Mycological Society* **68**, 421-427.
- Gardiner DM, Benfield AH, Stiller J, Stephen S, Aitken K, Liu C, Kazan K, 2017. A high-resolution genetic map of the cereal crown rot pathogen *Fusarium pseudograminearum* provides a near-complete genome assembly. *Molecular Plant Pathology* **19**, 217-216.
- Gargouri S, Mtat I, Kammoun LG, Zid M, Hajlaoui MR, 2011. Molecular genetic diversity in populations of *Fusarium pseudograminearum* from Tunisia. *Journal of Phytopathology* **159**, 306-313.
- Hansen FT, Gardiner DM, Lysøe E, Fuertes PR, Tudzynski B, Wiemann P, Sondergaard TE, Giese H, Brodersen DE, Sørensen JL, 2015. An update to polyketide synthase and non-ribosomal synthetase genes and nomenclature in *Fusarium*. *Fungal Genetics and Biology* **75**, 20-29.
- Ji LJ, Kong LX, Li QS, Wang LS, Chen D, Ma P, 2016. First report of *Fusarium pseudograminearum* causing Fusarium head blight of wheat in Hebei Province, China. *Plant Disease* **100**, 220-220.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM, 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Research* **27**, 722-736.

- Kristensen R, Torp M, Kosiak B, Holst-Jensen A, 2005. Phylogeny and toxigenic potential is correlated in *Fusarium* species as revealed by partial translation elongation factor 1 alpha gene sequences. *Mycological Research* **109**, 173-186.
- Lamprecht SC, Marasas WFO, Hardy MB, Calitz FJ, 2006. Effect of crop rotation on crown rot and the incidence of *Fusarium pseudograminearum* in wheat in the Western Cape, South Africa. *Australasian Plant Pathology* **35**, 419-426.
- Li HL, Yuan HX, Fu B, Xing XP, Sun BJ, Tang WH, 2012. First report of *Fusarium pseudograminearum* causing crown rot of wheat in Henan, China. *Plant Disease* **96**, 1065-1065.
- Lofgren LA, LeBlanc NR, Certano AK, Nachtigall J, LaBine KM, Riddle J, Broz K, Dong Y, Bethan B, Kafer CW, Kistler HC, 2018. *Fusarium graminearum*: pathogen or endophyte of North American grasses? *New Phytologist* **217**, 1203-1212.
- Moolhuijzen PM, Manners JM, Wilcox SA, Bellgard MI, Gardiner DM, 2013. Genome sequences of six wheat-infecting *Fusarium* species isolates. *Genome announcements* **1**.
- Murray GM, Brennan JP, 2009. Estimating disease losses to the Australian wheat industry. *Australasian Plant Pathology* **38**, 558-570.
- Murray GM, Brennan JP, 2010. Estimating disease losses to the Australian barley industry. *Australasian Plant Pathology* **39**, 85-96.
- O'Donnell K, Kistler HC, Tacke BK, Casper HH, 2000. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 7905-7910.
- O'Donnell K, Rooney AP, Proctor RH, Brown DW, McCormick SP, Ward TJ, Frandsen RJN, Lysøe E, Rehner SA, Aoki T, Robert V, Crous PW, Groenewald JZ, Kang S, Geiser DM, 2013. Phylogenetic analyses of RPB1 and RPB2 support a middle Cretaceous origin for a clade comprising all agriculturally and medically important fusaria. *Fungal Genetics and Biology* **52**, 20-31.
- Scott JB, Chakraborty S, 2006. Multilocus sequence analysis of *Fusarium pseudograminearum* reveals a single phylogenetic species. *Mycological Research* **110**, 1413-1425.
- Sieber CMK, Lee W, Wong P, Munsterkotter M, Mewes HW, Schmeitzl C, Varga E, Berthiller F, Adam G, Guldener U, 2014. The *Fusarium graminearum* genome reveals more secondary metabolite gene clusters and hints of horizontal gene transfer. *PloS one* **9**.
- Smedsgaard J, 1997. Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *Journal of Chromatography A* **760**, 264-270.

- Smiley RW, Gourlie JA, Easley SA, Patterson LM, 2005. Pathogenicity of fungi associated with the wheat crown rot complex in Oregon and Washington. *Plant Disease* **89**, 949-957.
- Summerell BA, Burgess LW, Backhouse D, Bullock S, Swan LJ, 2001. Natural occurrence of perithecia of *Gibberella coronicola* on wheat plants with crown rot in Australia. *Australasian Plant Pathology* **30**, 353-356.
- Sydenham EW, Marasas WFO, Thiel PG, Shephard GS, Nieuwenhuis JJ, 1991. Production of mycotoxins by selected *Fusarium graminearum* and *F. crookwellense* isolates. *Food Additives and Contaminants* **8**, 31-41.
- Sørensen JL, Sondergaard TE, 2014. The effects of different yeast extracts on secondary metabolite production in *Fusarium*. *International Journal of Food Microbiology* **170**, 55-60.
- Sørensen JL, Sondergaard TE, Covarelli L, Fuertes PR, Hansen FT, Frandsen RJN, Saei W, Lukassen MB, Wimmer R, Nielsen KF, Gardiner DM, Giese H, 2014a. Identification of the biosynthetic gene clusters for the lipopeptides fusaristatin A and W493 B in *Fusarium graminearum* and *F. pseudograminearum*. *Journal of Natural Products* **77**, 2619-2615.
- Sørensen LQ, Lysøe E, Larsen JE, Khorsand-Jamal P, Nielsen KF, Frandsen RJN, 2014b. Genetic transformation of *Fusarium avenaceum* by *Agrobacterium tumefaciens* mediated transformation and the development of a USER-Brick vector construction system. *BMC Molecular Biology* **15**, 15.
- Xu F, Song YL, Wang JM, Liu LL, Zhao K, 2017. Occurrence of *Fusarium* crown rot caused by *Fusarium pseudograminearum* on barley in China. *Plant Disease* **101**, 837-837.
- Zhang H, Gao S, Lercher MJ, Hu S, Chen W-H, 2012. EvolView, an online tool for visualizing, annotating and managing phylogenetic trees. *Nucleic Acids Research* **40**, W569-W572.

Figure legends

Figure 1. Structures of W493-B and fusaristatin A highlighting the reduced polyketide (black) and peptide (colored) parts.

Figure 2. Comparative analysis of the fusaristatin gene cluster and remnant fragments in *Fusarium*.

A. Illustration of the intact cluster in *F. pseudograminearum* CS5834 (BN849_0052030 - BN849_0052070) and seven other *Fusarium* species. Only five remnant fragments (R1-R5) are present in *F. pseudograminearum* CS3096, CS3220, CS3487, CS3270, CS3427 and RBG5266. **B.** Predicted amino acid sequence of regions corresponding to PKS6 fragments R2 - R4 in *F. pseudograminearum* strains CS3096 (lacks intact cluster) and CS5834 (has intact cluster). Amino acids are represented by standard single-letter abbreviations, and two letters stacked one on top of the other indicate a difference in the sequence of the two strains.

Figure 3. Molecular analyses of the *F. pseudograminearum* strains. **A.** Phylogenetic analyses of the 99 *F. pseudograminearum* strains (orange: Western Australia; red: South Australia; blue: New South Wales; purple: Queensland) and of selected genome sequenced *Fusarium* strains with *F. poae* strain 2516 as outgroup. Numbers indicate bootstrap values from 1000 replications. **B.** 1% agarose gels visualizing the PCR products for determining the presence (●) and absence (●) of *PKS6* of strains located in clade II. **C.** Multi-locus phylogeny of 16 *F. pseudograminearum* isolates with and without the fusaristatin gene cluster. Numbers indicate bootstrap values from 1000 replications.

Supplementary figure 1. Production of W493-B by *Fusarium pseudograminearum* strains collected in New South Wales (NSW), Queensland (QLD) and Western Australia (WA). Strains

436 from WA have been divided in fusaristatin producers (+) and nonproducers (-). The box plot
437 illustrate minimum and maximum; first and third quartile and mean peak areas.

438


439 **Supplementary Figure 2.** Alignment of *F. pseudograminearum* CS3096 and CS3894 in the region
440 where the fusaristatin gene cluster has been lost. A highlighted 100 bp region is present in CS3894,
441 but absent in CS3096.



Table 1. Description of genes in the fusaristatin cluster in *F. pseudograminearum* CS5834 and comparison (% identity on amino acid level) to *F. graminearum* NRRL 31084 and *F. avenaceum* Fa05001.

<i>F. pseudograminearum</i>	Length	Function	<i>F. graminearum</i>	<i>F. avenaceum</i>
BN849_0052030	4355 aa	Non-ribosomal peptide synthetase	FGSG_08209 (94%)	FAVG1_08708 (73%)
BN849_0052040	2554 aa	Polyketide synthase	FGSG_08208 (98%)	FAVG1_08709 (84%)
BN849_0052050	520 aa	Cytochrome P450 monooxygenase	FGSG_08207 (98%)	FAVG1_08710 (86%)
BN849_0052060	138 aa	Hypothetic protein	FGSG_08206 (96%)	FAVG1_08711 (79%)
BN849_0052070	511 aa	Aminotransferase	FGSG_08205 (96%)	FAVG1_08712 (78%)

Table 2. Production^a of W493-B and Fusaristatin A (Fst A) by *F. pseudograminearum* strains collected from New South Wales (NSW), Queensland (QLD), South Australia (SA) and Western Australia (WA).


Strain ^a	W493-B ^b	Fst A ^b	Location	State	Strain	W493-B	Fst A	Location	State
CS3096	•		Moree	NSW	CS7114	•		Bowenville	QLD
CS3164	•		Qurindi	NSW	CS7118	•		Marmadiaz	QLD
CS3166	•		Qurindi	NSW	CS7124	•		Hannaford	QLD
CS3173	•		Qurindi	NSW	CS7126	•		Hannaford	QLD
CS3184	•		Bladeville	NSW	CS7133	•		Toobeak	QLD
CS3220	•		Liverpool Plains	NSW	CS7139	•		Toobeak	QLD
CS3270	•		Liverpool Plains	NSW	CS7145	•		Wyaga	QLD
CS3293	•		Boggabri	NSW	CS7147	•		Wyaga	QLD
CS3319	•		Boggabri	NSW	CS7149	•		Warra	QLD
CS3361	•		Bellata	NSW	CS7153	•		Warra	QLD
CS3768	•		North Stat	NSW					
CS3784	•		North Stat	NSW	CS3891	•		Foolunga Street	SA
CS3941	•		Cooper Creek K	NSW	CS3894	•		Foolunga Street	SA
CS3950	•		Cooper Creek K	NSW	CS3900	•		Angus Valley	SA
CS3965	•		9 Miles Road	NSW	CS3907	•		Angus Valley	SA
CS3967	•		9 Miles Road	NSW					
CS3983	•		Livingstone Farm	NSW	CS5541	•	•	Stockdale	WA
CS3986	•		Livingstone Farm	NSW	CS5573	•	•	Stockdale	WA
CS7291	•		Nombi 1	NSW	CS5588	•	•	Tammin	WA
CS7302	•		Spring Ridge 1	NSW	CS5703	•	•	Tammin	WA
CS7305	•		Spring Ridge 1	NSW	CS5834	•	•	Tammin	WA
CS7311	•		Nombi 1	NSW	CS5877	•		Farm 3	WA
CS7313	•		Nowbi 1	NSW	CS5894	•	•	Jerramungub	WA
CS7319	•		Spring Ridge 2	NSW	CS5897	•	•	Jerramungub	WA
CS7344	•		Nowbi 2	NSW	CS7054	•	•	Lake Grace	WA
CS7350	•		Nowbi 2	NSW	CS7055	•		Boxwood Hill	WA
CS7358	•		Tambar Springs	NSW	CS7056	•	•	Boxwood Hill	WA
CS7374	•		Tambar Springs	NSW	CS7060	•	•	Lake Grace	WA
CS7385	•		Spring Ridge 3	NSW	CS7062	•	•	Lake Grace	WA
CS7391	•		Spring Ridge 3	NSW	CS7065	•		Mettler	WA
CS7405	•		Spring Ridge 4	NSW	CS7066	•	•	Wellstead	WA
CS7407	•		Bladeville	NSW	CS7069	•		Wellstead	WA
CS7420	•		Spring Ridge 5	NSW	CS7078	•		Lake Grace	WA
CS7427	•		Spring Ridge 5	NSW	CS7080	•		Lake Grace	WA
CS7436	•		Spring Ridge 2	NSW	CS7081	•	•	Carnamagh	WA
CS7453	•		Spring Ridge 6	NSW	CS7082	•		Lake King	WA
CS7460	•		Werris Creek	NSW	CS7084	•		Lake King	WA
CS7461	•		Werris Creek	NSW	CS7085	•	•	Lake King	WA
CS7463	•		Kelvin	NSW	CS7088	•		Lake King	WA
CS7464	•		Kelvin	NSW	CS7089	•		Grasspatch	WA
CS7465	•		Caroona 4	NSW	CS7090	•		Grasspatch	WA
CS7467	•		Caroona 4	NSW	CS7091	•		Grasspatch	WA
					CS7093	•		Grasspatch	WA
CS3002				QLD	CS7094	•		Grasspatch	WA
CS3427	•		Wilga Downs	QLD	CS7098	•		Grasspatch	WA
CS3438	•		Wilga Downs	QLD	CS7099	•		Salmon Gums	WA
CS3442	•		Coondiwindi	QLD	CS7100	•		Salmon Gums	WA
CS3744	•		Kentare	QLD	CS7104	•		Salmon Gums	WA
CS3752	•		Kentare	QLD	CS7105	•		Lake Grace	WA
CS3910	•		Westfield	QLD	CS7108	•	•	Lake Grace	WA
CS7113	•		Bowenville	QLD					




A**B**



B



C



Fusaristatin gene cluster (● present; ● absent)

Highlights

Ancestral *Fusarium pseudograminearum* strains produced fusaristatin

Fusaristatin-producing *F. pseudograminearum* are confined to Western Australia

Remnant fragments of fusaristatin cluster was found in non-producers