

The relationship between serum potassium concentrations and electrocardiographic characteristics in 163,547 individuals from primary care

Krogager, Maria Lukacs; Kragholm, Kristian Hay; Skals, Regitze Kuhr; Mortensen, Rikke Nørmark; Polcwiartek, Christoffer Bech; Graff, Claus; Kanter, Jørgen K.; Holst, Anders Gaarsdal; Søgaard, Peter; Pietersen, Adrian; Torp-Pedersen, Christian; Hansen, Steen Møller

Published in:

Book of abstracts from ESC Congress 2019

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Krogager, M. L., Kragholm, K. H., Skals, R. K., Mortensen, R. N., Polcwiartek, C. B., Graff, C., Kanter, J. K., Holst, A. G., Søgaard, P., Pietersen, A., Torp-Pedersen, C., & Hansen, S. M. (2019). The relationship between serum potassium concentrations and electrocardiographic characteristics in 163,547 individuals from primary care. In *Book of abstracts from ESC Congress 2019* Article P6562 <https://esc365.escardio.org/Congress/ESC-CONGRESS-2019/Poster-Session-7-Arrhythmias-Miscellaneous/200003-the-relationship-between-serum-potassium-concentrations-and-electrocardiographic-characteristics-in-163-547-individuals-from-primary-care>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: December 06, 2025

The relationship between serum potassium concentrations and electrocardiographic characteristics in 163,547 individuals from primary care

Authors:

ML Krogager¹, K Kragholm², RK Skals², RN Mortensen², C Polcwiartek², C Graff³, JB Nielsen⁴, JK Kanter⁵, AG Holst⁶, P Soegaard², AH Pietersen⁷, C Torp-Pedersen², SM Hansen², ¹Aalborg University Hospital, Unit of Epidemiology and Biostatistics - Aalborg - Denmark, ²Aalborg University Hospital - Aalborg - Denmark, ³Aalborg University - Aalborg - Denmark, ⁴Division of Cardiovascular Medicine, Department of Internal Medicine - Michigan - United States of America, ⁵University of Copenhagen, Laboratory of experimental cardiology - Copenhagen - Denmark, ⁶Rigshospitalet - Copenhagen University Hospital, Cardiology - Copenhagen - Denmark, ⁷Hillerød Hospital, Cardiology - Hillerød - Denmark,

Topic(s):

Arrhythmias, General – Prevention

Citation:

Funding Acknowledgements:

None

Aims

Potassium disturbances are common and associated with increased morbidity and mortality, even in patients without prior cardiovascular disease. We examined six electrocardiographic (ECG) measures and their association to serum potassium levels.

Methods and results

From the Copenhagen General Practitioners' Laboratory, we identified 163,547 individuals aged ≥16 years with a first available ECG and a concomitant serum potassium measurement during 2001-2011. Restricted cubic splines curves showed a non-linear relationship between potassium and the Fridericia corrected QT (QTcF) interval, T-wave amplitude, morphology combination score (MCS), PR interval, P-wave amplitude and duration. Therefore, potassium was stratified in two intervals K: 2.0-4.1 mmol/L and 4.2-6.0 mmol/L for further analyses. Within the low potassium range, we observed: QTcF was 12.5 ms longer for each mmol/L decrease in potassium ($p<0.0001$); T-wave amplitude was 55.1 μ V lower for each mmol/L decrease in potassium ($p<0.0001$); and MCS was 0.09 higher per mmol/L decrease in potassium ($p<0.001$). Moreover, P-wave duration and PR interval were prolonged by 3.1 and 3.3 ms for each mmol/L decrease in potassium ($p<0.0001$), respectively. Within the lowest potassium range (2.0-4.1 mmol/L) P-wave amplitude was 10.7 μ V higher for each mmol/L decrease in potassium ($p<0.0001$). Within the high potassium range associations with the above-mentioned ECG parameters were much weaker.

Conclusion

The association of potassium with six commonly measured ECG parameters was non-linear. Strong associations between ECG abnormalities and potassium were seen among individuals with lower potassium levels (≤ 4.1 mmol/L).

