



Neuromuscular Exercises Improve Shoulder Function More Than Standard Care Exercises in Patients With a Traumatic Anterior Shoulder Dislocation

A Randomized Controlled Trial

Eshoj, Henrik Rode; Rasmussen, Sten; Frich, Lars Henrik; Hvass, Inge; Christensen, Robin; Boyle, Eleanor; Jensen, Steen Lund; Søndergaard, Jens; Søgaard, Karen; Juul-Kristensen, Birgit

Published in:

Orthopaedic Journal of Sports Medicine

DOI (link to publication from Publisher):

[10.1177/2325967119896102](https://doi.org/10.1177/2325967119896102)

Creative Commons License

CC BY-NC-ND 4.0

Publication date:

2020

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Eshoj, H. R., Rasmussen, S., Frich, L. H., Hvass, I., Christensen, R., Boyle, E., Jensen, S. L., Søndergaard, J., Søgaard, K., & Juul-Kristensen, B. (2020). Neuromuscular Exercises Improve Shoulder Function More Than Standard Care Exercises in Patients With a Traumatic Anterior Shoulder Dislocation: A Randomized Controlled Trial. *Orthopaedic Journal of Sports Medicine*, 8(1), Article 2325967119896102.

<https://doi.org/10.1177/2325967119896102>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Neuromuscular Exercises Improve Shoulder Function More Than Standard Care Exercises in Patients With a Traumatic Anterior Shoulder Dislocation

A Randomized Controlled Trial

Henrik Rode Eshoj,^{*†} PhD, Sten Rasmussen,^{‡§} MD, PhD, Lars Henrik Frich,^{||¶} MD, PhD, Inge Hvass,[#] MD, PhD, Robin Christensen,^{**††} PhD, Eleanor Boyle,^{*} PhD, Steen Lund Jensen,^{‡‡} MD, PhD, Jens Søndergaard,^{§§} MD, PhD, Karen Søgaard,^{*¶} PhD, and Birgit Juul-Kristensen,^{*|||} PhD

Investigation performed at University of Southern Denmark, Odense, Denmark

Background: There is an important gap in knowledge about the effectiveness of nonoperative treatment (exercise) for patients with traumatic primary and recurrent anterior shoulder dislocations (ASDs).

Purpose/Hypothesis: The purpose of this study was to assess the efficacy and safety of physical therapist-supervised, shoulder instability neuromuscular exercise (SINEX) versus self-managed, home-based, standard care shoulder exercise (HOMEX) in patients with traumatic ASDs. The hypothesis was that SINEX would have a larger effect and fewer adverse events compared with HOMEX.

Study Design: Randomized controlled trial; Level of evidence, 2.

Methods: A total of 56 participants with radiographically verified, trauma-initiated primary or recurrent ASDs and self-reported decreased shoulder function were randomized to 12 weeks of either SINEX or HOMEX. The SINEX program consisted of 7 exercises, individually progressing from basic (2 × 20 repetitions each day) to elite (2 × 10 repetitions, 3 times weekly). The HOMEX program included 5 shoulder exercises performed 3 times weekly (2 × 10 repetitions). The primary outcome was the Western Ontario Shoulder Instability Index (WOSI) score, ranging from 0 (best possible) to 2100. The between-group minimal clinically important difference at 12 weeks was 250 points. Secondary outcomes included WOSI subdomain scores, patient-reported ratings of kinesiophobia and pain, objective shoulder function, patient satisfaction, and number of adverse events.

Results: The between-group mean difference in the WOSI total score at 12 weeks significantly favored SINEX over HOMEX (−228.1 [95% CI, −430.5 to −25.6]). SINEX was furthermore superior to HOMEX in most of the secondary outcomes (3/4 subdomains of the WOSI and pain level during the past 7 days as well as clinical signs of anterior shoulder instability). Also, although not statistically significant, less than half the proportion of the SINEX patients compared with the HOMEX patients (3/27 [11%] vs 6/24 [25%], respectively; $P = .204$) underwent or were referred for shoulder stabilizing surgery. Satisfaction with both exercise programs was high, and no serious adverse events were reported.

Conclusion: Neuromuscular shoulder exercise (SINEX) was superior to standard care exercise (HOMEX) in patients with traumatic ASDs. Further long-term follow-ups on treatment effects are needed.

Registration: NCT02371928 (ClinicalTrials.gov identifier).

Keywords: shoulder; traumatic; instability; neuromuscular exercise; physical therapy

The Orthopaedic Journal of Sports Medicine, 8(1), 2325967119896102
DOI: 10.1177/2325967119896102
© The Author(s) 2020

A traumatic anterior shoulder dislocation (ASD) is a common athletic injury among young, active patients.³⁴ The annual incidence of primary traumatic ASDs is between 11.2 and 26.2 per 100,000 people.^{30,32,52} Patients with primary traumatic ASDs are prone to recurrent ASDs because

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

of lesions of the glenoid labrum, an important stabilizer of the shoulder.^{31,42} This may lead to recurrent ASDs and symptoms for longer periods,³ reduced participation in physical activity, and decreased shoulder-related quality of life.^{16,37,44}

Diverging strategies exist as to the optimum treatment for this group. For young, active male patients with primary traumatic ASDs, 2 systematic reviews have concluded that shoulder stabilizing surgery with postsurgical shoulder rehabilitation was superior to nonsurgical standard care treatment only in reducing the risk of recurrent ASDs.^{19,34} These systematic reviews included 4 randomized controlled trials (RCTs) with rehabilitation of only primary traumatic ASDs,^{5,22,27,50} and these studies had several limitations: limited description of the rehabilitation protocol and lack of short-term treatment effect measures. Moreover, the primary endpoint was objectively registered anterior shoulder redislocation, which usually does not occur until 12 to 24 months after a primary traumatic ASD.⁴⁰ Therefore, from RCTs, little is known about patients with primary and recurrent traumatic ASDs on short-term and subjective outcomes of shoulder symptoms and function.²⁵

Furthermore, in qualitative, cross-sectional, and cohort studies, it was reported that many patients (~50%) with a primary traumatic ASD who had been treated nonsurgically did not experience recurrent ASDs, whereas shoulder function in general was poor.²¹ Also, it was reported that some patients who were treated with surgery and rehabilitation reported poor shoulder function, a fear of reinjuries, and unstable shoulders for long periods.^{16,44} This would tend to indicate that nonsurgical standard care treatment is only suboptimal and that initial surgery is not ideal for all patients with traumatic ASDs (primary or recurrent). In addition, as not all patients are candidates for surgery (surgeon's or one's own decision), an efficient and evidence-based standardized physical therapy regimen for the nonsurgical management of traumatic ASDs is needed. Such a regimen has not yet been established, as also seen in treatment recommendations within this area, which are only weakly substantiated as described because of the absence of RCT studies that compare the effect of nonsurgical exercise programs.²⁵ Traditionally, nonsurgical standard care treatment for patients with traumatic ASDs

consists of closed reduction followed by immobilization in a shoulder sling, and, if provided, some type of physical therapy (eg, shoulder range of movement exercises and low load strengthening of the rotator cuff muscles).^{5,22,27} However, as a traumatic ASD often leads to loss of mechanical stability and deficits to the global neuromuscular and proprioceptive systems, in addition to weakness of the rotator cuff muscles,^{9,10,20,29,43} a greater focus on neuromuscular exercises that enhance compensatory functional shoulder stability may seem relevant. While other studies on patients with traumatic knee instability (anterior cruciate ligament tears) have shown a large effect for progressive neuromuscular exercise programs,^{11,17} such programs do not exist for patients with traumatic ASDs.⁵³ This is why it seems reasonable to also develop and evaluate treatment effects of such a program for patients with traumatic ASDs.³⁶

Therefore, the purpose of this RCT was to evaluate the efficacy and safety of a nonsurgical, supervised, progressive shoulder instability neuromuscular exercise (SINEX) program compared with a self-managed, home-based, standard care shoulder exercise program (HOMEX) in patients with traumatic ASDs. The hypothesis was that SINEX would (1) have a larger effect than HOMEX and (2) not induce more adverse events than HOMEX.

METHODS

Trial Design

A multicenter, randomized, assessor-blinded, controlled clinical superiority trial was conducted using a parallel (1:1) group design. All patients gave written informed consent, and the study fulfilled the principles of the Declaration of Helsinki.⁵¹ The study conformed to the CONSORT 2010 statement.⁴¹ Details of the trial design and methods have been published elsewhere.¹⁵ The statistical analysis plan is provided as supplementary material (see Appendix 1, available as supplemental material). Deviations to the study protocol¹⁵ were as follows: Initially, patients had to have clinical signs of anterior shoulder instability with positive findings on a minimum of 2 of 3 clinical tests (apprehension, relocation, surprise). However, this criterion was

^{|||}Address correspondence to Birgit Juul-Kristensen, PhD, Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark (email: bjuul-kristensen@health.sdu.dk).

*Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark.

[†]Quality of Life Research Center, Department of Haematology, Odense University Hospital, Odense, Denmark.

[‡]Department of Clinical Medicine, Aalborg University, Aalborg, Denmark.

[§]Orthopaedic Research Unit, Aalborg University Hospital, Aalborg, Denmark.

^{||}Department of Orthopaedic Surgery and Traumatology, Odense University Hospital, Odense, Denmark.

[¶]Department of Clinical Research, University of Southern Denmark, Odense, Denmark.

[#]Shoulder Section, Department of Orthopaedic Surgery, South-West Jutland Hospital, Esbjerg, Denmark.

^{**}Musculoskeletal Statistics Unit, The Parker Institute, Bispebjerg and Frederiksberg Hospital, University of Copenhagen, Copenhagen, Denmark.

^{††}Rheumatology Research Unit, Department of Clinical Research, University of Southern Denmark, Odense, Denmark.

^{‡‡}Shoulder Section, Department of Orthopaedic Surgery, Aalborg University Hospital, Farsoe, Denmark.

^{§§}Research Unit of General Practice, Department of Public Health, University of Southern Denmark, Odense, Denmark.

Final revision submitted October 15, 2019; accepted October 25, 2019.

The authors declared that there are no conflicts of interest in the authorship and publication of this contribution. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval for this study was obtained from the ethics committee for the Region of Southern Denmark (project No. S-20140093).

removed from inclusion criteria because many of the patients reported poor shoulder function without having clinical signs of anterior shoulder instability. However, the clinical tests were retained as treatment effect measures. Furthermore, the objective measurement of prone-lying shoulder stability (using a Nintendo Wii Balance Board), originally included in the objective outcome test battery,¹⁵ was withdrawn because of low concurrent validity compared with the gold standard (force platform).¹⁴

Setting and Participants

Patients were recruited from 3 orthopaedic shoulder units in the Regions of Southern and Northern Denmark from March 2015 to March 2017, with the final deadline for enrollment set at March 31, 2017.¹⁵ Eligible patients were men and women between the ages of 18 and 39 years with trauma-initiated, unidirectional anterior shoulder instability.¹⁸ Inclusion criteria were radiographically verified acute primary or recurrent ASDs (using a radiograph that showed the actual dislocation, which is standard procedure before manual reduction in Denmark) and self-reported decreased ability to perform shoulder movements during daily activities in the previous 7 days. Patients were assessed for eligibility to participate in the RCT within 3 to 6 weeks after their current shoulder injury. This screening window was chosen to ensure that all patients had completed initial immobilization of the shoulder after reduction. Exclusion criteria were humeral head fractures and/or bony Bankart lesions requiring early surgery (evaluated by orthopaedic surgeons), prior surgery in the affected shoulder joint, more than 5 ASDs in the patient's lifetime, suspected competing diagnoses (such as rheumatoid arthritis, cancer, neurological disorders, fibromyalgia, psychiatric diseases), sensory and motor deficits in the neck and shoulder, current pregnancy, inability to speak or write Danish, and/or inability to attend 12 weeks of supervised shoulder training.

Stratified Randomization and Blinding

Patients were randomly assigned to either SINEX or HOMEX with a 1:1 allocation per study site, stratified according to injury status: that is, a primary (first-time) or recurrent (second- to fifth-time) ASD. A computer-generated randomization sequence was produced using PROC PLAN (SAS Institute) to generate the schedules before any patient was enrolled, allocating patients in permuted blocks of 2, 4, or 6 to the SINEX or HOMEX program. At each trial site, these (stratified) lists were sequentially numbered in opaque, concealed envelopes. A total of 2 outcome assessors, blinded to treatment allocation, performed all measurements according to procedures described previously.¹⁵

Interventions

As standard care treatment varies, the HOMEX program was developed to reflect the core similarity of standard care packages while still representing a realistic treatment option. The SINEX program was developed to accommodate shoulder mechanical loss and proprioceptive impairment

subsequent to trauma-initiated primary or recurrent ASDs.^{9,10,20,29,43}

SINEX Program. A cornerstone of the neuromuscular exercises was movement quality through guided supervision from physical therapists. The neuromuscular exercises included strength, coordination, balance, and proprioception, which were integrated simultaneously into various body positions, enhancing compensatory functional shoulder stability. This is in contrast with the standard care program (HOMEX) that relied primarily on strength training to increase muscle mass. The SINEX group received 12 weeks of individually tailored, supervised sessions of progressive shoulder exercise in addition to functional kinetic chain exercise. The SINEX program included 7 exercises targeting the glenohumeral and scapular muscles. Each exercise has 7 progression levels (basic to elite), with exercises at the basic level performed every day (2×20 repetitions) and exercises at the elite level performed 3 times weekly (2×10 repetitions). Exercises followed general strength training principles, with basic and elite levels referring to low and high load exercises. All patients in the SINEX group had access to online exercise instructions and videos through the physical therapy website (digifys.com).

Supervised sessions were provided throughout the 12 weeks, lasting approximately 45 minutes each. Supervised sessions were offered twice a week for the first 2 weeks and then once a week for the remaining 10 weeks, summing up to a maximum of 14 sessions. The amount of supervised sessions needed was decided by physical therapists based on the movement control and capability of the individual patient. However, to satisfactorily complete the SINEX program, attendance of at least 7 (50%) supervised sessions (of 14 possible) was required in addition to completion of at least two-thirds (66%) of the planned home-based exercises (self-reported training diary). A full description of the SINEX program is provided as supplementary material (see Appendix 2, available as supplemental material).

HOMEX Program. The HOMEX group received 1 introductory supervised physical therapy session, including a leaflet with photographs and descriptions of exercises. Patients were instructed not to perform exercises that exceeded their pain limit and that provoked shoulder pain. The HOMEX program consisted of active exercises for the rotator cuff and scapular muscles using elastic bands and 1 exercise for mobility/coactivation of the scapular and core stability muscles. Patients had to perform the exercises for 12 weeks, 3 times weekly (2×10 repetitions). After 6 weeks, patients received a telephone call from a physical therapist to ascertain the appropriate progression of and compliance with the exercises. Compliance with the HOMEX program was reached, with a minimum of two-thirds (66%) of the planned home training completed (via self-reported training diary). Further details of the HOMEX program are provided as supplementary material (see Appendix 3, available as supplemental material).

Outcome Measures

The primary endpoint was the change in the total score of the patient-reported Western Ontario Shoulder Instability

Index (WOSI) from baseline to the 12-week follow-up. The WOSI covers 21 items, each ranging from 0 to 100, with higher scores representing worse shoulder-related quality of life.²⁶ A reliable and validated Danish version of the WOSI was used.¹²

Secondary patient-reported outcomes were the sub-domain scores (physical symptoms, sport function, lifestyle, emotions) of the WOSI collected at baseline and weeks 4, 8, and 12. Further secondary patient-reported outcome measures included the Tampa Scale of Kinesiophobia,^{35,47} the EuroQol 5-Dimensions questionnaire,³⁸ and the Patient-Specific Functional Scale⁷ as well as pain intensity now, average pain intensity in the past 24 hours, and average pain intensity in the previous 7 days using the Numeric Pain Rating Scale (NPRS).⁸ Objective secondary outcome measures included the Constant-Murley score (CMS), including the CMS subdomains of range of motion and isometric shoulder abduction strength (in kg) with the use of a dynamometer,² blindfolded shoulder joint reposition sense (JRS) testing within low ranges of shoulder flexion and abduction (0° to $60^\circ \pm 10^\circ$) with the use of a laser beam,⁴⁵ and clinical tests for anterior shoulder instability¹³ as well as the Beighton score for the classification of generalized joint hypermobility.^{4,24} Patients further reported their impression of recovery at week 12 using the global perceived effect (GPE) scale and responded to questions on exercise-related adverse events and other shoulder-related issues after participation.

Statistical Analysis

To reduce bias in the interpretation of the primary outcome, a blinded interpretation framework was used (see Appendix 4, available as supplemental material).²³ All analyses followed the intention-to-treat (ITT) principle; missing data were replaced using nonresponder imputation (ie, baseline observation carried forward technique)³³ because of the assumption that those who dropped out returned to their baseline WOSI score (ie, null imputation).⁴⁹

Simple descriptive statistics of either group means with standard deviations or within-group numbers and proportions were used. Change values were calculated, and results are presented as the estimated adjusted between-group mean difference of change values, besides adjusted risk differences, with 95% CIs and associated *P* values. Analysis of covariance (ANCOVA) was used to analyze the mean change and self-reported data of continuous outcomes, presented as estimates of least squares means with 95% CIs. The model included treatment group, study center, sex, and injury status (primary or recurrent ASD) as fixed effects, adjusting each patient's dependent outcome for his or her baseline score, with baseline as a covariate. For the categorical outcomes (eg, clinical test findings, adverse events), logistic regression analysis was used, presenting probability estimates with 95% CIs using the same fixed effects and covariates as in the ANCOVA above. For the categorical outcomes on self-reported data and adverse events at week 12, data are presented as percentages, with differences between groups determined by a chi-square test or Fisher exact test.

For the longitudinal part of the trial (no imputation for missing data needed), a linear mixed model with repeated measures of the WOSI (weeks 4, 8, and 12) was performed to test the difference and trajectory over time between the 2 groups (interaction: group \times time) with the same fixed effects and covariates as in the ANCOVA. For sensitivity and exploratory purposes, per-protocol analysis was performed, which included only the participants who had acceptable exercise compliance.¹⁵

The study was powered to detect a difference of at least 250 points as the minimal clinically important difference (MCID) between SINEX and HOMEX for the primary outcome.²⁸ To achieve 90% power to show a clinically relevant between-group difference in favor of SINEX, with a 2-sided type I error rate of 5%, 36 patients per group were necessary.

All statistical analyses were performed with SPSS (Version 24.0; IBM). *P* values $<.05$ (2-sided) were considered statistically significant.

Patient Involvement

Patients with traumatic ASDs provided feedback about the neuromuscular exercises during the development and design of the SINEX program. After consenting to participate in the study, the patients were asked whether they wanted to receive a letter explaining the results of this trial. Accordingly, a letter in lay terms will be forwarded to those interested when the study findings are published.

RESULTS

Participant Characteristics

A total of 130 patients were screened for eligibility. Of these, 56 patients fulfilled the eligibility criteria, comprising the ITT population (Figure 1), and were randomized to either the SINEX or the HOMEX group. A total of 51 of 56 (93%) completed the 12-week follow-up assessment (SINEX: $n = 27$; HOMEX: $n = 24$). At baseline, the groups were comparable in demographics and clinical characteristics (except for the WOSI lifestyle and NPRS within the past 24 hours in which the HOMEX group had poorer scores). The mean age was 25.8 ± 5.8 years and 26.2 ± 6.4 years for the HOMEX and SINEX groups, respectively, and a majority of the patients were male (49/56; 88%). Baseline characteristics are described in Tables 1 and 2.

For those returning diaries after completing the home exercise ($n = 18$ patients in each group), the overall adherence was a mean of 71% for SINEX and 79% for HOMEX. However, compliance with SINEX (at least 7/14 supervised exercise sessions and at least 66% of the scheduled home-based training) corresponded to 12 of 28 (43%) patients attending a mean of 9 supervised sessions and 82% of the scheduled home-based training. Compliance with HOMEX (at least 66% of the scheduled home-based training) corresponded to 15 of 28 (54%) patients performing a mean of 87% of the home-based training. Therefore, the per-protocol




Figure 1. Flowchart of participants with traumatic anterior shoulder dislocations (ASDs) in the study. A total of 49 patients did not meet the inclusion criteria: 11 did not meet the age requirements (<18 years: n = 5; >39 years: n = 6), 11 had no radiographic verification of a shoulder dislocation, 17 reported not having any problems with shoulder movements during daily activities in the previous 7 days, 2 patients had a bony Bankart lesion or humeral head fracture, 1 patient had experienced more than 5 ASDs in his or her lifetime, 2 patients had suspected competing diagnoses (sclerosis: n = 1; unknown neurological disorder: n = 1), 1 patient was pregnant, 3 patients had no clinical signs of anterior shoulder instability, and 1 patient was not able to speak and write Danish. HOMEX, self-managed, home-based, standard care shoulder exercise program; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise.

population was composed of 12 and 15 patients for the SINEX and HOMEX groups, respectively.

Primary Outcome

The SINEX group had a significantly greater improvement on the WOSI total, although marginally below the cutoff for the MCID, compared with the HOMEX group in the change from baseline to the 12-week follow-up (between-group

mean difference, -228.1 [95% CI, -430.5 to -25.6]; *P* = .028) (Table 3). The mean change in the WOSI total score was 655.3 (95% CI, 457.5 to 853.0) in the SINEX group and 427.2 (95% CI, 245.9 to 608.6) in the HOMEX group.

Secondary Outcomes

On 3 of 4 WOSI subdomains and in pain during the previous 7 days, significantly larger improvements were seen in

TABLE 1
Baseline Demographic Information^a

	SINEX Group (n = 28)	HOMEX Group (n = 28)
Male sex	26 (93)	23 (82)
Age, mean ± SD, y	26.2 ± 6.4	25.8 ± 5.8
Weight, mean ± SD, kg	86.0 ± 19.0	80.8 ± 13.4
Height, mean ± SD, cm	180.8 ± 8.0	178.3 ± 8.0
Educational level		
University	7 (25)	10 (36)
College/technical school	12 (43)	12 (43)
Below high school	5 (18)	2 (7)
No formal education	4 (14)	4 (14)
Occupational status		
Full-time employed	16 (57)	15 (54)
Part-time employed	2 (7)	0 (0)
Student	9 (32)	10 (36)
Unemployed/retired	0 (0)	0 (0)
On sick leave	1 (4)	3 (11)
Dominant arm, right	25 (89)	26 (93)
Injured shoulder, right	16 (57)	12 (43)
Injury mechanism		
Fell on arm	13 (46)	15 (54)
Arm was pulled	4 (14)	3 (11)
External force to shoulder	2 (7)	1 (4)
Other ^b	9 (32)	9 (32)
Traumatic anterior shoulder dislocation(s) on injured side		
1	18 (64)	19 (67)
2	3 (11)	7 (25)
3	4 (14)	1 (4)
4	2 (7)	1 (4)
5	1 (4)	0 (0)
Previous shoulder treatment(s) for current shoulder injury	8 (29)	9 (32)
Exercise treatment	5 (18)	9 (32)
Passive treatment	1 (4)	1 (4)
Chiropractic	1 (4)	0 (0)
Analgesic medication (medically prescribed)	2 (7)	4 (14)
Currently physically active	23 (82)	22 (79)
Physically active ≥4 h/wk	7 (25)	6 (21)

^aData are reported as n (%) unless stated otherwise. HOMEX, self-managed, home-based, standard care shoulder exercise program; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise.

^bDuring athletic activities (soccer, gymnastics, fun wrestling, motocross).

favor of the SINEX group (Table 3). Further significant improvements in favor of SINEX were seen in clinical signs of anterior shoulder instability with the apprehension test and in CMS strength of the noninjured shoulder (Table 3). Also, less than half of the patients (33% and 46%, respectively) in the SINEX and HOMEX groups returned to pre-injury levels of sport (Table 4). At the same time, the mean ratings of actual shoulder function and ability to perform sport/leisure activities on the GPE scale were rated “improved” to “much improved,” with a statistically

TABLE 2
Baseline Outcome Measure Scores^a

	SINEX Group (n = 28)	HOMEX Group (n = 28)
WOSI, ^b total (range, 0-2100)	970.2 ± 346.9	1145.5 ± 376.2
Physical symptoms (range, 0-1000)	351.8 ± 160.0	406.8 ± 206.2
Sport function (range, 0-400)	212.4 ± 83.4	259.8 ± 93.1
Lifestyle (range, 0-400)	205.2 ± 88.4	255.8 ± 86.0
Emotions (range, 0-300)	200.9 ± 62.4	223.1 ± 67.8
TSK ^b (range, 17-68)	43.3 ± 5.1	42.9 ± 4.0
High fear of movement and reinjury (TSK ≥37), n (%)	26 (93)	26 (93)
Pain intensity for current shoulder injury ^b (range, 0-10)		
Currently	2.4 ± 2.0	3.5 ± 2.5
Mean during past 24 h	2.6 ± 1.9	4.0 ± 2.1
Mean during previous 7 d	3.7 ± 1.8	4.6 ± 1.8
EQ-5D VAS ^c (range, 0-100)	74.8 ± 15.5	73.7 ± 18.4
PSFS ^c (range, 0-10)	4.3 ± 2.5	4.5 ± 2.1
Generalized joint hypermobility ^b (range, 0-9; positive ≥5), n (%)	3 (11)	2 (7)
Positive clinical test findings, n (%)		
Apprehension	28 (100)	26 (93)
Relocation	25 (89)	21 (75)
Surprise	23 (82)	22 (79)
CMS, total ^c (range, 0-100)	72.6 ± 15.3	67.6 ± 20.7
Range of motion ^c (range, 0-40)	30.6 ± 8.0	28.0 ± 9.7
Strength, ^d kg		
Injured shoulder	17.7 ± 6.8	18.1 ± 8.3
Noninjured shoulder	26.2 ± 7.4	25.8 ± 6.9
Shoulder JRS (mean absolute error), ^e cm		
Flexion	5.2 ± 6.2	5.5 ± 4.2
Abduction	5.5 ± 6.6	6.4 ± 4.8

^aData are reported as mean ± SD unless stated otherwise. CMS, Constant-Murley score; EQ-5D VAS, EuroQol 5-Dimensions questionnaire visual analog scale; HOMEX, self-managed, home-based, standard care shoulder exercise program; JRS, joint reposition sense; PSFS, Patient-Specific Functional Scale; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise; TSK, Tampa Scale of Kinesiophobia; WOSI, Western Ontario Shoulder Instability Index.

^bHigher scores reflect worse status.

^cLower scores indicate better status.

^dIsometric shoulder strength at 90° of shoulder abduction in the scapular plane with the use of a dynamometer.

^eProprioceptive repositioning testing at 60° ± 10° of shoulder flexion and abduction with the use of a laser beam.

significant difference in favor of the SINEX group compared with the HOMEX group (6.2 vs 5.5 [$P = .012$] and 5.8 vs 4.9 [$P = .025$], respectively) (Table 4).

Although not statistically significant, only half of the patients in the SINEX group (3/27; 11%) compared with the HOMEX group (6/24; 25%) underwent or were referred for shoulder stabilizing surgery (Table 4). In the multilevel analyses, there was no difference between groups over time for the WOSI total score and for each of the WOSI

TABLE 3
Changes in Outcome Measure Scores Between Baseline and Week 12 in Intention-to-Treat Population^a

	Mean Change From Baseline			P Value
	SINEX Group (n = 28)	HOMEX Group (n = 28)	Difference, HOMEX vs SINEX	
WOSI, ^b total	655.3 (457.5 to 853.0)	427.2 (245.9 to 608.6)	-228.1 (-430.5 to -25.6)	.028
Physical symptoms	175.8 (112.8 to 238.8)	83.3 (18.1 to 148.6)	-92.5 (-181.0 to -3.9)	.041
Sport function	185.2 (139.5 to 230.9)	128.0 (87.3 to 168.1)	-57.5 (-106.5 to -8.5)	.022
Lifestyle	154.1 (109.2 to 199.0)	104.3 (67.7 to 144.9)	-49.8 (-95.8 to -3.8)	.034
Emotions	128.8 (89.4 to 168.3)	101.0 (66.1 to 136.0)	-27.8 (-69.5 to 13.9)	.187
TSK ^b	3.1 (1.3 to 4.8)	2.2 (0.4 to 4.0)	0.8 (-3.4 to 1.6)	.485
High fear of movement and reinjury at baseline (TSK ≥37) and low fear of movement and reinjury at follow- up (TSK <37), n (%)	9 (32.1)	6 (21.4)	-10.7 (-43.2 to 15.9)	.370
Pain intensity of current shoulder injury ^b				
Currently	2.4 (1.5 to 3.3)	1.9 (1.1 to 2.8)	0.5 (-1.4 to 0.5)	.302
Mean in past 24 h	2.1 (1.1 to 3.2)	1.7 (0.8 to 2.7)	-0.4 (-1.5 to 0.6)	.453
Mean in previous 7 d	3.3 (2.4 to 4.3)	2.2 (1.4 to 3.1)	-1.1 (-2.1 to -0.9)	.033
EQ-5D VAS ^c	-13.9 (-20.4 to -7.4)	-8.6 (-14.4 to -2.8)	5.3 (-1.6 to 12.1)	.127
PSFS ^c	-2.5 (-3.6 to 1.4)	-1.3 (-2.5 to -0.2)	1.2 (-0.3 to 2.7)	.124
Positive clinical test findings at baseline and negative at follow-up, n (%)				
Apprehension	18 (64.3)	8 (28.6)	-35.7 (-62.2 to -9.6)	.012
Relocation	14 (50.0)	8 (28.6)	-21.4 (-49.3 to 4.4)	.083
Surprise	15 (53.6)	10 (35.7)	-17.9 (-44.4 to 8.3)	.180
CMS, total ^c	-16.3 (-22.0 to -10.6)	-13.0 (-18.9 to -7.2)	3.2 (-4.8 to 11.3)	.427
Range of motion ^c	-7.3 (-10.0 to -4.6)	-3.5 (-6.2 to -0.8)	3.8 (-0.1 to 7.7)	.057
Strength, ^d kg				
Injured shoulder	2.5 (-5.5 to 0.5)	2.0 (-4.8 to 0.6)	0.5 (-2.5 to 3.4)	.762
Noninjured shoulder	1.5 (0.6 to 2.4)	0.2 (-0.9 to 1.1)	-1.3 (-2.6 to 0.0)	.049
Shoulder JRS (mean absolute error), ^e cm				
Flexion	2.2 (-0.2 to 4.6)	2.3 (0.1 to 4.5)	0.1 (-2.2 to 2.4)	.921
Abduction	1.5 (-0.5 to 3.4)	0.4 (-1.7 to 2.4)	-1.1 (-3.8 to 1.6)	.422

^aData are reported as mean (95% CI) unless stated otherwise. *P* values for continuous and dichotomous outcomes were calculated with analysis of covariance and logistic regression, respectively, using treatment type, study center, sex, and injury status (primary or recurrent anterior shoulder dislocation) as fixed effects and baseline values as covariates. Bolded *P* values indicate statistical significance. CMS, Constant-Murley score; EQ-5D VAS, EuroQol 5-Dimensions questionnaire visual analog scale; HOMEX, self-managed, home-based, standard care shoulder exercise program; JRS, joint reposition sense; PSFS, Patient-Specific Functional Scale; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise; TSK, Tampa Scale of Kinesiophobia; WOSI, Western Ontario Shoulder Instability Index.

^bHigher scores reflect worse status.

^cLower scores indicate better status.

^dIsometric shoulder strength at 90° of shoulder abduction in the scapular plane with the use of a dynamometer.

^eProprioceptive repositioning testing at 60° ± 10° of shoulder flexion and abduction with the use of a laser beam.

subdomain scores (baseline and weeks 4, 8, and 12), which was only shown for the WOSI total (Figure 2).

Adverse Events

Few patients reported recurrent ASDs at the 12-week follow-up, with no significant difference between groups (Table 4). In addition, a large proportion (85% and 83%, respectively) in the SINEX and HOMEX groups reported "very little or not at all" on recurrent anterior shoulder instability or subluxations, with no significant difference between groups (Table 4). Short-lasting adverse events at 12 weeks were few, with no significant group differences (Table 5). The most common adverse events were exercise-induced shoulder pain, soreness, and muscle fatigue (SINEX: n = 10; HOMEX: n = 10).

Sensitivity and Per-Protocol Analyses

Patients who did and did not complete the follow-up assessments (5 patients in total) did not differ on all baseline variables (data not presented), except for employment now in which those lost to follow-up had a lower rate of full-time employment and a higher rate of sick leave. The per-protocol analysis confirmed the ITT analysis, with further significant group differences in favor of SINEX (data not presented).

DISCUSSION

Although the MCID was not reached, this multicenter RCT involving patients with traumatic ASDs showed the

TABLE 4
Self-reported Data at 12-Week Follow-up^a

	SINEX Group (n = 27)	HOMEX Group (n = 24)	P Value
Currently employed or a student			.177
Yes	26 (96)	20 (83)	
Yes, but on sick leave because of current shoulder injury	0 (0)	2 (8)	
No, because of current shoulder injury	0 (0)	1 (4)	
No, for reasons other than current shoulder injury	1 (4)	1 (4)	
Time away from work/education because of current shoulder injury			.184
None	20 (74)	12 (50)	
1-9 d	2 (7)	7 (29)	
10-24 d	3 (11)	2 (8)	
≥25 d	2 (7)	3 (13)	
Use of painkillers (<4 d/wk) because of current shoulder injury			.503
None	24 (89)	23 (96)	
1-4 pills/d	2 (7)	1 (4)	
5-8 pills/d	1 (4)	0 (0)	
Received other treatment(s)			
No other treatment	24 (89)	21 (88)	.865
Physical therapy	1 (4)	2 (8)	.472
Corticosteroid injection	0 (0)	1 (4)	.998
Massage	1 (4)	2 (8)	.494
Other	1 (4)	2 (8)	.494
Rated recurrent anterior shoulder dislocation(s) (yes)	3 (11)	2 (8)	≥.999
Rated recurrent anterior shoulder instability or subluxation(s) (very little/not at all)	23 (85)	20 (83)	.939
Rated extent of impact from actual shoulder injury on number of duties performed at work/school within past 4 wk (very little/not at all)	22 (82)	21 (88)	.600
Rated extent of impact from actual shoulder injury on quality of duties at work/school within past 4 wk (very little/not at all)	22 (82)	20 (83)	.738
Referred to or underwent shoulder stabilizing surgery (yes)	3 (11)	6 (25)	.204
Rated actual shoulder condition as satisfying (to some extent/to a large degree)	22 (82)	18 (75)	.609
Rated current shoulder exercise treatment to have failed (very little/not at all)	26 (96)	20 (83)	.237
Desire to undergo shoulder surgery because of actual shoulder function			.629
No	15 (56)	15 (65)	
No, because of fear of surgery	6 (22)	2 (9)	
Yes	6 (22)	6 (26)	
Rated having returned to preinjury level of sport (to some extent/to a large degree)	9 (33)	11 (46)	.240
Global perceived effect scale (range, 1-7; 7 = very much improved), mean ± SD			
Actual shoulder function	6.2 ± 0.3	5.5 ± 0.3	.012
Ability to perform activities of daily living	5.8 ± 0.3	5.2 ± 0.3	.051
Ability to perform sport/leisure activities	5.8 ± 0.4	4.9 ± 0.4	.025
Shoulder-related quality of life	5.4 ± 0.3	4.8 ± 0.3	.125

^aData are reported as n (%) unless stated otherwise. Bolded P values indicate statistical significance. HOMEX, self-managed, home-based, standard care shoulder exercise program; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise.

superior effects of supervised neuromuscular shoulder exercise versus standard care shoulder exercise after 12 weeks of nonsurgical treatment with regard to the primary outcome of patient-reported shoulder function and quality of life. Further superior effects of SINEX were seen in a number of key secondary outcomes, and although not statistically significant, less than half of the patients in the SINEX group (3/27) compared with the HOMEX group (6/24) underwent or were referred for shoulder stabilizing surgery at the 12-week follow-up. Satisfaction with both exercise programs was high, and no serious adverse events were reported.

We are not aware of any other RCTs investigating the effectiveness of a nonsurgical, physical therapist-supervised progressive exercise regimen in patients with

traumatic ASDs.²⁵ Only 1 other study (noncontrolled cohort study) has evaluated the effect of a 12-week nonsurgical progressive shoulder rehabilitation program in patients with traumatic ASDs (primary dislocations only).¹ That study found that 15 of 20 (75%) patients had no clinical sign of anterior shoulder instability (negative apprehension test finding),¹ which is in line with the current results of 18 of 28 (64%) patients in the SINEX group. Regarding returning to preinjury levels of sport, 15 of 20 (75%) patients in the study by Aronen and Regan¹ returned to unrestricted athletic participation at the 12-week follow-up, which included the sport that caused their initial dislocation. This is in contrast with the current findings in which a smaller proportion of patients in both groups (SINEX, 9/27 [33%];




Figure 2. Western Ontario Shoulder Instability (WOSI) total score at baseline and weeks 4, 8, and 12 for the SINEX versus HOMEX groups among patients with traumatic anterior shoulder dislocations. The graphs illustrate the results from the intention-to-treat population. Data points represent least squares means. (A) Nonresponder imputation (SINEX, n = 28; HOMEX, n = 28). (B) As observed (SINEX, n = 28, 26, 18, and 27, respectively; HOMEX, n = 28, 25, 22, and 25, respectively). (C) Per-protocol analysis (SINEX, n = 12, 10, 10, and 12, respectively; HOMEX, n = 15, 14, 15, and 15, respectively). HOMEX, self-managed, home-based, standard care shoulder exercise program; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise.

HOMEX, 11/24 [46%]) returned to their preinjury level of sport at the 12-week follow-up. One explanation for this discrepancy may be the difference in populations (occupational¹ vs general and not necessarily athletic population [no sport to return to] and first-time¹ vs first-time or recurrent ASD). Another reason may be the different definitions

TABLE 5
Adverse Events From Baseline to Week 12^a

	SINEX Group (n = 27)	HOMEX Group (n = 23)	P
Issues related to shoulder because of shoulder exercise program			
Pain	3 (11)	2 (9)	.766
Soreness	6 (22)	4 (17)	.651
Muscle fatigue	1 (4)	4 (17)	.130
Other (eg, skin irritation)	1 (4)	1 (4)	.882
At least 1 of the above	8 (30)	8 (35)	.668
Sought health care from general practitioner because of shoulder-related issues	2 (7)	1 (4)	.773
Sought health care from orthopaedic surgeon because of shoulder-related issues	0 (0)	2 (9)	.998
Issues related to other body parts because of shoulder exercise program	3 (11)	4 (17)	.335

^aData are reported as n (%). Includes any specific and serious adverse events. P values were calculated with the Fisher exact test. HOMEX, self-managed, home-based, standard care shoulder exercise program; SINEX, nonsurgical, supervised, progressive shoulder instability neuromuscular exercise.

of return to sport (“being allowed to participate in unrestricted athletic participation”¹ and “returning to preinjury level of sport”). Nonetheless, the superior benefit of SINEX in the current study was further consistent with the results of the GPE scale in which the mean rating (7-point Likert scale; 7 = best) for capability to perform sport/leisure activities was higher in the SINEX group than in the HOMEX group (5.8 vs 4.9, respectively; P = .025) at the 12-week follow-up. This corresponds to “much improved” from only “improved.” Thereby, the current results seem to reflect the conclusion of Aronen and Regan, stating “that a progressive neuromuscular program substantially improves the likelihood of full return to activity.”

In agreement with the hypothesis, the SINEX program with supervised physical therapy guidance in neuromuscular shoulder exercises had a larger effect compared with the HOMEX program and was safe to perform. However, with the 95% CI of the primary outcome including the cutoff of 250 points for the WOSI, prespecified as a clinically relevant MCID, a true between-group difference in favor of SINEX cannot be ruled out. However, while previous studies have recommended changes ranging from 210 to 400 of 2100 (10%-19%) for the WOSI total score,^{6,28,46} one might question the current MCID threshold, as it is at the lower end of the suggested changes (228.1). However, despite the fact that both groups were expected to improve, regardless of treatment allocation, a large between-group treatment effect (eg, MCID of 400/2100 [19%]) was not anticipated. Thus, the current MCID of at least 250 of 2100 (12%) points for the WOSI total was pragmatically chosen based on the existing literature and within the common rule of thumb of a minimum of 10% difference for interpreting MCIDs in

patient-reported outcomes.³⁹ Furthermore, most of the secondary patient-reported outcome measures supported the primary finding in the current study but not the objective outcome measures of the CMS and JRS. The reason may be that the CMS is not specifically designed to evaluate shoulder instability⁴⁸ and that JRS testing, for reliability reasons,⁴⁵ was performed in low-range positions only (<60° of shoulder flexion/abduction), thus not able to capture the proprioceptive changes achieved during the SINEX program predominantly including overhead exercises.

Finally, it should be noted that the long-term effects regarding preventing recurrent ASDs, avoiding surgery, and returning to sport, among others, of the current treatments are unknown. In addition, it is unknown how much added cost attributed to physical therapy there is associated with administering SINEX, but a long-term follow-up at 1 and 2 years and cost-benefit analysis will be performed.

Limitations and Strengths

Although the sample size and adherence were relatively low, which increased the risk of type II errors, the current data showed a significant and nearly clinically relevant difference in favor of SINEX. To further support these findings, both the sensitivity and per-protocol analyses confirmed the current ITT analysis.

Another limitation is that the number of glenohumeral Hill-Sachs lesions and anterior bone loss in each group are unknown. However, the nature of the randomized design ensures that these characteristics were equally distributed between the 2 groups, thereby limiting any potential bias. Also, the decision to enroll patients with primary as well as recurrent ASDs may seem controversial, as they may have had different starting points for treatment. However, a recent cross-sectional analysis revealed that patients with primary and recurrent ASDs present with equally poor shoulder function and a high fear of reinjuries, thereby indicating equal indications for receiving treatment regardless of the number of previous dislocations.¹⁶ Furthermore, although stratification may be seen as a limitation, the use of this technique ensured that the number of patients with primary and recurrent ASDs were equally distributed between the 2 groups, thereby limiting any potential bias.

In some countries, it may be that a home-based program is not the standard care for patients with traumatic ASDs, which is why the generalizability of the current findings may be limited. However, the current HOMEX program was designed to reflect the core similarity of standard care packages across most hospitals in Denmark. Because of the design, it was not possible to blind the patients and the treating physical therapists; yet, because all outcome assessments and analyses of the primary outcome were performed blinded and all WOSI scores showed a consistent pattern in favor of SINEX, this is not considered a major limitation.

It is also a limitation that we cannot conclude whether it is the neuromuscular aspect, the difference in supervision, or a combination of the two that makes up the current treatment effect. Finally, because of technical errors having wrongly allocated 2 patients in the analysis performed after the consensus agreement on blinded data interpretation

(see Appendix 4, available as supplemental material), these data could not be used entirely as intended. However, the revised analysis further confirmed the findings in the blinded interpretation document, and the topics suggested as points for discussion in the consensus agreement could still be used.

One strength of this study is the detailed and published protocol including blinded analysis and interpretation of the primary outcome. Furthermore, this study developed and used a standardized, individualized, physical therapist-supervised neuromuscular shoulder exercise program targeting both primary and recurrent ASDs. This study therefore complies with recent recommendations for further research, pointing to the need for more studies to refine an ideal physical therapy regimen for nonsurgical management after primary and recurrent ASDs.²⁵

Another strength is that only 2 outcome examiners were used, which is why assessment variability was kept to a minimum for the objectively measured outcomes. Also, the rate of loss to follow-up was low and equal in both groups.

Finally, it is a strength that the control group (HOMEX) received a realistic and active treatment approach that is often used today and expected to induce a true treatment effect. Still, SINEX managed to improve patients more.

CONCLUSION

Ultimately, 12 weeks of neuromuscular shoulder exercise (SINEX) was superior to a standard care home exercise program (HOMEX) in improving patient-reported shoulder function in patients with traumatic ASDs, although the improvement did not reach the MCID. As SINEX showed the most benefit, this exercise program could potentially be the first-choice option in patients with traumatic ASDs who are not candidates for early shoulder stabilizing surgery. Further long-term treatment effects on patient-reported outcomes and shoulder redislocations as well as socioeconomic evaluations of SINEX are needed to explore the potential for nonsurgical rehabilitation of this complex patient group.

ACKNOWLEDGMENT

The authors thank the staff involved in prescreening the patients at the Odense University Hospital, Aalborg University Hospital, and South-West Jutland Hospital; Annie Gam Petersen for the recruitment of study participants; Odense University Hospital, Aalborg University Hospital, and South-West Jutland Hospital for the use of their facilities for treatment and outcome assessments; Elizabeth Andreassen for help with outcome assessments and data collection; physical therapists Lotte Mejlvig Nielsen, Annelene Houen Larsen, Marianne Gregersen, Gitte Poulsen, Sarah Tofte Benzon, and Mette Nybo Bjerregaard for the instruction and treatment of study participants; Anne Marie Rosager for assisting with administrative tasks; Suzanne Capell for academic editing; and the study participants who made this trial possible.

SUPPLEMENTAL MATERIAL

Appendices 1-4 for this article are available at <http://journals.sagepub.com/doi/suppl/10.1177/2325967119896102>

REFERENCES

- Aronen JG, Regan K. Decreasing the incidence of recurrence of first time anterior shoulder dislocations with rehabilitation. *Am J Sports Med.* 1984;12(4):283-291.
- Ban I, Troelsen A, Christiansen DH, et al. Standardised test protocol (Constant score) for evaluation of functionality in patients with shoulder disorders. *Dan Med J.* 2013;60(4):A4608.
- Bankart AS, Cantab MC. Recurrent or habitual dislocation of the shoulder-joint. *Clin Orthop Relat Res.* 1993;291:3-6.
- Beighton P, Solomon L, Soskolne CL. Articular mobility in an African population. *Ann Rheum Dis.* 1973;32(5):413-418.
- Bottoni CR, Wilckens JH, DeBerardino TM, et al. A prospective, randomized evaluation of arthroscopic stabilization versus nonoperative treatment in patients with acute, traumatic, first-time shoulder dislocations. *Am J Sports Med.* 2002;30(4):576-580.
- Cacchio A, Paoloni M, Griffin SH, et al. Cross-cultural adaptation and measurement properties of an Italian version of the Western Ontario Shoulder Instability Index (WOSI). *J Orthop Sports Phys Ther.* 2012;42(6):559-567.
- Donnelly C, Carswell A. Individualized outcome measures: a review of the literature. *Can J Occup Ther.* 2002;69(2):84-94.
- Downie WW, Leatham PA, Rhind VM, et al. Studies with pain rating scales. *Ann Rheum Dis.* 1978;37(4):378-381.
- Edouard P, Degache F, Beguin L, et al. Rotator cuff strength in recurrent anterior shoulder instability. *J Bone Joint Surg Am.* 2011;93(8):759-765.
- Edouard P, Gasq D, Calmels P, et al. Sensorimotor control deficiency in recurrent anterior shoulder instability assessed with a stabilometric force platform. *J Shoulder Elbow Surg.* 2014;23(3):355-360.
- Eitzen I, Moksnes H, Snyder-Mackler L, et al. A progressive 5-week exercise therapy program leads to significant improvement in knee function early after anterior cruciate ligament injury. *J Orthop Sports Phys Ther.* 2010;40(11):705-721.
- Eshoj H, Bak K, Blond L, et al. Translation, adaptation and measurement properties of an electronic version of the Danish Western Ontario Shoulder Instability Index (WOSI). *BMJ Open.* 2017;7(7):e014053.
- Eshoj H, Ingwersen KG, Larsen CM, et al. Intertester reliability of clinical shoulder instability and laxity tests in subjects with and without self-reported shoulder problems. *BMJ Open.* 2018;8(3):e018472.
- Eshoj H, Juul-Kristensen B, Jorgensen RG, et al. Reproducibility and validity of the Nintendo Wii Balance Board for measuring shoulder sensorimotor control in prone lying. *Gait Posture.* 2017;52:211-216.
- Eshoj H, Rasmussen S, Frich LH, et al. A neuromuscular exercise programme versus standard care for patients with traumatic anterior shoulder instability: study protocol for a randomised controlled trial (the SINEX study). *Trials.* 2017;18(1):90.
- Eshoj H, Rasmussen S, Frich LH, et al. Patients with non-operated traumatic primary or recurrent anterior shoulder dislocation have equally poor self-reported and measured shoulder function: a cross-sectional study. *BMC Musculoskelet Disord.* 2019;20(1):59.
- Frobell RB, Roos EM, Roos HP, et al. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med.* 2010;363(4):331-342.
- Gerber C, Nyffeler RW. Classification of glenohumeral joint instability. *Clin Orthop Relat Res.* 2002;400:65-76.
- Handoll HH, Almayyah MA, Rangan A. Surgical versus non-surgical treatment for acute anterior shoulder dislocation. *Cochrane Database Syst Rev.* 2004;1:CD004325.
- Hayes K, Callanan M, Walton J, et al. Shoulder instability: management and rehabilitation. *J Orthop Sports Phys Ther.* 2002;32(10):497-509.
- Hovelius L, Rahme H. Primary anterior dislocation of the shoulder: long-term prognosis at the age of 40 years or younger. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(2):330-342.
- Jakobsen BW, Johannsen HV, Suder P, et al. Primary repair versus conservative treatment of first-time traumatic anterior dislocation of the shoulder: a randomized study with 10-year follow-up. *Arthroscopy.* 2007;23(2):118-123.
- Järvinen TL, Sihvonen R, Bhandari M, et al. Blinded interpretation of study results can feasibly and effectively diminish interpretation bias. *J Clin Epidemiol.* 2014;67:769-772.
- Juul-Kristensen B, Rogind H, Jensen DV, et al. Inter-examiner reproducibility of tests and criteria for generalized joint hypermobility and benign joint hypermobility syndrome. *Rheumatology.* 2007;46(12):1835-1841.
- Kavaja L, Lahdeoja T, Malmivaara A, et al. Treatment after traumatic shoulder dislocation: a systematic review with a network meta-analysis. *Br J Sports Med.* 2018;52(23):1498-1506.
- Kirkley A, Griffin S, McLintock H, et al. The development and evaluation of a disease-specific quality of life measurement tool for shoulder instability: the Western Ontario Shoulder Instability Index (WOSI). *Am J Sports Med.* 1998;26(6):764-772.
- Kirkley A, Griffin S, Richards C, et al. Prospective randomized clinical trial comparing the effectiveness of immediate arthroscopic stabilization versus immobilization and rehabilitation in first traumatic anterior dislocations of the shoulder. *Arthroscopy.* 1999;15(5):507-514.
- Kirkley A, Werstine R, Ratjek A, et al. Prospective randomized clinical trial comparing the effectiveness of immediate arthroscopic stabilization versus immobilization and rehabilitation in first traumatic anterior dislocations of the shoulder: long-term evaluation. *Arthroscopy.* 2005;21(1):55-63.
- Lephart SM, Warner JJ, Borsa PA, et al. Proprioception of the shoulder joint in healthy, unstable, and surgically repaired shoulders. *J Shoulder Elbow Surg.* 1994;3(6):371-380.
- Leroux T, Wasserstein D, Veillette C, et al. Epidemiology of primary anterior shoulder dislocation requiring closed reduction in Ontario, Canada. *Am J Sports Med.* 2014;42(2):442-450.
- Levine WN, Flatow EL. The pathophysiology of shoulder instability. *Am J Sports Med.* 2000;28(6):910-917.
- Liavaag S, Svenningsen S, Reikeras O, et al. The epidemiology of shoulder dislocations in Oslo. *Scand J Med Sci Sports.* 2011;21(6):e334-e340.
- Little RJ, D'Agostino R, Cohen ML, et al. The prevention and treatment of missing data in clinical trials. *N Engl J Med.* 2012;367(14):1355-1360.
- Longo UG, Loppini M, Rizzello G, et al. Management of primary acute anterior shoulder dislocation: systematic review and quantitative synthesis of the literature. *Arthroscopy.* 2014;30(4):506-522.
- Mintken PE, Cleland JA, Whitman JM, et al. Psychometric properties of the Fear-Avoidance Beliefs Questionnaire and Tampa Scale of Kinesiophobia in patients with shoulder pain. *Arch Phys Med Rehabil.* 2010;91(7):1128-1136.
- Monk AP, Garfjeld Roberts P, Logishetty K, et al. Evidence in managing traumatic anterior shoulder instability: a scoping review. *Br J Sports Med.* 2015;49(5):307-311.
- Olds M, Ellis R, Donaldson K, et al. Risk factors which predispose first-time traumatic anterior shoulder dislocations to recurrent instability in adults: a systematic review and meta-analysis. *Br J Sports Med.* 2015;49(14):913-922.
- Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol Group. *Ann Med.* 2001;33(5):337-343.
- Ringash J, O'Sullivan B, Bezjak A, et al. Interpreting clinically significant changes in patient-reported outcomes. *Cancer.* 2007;110(1):196-202.
- Robinson CM, Howes J, Murdoch H, et al. Functional outcome and risk of recurrent instability after primary traumatic anterior shoulder dislocation in young patients. *J Bone Joint Surg Am.* 2006;88(11):2326-2336.

41. Schulz KF, Altman DG, Moher D, et al. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ*. 2010;340:C332.
42. Simonet WT, Cofield RH. Prognosis in anterior shoulder dislocation. *Am J Sports Med*. 1984;12(1):19-24.
43. Smith RL, Brunolli J. Shoulder kinesthesia after anterior glenohumeral joint dislocation. *Phys Ther*. 1989;69(2):106-112.
44. Tjong VK, Devitt BM, Murnaghan ML, et al. A qualitative investigation of return to sport after arthroscopic Bankart repair: beyond stability. *Am J Sports Med*. 2015;43(8):2005-2011.
45. Vafadar AK, Cote JN, Archambault PS. Interrater and intrarater reliability and validity of 3 measurement methods for shoulder-position sense. *J Sport Rehabil*. 2016;25(1).
46. van der Linde JA, van Kampen DA, van Beers L, et al. The responsiveness and minimal important change of the Western Ontario Shoulder Instability Index and Oxford Shoulder Instability Score. *J Orthop Sports Phys Ther*. 2017;47(6):402-410.
47. Vlaeyen JW, Kole-Snijders AM, Boeren RG, et al. Fear of movement/ (re)injury in chronic low back pain and its relation to behavioral performance. *Pain*. 1995;62(3):363-372.
48. Vrotsou K, Avila M, Machon M, et al. Constant-Murley score: systematic review and standardized evaluation in different shoulder pathologies. *Qual Life Res*. 2018;27(9):2217-2226.
49. White IR, Horton NJ, Carpenter J, et al. Strategy for intention to treat analysis in randomised trials with missing outcome data. *BMJ*. 2011;342:D40.
50. Wintzell G, Haglund-Akerlind Y, Nowak J, et al. Arthroscopic lavage compared with nonoperative treatment for traumatic primary anterior shoulder dislocation: a 2-year follow-up of a prospective randomized study. *J Shoulder Elbow Surg*. 1999;8(5):399-402.
51. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2000;284(23):3043-3045.
52. Zacchilli MA, Owens BD. Epidemiology of shoulder dislocations presenting to emergency departments in the United States. *J Bone Joint Surg Am*. 2010;92(3):542-549.
53. Zech A, Hubscher M, Vogt L, et al. Neuromuscular training for rehabilitation of sports injuries: a systematic review. *Med Sci Sports Exerc*. 2009;41(10):1831-1841.