

Endoscopic ultrasound-guided pleural biopsy in the hands of the pulmonologist

Bibi, Rana; Bodtger, Uffe; Nessar, Rafi; Jensen, Henrik K; Skovgaard Christiansen, Ida; Clementsen, Paul F

Published in:
Respirology case reports

DOI (link to publication from Publisher):
[10.1002/rcr2.517](https://doi.org/10.1002/rcr2.517)

Creative Commons License
CC BY-NC-ND 4.0

Publication date:
2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Bibi, R., Bodtger, U., Nessar, R., Jensen, H. K., Skovgaard Christiansen, I., & Clementsen, P. F. (2020). Endoscopic ultrasound-guided pleural biopsy in the hands of the pulmonologist. *Respirology case reports*, 8(2), 1-3. Article e00517. <https://doi.org/10.1002/rcr2.517>

General rights


Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Endoscopic ultrasound-guided pleural biopsy in the hands of the pulmonologist

Rana Bibi¹, Uffe Bodtger^{2,3,4}, Rafi Nessar³, Henrik K. Jensen¹, Ida Skovgaard Christiansen³  & Paul F. Clementsen^{3,5,6}

¹Department of Respiratory Medicine, Aalborg University Hospital, Aalborg, Denmark.

²Department of Respiratory Medicine, Naestved Hospital, Naestved, Denmark.

³Department of Internal Medicine, Zealand University Hospital, Roskilde, Denmark.

⁴Institute of Regional Health Research, University of Southern Denmark, Odense, Denmark.

⁵Copenhagen Academy for Medical Education and Simulation, Capital Region of Denmark and University of Copenhagen, Copenhagen, Denmark.

⁶Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.

Keywords

EBUS, endoscopic ultrasound, EUS-B, pleura, pleural metastasis.

Correspondence

Ida Skovgaard Christiansen, Department of internal Medicine, Zealand University Hospital, Roskilde, Sygehusvej 10, 4000 Roskilde, Denmark. E-mail: ida.skovgaard@gmail.com

Received: 3 November 2019; Revised: 29 November 2019; Accepted: 13 December 2019; Associate Editor: Jonathan Williamson.

Respirology Case Reports, 8 (2), 2020, e00517

doi: 10.1002/rcr2.517

Abstract

The use of the endobronchial ultrasound (EBUS) endoscope in the oesophagus, the so-called EUS-B procedure, for the diagnosis and staging of thoracic malignancy is quickly gaining ground. Pleural lesions located close to the oesophagus can be inaccessible to transthoracic biopsy and endoscopic procedures can be the only option. We here present two cases demonstrating that EUS-B-guided fine needle aspiration (EUS-B-FNA) of pleural lesions is possible. The first case demonstrates a EUS-B-FNA with malignant mesothelioma of a pleural lesion in a 70-year-old patient. In the second case, EUS-B-FNA diagnosed a pleural metastasis from adenoid cystic adenocarcinoma in a 75-year-old-patient. In conclusion, we hereby demonstrate that EUS-B-FNA from pleural lesions is feasible and appears to be safe.

Introduction

Endoscopic ultrasound (EUS) via the oesophagus can be performed either with the use of a conventional gastrointestinal endoscope (EUS-fine needle aspiration (FNA)) or by using the endobronchial ultrasound (EBUS) endoscope in the oesophagus (EUS-B-FNA) [1,2]. The use of EUS-B is quickly gaining ground. With EUS-B-FNA, it is possible to biopsy lesions in the lungs, mediastinum, liver, and left adrenal gland [1–3].

Case Report

Case 1

A 70-year-old male presented with left pleural thickening with metabolically active areas close to the oesophagus at positron emission tomography-computed tomography (PET-CT) (Fig. 1). Ten years earlier, he was treated for a left-sided malignant pleural mesothelioma (MPM) with partial pericardiectomy, pneumonectomy, and adjuvant chemotherapy. We biopsied the pleural thickening 40 cm from the

teeth row with EUS-B-FNA under conscious sedation using a flexible EBUS endoscope (Olympus XBF-UC180F; Olympus Medical Systems Europe, Ltd., Hamburg, Germany) and a 22 Gauge needle. Three needle passes were performed. Rapid onsite evaluation was not available. The smears and cell blocks showed an abundance of material consisting of rounded cells with medium to ample cytoplasm, and centrally located, rounded to oval, moderately hyperchromic nuclei with distinct nucleolus. The cells were positive for calretinin, CK5, and D240; negative for EP4, CEA, and desmin; and considered diagnostic for MPM of epithelioid type. The patient was referred for palliative treatment.

Case 2

A 75-year-old woman presented with thickening of the right pleura, and PET-CT suggested pleural metastases (Fig. 2A). Seven years earlier, she was diagnosed with adenoid cystic adenocarcinoma in the left glandular parotid region and the




Figure 1. Positron emission tomography-computed tomography (PET-CT) showing thickened and metabolic active left pleural lesions close to the oesophagus, arrow indicates site of biopsy.

tumour was removed surgically. We biopsied the pleural thickening 32 cm from the teeth row with EUS-B-FNA under conscious sedation (Fig. 2B). Three needle passes were performed. The smears and cell blocks showed an abundance of material, predominantly of malignant cells in pseudoglandular or cribriform formations positive for CK7, P63, P40, ASMA, calpotin, and CD117, and negative for EMA and S-100. The biopsies were considered diagnostic for metastasis from the adenoid cystic adenocarcinoma. The patient was offered experimental chemotherapy.

Discussion

The diagnosis and staging of lung cancer and MPM by EUS-FNA is well described [4]. Complications are rare, but oesophageal perforation has been reported in one patient in a cohort of 32 patients with suspected malignant mesothelioma biopsied with EUS-FNA [4]; no complications were observed in either of the two presented cases. Compared to EUS-B, EUS has many advantages, the ultrasonic window angle is larger, the ultrasonic image has higher resolution, and needle




Figure 2. (A) Positron emission tomography-computed tomography (PET-CT) showing thickened and metabolically active right pleural lesions close to the oesophagus. (B) EUS-B-FNA of the pleural lesion.

manoeuvrability is better. However, these benefits are outweighed as pulmonologists are traditionally reluctant to use the EUS endoscope [5], but are increasingly familiar with performing the EUS-B procedures [1–5]. There are obvious logistic and practical advantages in encouraging this, as the pulmonologist in the same session can combine sampling by EUS-B, EBUS, and bronchoscopy in patients suspected of thoracic malignancy including pleural lesions—especially if these are located close to the oesophagus and, therefore, inaccessible to transthoracic biopsy.

Disclosure Statement

Appropriate written informed consent was obtained for publication of this case report and accompanying images.

References

1. Vilmann P, Clementsen PF, Colella S, et al. 2015. Combined endobronchial and esophageal endosonography for the diagnosis and staging of lung cancer: European Society of Gastrointestinal Endoscopy (ESGE) guideline, in cooperation with the European Respiratory Society (ERS) and the

- European Society of Thoracic Surgeons (ESTS). *Endoscopy* 47(6):545–559.
2. Crombag LMMJ, Szlubowski A, Stigt JA, et al. 2017. EUS-B-FNA vs conventional EUS-FNA for left adrenal gland analysis in lung cancer patients. *Lung Cancer* 108(1):38–44.
 3. Korevaar DA, Colella S, Spijker R, et al. 2017. Esophageal endosonography for the diagnosis of intrapulmonary tumors: a systematic review and a meta-analysis. *Respiration* 93(2):126–137.
 4. Tournoy KG, Burgers SA, Annema JT, et al. 2008. Transesophageal endoscopic ultrasound with fine needle aspiration in the preoperative staging of malignant pleural mesothelioma. *Clin. Cancer Res.* 14(19):6259–6263.
 5. Vilmann P, and Clementsen PF. 2015. Combined EUS and EBUS are complementary methods in lung cancer staging: do not forget the esophagus. *Endosc. Int. Open* 3(4): E300–E301.