

## **DNA methylation biomarkers in peripheral blood of patients with head and neck squamous cell carcinomas. A systematic review**

Danstrup, Christian Sander; Marcussen, Mette; Pedersen, Inge Søkilde; Jacobsen, Henrik; Dybkær, Karen; Gaihede, Michael

*Published in:*  
PLOS ONE

*DOI (link to publication from Publisher):*  
[10.1371/journal.pone.0244101](https://doi.org/10.1371/journal.pone.0244101)

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

Danstrup, C. S., Marcussen, M., Pedersen, I. S., Jacobsen, H., Dybkær, K., & Gaihede, M. (2020). DNA methylation biomarkers in peripheral blood of patients with head and neck squamous cell carcinomas. A systematic review. *PLOS ONE*, 15(12), Article e0244101. <https://doi.org/10.1371/journal.pone.0244101>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.




RESEARCH ARTICLE

# DNA methylation biomarkers in peripheral blood of patients with head and neck squamous cell carcinomas. A systematic review

Christian Sander Danstrup<sup>1,2\*</sup>, Mette Marcussen<sup>2</sup>, Inge Søkilde Pedersen<sup>2,3,4</sup>, Henrik Jacobsen<sup>1</sup>, Karen Dybkær<sup>2,4,5</sup>, Michael Gaihede<sup>1,4</sup>

**1** Department of Otorhinolaryngology–Head & Neck Surgery and Audiology, Aalborg University Hospital, Aalborg, Denmark, **2** Clinical Cancer Research Center, Aalborg University Hospital, Aalborg, Denmark, **3** Department of Molecular Diagnostics, Aalborg University Hospital, Aalborg, Denmark, **4** Department of Clinical Medicine, Aalborg University, Aalborg, Denmark, **5** Department of Hematology, Aalborg University Hospital, Aalborg, Denmark

\* [csd@rn.dk](mailto:csd@rn.dk)



## OPEN ACCESS

**Citation:** Danstrup CS, Marcussen M, Pedersen IS, Jacobsen H, Dybkær K, Gaihede M (2020) DNA methylation biomarkers in peripheral blood of patients with head and neck squamous cell carcinomas. A systematic review. PLoS ONE 15(12): e0244101. <https://doi.org/10.1371/journal.pone.0244101>

**Editor:** Mariana Brait, Johns Hopkins University, UNITED STATES

**Received:** April 24, 2020

**Accepted:** December 2, 2020

**Published:** December 17, 2020

**Copyright:** © 2020 Danstrup et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its [Supporting Information files](#).

**Funding:** This study has been supported by Department of Clinical Medicine, Aalborg University and Department of Otorhinolaryngology – Head & Neck Surgery and Audiology, Aalborg University Hospital, Denmark. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract

### Background

Head and neck squamous cell carcinomas (HNSCC) are often diagnosed in advanced stages. In search of new diagnostic tools, focus has shifted towards the biological properties of the HNSCC, and the number of different biomarkers under investigation is rapidly growing.

### Objectives

The objective was to review the current literature regarding aberrantly methylated DNA found in peripheral blood plasma or serum in patients with HNSCC and to evaluate the diagnostic accuracy of these changes.

### Methods

The inclusion criteria were clinical studies involving patients with verified HNSCC that reported findings of aberrantly methylated DNA in peripheral blood serum or plasma. We systematically searched PubMed, OVID Embase and Cochrane Library. In addition to the search, we performed forward and backward chaining in references and Web of Science. The protocol was registered in PROSPERO: CRD42019135406. Two authors independently extracted data. The quality and the risk of bias of the included studies were assessed by the QUADAS-2 tool.

### Results

A total of 1,743 studies were found eligible for screening, while ultimately seven studies were included. All studies were found to have methodological weaknesses, mainly

**Competing interests:** The authors have declared that no competing interests exist.

concerning patient selection bias. The best individual marker of HNSCC was Septin 9 in plasma with a sensitivity of 57% and a specificity of 95%.

## Conclusions

None of the aberrantly methylated genes found in the retrieved studies are applicable as single diagnostic markers for HNSCC and the best gene-panels still lack diagnostic accuracy. Future studies may benefit from newer sequencing techniques but validation studies with well-designed cohorts are also needed in the process of developing epigenetic based diagnostic tests for HNSCC.

## Introduction

Head and neck cancer represents 4% of all malignancies with more than 650,000 new cases worldwide each year, where the majority of these cancers are squamous cell carcinomas (HNSCC) [1–3]. The overall 5 year survival of the patients is 60%, and it has improved only moderately over the last decades [4–6]. A major reason for this is probably that patients with HNSCC often present at later stages of their disease (III–IV as defined by the American Joint Committee on Cancer (AJCC) cancer staging manual [7]). Survival is inversely related to disease stage [8], and for patients with oro- and hypopharyngeal cancers, 72–75% present with late stages, with a 5 year survival rate below 50% [9, 10]. Smoking and alcohol consumption are major risk factors in HNSCC [11], however, the role of human papillomavirus (HPV) has within the last 10 years been increasingly recognized as a factor, particularly in the development of oropharyngeal cancer [12]. The HPV-related oropharyngeal carcinomas have a better prognosis and show better response to treatment [13]. The differences between HPV-related and non HPV-related oropharyngeal carcinomas have caused that separate staging systems were established in the eighth edition of AJCC cancer staging manual [7].

In search of tools for earlier diagnosis, improved prognostic evaluation and detection of recurrent disease, focus has shifted towards the biological properties of HNSCC, and the number of biomarkers under investigation is rapidly growing. Initially, proteins and later changes in DNA, RNA (microRNA), and circulating tumor cells have been investigated. A particular evolving branch of DNA-research is epigenetic changes, where DNA-methylation represents one of the most studied modifications. This consists of the addition of a methyl (CH<sub>3</sub>) residue in a cytosine base proceeding a guanosine base, known as a CpG dinucleotide; in normal human DNA 3–6% of all cytosines are methylated [14].

Most CpG dinucleotides are found in CpG-rich regions, known as CpG islands [15]. 55–75% of human genes have been reported to have CpG rich promoters, and these are predominantly unmethylated in normal tissue [16, 17]. In the development of cancers, aberrant methylation of promoter-regions is of particular interest, as hypermethylation of tumor-suppressor genes may result in downregulation of the activity of the tumor suppressor function, while hypomethylation of promoter regions in oncogenes may result in an increased gene expression [18–21].

Cell-free DNA (cfDNA) from solid tumors has been demonstrated in the bloodstream. While the exact mechanism behind such release of cfDNA is still debated, apoptotic and necrotic cells are thought to be the origin [22, 23]. If these DNA-fragments can be detected and their methylation-status determined, this holds the potential of becoming an important tool in early demonstration and diagnosis of malignant solid tumors. In several types of malignant solid tumors, but not HNSCC, commercial tests based on aberrant methylation are

already available (e.g. the Epi proColon<sup>®</sup> and Cologuard<sup>®</sup> for colorectal-cancer screening and the Epi proLung<sup>®</sup> in detection of lung cancer) [24–29].

Based on this background, the primary objective of this paper was to review the current literature of the usage of aberrantly methylated DNA found in the peripheral blood plasma or serum of patients with HNSCC and evaluate the diagnostic accuracy of these changes. A second objective was to evaluate the potential prognostic value of such findings.

## Materials and methods

We conducted a systematic review according to the Preferred Reporting Items for Systematic review and Meta-analysis (PRISMA) guidelines ([S1 Checklist](#)) [30]. The protocol was registered in the PROSPERO database for systematic reviews with registration number: CRD42019135406 [31].

### Inclusion criteria

Clinical studies involving patients with histopathologically verified HNSCC that reported findings of aberrantly methylated DNA in peripheral blood serum or plasma were eligible.

### Exclusion criteria

Studies were excluded, if they were: (1) performed in animals; (2) only describing aberrantly methylated DNA in solid tumors; (3) published only as conference abstracts; (4) published only in a review; or (5) published in a language other than English.

### Search strategy

In June 2019, we performed a systematic literature search in OVID Embase, PubMed and Cochrane Library. No limits were placed on the publication dates in our search. In addition to the search, we performed forward and backward chaining in references and Web of Science. The search strategy was structured by one reviewer (C.D.) in collaboration with Pernille Gaardsted, research librarian at the Medical Library, Aalborg University Hospital. An updated search was performed in March 2020. The full search strategy appears in [S1 Table](#).

### Study selection

Papers found according to our search strategy were imported into the Covidence online software (Covidence, Melbourne AUS for data management and literature screening [32]). After duplicate removal, one reviewer (C.D.) screened titles and abstracts of all potential studies using the inclusion criteria. Then all full text articles were assessed for eligibility by two independent reviewers (C.D., M.M.). Any disagreement was resolved by consensus. A flowchart showing the study selection process in according to PRISMA was generated ([Fig 1](#)).

### Data collection process

Two authors (C.D., M.M.) independently extracted data and assessed the risk of bias in the included studies. Data extraction included the name of first author, year of publication, number of HNSCC cases, number of controls, medium examined (plasma/serum), type of methylation analysis used, diagnostic performance, prognostic value of the methylation status in HNSCC patients measured by recurrence free survival and overall survival. Any disagreement was resolved by consensus.

Due to the heterogeneity of the included studies a meta-analysis was not considered meaningful and a structured narrative synthesis was performed.




Fig 1. PRISMA flow diagram. From Moher et al. [30].

<https://doi.org/10.1371/journal.pone.0244101.g001>

### Quality assessment and assessment of the risk of bias

The quality and the risk of bias of the included studies were assessed by the QUADAS-2 tool [33]. The tool is designed to assess the quality of primary diagnostic accuracy studies. The following domains were evaluated: Patient selection, index test, reference standard, flow and timing.

### Results

A total of 2428 studies were imported into Covidence. After removal of duplicates by Covidence (685) a total of 1743 studies were eligible for screening according to the inclusion

criteria. The screening process resulted in the removal of 1705 papers (incl. two duplicates found during the screening) and left 38 papers for assessment. The full text assessment resulted in the further exclusion of 31 studies and the inclusion of a total of 7 studies for final analysis [34–40] (Fig 1). The reference lists of the included articles were screened for additional relevant articles, but none met the inclusion criteria.

The earliest study was performed in 2000 [40], and the latest studies performed in 2017 [37, 38]. Six of the seven studies used gene panels based on results derived from earlier published studies of HNSCC or other malignant tumors (e.g. lung, breast and colorectal cancers), while Mydlarz et al. chose a panel based on own earlier findings in saliva [35]. All studies evaluated the methylation status of the primary tumor and of peripheral blood, serum, or plasma with focus on a limited number of genes. None of the investigators performed a broad or genome-wide analysis. The studies show a large variation in included number of cases and controls. Thus, 425 HNSCC patients were included in the study by Schröck et al. [37], contrasting the 17 patients with oral cancer described by Nakahara et al. [39].

It is noteworthy that studies [37, 38] are from the same research group and that studies [34, 35, 40] are also from one group.

## Results from studies of methylation in serum

In 2008, Carvalho et al. [34] published a study on evaluation of promoter hypermethylation in body fluids as a diagnostic tool in patients with HNSCC and healthy controls using quantitative methylation-specific polymerase chain reaction (qMSP). They examined a panel of 21 genes that previously were described as hypermethylated in HNSCC or other solid tumors. Six genes found in serum were significantly associated with HNSCC. Hypermethylation of each gene was treated as a binary variable (methylation versus no methylation) and each gene was evaluated separately and combined in a panel of the best possible combinations regarding sensitivity and specificity (Table 1).

The performances of the selected panels were explored using multivariable logistic regression and receiver operating characteristic (ROC) curves. Internal validation of the logistic regression was done by using an approximation to the leave-one-out jackknife procedure. Of the six genes only one, Hypermethylated in Cancer 1 (HIC1), was considered useful as a diagnostic single-gene-marker with a sensitivity of 31.4% and a specificity of 92.5%. No combinations of the selected genes were able to surpass the specificity of HIC1. A panel of five genes showed a sensitivity of 87.2% and a specificity of 42.3%.

In their study, Mydlarz et al. [35] evaluated 100 HNSCC patients and 50 controls using qMSP to detect hypermethylated Endothelin Receptor Type B (EDNRB), Deleted in Colorectal Cancer (DCC) and Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A). Detection of hypermethylation was interpreted in a binary fashion without any cut-off as no methylation was found in normal serum. In the HNSCC group, genes were hypermethylated in 10%, 2% and 1%, respectively. No hypermethylation of the specific genes could be demonstrated in the control group. There was no significant association between serum positivity for hypermethylated genes and local recurrence free and overall survival (Table 2).

In another study, Nakahara et al. [39] examined hypermethylation of CDKN2A in biopsies and serum of 17 oral squamous cell carcinoma (OSCC) patients and eight healthy controls using MSP. Methylation was treated as a binary outcome. They found a sensitivity of 35% and a specificity of 100% in serum. The authors performed pre- and post-treatment serum analyses, and all patients with aberrant methylation in serum before treatment were non-methylated at two months post-treatment follow up. Four patients had recurrence and three of these

Table 1. Diagnostic value of DNA methylation in plasma or serum.

Gene	Name	Cases/ controls	Concordance rate between methylation in tumor and blood	Anatomical site	Stage	Specimen	Sensitivity	Specificity	AUC	Method	Reference
CDH1	Cadherin 1	62/320		All	NA	Serum	32%	73%	0.53	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
CCND2	Cyclin D2	47/284		All	NA	Serum	6%	98%	0.52	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
TGFBR2	Transforming Growth Factor Beta Receptor 2	37/134		All	NA	Serum	8%	94%	0.51	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
TIMP3	TIMP Metalloproteinase Inhibitor 3	50/296		All	NA	Serum	10%	95%	0.52	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
HIC1	Hypermethylated in Cancer 1	70/373		All	NA	Serum	31%	93%	0.62	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
PGP9.5	Neuron Cytoplasmatic Protein 9.5	52/203		All	NA	Serum	8%	98%	0.53	qMSP <sup>b</sup>	Carvalho et al. [34] (2008)
SEPT9 <sup>1</sup>	Septin 9	137/170	-	All	All	Plasma	62%	92%	0.79	quasi-digital PCR	De Vos et al. [38] (2017)
SHOX2 <sup>1</sup>	Short Stature Homeobox 2	137/170	-	All	All	Plasma	53%	87%	0.78	quasi-digital PCR	De Vos et al. [38] (2017)
SEPT9 <sup>2</sup>	Septin 9	141/102	-	All	All	Plasma	55%	90%	0.75	quasi-digital PCR	De Vos et al. [38] (2017)
SHOX2 <sup>2</sup>	Short Stature Homeobox 2	141/102	-	All	All	Plasma	43%	95%	0.77	quasi-digital PCR	De Vos et al. [38] (2017)
EDNRB	Endothelin Receptor Type B	100/50	-	All	All	Serum	10%	100%		qMSP <sup>b</sup>	Mydlarz et al. [35] (2016)
DCC	Deleted in Colo-rectal Cancer	100/50	-	All	All	Serum	2%	100%		qMSP <sup>b</sup>	Mydlarz et al. [35] (2016)
CDKN2A	Cyclin-Dependent Kinase Inhibitor 2A	100/50	-	All	All	Serum	1%	100%		qMSP <sup>b</sup>	Mydlarz et al. [35] (2016)
CDKN2A	Cyclin-Dependent Kinase Inhibitor 2A	17/8	65%	Oral	All	Serum	35%	100%		MSP <sup>a</sup>	Nakahara et al. [39] (2005)
CDKN2A	Cyclin-Dependent Kinase Inhibitor 2A	50/0	27%	All	All	Serum	4%	-		MSP <sup>a</sup>	Sanchez-Céspedes et al. [40] (2000)
MGMT	O <sup>6</sup> -Methylguanine-DNA-Methyltransferase	50/0	33%	All	All	Serum	7%	-		MSP <sup>a</sup>	Sanchez-Céspedes et al. [40] (2000)
DAPK	Death-Associated Protein Kinase	50/0	18%	All	All	Serum	2%	-		MSP <sup>a</sup>	Sanchez-Céspedes et al. [40] (2000)

(Continued)

Table 1. (Continued)

Gene	Name	Cases/ controls	Concordance rate between methylation in tumor and blood	Anatomical site	Stage	Specimen	Sensitivity	Specificity	AUC	Method	Reference
GSTP1	Glutathione S-transferase p1	50/0	0%	All	All	Serum	0%	-		MSP <sup>a</sup>	Sanchez-Cespedes et al. [40] (2000)
SHOX2 <sup>1</sup>	Short Stature Homeobox 2	137/122	-	All	All	Plasma	50%	95%	0.80	qPCR	Schröck et al. [37] (2017)
SEPT9 <sup>1</sup>	Septin 9	137/122	-	All	All	Plasma	57%	95%	0.79	qPCR	Schröck et al. [37] (2017)
SHOX2 <sup>2</sup>	Short Stature Homeobox 2	141/102	-	All	All	Plasma			0.79	qPCR	Schröck et al. [37] (2017)
SEPT9 <sup>2</sup>	Septin 9	141/102	-	All	All	Plasma			0.74	qPCR	Schröck et al. [37] (2017)
CDKN2A	Cyclin-Dependent Kinase Inhibitor 2A	20/24	49%	All	All	Plasma	65% (13/20)	67% (16/24)		Real time PCR	Wong et al. [36] (2003)
CDKN2B	Cyclin-Dependent Kinase Inhibitor 2B	20/24	60%	All	All	Plasma	60% (12/20)	50% (12/24)		Real time PCR	Wong et al. [36] (2003)

a: Methylation specific PCR.

b: Quantitative methylation specific PCR.

c: Combined bisulfite restriction analysis.

1: Training Cohort.

2: Testing cohort.

\*The test is between OSCC and OP. No healthy controls.

<https://doi.org/10.1371/journal.pone.0244101.t001>

patients, had aberrant methylation in the peripheral blood. In addition, all three had aberrant methylation in both tumor and serum prior to treatment.

Using MSP for either methylated or unmethylated sequences, Sanchez-Cespedes et al. [40] published a study in 2000 examining aberrant methylation of four genes, CDKN2A, O<sup>6</sup>-methylguanine-DNA-methyltransferase (MGMT), Death-associated protein kinase (DAPK), Glutathione S-transferase P1 (GSTP1). Out of 95 HNSCC tumor and serum samples, 52 patients had aberrant methylation in the primary tumor. MSP was unsuccessful in two cases and therefore only 50 serum samples were evaluated. Twenty-one of 50 patients had the same methylation changes in the serum as in the primary tumor. CDKN2A was positive in eight serum samples, MGMT in 14 and DAPK in three. GSTP1 hypermethylation was not found in any tumor or serum sample. As control subjects, they analyzed serum of 25 patients, who did not have aberrant methylation in the primary tumor. None of the serum samples in the control group, showed aberrant methylation of the four genes. Seven of the patients had a new serum sample post-treatment. None of the methylation-negative patients had converted, however, one DAPK-positive patient had become negative, whereas two patients still were methylation-positive 5–6 months post-treatment.

Table 2. Prognostic value of DNA methylation in plasma or serum.

Gene(s)	Name	Anatomical site	Stage	Local recurrence-free survival (hazard ratio)	Overall survival (hazard ratio)	Method	Reference
EDNRB/DCC/CDKN2A		All	All	1.35 <sup>c</sup> (0.17–10.5)	1.35 <sup>c</sup> (0.41–4.43)	qMSP	Mydlarz et al. [35]
SEPT9 <sup>1</sup>	Septin 9	All	All	-	5.27 <sup>a</sup> (2.03–13.68)	qPCR	Schröck et al. [37]
SHOX2 <sup>1</sup>	Short statue Homeobox 2	All	All	-	2.32 <sup>a</sup> (1.12–4.83)	qPCR	Schröck et al. [37]
SEPT9 <sup>1</sup>	Septin 9	All	All	1.32 <sup>b</sup> (1.09–1.59)	1.23 <sup>b</sup> (1.07–1.41)	qPCR	Schröck et al. [37]
SEPT9 <sup>2</sup>	Septin 9	All	All	-	2.78 <sup>a</sup> (1.16–6.67)	qPCR	Schröck et al. [37]
SHOX2 <sup>2</sup>	Short statue Homeobox 2	All	All	-	2.50 <sup>a</sup> (1.12–5.60)	qPCR	Schröck et al. [37]
SEPT9 <sup>1</sup>	Septin 9	All	All	-	1.90 <sup>d</sup> (1.43–2.69)	quasi-digital PCR	De Vos et al. [38]
SHOX2 <sup>1</sup>	Short statue Homeobox 2	All	All	-	1.85 <sup>d</sup> (1.18–2.91)	quasi-digital PCR	De Vos et al. [38]
SEPT9 <sup>2</sup>	Septin 9	All	All	-	1.67 <sup>d</sup> (1.13–2.45)	quasi-digital PCR	De Vos et al. [38]
SHOX2 <sup>2</sup>	Short statue Homeobox 2	All	All	-	1.71 <sup>d</sup> (1.05–2.80)	quasi-digital PCR	De Vos et al. [38]

a: Univariate cox proportional hazards analysis of the 129 cases with positive serum samples using the introduced cut-off value.

b: Multivariate cox proportional hazards analysis of the 129 cases using the DNA methylations levels as a continuous variable.

c: Univariate proportional hazards analysis.

d: Univariate cox proportional hazards analysis.

1: Training Cohort.

2: Testing Cohort.

<https://doi.org/10.1371/journal.pone.0244101.t002>

## Results from studies of methylation in plasma

Short Statute Homeobox 2 (SHOX2) and Septin 9 (SEPT9) have been examined by Schröck et al. and De Vos et al. [37, 38]. First, a *training cohort* with 284 patients with HNSCC and 122 matched healthy controls and a *test cohort* with 141 patients with HNSCC and 102 matched healthy individuals were tested for these two biomarkers with triplex quantitative PCR. Only 137 of 284 patients with HNSCC in the *training cohort*, had baseline blood plasma samples available. The authors introduced a methylation cut-off value to dichotomize the methylation values and levels below these values were considered sporadic background methylation. Matched tumor samples and pretherapeutic methylation in plasma from 55 patients were analyzed [37]. SHOX2 methylation in plasma correlated significantly with tissue levels (Spearman's  $p = 0.36$ ,  $P = 0.007$ ), while SEPT9 showed a suggestive trend (Spearman's  $p = 0.25$ ,  $P = 0.067$ ). ROC curves have been computed as a measure of diagnostic accuracy and in the *training cohort* SHOX2/SEPT9 sensitivity and specificity were reported to be 59% and 96%, respectively (Table 1). Using univariate Cox proportional hazard analysis for both cohorts, the authors found a significantly higher risk of death for methylation positive patients for both markers [37, 38] (Table 2).

In their second study [38], the team evaluated three different algorithms to assess methylated cfDNA; Relative quantification, Absolute quantification and Quasi-Digital PCR. They used the same two markers (SEPT9 and SHOX2) and almost the same training and testing cohort as in their previous study [37], only adding 48 extra controls to the *training cohort*. All three

Table 3. Risk of bias and applicability concerns.

Study	RISK OF BIAS				APPLICABILITY CONCERNS		
	PATIENT SELECTION	INDEX TEST	REFERENCE STANDARD	FLOW AND TIMING	PATIENT SELECTION	INDEX TEST	REFERENCE STANDARD
Carvalho et al. [34]	High	High	Low	Unclear	Low	Low	Low
Mydlarz et al. [35]	Unclear	High	Low	Unclear	Low	Low	Low
Nakahara et al. [39]	High	High	Low	Unclear	Unclear	Low	Low
Sanchez-Cespedes et al. [40]	Unclear	High	Low	Unclear	High	Unclear	Low
Schröck et al. [37]	High	High	Low	Unclear	Low	Low	Low
De Vos et al. [38]	High	High	Low	Unclear	Low	Low	Low
Wong et al. [36]	High	High	Low	Unclear	Unclear	Low	Low

<https://doi.org/10.1371/journal.pone.0244101.t003>

algorithms showed similar Area Under the Curve (AUC) in the ROC curves (Performance for Quasi-Digital PCR can be found in Table 1). In prognostic evaluation in both cohorts, all three algorithms also showed significantly increased hazard ratios for methylation positive patients (Performance for Quasi-Digital PCR can be found in Table 2).

Finally, Wong and colleagues [36], used real time PCR to examine promoter methylation of the p16 and p15 genes in plasma. Methylated p16 was found in 65% of the patients with HNSCC in contrast to 20% of the healthy controls. Methylated p15 was seen in the plasma of 60% of the patients with HNSCC and in 50% of the healthy controls. The mean concentration of methylated p16 and p15 was significantly higher in HNSCC patients compared to healthy controls ( $p = 0.016$  and  $0.0037$  respectively), but no cut-off values for background methylation was defined.

### Risk of bias

The QUADAS-2 evaluation of the included studies showed that all studies had a high or unclear risk of bias related to patient selection. None of the studies explicitly described how patients or controls were enrolled, e.g. consecutively or randomly or other criteria. Furthermore, all the studies were case control by design. All index-tests were interpreted with the knowledge of the reference standard in the HNSCC groups. None of the included studies described the precise flow and timing regarding time from diagnosis to biopsy and blood sampling. (An overview of the QUADAS-2 evaluation appears in Table 3).

### Applicability concerns

There were no concerns regarding the condition as defined by the reference standard in any of the included studies. One study [40], was graded as unclear concerning the index test, as no healthy control group was included.

Two studies [27, 30], did not report any information on the control groups, and they were judged as unclear regarding patient selection. As no healthy control group was defined in the study by Sanchez-Cespedes et al. [40], the concern regarding patient selection was high. The remaining four studies [34, 35, 37, 38], were judged as low as they had matched control groups and provided tables with demographic information regarding age, gender, tobacco- and alcohol consumption. (An overview of the applicability concerns is given in Table 3)

### Discussion

Epigenetic changes, including aberrant DNA methylation, in human cancers was initially described more than thirty years ago [41]. With the development of new laboratory techniques

and assays, the ability to examine body fluids, especially blood, for circulating tumor cells or tumor DNA has inspired the search of new diagnostic and prognostic tools in cancer. The objective of this review was to provide an overview of aberrantly methylated DNA found in plasma or serum of peripheral blood and their diagnostic and prognostic potential in patients with HNSCC.

The majority of the studies, we retrieved in this systematic review, described both sensitivity and specificity of the different aberrantly methylated genes, whereas the study by Sanchez-Cespedes et al. [40] did not include a healthy control group, so that specificity could not be calculated.

Not all included studies presented both diagnostic and prognostic data, as only three studies [35, 37, 38], described follow-up, and therefore, they were able to calculate hazard ratios on local recurrence free survival and overall survival.

Both single genes and gene-panels were evaluated. The best individual marker was found by Schröck et al. [37] as detected hypermethylated SEPT9 in plasma had a 57% sensitivity and 95% specificity. Hypermethylated SEPT9 was also assessed in studies of other solid tumors, e.g. the Epi proColon<sup>®</sup> that is a commercially available blood-based DNA hypermethylation test screening for colorectal cancer. As a single marker, the assay reached a sensitivity of 48.2% and specificity of 91.5% in a large prospective study with 6874 asymptomatic patients scheduled for colorectal cancer screening [42].

When combining both SHOX2 and SEPT9 ( $\text{Mean}_{\text{Shox2/SEPT9}}$ ), Schröck et al. [37] demonstrated that the sensitivity and specificity can be improved to 59% and 96%, respectively. In the study by Carvalho et al. [34] the best combined gene panel generated a sensitivity of 87% and a specificity of 42%. Their study was limited by the fact that not all genes were analyzed in all HNSCC- and control samples.

### Anatomical subsite, stage and HPV status

The anatomical subsite and HNSCC tumor stage may also play a role in the release of tumor DNA, and hence, the methylation-findings in peripheral blood. Thus, tumors in subsites with rich lymphatic drainage and blood supply may possibly be detected earlier. For example, the lymphatic drainage and blood supply of the larynx is remarkably different to other subsites due to the dual embryological origin [43]. Tumor size may also play a role, as small T1 tumors of the larynx will present a small cell-turnover compared to a large invasive T4b supraglottic tumor invading the prevertebral space.

The incidence of HPV-related HNSCC has been increasing during the last decades and is now described as an independent subtype with a distinct morphological profile and occurs primarily in the oropharynx of younger patients without tobacco and alcohol as co-factors [44, 45]. It has been shown that expression of the p16 protein is highly correlated to HPV in oropharyngeal cancer where the protein is upregulated in contrast to non HPV-related HNSCC. Due to this strong correlation, immunohistochemical analysis of p16 are used as a surrogate marker of HPV-related HNSCC [46].

This etiological linkage and role of HPV should be taken into consideration when evaluating methylation profiles as these may differ between HPV-related and non HPV-related HNSCC. This has been addressed by Delgi Esposti et al. [47]. Using pooled genome-wide analyses they found that HPV infection affects the DNA methylation in HNSCC across different anatomical subsites.

Of the seven studies included, only Mydlarz et al. [35] reported the HPV-status in their HNSCC patients, but did not mention the method used to determine HPV-status. They described the HPV- status related to survival but not in relation to the biomarkers examined.

Five studies [34–36, 39, 40] evaluated methylation of the p16 promoter, but none described or discussed the findings in relation to HPV and potential differences between HPV-related and non HPV-related HNSCC.

The mechanism of DNA release into the circulation is not fully clarified. However, it is suggested that circulating DNA is derived from apoptotic or neoplastic cells or from tumor cells that are not capable of generating metastases [22, 23, 48]. Carvalho et al. [34] did not report the tumor size and stage of the cancer patients examined, while Wong et al. [36] only reported methylation status of the primary tumor in relation to tumor size. All positive samples in the study by Mydlarz et al. [35] were Stage IV cancers and Schröck et al. [37] found a strong correlation between the level of the biomarkers in plasma and tumor- and nodal-category.

This should be taken into consideration when evaluating blood-based biomarkers as screening tools for HNSCC and if possible, anatomical subsite, HPV status, tumor size and stage should be reported when evaluating the potential diagnostic and prognostic biomarkers.

### Examination of plasma versus serum—flow and timing

The use of plasma and serum was almost equally distributed as four studies [34, 35, 39, 40], evaluated methylation in serum, and three studies [36–38] evaluated methylation in plasma.

In general, the amount of cell free DNA is higher in serum compared to plasma (2–24 times). Therefore, serum is suggested as the best source for evaluation, however, Jung et al. [49] examined the changes in the concentration of DNA in serum and plasma during storage and handling of blood samples. They found no change in DNA concentration in plasma stored 8h at room temperature, but time delay and storage temperature had significant impact (3.8–4.8 times higher for samples stored 2–8h at room temperature) on the concentration of DNA in serum.

Of the included studies, only Schröck et al. [37] and De Vos et al. [38] explicitly described the plasma preparation regarding timing, as plasma were prepared within 2h after sampling.

None of the included studies described the flow and timing between the biopsy (histologically verifying the diagnosis) and the blood samples. If a large biopsy in a vascularized field is taken, it may cause a release of tumor-cells in the bloodstream, and hence, an elevation of cfDNA in the blood sample afterwards. Henriksen et al. demonstrated that total cfDNA levels were elevated up to four weeks after surgery in patients undergoing colorectal cancer surgery [50]. Therefore, when studying biomarkers potentially released from tumor tissue, the study should preferably include a measurement of the DNA methylation status before (and after) the primary biopsy with documented flow and timing, or at least describe the flow of sample collection.

### Cut-off values and background methylation

Only two studies [35, 39] did not find any aberrant methylation of the evaluated biomarkers in the healthy control groups. The fact that the other studies find aberrant methylation in healthy controls suggest that other factors may contribute to the methylation-level in general. This issue was only addressed by Schröck and others [37] and defined as sporadic background methylation. Therefore, a methylation level cutoff was introduced.

Wong et al. [36] evaluated the resection-margins of 29 smoking patients with HNSCC. They found that methylation-levels of the p15 promoter were significantly higher in resection margins (and histopathologic healthy tissue) in chronic smokers compared to non-smokers. These findings have also been documented by others [51, 52]. When evaluating a biomarker as a screening or diagnostic tool, it should be evaluated in a group of people at risk. Matched controls regarding age, gender, use of tobacco, alcohol and general health-status should be

preferred. Of the included studies, two failed to report demographics of their control patients, hence introducing potential bias in the patient selection [36, 39].

### Reflections of tumor-methylation in peripheral blood–concordance rate

All the included studies describe that methylation analyses of the primary tumors have been performed, but not all described the concordance between tumor- and the serum-methylation.

Using Spearman correlation, Schröck et al. analyzed 55 matched tumor and blood samples and found a significant correlation for SHOX2 but not for SEPT9. Sanchez-Cespedes et al. [40] found that 42% of the examined patients had the same methylation changes in serum and the primary tumor.

In the study of aberrantly methylated CDKN2A by Nakahara et al. [39], 54.5% of the patients had matching tumor and serum changes. In a study of CDKN2A in 94 colorectal cancer patients, a concordance rate of 30% was found [53], and in a study of primary lung cancer (using MSP) by Ooki et al., the concordance rate was between 20–56% in a six-gene panel [54].

None of the included studies had a perfect concordance rate of methylation in tumor and plasma/ serum, which raises the question, why that is so. HNSCC are highly heterogeneous tumors and different methylation-profiles may exist even within the tumor. Thus, different biopsy-sites may yield different methylation-profiles. If different methylation-profiles exists, this may also be reflected in the DNA released to the circulation.

### Prognostic value of aberrant blood methylation in HNSCC and post-treatment surveillance

Several studies have evaluated the prognostic value of aberrant methylation profiles in the primary tumors of HNSCC [55–59]. Of the included studies, only three [35, 37, 38] had follow-up, thus describing local recurrence free- and overall survival. Mydlarz et al. [35] did not find a significant association between local recurrence or overall survival and promoter hypermethylation of the three genes examined (Table 2).

Schröck et al. [37] performed univariate proportional hazards analyses on both cohorts and multivariate analyses on their *training cohort*. In the univariate analyses, patients with positive plasma levels of either SEPT9 or SHOX2 were at higher risk of death compared to the methylation negative patients. In multivariate analyses of the 129 patients in their *training cohort*, SEPT9 proved to be an independent prognostic factor regarding local recurrence free survival (HR: 1.32 CI:1.09–1.59) and overall survival (HR:1.23 CI: 1.07–1.41) (Table 2). De Vos et al. [38] found both SEPT9 and SHOX2 to be significant prognostic factors regarding overall survival in all tested quantification algorithms (Table 2).

Post-treatment evaluation of aberrant methylated DNA was only reported in two of the seven studies included [39, 40]. Nakahara et al. found that all six patients with positive tests with regard to pre-treatment aberrant methylated DNA had converted to negative tests two months after treatment; four of these patients were later diagnosed with recurrent disease, and three had converted back to positive tests [39]. Sanchez-Cespedes et al. reported from post-treatment samples that in three patients with positive pre-treatment tests, only one patient became negative after treatment, while in two patients' tests were still positive at five to six months of follow-up. It was not reported, whether these patients later had recurrent disease [40]. The role of post-treatment profiles of aberrant methylated DNA in HNSCC remains unclear. Symonds et al. reported conversion of positive to negative tests in 35 of 47 patients with colorectal-cancer [60]. Thus, it seems relevant to pursue this aspect in patients with HNSCC as it may reveal an important method for surveillance.

## Methods used in methylation analyses

The included studies use different methods in their methylation analyses. Methylation-specific PCR (MSP) was one of the earliest methods used in methylation studies and was initially described by Herman et al. in 1996 [61]. The method uses primers to recognize methylated- and unmethylated CpG's. The method is cost-effective, but has a disadvantage of false positives results [62]. MSP have been used in two studies in this review, which are also two of the oldest studies (Sanchez-Cespedes et al. in 2000 and Nakahara et al. in 2005).

Of more recent methods, Yokoi et al. named the quantitative methylation specific PCR (qMSP) as the “gold standard” of methylation analyses in a review from 2017 [63]. qMSP has been used in another two of the included studies [34, 35]. The group with Schröck and De Vos used triplex quantitative PCR (qPCR) and Wong et al. used real time PCR [36].

There are several challenges in the detection and quantification of cfDNA. The amount of circulating cfDNA from tumor is generally very limited and have a short half-life. Therefore, the methods used are particularly susceptible to timing and workflow, and optimal standardized methods are needed in order to obtain a high performance.

All the included studies used targeted assays covering only a few numbers of known promoters. Recent technological progress has enabled whole genome methylation analyses e.g. Methylated DNA immunoprecipitation sequencing (MeDIPseq), whole-genome bisulfite sequencing or the Infinium MethylationEpic Array [64, 65]. Each method has its advantages but common for all are the genome wide coverage compared to targeted assays.

Using comprehensive integrative molecular analyses of 10.000 specimens of primary tumors across 33 types of cancer in The Cancer Genome Atlas, Hoadley et al. [66] showed that the influence of cell type was evident in DNA-methylation based clustering. Using MeDIP-seq, Shen et al. [67] demonstrated detection of early-stage pancreatic cancer through cfDNA in plasma but also the potential of multi-cancer classification using the same protocol in a cohort of 189 plasma samples from 7 different tumor types. Early-stage detection across >50 different cancers, including HNSCC by analyzing methylation signatures in cfDNA with advanced computational biology has also recently been described by Liu et al. [68].

Ideally, each subtype of cancer presents a unique, detectable cfDNA signature and a genome-wide analysis with improved bioinformatics would be able to detect the cancer present at an early stage and the cell or tissue of origin.

## Methods used in bisulfite conversion

Despite using different PCR-methods, all the included studies use bisulfite treatment of purified DNA. Efficient bisulfite conversion of DNA prior to PCR or any other methylation analysis is an important step. Incomplete conversion may result in overestimation of methylation. Furthermore, bisulfite treatment may result in degradation and loss of DNA, especially when using samples of less than 200 ng DNA. In these cases, more than 95% of the bisulfite-treated DNA is lost, if using standard procedures [69]. However high recovery can be achieved through a rapid bisulfite-treatment protocol [70]. The challenge regarding efficient bisulfite conversion has also been addressed by Ørntoft and colleagues [71]. They compared 12 different kits for bisulfite conversion of circulating cfDNA and found that mean recovery ranged between 9 and 32% and a bisulfite conversion efficiency of 97–99.9%. Based on their findings, they recommend that an integrated bisulfite conversion efficiency control should be integrated in the protocol of studies examining methylation markers in cfDNA.

None of the included studies in this review had a description of DNA recovery and bisulfite conversion efficiency in their methods, nor mentioned the bisulfite treatment as a crucial step in their discussion.

## Conclusion

None of the examined genes found in this literature review are applicable as single diagnostic markers for HNSCC, and the best gene-panels still lack diagnostic accuracy. All the included studies evaluated predefined candidate genes based on earlier studies, and thus, future studies evaluating diagnostic methylation-markers in HNSCC may benefit from the use of whole genome sequencing, new microarray technology, and improved bioinformatics algorithms. Methylation levels seem to correlate with tumor size, stage, and nodal involvement so the diagnostic challenge of HNSCC may remain in small tumors with none- or uncharacteristic symptoms, but new panels with refined computational biology may be able to overcome this challenge.

Future evaluation of cfDNA methylation in peripheral blood could have a diagnostic as well as prognostic value and could be used for post-treatment surveillance. This would potentially improve the survival rate and reduce the treatment related morbidity in these patients by enabling early detection and less invasive procedures in the post-treatment follow-up care.

In conclusion, further studies are needed that ideally include thorough documentation of laboratory methods used and genome-wide analyses of single anatomical sites. This also includes analysis of correlation between methylation in tumor and peripheral blood, the HPV status, matched cases and controls, and sample size calculations. Altogether, these aspects may contribute to an improvement of HNSCC treatment regarding survival and the overall quality of life for these patients.

## Supporting information

**S1 Checklist. PRISMA 2009 checklist.**

(DOC)

**S1 Table. Search algorithms in PubMed and OVID EMBASE.**

(DOCX)

## Acknowledgments

The authors would like to thank Pernille Gaarsted, research librarian at the Medical Library, Aalborg University Hospital for her valuable time and help to structure the literature search.

## Author Contributions

**Conceptualization:** Christian Sander Danstrup, Inge Søkilde Pedersen, Karen Dybkær, Michael Gaihede.

**Formal analysis:** Christian Sander Danstrup, Mette Marcussen.

**Investigation:** Christian Sander Danstrup, Mette Marcussen.

**Methodology:** Mette Marcussen, Henrik Jacobsen, Michael Gaihede.

**Project administration:** Christian Sander Danstrup.

**Resources:** Henrik Jacobsen.

**Supervision:** Mette Marcussen, Inge Søkilde Pedersen, Henrik Jacobsen, Karen Dybkær, Michael Gaihede.

**Writing – original draft:** Christian Sander Danstrup.

**Writing – review & editing:** Mette Marcussen, Inge Søkilde Pedersen, Henrik Jacobsen, Karen Dybkær, Michael Gaihede.

## References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin.* 2020; 70: 7–30. <https://doi.org/10.3322/caac.21590> PMID: 31912902
2. Gatta G, Botta L, Sánchez MJ, Anderson LA, Pierannunzio D, Licitra L, et al. Prognoses and improvement for head and neck cancers diagnosed in Europe in early 2000s: The EURO CARE-5 population-based study. *Eur J Cancer.* 2015; 51: 2130–2143. <https://doi.org/10.1016/j.ejca.2015.07.043> PMID: 26421817
3. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018; 68: 394–424. <https://doi.org/10.3322/caac.21492> PMID: 30207593
4. Braakhuis BJM, Leemans CR, Visser O. Incidence and survival trends of head and neck squamous cell carcinoma in the Netherlands between 1989 and 2011. *Oral Oncol.* 2014; 50: 670–675. <https://doi.org/10.1016/j.oraloncology.2014.03.008> PMID: 24735546
5. Pulte D, Brenner H. Changes in Survival in Head and Neck Cancers in the Late 20th and Early 21st Century: A Period Analysis. *Oncologist.* 2010; 15: 994–1001. <https://doi.org/10.1634/theoncologist.2009-0289> PMID: 20798198
6. Jakobsen KK, Grønhoj C, Jensen DH, Karnov KKS, Agander TK, Specht L, et al. Increasing incidence and survival of head and neck cancers in Denmark: a nation-wide study from 1980 to 2014. *Acta Oncol (Madr).* 2018; 57: 1143–1151. <https://doi.org/10.1080/0284186X.2018.1438657> PMID: 29447088
7. Amin MB, Edge SB, Greene FL. *AJCC Cancer Staging Manual.* 8th. ed. New York: Springer; 2017.
8. Howlader N, Noone A, Krapcho M, Miller D, Brest A, Yu M, et al. SEER Cancer Statistics. SEER Cancer Stat Rev 1975–2017. 2020. Available: [https://seer.cancer.gov/csr/1975\\_2017/sections.html](https://seer.cancer.gov/csr/1975_2017/sections.html)
9. Thompson-Harvey A, Yetukuri M, Hansen AR, Simpson MC, Adjei Boakye E, Varvares MA, et al. Rising incidence of late-stage head and neck cancer in the United States. *Cancer.* 2020; 126: 1090–1101. <https://doi.org/10.1002/cncr.32583> PMID: 31722124
10. Janz TA, Graboyes EM, Nguyen SA, Ellis MA, Neskey DM, Harruff EE, et al. A Comparison of the NCDB and SEER Database for Research Involving Head and Neck Cancer. *Otolaryngol—Head Neck Surg (United States).* 2019; 160: 284–294. <https://doi.org/10.1177/0194599818792205> PMID: 30129822
11. Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Susan S, et al. Smoking and Drinking in Relation to Oral and Pharyngeal Cancer. *Cancer Res.* 1988; 48: 3282–3287. PMID: 3365707
12. Lassen P. The role of Human papillomavirus in head and neck cancer and the impact on radiotherapy outcome. *Radiother Oncol.* 2010; 95: 371–380. <https://doi.org/10.1016/j.radonc.2010.04.022> PMID: 20493569
13. Fakhry C, Westra WH, Li S, Cmelak A, Ridge JA, Pinto H, et al. Improved survival of patients with human papillomavirus-positive head and neck squamous cell carcinoma in a prospective clinical trial. *J Natl Cancer Inst.* 2008; 100: 261–9. <https://doi.org/10.1093/jnci/djn011> PMID: 18270337
14. Mulero-Navarro S, Esteller M. Epigenetic biomarkers for human cancer: The time is now. *Critical Reviews in Oncology/Hematology. Crit Rev Oncol Hematol;* 2008. pp. 1–11. <https://doi.org/10.1016/j.critrevonc.2008.03.001> PMID: 18430583
15. Bird AP. CpG-Rich islands and the function of DNA methylation. *Nature.* 1986; 321: 209–213. <https://doi.org/10.1038/321209a0> PMID: 2423876
16. Antequera F, Bird A. Number of CpG islands and genes in human and mouse. *Proc Natl Acad Sci U S A.* 1993; 90: 11995–11999. <https://doi.org/10.1073/pnas.90.24.11995> PMID: 7505451
17. Bestor TH, Edwards JR, Boulard M. Notes on the role of dynamic DNA methylation in mammalian development. *Proc Natl Acad Sci U S A.* 2015; 112: 6796–6799. <https://doi.org/10.1073/pnas.1415301111> PMID: 25368180
18. Hasegawa M, Nelson HH, Peters E, Ringstrom E, Posner M, Kelsey KT. Patterns of gene promoter methylation in squamous cell cancer of the head and neck. *Oncogene.* 2002; 21: 4231–4236. <https://doi.org/10.1038/sj.onc.1205528> PMID: 12082610
19. Lu YF, Yu JR, Yang Z, Zhu GX, Gao P, Wang H, et al. Promoter hypomethylation mediated upregulation of MicroRNA-10b-3p targets FOXO3 to promote the progression of esophageal squamous cell carcinoma (ESCC) 11 Medical and Health Sciences 1112 Oncology and Carcinogenesis. *J Exp Clin Cancer Res.* 2018;37. <https://doi.org/10.1186/s13046-018-0689-3> PMID: 29475441

20. Sun W, Liu Y, Glazer CA, Shao C, Bhan S, Demokan S, et al. TKTL1 is activated by promoter hypomethylation and contributes to head and neck squamous cell carcinoma carcinogenesis through increased aerobic glycolysis and HIF1 $\alpha$  stabilization. *Clin Cancer Res*. 2010; 16: 857–866. <https://doi.org/10.1158/1078-0432.CCR-09-2604> PMID: 20103683
21. Basu B, Chakraborty J, Chandra A, Katarkar A, Baldevbhai JRK, Dhar Chowdhury D, et al. Genome-wide DNA methylation profile identified a unique set of differentially methylated immune genes in oral squamous cell carcinoma patients in India. *Clin Epigenetics*. 2017; 9: 13. <https://doi.org/10.1186/s13148-017-0314-x> PMID: 28174608
22. Jahr S, Hentze H, Englisch S, Hardt D, Fackelmayer FO, Hesch RD, et al. DNA fragments in the blood plasma of cancer patients: Quantitations and evidence for their origin from apoptotic and necrotic cells. *Cancer Res*. 2001; 61: 1659–1665. PMID: 11245480
23. Diehl F, Li M, Dressman D, He Y, Shen D, Szabo S, et al. Detection and quantification of mutations in the plasma of patients with colorectal tumors. *Proc Natl Acad Sci U S A*. 2005; 102: 16368–16373. <https://doi.org/10.1073/pnas.0507904102> PMID: 16258065
24. Issa IA, NouredDine M. Colorectal cancer screening: An updated review of the available options. *World Journal of Gastroenterology*. Baishideng Publishing Group Co., Limited; 2017. pp. 5086–5096. <https://doi.org/10.3748/wjg.v23.i28.5086> PMID: 28811705
25. Imperiale TF, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Lidgard GP, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med*. 2014; 370: 1287–1297. <https://doi.org/10.1056/NEJMoa1311194> PMID: 24645800
26. Weiss G, Schlegel A, Kottwitz D, König T, Tetzner R. Validation of the SHOX2/PTGER4 DNA Methylation Marker Panel for Plasma-Based Discrimination between Patients with Malignant and Nonmalignant Lung Disease. *J Thorac Oncol*. 2017; 12: 77–84. <https://doi.org/10.1016/j.jtho.2016.08.123> PMID: 27544059
27. Partin AW, Van Neste L, Klein EA, Marks LS, Gee JR, Troyer DA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol*. 2014; 192: 1081–1087. <https://doi.org/10.1016/j.juro.2014.04.013> PMID: 24747657
28. Mancini M, Righetto M, Zumerle S, Montopoli M, Zattoni F. The Bladder EpiCheck Test as a Non-Invasive Tool Based on the Identification of DNA Methylation in Bladder Cancer Cells in the Urine: A Review of Published Evidence. *Int J Mol Sci*. 2020; 21: 6542. <https://doi.org/10.3390/ijms21186542> PMID: 32911596
29. van Kessel KEM, Beukers W, Lurkin I, Ziel-van der Made A, van der Keur KA, Boormans JL, et al. Validation of a DNA Methylation-Mutation Urine Assay to Select Patients with Hematuria for Cystoscopy. *J Urol*. 2017; 197: 590–595. <https://doi.org/10.1016/j.juro.2016.09.118> PMID: 27746284
30. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*. 2009. <https://doi.org/10.1371/journal.pmed.1000097> PMID: 19621072
31. Prospero. Available: <https://www.crd.york.ac.uk/prospero/>
32. Covidence review tool. Available: <https://www.covidence.org>
33. University of Bristol. QUADAS. Available: <http://www.bristol.ac.uk/population-health-sciences/projects/quadas/>
34. Carvalho AL, Jeronimo C, Kim MM, Henrique R, Zhang Z, Hoque MO, et al. Evaluation of promoter hypermethylation detection in body fluids as a screening/diagnosis tool for head and neck squamous cell carcinoma. *Clin Cancer Res*. 2008; 14: 97–107. <https://doi.org/10.1158/1078-0432.CCR-07-0722> PMID: 18172258
35. Mydlarz WK, Hennessey PT, Wang H, Carvalho AL, Califano JA. Serum biomarkers for detection of head and neck squamous cell carcinoma. *Head Neck*. 2016; 38: 9–14. <https://doi.org/10.1002/hed.23842> PMID: 24995714
36. Wong TS, Man MWL, Lam AKY, Wei WI, Kwong YL, Yuen APW. The study of p16 and p15 gene methylation in head and neck squamous cell carcinoma and their quantitative evaluation in plasma by real-time PCR. *Eur J Cancer*. 2003; 39: 1881–1887. [https://doi.org/10.1016/s0959-8049\(03\)00428-3](https://doi.org/10.1016/s0959-8049(03)00428-3) PMID: 12932666
37. Schröck A, Leisse A, De Vos L, Gevensleben H, Dröge F, Franzen A, et al. Free-circulating methylated DNA in blood for diagnosis, staging, prognosis, and monitoring of head and neck squamous cell carcinoma patients: An observational prospective cohort study. *Clin Chem*. 2017; 63: 1288–1296. <https://doi.org/10.1373/clinchem.2016.270207> PMID: 28515105
38. de Vos L, Gevensleben H, Schröck A, Franzen A, Kristiansen G, Bootz F, et al. Comparison of quantification algorithms for circulating cell-free DNA methylation biomarkers in blood plasma from cancer patients. *Clin Epigenetics*. 2017; 9: 125. <https://doi.org/10.1186/s13148-017-0425-4> PMID: 29213339

39. Nakahara Y, Shintani S, Mihara M, Hino S, Hamakawa H. Detection of p16 promoter methylation in the serum of oral cancer patients. *Int J Oral Maxillofac Surg*. 2006; 35: 362–365. <https://doi.org/10.1016/j.ijom.2005.08.005> PMID: 16298513
40. Sanchez-Cespedes M, Esteller M, Wu L, Nawroz-Danish H, Yoo GH, Koch WM, et al. Gene promoter hypermethylation in tumors and serum of head and neck cancer patients. *Cancer Res*. 2000; 60: 892–895. PMID: 10706101
41. Feinberg AP, Vogelstein B. Hypomethylation of ras oncogenes in primary human cancers. *Biochem Biophys Res Commun*. 1983; 111: 47–54. [https://doi.org/10.1016/s0006-291x\(83\)80115-6](https://doi.org/10.1016/s0006-291x(83)80115-6) PMID: 6187346
42. Church TR, Wandell M, Lofton-Day C, Mongin SJ, Burger M, Payne SR, et al. Prospective evaluation of methylated SEPT9 in plasma for detection of asymptomatic colorectal cancer. *Gut*. 2014; 63: 317–25. <https://doi.org/10.1136/gutjnl-2012-304149> PMID: 23408352
43. Armstrong WB, Netterville JL. Anatomy of the larynx, trachea, and bronchi. *Otolaryngologic Clinics of North America*. 1995. pp. 685–699. PMID: 7478631
44. Lassen P, Primdahl H, Johansen J, Kristensen CA, Andersen E, Andersen LJ, et al. Impact of HPV-associated p16-expression on radiotherapy outcome in advanced oropharynx and non-oropharynx cancer. *Radiother Oncol*. 2014; 113: 310–316. <https://doi.org/10.1016/j.radonc.2014.11.032> PMID: 25544647
45. Westra WH. The Morphologic Profile of HPV-Related Head and Neck Squamous Carcinoma: Implications for Diagnosis, Prognosis, and Clinical Management. *Head and Neck Pathology*. *Head Neck Pathol*; 2012. pp. 48–54. <https://doi.org/10.1007/s12105-011-0288-5> PMID: 21877232
46. El-Naggar AK, Westra WH. P16 expression as a surrogate marker for HPV-related oropharyngeal carcinoma: A guide for interpretative relevance and consistency. *Head and Neck*. *Head Neck*; 2012. pp. 459–461. <https://doi.org/10.1002/hed.21974> PMID: 22180304
47. Degli Esposti D, Sklias A, Lima SC, Beghelli-de la Forest Divonne S, Cahais V, Fernandez-Jimenez N, et al. Unique DNA methylation signature in HPV-positive head and neck squamous cell carcinomas. *Genome Med*. 2017; 9: 33. <https://doi.org/10.1186/s13073-017-0419-z> PMID: 28381277
48. Sidransky D. Emerging molecular markers of cancer. *Nature Reviews Cancer*. *European Association for Cardio-Thoracic Surgery*; 2002. pp. 210–219. <https://doi.org/10.1038/nrc755> PMID: 11990857
49. Jung M, Klotzek S, Lewandowski M, Fleischhacker M, Jung K. Changes in concentration of DNA in serum and plasma during storage of blood samples. *Clin Chem*. 2003; 49: 1028–9. <https://doi.org/10.1373/49.6.1028> PMID: 12766024
50. Henriksen, T.V.; Reinert, T.; Sethi, H.; Gögenur I. The effect of surgical trauma on postoperative circulation free DNA levels in patients with colorectal cancer—implications for studies of circulation tumor DNA. *Odense, Denmark*; 2019.
51. Queimado LV, Raju R, Cheng L, Ramachandran I, Brabcova E, Naifeh M, et al. Abstract 4299: Down-regulation of Wnt inhibitory factor 1 is an early event in oropharyngeal cancer associated with tobacco-use. *Cancer Res*. 2013; 73: 4299–4299. <https://doi.org/10.1158/1538-7445.am2013-4299>
52. Chang HW, Ling GS, Wei WI, Yuen APW. Smoking and drinking can induce p15 methylation in the upper aerodigestive tract of healthy individuals and patients with head and neck squamous cell carcinoma. *Cancer*. 2004; 101: 125–132. <https://doi.org/10.1002/cncr.20323> PMID: 15221997
53. Nakayama H, Hibi K, Taguchi M, Takase T, Yamazaki T, Kasai Y, et al. Molecular detection of p16 promoter methylation in the serum of colorectal cancer patients. *Cancer Lett*. 2002; 188: 115–119. [https://doi.org/10.1016/s0304-3835\(01\)00839-4](https://doi.org/10.1016/s0304-3835(01)00839-4) PMID: 12406556
54. Ooki A, Maleki Z, Tsay JCJ, Goparaju C, Brait M, Turaga N, et al. A panel of novel detection and prognostic methylated DNA markers in primary non-small cell lung cancer and serum DNA. *Clin Cancer Res*. 2017; 23: 7141–7152. <https://doi.org/10.1158/1078-0432.CCR-17-1222> PMID: 28855354
55. Hayashi M, Wu G, Roh JL, Chang X, Li X, Ahn J, et al. Correlation of gene methylation in surgical margin imprints with locoregional recurrence in head and neck squamous cell carcinoma. *Cancer*. 2015; 121: 1957–1965. <https://doi.org/10.1002/cncr.29303> PMID: 25773145
56. Misawa K, Mochizuki D, Imai A, Endo S, Mima M, Misawa Y, et al. Prognostic value of aberrant promoter hypermethylation of tumor-related genes in early-stage head and neck cancer. *Oncotarget*. 2016; 7: 26087–26098. <https://doi.org/10.18632/oncotarget.8317> PMID: 27027429
57. Roh JL, Wang XV, Manola J, Sidransky D, Forastiere AA, Koch WM. Clinical correlates of promoter hypermethylation of four target genes in head and neck cancer: A cooperative group correlative study. *Clin Cancer Res*. 2013; 19: 2528–2540. <https://doi.org/10.1158/1078-0432.CCR-12-3047> PMID: 23444219

58. Virani S, Bellile E, Bradford CR, Carey TE, Chepeha DB, Colacino JA, et al. NDN and CD1A are novel prognostic methylation markers in patients with head and neck squamous carcinomas. *BMC Cancer*. 2015; 15: 825. <https://doi.org/10.1186/s12885-015-1806-8> PMID: 26518708
59. Zuo C, Ai L, Ratliff P, Suen JY, Hanna E, Brent TP, et al. O6-methylguanine-DNA methyltransferase gene: Epigenetic silencing and prognostic value in head and neck squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev*. 2004; 13: 967–975. PMID: 15184253
60. Symonds EL, Pedersen SK, Murray DH, Jedi M, Byrne SE, Rabbitt P, et al. Circulating tumour DNA for monitoring colorectal cancer—a prospective cohort study to assess relationship to tissue methylation, cancer characteristics and surgical resection. *Clin Epigenetics*. 2018; 10. <https://doi.org/10.1186/s13148-018-0500-5> PMID: 29796114
61. Herman JG, Graff JR, Myöhänen S, Nelkin BD, Baylin SB. Methylation-specific PCR: A novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci U S A*. 1996; 93: 9821–9826. <https://doi.org/10.1073/pnas.93.18.9821> PMID: 8790415
62. Kristensen LS, Hansen LL. PCR-based methods for detecting single-locus DNA methylation biomarkers in cancer diagnostics, prognostics, and response to treatment. *Clin Chem*. 2009; 55: 1471–83. <https://doi.org/10.1373/clinchem.2008.121962> PMID: 19520761
63. Yokoi K, Yamashita K, Watanabe M. Analysis of DNA methylation status in bodily fluids for early detection of cancer. *International Journal of Molecular Sciences*. MDPI AG; 2017. <https://doi.org/10.3390/ijms18040735> PMID: 28358330
64. Vidal E, Sayols S, Moran S, Guillaumet-Adkins A, Schroeder MP, Royo R, et al. A DNA methylation map of human cancer at single base-pair resolution. *Oncogene*. 2017; 36: 5648–5657. <https://doi.org/10.1038/onc.2017.176> PMID: 28581523
65. Clark C, Palta P, Joyce CJ, Scott C, Grundberg E, Deloukas P, et al. A Comparison of the Whole Genome Approach of MeDIP-Seq to the Targeted Approach of the Infinium HumanMethylation450 BeadChip® for Methylome Profiling. *PLoS One*. 2012; 7. <https://doi.org/10.1371/journal.pone.0050233> PMID: 23209683
66. Hoadley KA, Yau C, Hinoue T, Wolf DM, Lazar AJ, Drill E, et al. Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. *Cell*. 2018; 173: 291–304.e6. <https://doi.org/10.1016/j.cell.2018.03.022> PMID: 29625048
67. Shen SY, Singhania R, Fehringer G, Chakravarthy A, Roehrl MHA, Chadwick D, et al. Sensitive tumour detection and classification using plasma cell-free DNA methylomes. *Nature*. 2018; 563: 579–583. <https://doi.org/10.1038/s41586-018-0703-0> PMID: 30429608
68. Liu MC, Oxnard GR, Klein EA, Swanton C, Seiden M V., Liu MC, et al. Sensitive and specific multi-cancer detection and localization using methylation signatures in cell-free DNA. *Ann Oncol*. 2020; 31: 745–759. <https://doi.org/10.1016/j.annonc.2020.02.011>
69. Tanaka K, Okamoto A. Degradation of DNA by bisulfite treatment. *Bioorganic Med Chem Lett*. 2007; 17: 1912–1915. <https://doi.org/10.1016/j.bmcl.2007.01.040> PMID: 17276678
70. Pedersen IS, Krarup HB, Thorlacius-Ussing O, Madsen PH. High recovery of cell-free methylated DNA based on a rapid bisulfite-treatment protocol. *BMC Mol Biol*. 2012; 13: 12. <https://doi.org/10.1186/1471-2199-13-12> PMID: 22448717
71. Worm Ørntoft MB, Jensen SØ, Hansen TB, Bramsen JB, Andersen CL. Comparative analysis of 12 different kits for bisulfite conversion of circulating cell-free DNA. *Epigenetics*. 2017; 12: 626–636. <https://doi.org/10.1080/15592294.2017.1334024> PMID: 28557629