

The Psychosocial Work Environment and Perceived Stress among Seniors with Physically Demanding Jobs

The SeniorWorkingLife Study

Vinstrup, Jonas; Meng, Annette; Sundstrup, Emil; Andersen, Lars L.

Published in:
International Journal of Environmental Research and Public Health

DOI (link to publication from Publisher):
[10.3390/ijerph18147437](https://doi.org/10.3390/ijerph18147437)

Creative Commons License
CC BY 4.0

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Vinstrup, J., Meng, A., Sundstrup, E., & Andersen, L. L. (2021). The Psychosocial Work Environment and Perceived Stress among Seniors with Physically Demanding Jobs: The SeniorWorkingLife Study. *International Journal of Environmental Research and Public Health*, 18(14), Article 7437.
<https://doi.org/10.3390/ijerph18147437>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.



Article

The Psychosocial Work Environment and Perceived Stress among Seniors with Physically Demanding Jobs: The SeniorWorkingLife Study

Jonas Vinstrup ^{1,*} , Annette Meng ¹ , Emil Sundstrup ¹ and Lars L. Andersen ^{1,2}

¹ National Research Centre for the Working Environment, 2100 Copenhagen, Denmark; ame@nfa.dk (A.M.); ESU@nfa.dk (E.S.); LLA@nfa.dk (L.L.A.)

² Sport Sciences, Department of Health Science and Technology, Aalborg University, 9220 Aalborg, Denmark

* Correspondence: jov@nfa.dk; Tel.: +45-3916-5200; Fax: +45-3916-5201

Abstract: Background: Poor psychosocial work conditions are known to foster negative health consequences. While the existing literature on this topic focus mainly on white-collar workers, the influence of different aspects of the psychosocial work environment in physically demanding jobs remain understudied. Likewise, senior workers represent a population of the workforce at increased risk of adverse health outcomes and premature exit from the labour market. This study investigates the association between psychosocial work factors and perceived stress among the senior work force. Methods: Utilizing cross-sectional findings, this study reports associations between psychosocial factors (organizational justice, cooperation and collegial support, decision latitude, clarity of tasks, and quality of leadership) and the outcome of perceived stress quantified by Cohen's Perceived Stress Scale (CPSS). Currently employed senior workers with physically demanding jobs were included in the analyses (n = 3386). Associations were modeled using general linear models with weights to make the estimates representative. Results: For all individually adjusted psychosocial variables, the category of "good" was consistently associated with lower stress scores compared to the categories of both "moderate" and "poor" (all $p < 0.0001$). Likewise, in the mutually adjusted analysis, the category of "good" was statistically different from "poor" for all included variables, while the category of "moderate" remained different from "poor" for "clarity of tasks", "cooperation and collegial support", and "decision latitude". Conclusions: Among senior workers with physically demanding jobs, poor ratings of organizational factors related to the psychosocial work environment are consistently associated with high stress scores. Blue-collar occupations focusing primarily on physical risk factors are recommended to increase awareness on psychosocial aspects that may be relevant to the local work environment.



Citation: Vinstrup, J.; Meng, A.; Sundstrup, E.; Andersen, L.L. The Psychosocial Work Environment and Perceived Stress among Seniors with Physically Demanding Jobs: The SeniorWorkingLife Study. *Int. J. Environ. Res. Public Health* **2021**, *18*, 7437. <https://doi.org/10.3390/ijerph18147437>

Academic Editor: Raphael M. Herr

Received: 10 May 2021

Accepted: 8 July 2021

Published: 12 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: psychological stress; workload; physical exertion; blue-collar workers; Cohen

1. Introduction

Since the first scientific article with the term "psychosocial work environment" embedded in the title was published in 1982 [1], this field of research has experienced exponential growth within academia and public health alike [2]. Likewise, the importance of psychosocial aspects within the local work environment has increasingly been recognized in the literature, giving rise to the notion that the physical demands and conditions are far from the be-all-end-all of a long, healthy, and productive work-life [3–7].

Among middle-aged workers, factors related to the psychosocial work environment have been shown to predict work exit by age 50. Specifically, low decision latitude, poor collegial support, and job insecurity double the risk of early work exit [8], whereas favorable psychosocial working conditions seem to significantly reduce this risk and may even facilitate working beyond pension age [9]. Recent results from a 20-year follow-up analysis of the Whitehall II study showed that high skill discretion and work-related social support

effectively diminish the risk of premature exit from the labor market [10]. Likewise, the inherent quality of a good psychosocial work environment is not only essential in retaining workers but has also shown to be of vital importance for the employee returning from sick leave. In short, even when adjusting for health complaints as well as current and expected future work ability, decision latitude and job promotion opportunities continue to predict return to work [11].

However, in the attempt to quantify the deleterious consequences of a poor psychosocial work environment, it is inherently difficult to separate these from the effects of stress. Alas, it is likely that psychosocial stress serves as a potent intermediate between a poor work environment and negative health outcomes [3,5,12–16]. For example, subjective ratings of the work environment have been found to correlate with biomarkers of stress [15], and a recent systematic review found psychosocial work stressors to be associated with an increased risk of all-cause mortality among the general working population [3]. Thus, differentiating between potential predictors and—perhaps more importantly—identifying specific aspects of the work environment associated with psychosocial stress seem of utmost importance and would enable targeted health strategies at the workplace. Additionally, while the majority of stress-related research has focused on white-collar workers, it is likely that the psychosocial work environment of their blue-collared counterparts have been disproportionately neglected [17]. Lastly, in addition to being more vulnerable to work-related adverse health outcomes [18], senior workers constitute a population of the workforce experiencing increased risk of premature exit from the labour market, of which namely seniors with physically demanding work have less opportunities in the workplace for supporting a prolonged work life [19].

Therefore, the present study sought to investigate associations between individual aspects of the psychosocial work environment and stress among seniors with physically demanding jobs. Since these job groups traditionally focus mainly on levels of physical exposure during work, the aim of this study was to emphasize aspects of the psychosocial work environment related to perceived stress.

2. Methods

2.1. Study Design and Population

Utilizing a cross-sectional study design, this study reports associations between occupational psychosocial factors and perceived stress. A large-scale questionnaire survey was dispatched in July 2018 to 30,000 Danes above the age of 50 (18,000 employed, 7000 unemployed, 3000 on voluntary early retirement, 2000 on disability pension), drawn as a probability sample at Statistics Denmark, and merged with national registers through individual social security numbers [20]. Publications from this research program can be found in PubMed using the term “SeniorWorkingLife”.

Initiated in 2018, the project investigates push and stay mechanisms among older workers and aspires to repeat the survey every 2–3 years. In the present analyses, 12,173 currently employed senior workers replied to the question about physical work demands (four categories from “sedentary” to “physically strenuous”). For the present article, we included only the two (self-reported) categories related to physically demanding jobs (i.e., “primarily walking/standing work with a lot of lifting or carrying” or “primarily physically strenuous heavy/fast work”) ($n = 3566$). Finally, we excluded those with depression, leaving a total sample size of 3386. The 10 largest job groups, in descending order, were: (1) nurse’s aides, (2) machine operators, (3) construction workers, (4) cleaners, (5) farmers and gardeners, (6) bricklayers and plumbers, (7) welders, (8) building and cleaning supervisors, (9) civil engineering labourers, and (10) carpenters and woodworkers.

The current study followed the STROBE recommendations for the reporting of cross-sectional studies [21] (ClinicalTrials Identifier: NCT03634410).

2.2. Predictors

Inspired by the Copenhagen Psychosocial Questionnaire (COPSOQ) [22] and adapted from the 2018 round of the Danish Work Environment Cohort Study (DWECS) [23], the present analyses include 12 individual items as predictors (Appendix A). These were rated on a 5-point Likert-scale (ranging from “never” to “always”) and subsequently merged into five categories related to different aspects of the psychosocial work environment. These were comprised of (1) organizational justice, (2) cooperation and collegial support, (3) decision latitude, (4) clarity of tasks, and (5) quality of leadership. For example, ratings to the questions “how often do you influence *how* you solve your work tasks?” and “how often do you influence *when* you solve your work tasks?”, represented in category 3, were averaged and converted into a 0–100 scale (i.e., poor; 0–50, moderate; 50–75, and good; 75–100). The weighted prevalence of each category is shown in Table 1. As different psychosocial factors are often related, we checked for multicollinearity ($r > 0.70$). The weighted Pearson correlation coefficient between the different factors ranged between 0.32–0.64, and therefore none of the factors were excluded from the mutually adjusted analysis.

Table 1. Demographics and weighted prevalence of work-related psychosocial factors.

	n	Weighted Percentage (%)	Weighted Mean (SD)
Age (mean and 95% CI)	3386		56.6 (5.3)
BMI (mean and 95% CI)	3279		26.6 (5.5)
Females	1371	43.6	
Smoking (yes)	769	22.0	
Organizational justice			
Poor (0–50)	1362	42.0	
Moderate (50–75)	1340	40.6	
Good (75–100)	652	17.4	
Cooperation and collegial support			
Poor (0–50)	466	13.6	
Moderate (50–75)	1418	43.7	
Good (75–100)	1476	42.7	
Decision latitude			
Poor (0–50)	729	21.6	
Moderate (50–75)	1276	37.4	
Good (75–100)	1362	41.0	
Clarity of tasks			
Poor (0–50)	508	14.6	
Moderate (50–75)	1892	56.8	
Good (75–100)	948	28.6	
Quality of leadership			
Poor (0–50)	1188	36.8	
Moderate (50–75)	1235	36.2	
Good (75–100)	922	27.0	
Level of physical activity during leisure time			
Sedentary	608	18.2	
Light intensity exercise > 4 h/week	2008	61.3	
Moderate intensity exercise > 4 h/week	646	18.9	
High intensity exercise several times/week	46	1.6	

Values are presented as absolute numbers (n), weighted means with standard deviations (SDs), and weighted percentages (%).

2.3. Outcome

Cohen’s Perceived Stress Scale (CPSS-10; scale 0–40) was used as a continuous outcome measure representing psychosocial stress; i.e., used interchangeably with “perceived stress” as per the consensus that psychological, sociological, and environmental factors all contribute to the feeling of stress [24,25]. Consisting of 10 questions, with each item rated on a 5-point Likert scale ranging from “never” to “almost always”, the scale shows

satisfactory validity, reliability, and internal consistency [26–28]. Likewise, the Danish consensus version of CPSS, used in the present study, exhibits cross-cultural adaptation and good psychometric properties [26].

2.4. Statistics

Associations were modelled using general linear models (Proc Glm, SAS) with CPSS-10 as the outcome (continuous variable, 0–40). The included psychosocial factors constituted the predictor variables. In model 1, each factor was entered separately (i.e., not mutually adjusted). Furthermore, this model was controlled for sex (from register), age (from register), highest attained education (from register), the type of physical work demands (from questionnaire), as well as lifestyle factors (from questionnaire: leisure time, physical activity, smoking habits, and body mass index). These covariates were chosen due to their potential (bi-directional) influence on psychosocial stress [29–32]. In model 2, all the psychosocial factors were entered simultaneously (i.e., mutually adjusted) and adjusted for the same control variables as model 1. In all analyses, model-assisted weights were applied to produce representative estimates. The weights were based on high-quality national registers at Statistics Denmark and took into account sex, age, occupational industry, highest completed education, family income, type, and origin [20]. Results were reported as least square means and 95% confidence intervals, as well as differences of least-square means and 95% confidence intervals.

3. Results

The current sample includes 3386 senior workers with physically demanding jobs, with Table 1 depicting demographics and weighted prevalence of the five included psychosocial variables related to the work environment.

Table 2 shows individually adjusted psychosocial variables and accompanying categories of “good” (reference), “moderate”, and “poor”. For all five psychosocial variables, the categories of “moderate” and “poor” were strongly associated with increased stress scores (all $p < 0.0001$). For example, for “organizational justice”, the categories of “good”, “moderate”, and “poor” corresponded to CPSS scores of 10.7 (95% CI 9.0–12.4), 12.2 (95% CI 10.5–13.9), and 14.4 (95% CI 12.8–16.1), respectively, with a similar pattern emerging from the remaining four variables.

Table 2. Model 1. Individually adjusted psychosocial variables and stress scores (0–40).

Psychosocial Variable	Stress Score		Difference between Means (95% CI)	p-Value
	Mean	95% CI		
Organizational justice				
Good	10.7	9.0–12.4		
Moderate	12.2	10.5–13.9	−1.5 (−2.0, −0.9)	<0.0001
Poor	14.4	12.8–16.1	−3.7 (−4.3, −3.2)	<0.0001
Cooperation and collegial support				
Good	11.2	9.5–12.9		
Moderate	13.4	11.7–15.1	−2.2 (−2.6, −1.9)	<0.0001
Poor	15.5	13.8–17.2	−4.3 (−4.9, −3.7)	<0.0001
Decision latitude				
Good	11.5	9.8–13.1		
Moderate	13.2	11.5–14.9	−1.7 (−2.2, −1.3)	<0.0001
Poor	15.2	13.5–16.9	−3.7 (−4.2, −3.2)	<0.0001
Clarity of tasks				
Good	10.2	8.5–11.9		
Moderate	12.7	11.1–14.4	−2.5 (−2.9, −2.1)	<0.0001
Poor	15.9	14.2–17.6	−5.7 (−6.3, −5.1)	<0.0001
Quality of leadership				
Good	10.8	9.1–12.5		
Moderate	12.4	10.7–14.1	−1.7 (−2.1, −1.2)	<0.0001
Poor	14.5	12.9–16.2	−3.8 (−4.3, −3.3)	<0.0001

Values are presented as means with 95% confidence intervals. Adjusted for age, sex, smoking, BMI, education, and level of physical activity during leisure time and work.

Likewise, Table 3 shows mutually adjusted associations between the included psychosocial factors and stress scores: With the exception of the category of “moderate” for “organizational justice” and “quality of leadership” ($p = 0.23$ and $p = 0.12$, respectively), the categories of both “moderate” and “poor” remained associated with increases in stress scores for “cooperation and collegial support”, “decision latitude”, and “clarity of tasks” (all $p \leq 0.02$).

Table 3. Model 2. Mutually adjusted psychosocial variables and stress scores (0–40).

Stress Score				
Psychosocial Variable	Mean	95% CI	Difference between Means (95% CI)	p-Value
Organizational justice				
Good	12.6	11.0–14.3		
Moderate	13.0	11.4–14.6	−0.3 (−0.9, 0.2)	0.23
Poor	13.4	11.8–15.0	−0.8 (−1.4, −0.1)	0.02
Cooperation and collegial support				
Good	12.1	10.5–13.8		
Moderate	13.0	11.3–14.6	−0.8 (−1.2, −0.4)	0.0003
Poor	13.9	12.2–15.5	−1.7 (−2.4, −1.1)	<0.0001
Decision latitude				
Good	12.3	10.7–14.0		
Moderate	12.9	11.3–14.5	−0.6 (−1.0, −0.1)	0.01
Poor	13.8	12.1–15.4	−1.4 (−2.0, −0.9)	<0.0001
Clarity of tasks				
Good	11.2	9.6–12.9		
Moderate	12.9	11.2–14.5	−1.6 (−2.1, −1.2)	<0.0001
Poor	14.9	13.2–16.5	−3.7 (−4.3, −3.0)	<0.0001
Quality of leadership				
Good	12.5	10.9–14.2		
Moderate	12.9	11.3–14.5	−0.4 (−0.9, 0.1)	0.12
Poor	13.5	11.9–15.2	−1.0 (−1.6, −0.4)	0.0008

Values are presented as means with 95% confidence intervals. Adjusted for age, sex, smoking, BMI, education, and level of physical activity during leisure time and work.

4. Discussion

This study reports associations between ratings of pre-defined aspects of the psychosocial work environment and stress scores, showing clear differences in both the individually and mutually adjusted analyses. Of note, the category of “clarity of tasks” shows the largest differences, exhibiting mean CPSS scores of 11.2, 12.9, and 14.9 for the categories of “good”, “moderate”, and “poor”, respectively, while the category of “cooperation and collegial support” exhibits scores of 12.1, 13.0, and 13.9, respectively. While the directionality cannot be established with certainty, these results highlight a potential mediating effect of stress on health-related outcomes commonly attributed directly to the psychosocial work environment, as well as the importance of differentiating between its inherent components.

Of note, the present analyses were performed on senior workers with physically demanding jobs. Among this population of the workforce, the majority of research is customarily done in relation to the aspects of the physical work environment [33–35]. While this common differentiation between job groups and their (assumed) differences in work-related stressors likely stems from outdated societal notions, this study infers that psychosocial stress is also a highly relevant topic among workers with physically demanding jobs. In fact, when comparing occupations based on the International Standard Classification of Occupations (ISCO), we have recently shown that workers with physically demanding jobs not only exhibit increased odds of musculoskeletal pain compared with more sedentary jobs, they are also likely to experience at least the same degree of psychosocial stress [36]. Therefore, while the inherent components of demanding physical work—likely to foster negative health and productivity outcomes [33,37,38]—are not to be

neglected, a growing body of evidence emphasizes the importance of including several aspects of the psychosocial work environment when informing policies [3,6,39]. In light of this, it is becoming increasingly clear that a broad array of very different organizational-, lifestyle-, and health-related factors relate to and indeed influence work ability [37,39–41]. Interestingly, while musculoskeletal pain—alone and in combination with high physical job demands—seems to be a strong predictor of poor work ability among the general working population [42] and senior workers alike [43,44], a range of psychosocial factors have been shown to modify these associations [39,45]. For example, in a nationally representative sample of middle-aged Finnish workers experiencing multi-site pain, 37% and 48% of men and women, respectively, report “good” (≥ 9 , 0–10 scale) work ability [39]. Following this, by utilizing data from the same cohort in a 7-year follow-up, Haukka et al. identify several work and lifestyle factors that influence the outcome of long-term sickness absence, ranging from “possibility to adjust workday length” to “no problems with working community or mental stress” [45]. Therefore, these and other recent results indicate that several organizational psychosocial factors serve as both predictors and protective determinants of work ability [46,47].

Summarily, while it is perspicuous that a plurality of physical and psychosocial risk factors exists within the work environment, the ability of the latter to modify the former and hereby directly influence work ability even among pain-ridden workers needs to be emphasized. The present study provides additional insight into the prevalence of psychosocial stress among workers with physically demanding jobs, highlighting the importance of deferring the prevailing focus on physical risk factors among this group of workers.

4.1. Perspectives

Considering the job demands–resources model by Demerouti and Bakker—an overarching framework with application across a wide range of occupational settings [48–51]—work-related stress is characterized by a mismatch between the inherent work demands and the adaptive resources of the individual, ultimately resulting in the neuroplastic, physiological, and behavioral changes associated with prolonged psychosocial stress [52,53]. This notion of allostatic (over)load and consequential homeostatic disruption echoes through all aspects of research on human stress syndromes, highlighting the fact that the brain does not discriminate between origins of various stressors [53]. However, creating impactful changes in the local work environment in the attempt to decrease the prevalence of stress is not always feasible. Specifically, among senior workers with physically demanding jobs, it is likely that intrinsic components of the work limit the potential for change and that some aspects of the psychosocial work environment are therefore easier to attune than others. For example, while individual “decision latitude” might be difficult to influence among most blue-collar jobs with predominantly predetermined tasks, a more suitable target for improvement may, for example, be “cooperation and collegial support”, as it relates less to the specific task at hand.

Therefore, while it is clear that individual demands, resources, and coping behavior modulate the stress response, local working environment policies would benefit from effectively identifying both positive and negative characteristics inherent to the occupational setting that are modifiable and may influence the well-being of the worker [54].

4.2. Strengths and Limitations

Limitations include the inherent risks associated with questionnaire surveys, including recall, non-response, and, most notably, common-method bias [55–57]. In effect, the directionality of the presented associations cannot, per the cross-sectional design of the study, be established, as it is likely that workers experiencing stress are more prone to rate aspects of their psychosocial work environment as being poor. However, likely in a bi-directional manner, the associations between perceived stress and work-related psychosocial factors remain convincing.

While based on self-reporting, a noteworthy strength includes the use of the Danish version of CPSS-10 for assessing psychosocial stress [26,27], and by utilizing a probability sample merged with national registers through social security numbers, it is likely that the presented results adequately represent the population of Danish senior workers.

5. Conclusions

Poor ratings of the psychosocial work environment are associated with higher stress scores among senior workers with physically demanding jobs. Blue-collar occupations focusing primarily on physical risk factors are recommended to adapt current policies to emphasize the importance of psychosocial risk factors relevant to the local work environment. Specifically, highlighting aspects related to collegial support and clarity of tasks may especially be important among this group of workers. Finally, these results signify the importance of recognizing perceived stress as a potential mediator of the negative health consequences attributed to the psychosocial work environment.

Author Contributions: Conceptualization, E.S. and L.L.A.; data curation, L.L.A.; formal analysis, J.V., A.M., E.S. and L.L.A.; funding acquisition, L.L.A.; methodology, J.V., A.M. and E.S.; project administration, J.V.; supervision, A.M. and L.L.A.; validation, J.V., A.M. and E.S.; visualization, J.V. and A.M.; writing—original draft, J.V.; writing—review and editing, A.M., E.S. and L.L.A. All authors have read and agreed to the published version of the manuscript.

Funding: Author L.L.A. obtained a grant from Trygffonden, Denmark.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: According to Danish law, questionnaire and register-based studies do not need approval by ethical and scientific committees, nor informed consent. All data has been de-identified by Statistics Denmark, will remain on the server of Statistics Denmark, and will be analyzed through remote access by the researchers.

Data Availability Statement: The authors encourage collaboration and use of data by other researchers. Data are stored on the server of Statistics Denmark, and researchers interested in using the data for scientific purposes should contact project leader Prof. Lars L. Andersen, lla@nfa.dk.

Acknowledgments: The authors thank the participants for engaging in the survey, as well as the researchers from the National Research Centre for the Working Environment, Aalborg University, Team Arbejdsliv, Statistics Denmark, and members of the Advisory Board for contributing to the design of the study.

Conflicts of Interest: There are no conflict of interest.

Appendix A. Individual Psychosocial Variables and Their Groupings

	Individual Questions	Category
1	How often are employees, who are affected by a given decision, heard?	Organizational justice
2	How often are all employees treated fairly at the workplace?	
3	How often do you and your colleagues help each other achieving the best possible result?	Cooperation and collegial support
4	At your workplace, how often are considerations taken towards employees with less energy (e.g., elderly or sick)?	
5	How often do you and your colleagues work together when problems arise that require solutions?	
6	How often do you have a say in how you complete your work tasks?	Decision latitude
7	How often do you have a say in when you complete your work tasks?	

	Individual Questions	Category
8	How often do you receive the information, guidance and instructions you need in order to do your job?	Clarity of tasks
9	How often do you know exactly what your work tasks are?	
10	How often are you exposed to conflicting demands at work?	
11	How often is your work recognized and valued by management?	Quality of leadership
12	How often do you receive the help and support you need from your immediate manager?	

References

1. Alfredsson, L.; Karasek, R.; Theorell, T. Myocardial infarction risk and psychosocial work environment: An analysis of the male Swedish working force. *Soc. Sci. Med.* **1982**, *16*, 463–467. [\[CrossRef\]](#)
2. Rugulies, R. What is a psychosocial work environment? *Scand. J. Work. Environ. Health* **2018**, *45*, 1–6. [\[CrossRef\]](#)
3. Taouk, Y.; Spittal, M.J.; Lamontagne, A.D.; Milner, A.J. Psychosocial work stressors and risk of all-cause and coronary heart disease mortality: A systematic review and meta-analysis. *Scand. J. Work. Environ. Health* **2019**, *46*, 19–31. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Heraclides, A.; Chandola, T.; Witte, D.R.; Brunner, E. Psychosocial Stress at Work Doubles the Risk of Type 2 Diabetes in Middle-Aged Women: Evidence from the Whitehall II Study. *Diabetes Care* **2009**, *32*, 2230–2235. [\[CrossRef\]](#)
5. Jood, K.; Karlsson, N.; Medin, J.; Pessah-Rasmussen, H.; Wester, P.; Ekberg, K.; K, J.; N, K.; J, M.; H, P.-R.; et al. The psychosocial work environment is associated with risk of stroke at working age. *Scand. J. Work. Environ. Health* **2017**, *43*, 367–374. [\[CrossRef\]](#)
6. Aronsson, G.; Theorell, T.; Grape, T.; Hammarström, A.; Hogstedt, C.; Marteinsdottir, I.; Skoog, I.; Träskman-Bendz, L.; Hall, C. A systematic review including meta-analysis of work environment and burnout symptoms. *BMC Public Health* **2017**, *17*, 264. [\[CrossRef\]](#)
7. Thorsen, S.V.; Jensen, P.H.; Bjørner, J.B. Psychosocial work environment and retirement age: A prospective study of 1876 senior employees. *Int. Arch. Occup. Environ. Health* **2016**, *89*, 891–900. [\[CrossRef\]](#)
8. Stansfeld, S.A.; Carr, E.; Smuk, M.; Clark, C.; Murray, E.; Shelton, N.; Head, J. Mid-life psychosocial work environment as a predictor of work exit by age 50. *PLoS ONE* **2018**, *13*, e0195495. [\[CrossRef\]](#) [\[PubMed\]](#)
9. Andersen, L.L.; Thorsen, S.V.; Larsen, M.; Sundstrup, E.; Boot, C.R.; Rugulies, R. Work factors facilitating working beyond state pension age: Prospective cohort study with register follow-up. *Scand. J. Work. Environ. Health* **2021**, *47*, 15–21. [\[CrossRef\]](#)
10. Fleischmann, M.; Carr, E.; A Stansfeld, S.; Xue, B.; Head, J. Can favourable psychosocial working conditions in midlife moderate the risk of work exit for chronically ill workers? A 20-year follow-up of the Whitehall II study. *Occup. Environ. Med.* **2017**, *75*, 183–190. [\[CrossRef\]](#) [\[PubMed\]](#)
11. Lau, B.; Shiryayeva, O.; Ruud, T.; Victor, M. What are they returning to? Psychosocial work environment as a predictor of returning to work among employees in treatment for common mental disorders: A prospective observational pre–post study. *PLoS ONE* **2019**, *14*, e0215354. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Harris, M.L.; Oldmeadow, C.; Hure, A.; Luu, J.; Loxton, D.; Attia, J. Stress increases the risk of type 2 diabetes onset in women: A 12-year longitudinal study using causal modelling. *PLoS ONE* **2017**, *12*, e0172126. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Arguelles, L.M.; Afari, N.; Buchwald, D.S.; Clauw, D.J.; Furner, S.; Goldberg, J. A twin study of posttraumatic stress disorder symptoms and chronic widespread pain. *Pain* **2006**, *124*, 150–157. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Vinstrup, J.; Jakobsen, M.D.; Andersen, L.L. Perceived Stress and Low-Back Pain Among Healthcare Workers: A Multi-Center Prospective Cohort Study. *Front. Public Health* **2020**, *8*, 297. [\[CrossRef\]](#)
15. Arnetz, J.; Sudan, S.; Goetz, C.; Counts, S.; Arnetz, B. Nurse Work Environment and Stress Biomarkers: Possible Implications for Patient Outcomes. *J. Occup. Environ. Med.* **2019**, *61*, 676–681. [\[CrossRef\]](#)
16. Vinstrup, J.; Jakobsen, M.D.; Calatayud, J.; Jay, K.; Andersen, L.L. Association of Stress and Musculoskeletal Pain With Poor Sleep: Cross-Sectional Study Among 3,600 Hospital Workers. *Front. Neurol.* **2018**, *9*, 968. [\[CrossRef\]](#)
17. Dédèlè, A.; Miškinytė, A.; Andrušaitytė, S.; Bartkutė, Ž. Perceived Stress among Different Occupational Groups and the Interaction with Sedentary Behaviour. *Int. J. Environ. Res. Public Health* **2019**, *16*, 4595. [\[CrossRef\]](#)
18. Jones, M.K.; Latreille, P.L.; Sloane, P.J.; Staneva, A.V. Work-related health risks in Europe: Are older workers more vulnerable? *Soc. Sci. Med.* **2013**, *88*, 18–29. [\[CrossRef\]](#) [\[PubMed\]](#)
19. Andersen, L.L.; Jensen, P.H.; Meng, A.; Sundstrup, E. Strong Labour Market Inequality of Opportunities at the Workplace for Supporting a Long and Healthy Work-Life: The SeniorWorkingLife Study. *Int. J. Environ. Res. Public Health* **2019**, *16*, 3264. [\[CrossRef\]](#)
20. Andersen, L.L.; Sundstrup, E. Study protocol for SeniorWorkingLife - push and stay mechanisms for labour market participation among older workers. *BMC Public Health* **2019**, *19*, 133. [\[CrossRef\]](#)

21. Von Elm, E.; Altman, D.G.; Egger, M.; Pocock, S.J.; Gøtzsche, P.C.; Vandenbroucke, J.P.; STROBE Initiative. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *J. Clin. Epidemiol.* **2008**, *61*, 344–349. [\[CrossRef\]](#)
22. Kristensen, T.S.; Hannerz, H.; Høgh, A.; Borg, V. The Copenhagen Psychosocial Questionnaire—A tool for the assessment and improvement of the psychosocial work environment. *Scand. J. Work. Environ. Health* **2005**, *31*, 438–449. [\[CrossRef\]](#) [\[PubMed\]](#)
23. National Research Centre for the Working Environment. Arbejdsmiljø og Helbred i Danmark 2018. 2018. Available online: <http://nfa.dk/-/media/NFA/Arbejdsmiljodata/Notat-om-OBmaal-AH2018.ashx?la=da> (accessed on 9 May 2021).
24. Cohen, S.; Gianaros, P.J.; Manuck, S.B. A Stage Model of Stress and Disease. *Perspect. Psychol. Sci.* **2016**, *11*, 456–463. [\[CrossRef\]](#)
25. Thoits, P.A. Stress and Health: Major Findings and Policy Implications. *J. Health Soc. Behav.* **2010**, *51*, S41–S53. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Eskildsen, A.; Dalgaard, V.L.; Nielsen, K.J.; Andersen, J.H.; Zachariae, R.; Olsen, L.R.; Järgensen, A.; Christiansen, D.H. Cross-cultural adaptation and validation of the Danish consensus version of the 10-item Perceived Stress Scale. *Scand. J. Work. Environ. Health* **2015**, *41*, 486–490. [\[CrossRef\]](#)
27. Cohen, S.; Kamarck, T.; Mermelstein, R. A global measure of perceived stress. *J. Health Soc. Behav.* **1983**, *24*, 385–396. [\[CrossRef\]](#)
28. Chiu, Y.-H.; Lu, F.J.-H.; Lin, J.-H.; Nien, C.-L.; Hsu, Y.-W.; Liu, H.-Y. Psychometric properties of the Perceived Stress Scale (PSS): Measurement invariance between athletes and non-athletes and construct validity. *PeerJ* **2016**, *4*, e2790. [\[CrossRef\]](#)
29. Cohen, S.; Lichtenstein, E. Perceived stress, quitting smoking, and smoking relapse. *Health Psychol.* **1990**, *9*, 466–478. [\[CrossRef\]](#)
30. Schultchen, D.; Reichenberger, J.; Mittl, T.; Weh, T.R.M.; Smyth, J.M.; Blechert, J.; Pollatos, O. Bidirectional relationship of stress and affect with physical activity and healthy eating. *Br. J. Health Psychol.* **2019**, *24*, 315–333. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Meule, A.; Reichenberger, J.; Blechert, J. Smoking, Stress Eating, and Body Weight: The Moderating Role of Perceived Stress. *Subst. Use Misuse* **2018**, *53*, 2152–2156. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Tajik, E.; Latiff, L.A.; Adznam, S.N.; Awang, H.; Siew, C.Y.; Abu Bakar, A.S. A study on level of physical activity, depression, anxiety and stress symptoms among adolescents. *J. Sports Med. Phys. Fit.* **2016**, *57*, 1382–1387.
33. Andersen, L.L.; Fallentin, N.; Thorsen, S.V.; Holtermann, A. Physical workload and risk of long-term sickness absence in the general working population and among blue-collar workers: Prospective cohort study with register follow-up. *Occup. Environ. Med.* **2016**, *73*, 246–253. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Villumsen, M.; Samani, A.; Jørgensen, M.B.; Gupta, N.; Madeleine, P.; Holtermann, A. Are forward bending of the trunk and low back pain associated among Danish blue-collar workers? A cross-sectional field study based on objective measures. *Ergonomics* **2014**, *58*, 246–258. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Bláfoss, R.; Sundstrup, E.; Jakobsen, M.D.; Brandt, M.; Bay, H.; Andersen, L.L. Physical workload and bodily fatigue after work: Cross-sectional study among 5000 workers. *Eur. J. Public Health* **2019**, *29*, 837–842. [\[CrossRef\]](#)
36. Vinstrup, J.; Sundstrup, E.; Andersen, L.L. Psychosocial stress and musculoskeletal pain among senior workers from nine occupational groups: Cross-sectional findings from the SeniorWorkingLife study. *BMJ Open* **2021**, *11*, e043520. [\[CrossRef\]](#)
37. Skovlund, S.V.; Bláfoss, R.; Sundstrup, E.; Andersen, L.L. Association between physical work demands and work ability in workers with musculoskeletal pain: Cross-sectional study. *BMC Musculoskelet. Disord.* **2020**, *21*, 166–168. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Vinstrup, J.; Jakobsen, M.D.; Madeleine, P.; Andersen, L.L. Physical Exposure during Patient Transfer and Risk of Back Injury & Low-Back Pain: Prospective Cohort Study. *BMC Musculoskelet. Disord.* **2020**, *21*, 715. [\[CrossRef\]](#)
39. Pensola, T.; Haukka, E.; Kaila-Kangas, L.; Neupane, S.; Leino-Arjas, P. Good work ability despite multisite musculoskeletal pain? A study among occupationally active Finns. *Scand. J. Public Health* **2016**, *44*, 300–310. [\[CrossRef\]](#)
40. Ezzatvar, Y.; Calatayud, J.; Andersen, L.L.; Vinstrup, J.; Alarcón, J.; Casaña, J. Dose-response association between multi-site musculoskeletal pain and work ability in physical therapists: A cross-sectional study. *Int. Arch. Occup. Environ. Health* **2020**, *93*, 863–870. [\[CrossRef\]](#) [\[PubMed\]](#)
41. Van den Berg, T.; Elders, L.; de Zwart, B.; Burdorf, A. The effects of work-related and individual factors on the Work Ability Index: A systematic review. *Occup. Environ. Med.* **2008**, *66*, 211–220. [\[CrossRef\]](#) [\[PubMed\]](#)
42. Bayattork, M.; Jakobsen, M.D.; Sundstrup, E.; Seidi, F.; Bay, H.; Andersen, L.L. Musculoskeletal pain in multiple body sites and work ability in the general working population: Cross-sectional study among 10,000 wage earners. *Scand. J. Pain* **2019**, *19*, 131–137. [\[CrossRef\]](#)
43. Skovlund, S.V.; Bláfoss, R.; Sundstrup, E.; Thomassen, K.; Andersen, L.L. Joint association of physical work demands and leg pain intensity for work limitations due to pain in senior workers: Cross-sectional study. *BMC Public Health* **2020**, *20*, 1741. [\[CrossRef\]](#)
44. Wilkie, R.; Blagojevic-Bucknall, M.; Jordan, K.P.; Pransky, G. Onset of Work Restriction in Employed Adults with Lower Limb Joint Pain: Individual Factors and Area-Level Socioeconomic Conditions. *J. Occup. Rehabil.* **2013**, *23*, 180–188. [\[CrossRef\]](#)
45. Haukka, E.; Ojajarvi, A.; Kaila-Kangas, L.; Leino-Arjas, P. Protective determinants of sickness absence among employees with multisite pain—A 7-year follow-up. *Pain* **2017**, *158*, 220–229. [\[CrossRef\]](#)
46. Saastamoinen, P.; Leino-Arjas, P.; Rahkonen, O.; Lahelma, E. Separate and combined associations of pain and emotional exhaustion with sickness absence. *Pain* **2016**, *157*, 186–193. [\[CrossRef\]](#)
47. Aronsson, V.; Toivanen, S.; Leineweber, C.; Nyberg, A. Can a poor psychosocial work environment and insufficient organizational resources explain the higher risk of ill-health and sickness absence in human service occupations? Evidence from a Swedish national cohort. *Scand. J. Public Health* **2019**, *47*, 310–317. [\[CrossRef\]](#)

-
48. Lesener, T.; Gusy, B.; Wolter, C. The job demands-resources model: A meta-analytic review of longitudinal studies. *Work Stress* **2019**, *33*, 76–103. [[CrossRef](#)]
 49. Van den Broeck, A.; Elst, T.V.; Baillien, E.; Sercu, M.; Schouteden, M.; De Witte, H.; Godderis, L. Job Demands, Job Resources, Burnout, Work Engagement, and Their Relationships: An Analysis Across Sectors. *J. Occup. Environ. Med.* **2017**, *59*, 369–376. [[CrossRef](#)]
 50. Bakker, A.; Demerouti, E. The Job Demands-Resources model: State of the art. *J. Manag. Psychol.* **2007**, *22*, 309–328. [[CrossRef](#)]
 51. Demerouti, E.; Bakker, A.B.; Nachreiner, F.; Schaufeli, W.B. The job demands-resources model of burnout. *J. Appl. Psychol.* **2001**, *86*, 499–512. [[CrossRef](#)]
 52. Bains, J.S.; Cusulin, J.I.W.; Inoue, W. Stress-related synaptic plasticity in the hypothalamus. *Nat. Rev. Neurosci.* **2015**, *16*, 377–388. [[CrossRef](#)] [[PubMed](#)]
 53. McEwen, B.S. Physiology and Neurobiology of Stress and Adaptation: Central Role of the Brain. *Physiol. Rev.* **2007**, *87*, 873–904. [[CrossRef](#)] [[PubMed](#)]
 54. Wadsworth, E.J.K.; Chaplin, K.S.; Smith, A.P. The work environment, stress and well-being. *Occup. Med.* **2010**, *60*, 635–639. [[CrossRef](#)]
 55. Cheung, K.L.; Klooster, P.M.T.; Smit, C.; De Vries, H.; Pieterse, M.E. The impact of non-response bias due to sampling in public health studies: A comparison of voluntary versus mandatory recruitment in a Dutch national survey on adolescent health. *BMC Public Health* **2017**, *17*, 276. [[CrossRef](#)]
 56. Choi, B.C.; Pak, A.W. A catalog of biases in questionnaires. *Prev. Chronic Dis.* **2004**, *2*, A13. [[PubMed](#)]
 57. Podsakoff, P.M.; MacKenzie, S.B.; Lee, J.Y.; Podsakoff, N.P. Common method biases in behavioral research: A critical review of the literature and recommended remedies. *J. Appl. Psychol.* **2003**, *88*, 879–903. [[CrossRef](#)] [[PubMed](#)]