

The Potential Application of Commercially Available Active Video Games to Cardiac Rehabilitation

Scoping Review

Sawa, Ryuichi; Saitoh, Masakazu; Morisawa, Tomoyuki; Takahashi, Tetsuya; Morimoto, Yuh; Kagiya, Nobuyuki; Kasai, Takatoshi; Dinesen, Birthe; Daida, Hiroyuki

Published in:
JMIR serious games

DOI (link to publication from Publisher):
[10.2196/31974](https://doi.org/10.2196/31974)

Creative Commons License
CC BY 4.0

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Sawa, R., Saitoh, M., Morisawa, T., Takahashi, T., Morimoto, Y., Kagiya, N., Kasai, T., Dinesen, B., & Daida, H. (2022). The Potential Application of Commercially Available Active Video Games to Cardiac Rehabilitation: Scoping Review. *JMIR serious games*, 10(1), Article e31974. <https://doi.org/10.2196/31974>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Review

The Potential Application of Commercially Available Active Video Games to Cardiac Rehabilitation: Scoping Review

Ryuichi Sawa¹, PT, MSc, PhD; Masakazu Saitoh¹, PT, MSc, PhD; Tomoyuki Morisawa¹, PT, MSc, PhD; Tetsuya Takahashi^{1,2}, PT, MSc, PhD; Yuh Morimoto³, PhD; Nobuyuki Kagiya^{2,4}, MD, PhD; Takatoshi Kasai^{2,4}, MD, PhD; Birthe Dinesen⁵, MSc, PhD; Hiroyuki Daida^{1,2,4}, MD, PhD

¹Department of Physical Therapy, Faculty of Health Science, Juntendo University, Tokyo, Japan

²Department of Digital Health and Telemedicine Research and Development, Faculty of Health Science, Juntendo University, Tokyo, Japan

³Faculty of Health Science, Juntendo University, Tokyo, Japan

⁴Department of Cardiovascular Biology and Medicine, Juntendo University Graduate School of Medicine, Tokyo, Japan

⁵Laboratory for Welfare Technologies - Telehealth & Telerehabilitation, Sport Sciences - Performance and Technology, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

Corresponding Author:

Hiroyuki Daida, MD, PhD

Department of Physical Therapy

Faculty of Health Science

Juntendo University

3-2-12 Hongo, Bunkyo-ku

Tokyo, 113-0033

Japan

Phone: 81 3 3813 3111

Email: daida@juntendo.ac.jp

Abstract

Background: Commercially available active video games (AVGs) have recently been used for rehabilitation in some specific patient populations but rarely in those with cardiovascular disease (CVD). Commercially available AVGs are designed to increase motivation for continuous play, which could be applicable to the long-term cardiac rehabilitation process.

Objective: The objective of this scoping review was to assess the effectiveness of AVG-induced physical exercise, safety management, and patient adherence by applying commercially available AVGs to cardiac rehabilitation.

Methods: Four databases (CINAHL, MEDLINE, PubMed, and SPORTDiscus) were searched for all years up to August 12, 2020. Articles were retained if they were written in English, included patients with CVD who were aged 18 years or older, and used AVGs as part of a physical exercise program. The included studies were then evaluated from the viewpoints of effectiveness as physical exercise, safety, and adherence management.

Results: Among 120 nonduplicate articles reviewed, 5 (4.2%) were eligible for inclusion, of which 3 (2.5%) were reported by the same research group. The AVG consoles used were Xbox Kinect and Nintendo Wii, and sports-related programs were adopted for the intervention. No adverse cardiac events occurred in the identified studies, and dropout rates tended to be low.

Conclusions: AVGs appear to be safe and feasible for promoting an active lifestyle in patients with CVD. However, the effectiveness of AVGs alone as a therapeutic exercise to improve physical function may be limited.

(*JMIR Serious Games* 2022;10(1):e31974) doi: [10.2196/31974](https://doi.org/10.2196/31974)

KEYWORDS

active video game; cardiac rehabilitation; physical exercise; rehabilitation; serious games; CVD; AVG; cardiovascular disease; exercise; safety; adherence

Introduction

Cardiovascular disease (CVD) is the most prevalent noncommunicable disease and the leading cause of death globally [1]. Despite the proven benefits for the improvement in mortality from cardiac rehabilitation and strong guideline recommendations [2-4], a variety of barriers that reduce adherence to cardiac rehabilitation remain, especially after hospital discharge [5-7]. In particular, barriers related to accessibility are expected to become the major barrier with progressive population aging. Symptoms associated with CVD lead to an excessively sedentary lifestyle, and this physical inactivity reinforces the tendency to be trapped in an age-related downward spiral toward frailty and disability [8,9]. Therefore, approaches that interrupt or help prevent falling into this spiral are urgently needed in cardiac rehabilitation.


Active video games (AVGs), which are games that require players to interact with objects within a virtual environment using some part of their body as the controller [10], have recently emerged because of the evolution of motion-sensing devices that are designed to detect and measure movements. During the past decade, video game companies have launched several commercially available AVG consoles (Figure 1). Owing to their appealing designs, AVGs can increase motivation and long-term engagement with physical exercise in the general population, including older adults [10-13]. A systematic review

has revealed the effectiveness of Nintendo Wii in the rehabilitation of adults with stroke [14]. However, only 6 studies were included in the review, suggesting that the evidence is still in the construction stage.

Therapeutic exercise is a core component in cardiac rehabilitation and aims to improve physical function such as exercise capacity and muscle strength through adequate exercise prescriptions [2]. On the other hand, because of the long-term rehabilitation process, practice guidelines consistently recommend comprehensive rehabilitation programs involving multiple components (eg, health education, advice on CVD risk reduction, physical activity, stress management) [3,4,15]. Furthermore, because of the aging population, it has become more important to provide multicomponent exercise programs as well as to increase patient motivation to engage in physical exercise. A previous systematic review reported that no studies integrated commercially available AVGs with multicomponent physical rehabilitation programs for motor function in patients with CVD [16].

Given this background, the objective of this scoping review was to explore the possibility of applying commercially available AVGs to cardiac rehabilitation. With this in mind, we assessed the available evidence in the literature for the physical effects of using AVGs in patients with CVD to examine ways to manage the safety of and adherence to such interventions.

Figure 1. Categories and characteristics of commercially available active video game consoles. LED: light-emitting diode; N/A: not available; RGB: red, green, blue; VR: virtual reality.



Methods

Design

We conducted a scoping review in accordance with the guidelines described by PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews) [17].

Search Strategy

A literature search of the electronic databases CINAHL, MEDLINE, PubMed, and SPORTDiscus was conducted for all years up to August 2020. The literature search was conducted using search terms covering medical and health science terminology (Multimedia Appendix 1).

Selection Process

Following the removal of duplicates, titles and abstracts were screened by at least 2 independent reviewers (TM, MS, YM, and RS) to exclude irrelevant articles as the first screening step. Articles were included if they focused on the influence of commercially available AVGs on patients with CVD aged 18 years or older. A commercially available AVG was defined as a digital game that video game companies launched into the entertainment market and that have controllers mounting motion-sensing devices for detecting bodily movements as signals. The inclusion criteria were articles that reported original quantitative data including case studies, studies with participants aged 18 years or older, studies that included participants with CVD, and studies that used a commercially available AVG as an intervention tool. Articles were ineligible if they were topic

news or an editorial, pictorial, perspective, letter, conference paper, or review. Those that reported qualitative data were also ineligible for this review. The exclusion criteria were studies that used commercially unavailable AVGs (research-based AVGs) and articles published in a language other than English. In the second screening step, the articles included from the first screening step were downloaded and screened again by the same independent reviewers using the same criteria as that used in the first screening. Disagreements arising between the reviewers at any stage of the study selection process were resolved through team discussions or by a third reviewer (TT) to reach consensus on whether the article met the inclusion criteria. Reference lists of relevant articles were also hand-searched to identify additional appropriate articles for inclusion in this review. Review articles were not included in this study, but were checked for potentially relevant references.

Data Extraction


Using a structured sheet in Excel (Microsoft), data were extracted by 1 reviewer (RS) and subsequently checked for accuracy by the other authors. The extracted data included details about the authors, year of publication, study design, location of the study, aims of the study, sample characteristics,

AVG console used in the intervention, AVG programs used in the intervention, details of the AVG intervention (eg, period, frequency, duration, intensity), and the outcome measures related to physical exercise. Physical exercise was defined as bodily movements to enhance or maintain physical activity. Additional data of interest included the study setting, supervision, delivery method for the AVG interventions (AVG alone or AVG with other programs), effectiveness of the AVG in terms of physical exercise, safety management, the number of adverse events, adherence management, the number of patients who dropped out, and reasons for dropout. Safety management was defined as efforts to prevent adverse events during the intervention period. Adherence was defined as performance of the intervention as planned, and efforts to manage patient adherence were extracted as adherence management. These data were grouped and arranged in Excel by the type of setting and the presence of supervision to report the study outcomes.

Results

A flowchart of the systematic screening process with the number of articles included or excluded at each stage is shown in [Figure 2](#).

Figure 2. A flow diagram of studies included according to PRISMA-ScR (Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews). AVG: active video game; CVD: cardiovascular disease.



From the initial 223 identified articles, 5 (4.2%) were retrieved and screened for eligibility. In terms of study design, 3 of the 5 (60%) articles reported randomized studies. The remaining 2/5 (40%) articles were a case study and a pre-post study. The details of the identified articles are shown in Table 1, and a summary of the intervention delivery modes and effectiveness is shown in Table 2. Table 3 shows the summary of adverse events and adherence across studies. One study investigated the impacts of adding AVG-based physical exercise to cardiac rehabilitation programs on acute hemodynamics in 27 outpatients with CVD or CVD risk [18]. That study was a cluster-randomized crossover trial and used the Xbox Kinect as an alternative tool for warm-up and conditioning sessions in center-based cardiac rehabilitation. Similar physiological acute hemodynamic responses to cardiac rehabilitation were found, with higher magnitudes of heart rate, respiratory rate, and the rate of perceived exertion during and after cardiac rehabilitation with the AVG. Another study conducted in 32 outpatients with CVD in a hospital reported that those who had been allocated into the AVG group showed significant improvements after a 6-week intervention period in physical activity measured as accelerometer arbitrary unit (AAU; median baseline, 255 AAU/min; end of program, 322 AAU/min; $P=.04$), whereas those allocated into the conventional cardiac rehabilitation group showed no change (median baseline, 225 AAU/min; end of program, 247 AAU/min; $P=.99$). Measures of energy expenditure also improved significantly in the AVG group compared with those in the conventional cardiac rehabilitation group [19]. These 2 studies took the exercise intensity of AVGs into consideration based on physiological measures such as heart rate. On the other hand, 3 studies reported by the same research group [20-22] focused on the effects of home-based

AVGs in patients with heart failure. The research group has published the results from a case study, a pilot study [22], and an open-label randomized study [20]. A total of 605 patients from multiple countries got the same instructions about the time and frequency for playing AVGs but were given no concrete numerical target regarding exercise intensity. Although the intervention through home-based AVGs was safe and feasible, it was not effective in improving the following outcomes: the 6-minute walk test, unilateral isotonic heel-lift, bilateral isometric shoulder abduction, unilateral isotonic shoulder flexion, exercise motivation, exercise self-efficacy, and physical activity.

The effectiveness of the AVG intervention was inconsistent. Another study conducted in the home setting reported that AVGs were ineffective in improving physical function [20], whereas studies in the hospital setting clarified the effectiveness of using AVGs as a supplementary tool for center-based, conventional cardiac rehabilitation programs [18,19]. The adherence rates and occurrence of adverse events showed similar trends in the identified studies. Regarding safety management, studies in the hospital setting controlled exercise intensity by monitoring the participants' physiological changes during the intervention, whereas those in the home setting provided safety guidelines and instructions for adapting AVG play to each patient's physical condition before the intervention. Furthermore, patients could call the research staff to ask questions during a given time period. Although myalgia and osteoarthritic knee pain occurred as musculoskeletal-related events in 2 studies, no cardiac-related adverse events were reported during the interventions. To manage patient adherence, the research staff conducted motivational calls or provided telephone guidance to each patient. The dropout rate seemed to be relatively low (0%-23%).

Table 1. Summary of study characteristics.

Study	Setting	Age, composition, and disease state	Active video game	Dose and duration	Outcome
Alves da Cruz et al, 2020 [18], cluster randomized controlled trial in Brazil	Hospital	<ul style="list-style-type: none"> N=27 48% female Age: 63.4 (12.71) years Status: CVD,^a CVD risk 	<ul style="list-style-type: none"> Console: Xbox Kinect Program: Just Dance 2015 (warm-up), Shape Up (conditioning) 	<ul style="list-style-type: none"> Period: 15 min (warm-up), 30 min, (conditioning) Frequency: Once Duration: 1 day Intensity: Based on heart rate reserve and rate of perceived exertion during intervention 	<ul style="list-style-type: none"> Hemodynamics: systolic blood pressure, diastolic blood pressure, respiratory rate, oxygen saturation, heart rate, rate of perceived exertion
Jaarsma et al, 2020 [20], open-label randomized study in multiple countries	Home	<ul style="list-style-type: none"> N=605 29% female Age: 67 (12) years Status: HF^b 	<ul style="list-style-type: none"> Console: Nintendo Wii Program: Nintendo Wii Sports 	<ul style="list-style-type: none"> Period: 30 min per day Frequency: 5 days a week Duration: 12 weeks Intensity: not reported 	<ul style="list-style-type: none"> EC^c: 6MWT^d Muscle function: unilateral isotonic heel-lift, bilateral isometric shoulder abductions, unilateral isotonic shoulder flexion EM^e: Questionnaire Exercise SE^f: Questionnaire PA^g: single question
Klompstra et al, 2013 [21], case study in Sweden	Home	<ul style="list-style-type: none"> N=1 0% female Age: 74 years Status: HF 	<ul style="list-style-type: none"> Console: Nintendo Wii Program: Nintendo Wii Sports 	<ul style="list-style-type: none"> Period: 15 min Frequency: everyday Duration: 12 weeks Intensity: not reported 	<ul style="list-style-type: none"> PA: accelerometer EC: 6MWT EM: questionnaire Exercise SE: questionnaire Perceived physical effort: Borg scale
Klompstra et al, 2014 [22], pilot study in Sweden	Home	<ul style="list-style-type: none"> N=32 31% female Age: 63 (19-88) years Status: HF 	<ul style="list-style-type: none"> Console: Nintendo Wii Program: Nintendo Wii Sports 	<ul style="list-style-type: none"> Period: 20 min per day Frequency: not recorded Duration: 12 weeks Intensity: not recorded 	<ul style="list-style-type: none"> EC: 6MWT PA: accelerometer
Ruivo et al, 2017 [19], pilot randomized controlled trial in Ireland	Hospital	<ul style="list-style-type: none"> N=32 18.7% female Age: 59.9 (10.2) years Status: CVD 	<ul style="list-style-type: none"> Console: Nintendo Wii Program: Nintendo Wii Sports 	<ul style="list-style-type: none"> Period: 1-hour sessions Frequency: twice a week Duration: 6 weeks Intensity: based on heart rate from a precardiac rehabilitation test 	<ul style="list-style-type: none"> EC: Bruce ramp protocol PA: accelerometer

^aCVD: cardiovascular disease.^bHF: heart failure.^cEC: exercise capacity.^d6MWT: 6-minute walk test.^eEM: exercise motivation.^fSE: self-efficacy.^gPA: physical activity.

Table 2. Summary of intervention delivery and effectiveness across studies.

Study	Supervision	AVG ^a alone/with other programs	Effectiveness	
			Results	Conclusions
Alves da Cruz et al, 2020 [18]	SV ^b	With cardiac rehabilitation program	<ul style="list-style-type: none"> Increased heart rate Increased respiratory rate Increased rate of perceived exertion 	<ul style="list-style-type: none"> Greater heart rate, respiratory rate, and rate of perceived exertion were observed during and 5 min after the AVG session
Jaarsma et al, 2020 [20]	No SV	AVG alone	<ul style="list-style-type: none"> No change^c in exercise capacity, muscle function, exercise motivation, exercise self-efficacy, or PA^d 	<ul style="list-style-type: none"> AVG was safe and feasible in patients with heart failure Not effective in improving outcomes
Klompstra et al, 2013 [21]	No SV	AVG alone	<ul style="list-style-type: none"> Increased PA Increased exercise motivation Increased exercise self-efficacy No change in perceived physical effort 	<ul style="list-style-type: none"> Further research is needed to generalize the results from the case study
Klompstra et al, 2014 [22]	No SV	AVG alone	<ul style="list-style-type: none"> Exercise capacity No change in PA 	<ul style="list-style-type: none"> AVG has the potential to increase exercise capacity in patients with heart failure
Ruivo et al, 2017 [19]	SV	With conventional program	<ul style="list-style-type: none"> Increased PA Increased energy expenditure per body weight 	<ul style="list-style-type: none"> Cardiac rehabilitation sessions with AVG are feasible and safe Significant improvement in PA and energy expenditure

^aAVG: active video game.^bSV: supervision.^cAfter correction for baseline and confounders.^dPA: physical activity.

Table 3. Summary of adverse events and adherence across studies.

Study	Adverse events		Adherence		
	Safety management	Number of adverse events	Adherence management	Dropout rate, n (%)	Reasons for dropout
Alves da Cruz et al, 2020 [18]	<ul style="list-style-type: none"> Control exercise intensity 	<ul style="list-style-type: none"> Cardiac, 0 Musculoskeletal, 0 	NR ^a	0	NR
Jaarsma et al, 2020 [20]	<ul style="list-style-type: none"> Instruction for adapting active video games Phone consultation 	<ul style="list-style-type: none"> Cardiac, 0 Musculoskeletal, 0 	<ul style="list-style-type: none"> Motivational calls 	71 (23%)	<ul style="list-style-type: none"> Medical-related issues Practical logistic reasons Refusal to continue Loss to follow-up Death
Klompstra et al, 2013 [21]	<ul style="list-style-type: none"> Safety guideline Phone call to heart failure nurse 	<ul style="list-style-type: none"> Cardiac, 0 Musculoskeletal, 0 	<ul style="list-style-type: none"> Remote guidance Follow-up visits 	0	NR
Klompstra et al, 2014 [22]	<ul style="list-style-type: none"> Safety guideline Phone call to heart failure nurse 	<ul style="list-style-type: none"> Cardiac, 0 Musculoskeletal, 1 (myalgia) 	<ul style="list-style-type: none"> Telephone guidance 	1 (3%)	<ul style="list-style-type: none"> Moved abroad for work
Ruivo et al, 2017 [19]	<ul style="list-style-type: none"> Playing active video games within individual target heart rate zones (55%-70%) Safety guideline Supervision at all times Monitoring by telemetry 	<ul style="list-style-type: none"> Cardiac, 0 Musculoskeletal, 1 (osteoarthritic knee pain) 	<ul style="list-style-type: none"> Motivational calls 	1 (6.3%)	<ul style="list-style-type: none"> Returned to work

^aNR: not reported.

Discussion

Overall Findings

This scoping review aimed to explore the possibility of applying commercially available AVGs to cardiac rehabilitation. Only 5 articles with commercially available AVGs were identified. All identified studies involved AVG interventions with a fixed time and frequency; however, exercise intensity was ill-defined in studies performed in the home setting. AVG interventions for patients with CVD may be safe and feasible regardless of the setting. On the other hand, some challenges seem to remain for the application of commercially available AVGs as therapeutic exercise tools in cardiac rehabilitation from the perspective of the balance between effectiveness and safety management, especially in terms of controlling exercise intensity.

Effectiveness

Although 4 identified studies provided interventions involving AVGs in a given period to patients with CVD, the effectiveness of the interventions in improving physical function was inconsistent. Conventional cardiac rehabilitation programs recommend that therapeutic exercise be generally composed of a combination of 2 types of exercise: aerobic and resistance [23]. Aerobic exercise is defined as any activity that uses large

muscle groups, can be maintained continuously, and is rhythmic in nature [24], whereas resistance exercise is expected to enhance the strength of major muscle groups by contracting muscles against external resistance [24]. It is essential for AVGs to require both types of exercise during play for adaptation to therapeutic exercise in cardiac rehabilitation. Regarding the game programs, sports-related programs that cause whole body movement during play, such as baseball and golf, were used. Several studies have reported the exercise intensity achieved during play for each game program in older people and patients after chronic stroke [25-27] but not in patients with CVD. Two randomized studies used an AVG as an intervention tool, but the results were inconsistent. An international multicenter study by Jaarsma et al [20] investigated the effects of access to a home-based AVG (Nintendo Wii) in patients with heart failure on submaximal aerobic exercise capacity, as assessed by the 6-minute walk test (6MWT), and muscle function, as measured by unilateral isotonic heel-lift, bilateral isometric shoulder abduction, and unilateral isotonic shoulder flexion. The treatment effects in the 6MWT were not significant at 3 months (4.3 m; 95% CI -6.9 to 15.5), 6 months (1.8 m; 95% CI -10.3 to 14), or 12 months (6.8 m; 95% CI -7.1 to 20.7) after correcting for baseline 6MWT results and confounders. Regarding muscle function, no outcomes were significantly improved except for left heel rise at 6 months. On the other hand, another study reported significant improvement in daily

physical activity and related energy expenditure per body weight after a 6-week intervention compared with that in the control group [19]. In that study, Nintendo Wii was used as the supplementary tool for the center-based, conventional cardiac rehabilitation program. The results suggest that exercise induced by AVGs without supervision is not sufficient to improve physical function. From the perspective of exercise prescription, it is essential to clarify the exercise intensity of each game program in specific subjects. To implement more effective therapeutic exercises in home-based cardiac rehabilitation, further laboratory-based studies are needed in which exercise intensity is measured in patients with CVD while they play AVGs. Additionally, the AVG consoles used in the identified studies were Xbox Kinect and Nintendo Wii, neither of which provides external resistance to cause muscle contractions. Therefore, it might be better to use AVGs in combination with resistance training. A device that can detect muscle contractions as a signal for controlling AVGs also needs to be developed. Commercially available AVGs are designed to encourage players to engage in physical activity continuously; therefore, AVG-based exercise interventions in the home setting may be applicable to patients with CVD from the perspective of promoting safe physical activity.

Safety Management

Regarding safety management, the identified studies prepared safety guidelines prior to the interventions. Patients could call an instructor or nurse at a given time in studies conducted in the home setting. In addition, in the identified studies, no cardiac-related adverse events were reported. The dropout rates were relatively low compared with those with conventional cardiac rehabilitation [28], partially because of the appealing design of commercially available AVGs such as the Nintendo Wii and Xbox Kinect for increasing motivation and long-term engagement to play the game. The challenges associated with center-based cardiac rehabilitation programs are low participation and high premature dropout rates [28-30]. A variety of barriers that can be characterized at 3 interrelated levels—patient, provider, and health care system—have been reported in previous studies [6,31-33]. Older age, lower socioeconomic status, schedule conflicts, disinterest in attending a program, and comorbidities were included as barriers with respect to patients' characteristics. Insufficient physician knowledge about the effectiveness of cardiac rehabilitation and inappropriate referrals to cardiac rehabilitation are often described as barriers at the provider level. System-level barriers to cardiac rehabilitation include transportation problems and the limited availability of cardiac rehabilitation programs for outpatients. Potential approaches to overcome these challenges include home-based cardiac rehabilitation and cardiac telerehabilitation [34-36]. Recent systematic reviews have concluded that home- and center-based cardiac rehabilitation have similar benefits in terms of clinical events, exercise capacity, and health-related quality of life among patients after myocardial infarction or coronary revascularization or among patients with heart failure [37,38]. Furthermore, another systematic review concluded that multidisciplinary or exercise-based cardiac telerehabilitation is safe and cost-effective and can be an alternative option to center-based cardiac rehabilitation in patients with coronary

artery disease and chronic heart failure [35]. As AVG interventions seem to be safe and feasible for keeping patients with CVD active at home, home-based cardiac rehabilitation and cardiac telerehabilitation in conjunction with AVGs could help such patients achieve and maintain a more active lifestyle.

Future Perspectives

AVGs are rapidly evolving with improvements in motion-sensing technologies, and 2 such AVG consoles, Nintendo Wii and Microsoft Kinect, were used as intervention tools for patients with CVD in the identified studies. Nintendo Wii, released in 2006, uses motion sensing technology in the primary controller to enable users to control game actions through their arm gestures. Nintendo Wii was the first commercially available video game console to induce body movement in the real world to manipulate objects or play sports in a virtual world. Following the Nintendo Wii, Microsoft launched Kinect in 2010 as a line of motion-sensing input devices. Microsoft Kinect, which was introduced to replace traditional game controllers, enables users to play games by using signals from whole-body gestures. A systematic review, published in 2016 [16] on the effectiveness of commercial AVGs for physical rehabilitation of motor function reported that almost 80% of included studies ($n=126$) used Nintendo Wii as the AVG console, which was a far higher rate than the rate at which Microsoft Kinect was used. The evolution seen in the past decade in motion-sensing systems to detect human body movement has accelerated the integration of commercial AVGs into therapeutic exercise for older people [10-13], patients after stroke [14], and patients with multiple sclerosis [39]. However, the application of AVGs in patients with CVD has yet to be addressed, partially because current AVG consoles do not induce resistance exercise. There could be 2 options to break through this barrier: the use of electromyographic signals and the development of a muscle contraction-induced controller. Some video games have introduced an electromyographic system to detect muscle contractions in the research setting [40,41] but none in the commercial setting. AVG consoles incorporating an electromyographic system in combination with a motion-sensing system could accelerate the application of AVGs in therapeutic exercise. Regarding the development of controllers that induce muscle contractions, Ring Fit Adventure, launched by Nintendo in 2017, is an action role-playing game for the Nintendo Switch that consists of 2 physical components as accessories: The Leg Strap and Ring-Con. The Leg Strap affixes the original controller to the user's thigh to detect leg movement, and Ring-Con is a ring that includes a strain sensor that can detect the bending of the ring by using the original controller as a logger. These accessories enable users to engage in both aerobic and resistance exercise during gameplay. Isometric resistance exercises usually require breath-holding, which elevates blood pressure. Speech recognition systems have the potential for safety management by not requiring users to hold their breath while performing resistance exercises even remotely. However, to our knowledge, no studies have investigated exercise intensity during gameplay using Ring Fit Adventure, which could be the first step in clarifying exercise intensity for its application as a tool in safe therapeutic exercises as a part of cardiac rehabilitation. Concurrently, the feasibility

of the combined use of AVGs and other devices for monitoring muscle contractions should be investigated to enable safe therapeutic exercise in home-based cardiac rehabilitation.

Subjects in identified articles in this review spanned a wide range of generations, including older generations. With increasing age, digital divide—the gap between individuals who have access to modern information and communication technology, including digital devices, and those who lack access—should be considered. Older generations usually do not have the opportunity or knowledge to use digital devices and therefore could not utilize them properly. In an identified study investigated in the home setting [22], the instructor demonstrated the AVG play at the patients' home and also conducted a 1-hour introduction session at the hospital. Proper introduction of AVGs is essential for its application in therapeutic exercise, especially in older adults.

Limitations

Commercially available AVGs could potentially be applied to cardiac rehabilitation; however, the evidence obtained in this review should be interpreted with caution because research in this field is rapidly growing. In addition, this scoping review

excluded studies that used commercially available AVG consoles with programs designed for research, which could offer hints for the next step toward the smooth application of commercially available AVGs in cardiac rehabilitation.

Conclusions

The results of this scoping review suggest that evidence remains lacking for the application of commercially available AVGs in cardiac rehabilitation for patients with CVD in both laboratory-based and clinical studies. Commercially available AVGs may be suitable as a tool for promoting an active lifestyle in patients with CVD even in the home setting. However, commercially available AVGs should be used with caution as an alternative to therapeutic exercise. The need for remotely accessible cardiac rehabilitation programs, including therapeutic exercises, is increasing because of the aging population. Further studies are needed to investigate the appropriate frequency, intensity, and duration of time of commercially available AVGs for effective therapeutic exercises in patients with CVD. In addition, the development of devices that can detect muscle contractions when AVGs are played is recommended for the addition of resistance exercise and institution of more effective therapeutic exercises.

Conflicts of Interest

TT, NK, TK, and HD are affiliated with a department funded by Philips Japan; Asahi Kasei Corporation; Inter Reha Co, Ltd and Toho Holdings Co, Ltd based on collaborative research agreements.

Multimedia Appendix 1

Literature search terminology and number of records included from databases.

[DOCX File, 23 KB-Multimedia Appendix 1]

References

1. Cardiovascular diseases (CVDs). World Health Organization. URL: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)) [accessed 2021-03-16]
2. Dalal H, Doherty P, Taylor R. Cardiac rehabilitation. *BMJ* 2015 Sep 29;351:h5000 [FREE Full text] [doi: [10.1136/bmj.h5000](https://doi.org/10.1136/bmj.h5000)] [Medline: [26419744](https://pubmed.ncbi.nlm.nih.gov/26419744/)]
3. Wenger N. Current status of cardiac rehabilitation. *J Am Coll Cardiol* 2008 Apr 29;51(17):1619-1631 [FREE Full text] [doi: [10.1016/j.jacc.2008.01.030](https://doi.org/10.1016/j.jacc.2008.01.030)] [Medline: [18436113](https://pubmed.ncbi.nlm.nih.gov/18436113/)]
4. Ambrosetti M, Abreu A, Corrà U, Davos C, Hansen D, Frederix I, et al. Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the Secondary Prevention and Rehabilitation Section of the European Association of Preventive Cardiology. *Eur J Prev Cardiol* 2020 Apr 07;2047487320913379. [doi: [10.1177/2047487320913379](https://doi.org/10.1177/2047487320913379)] [Medline: [33611446](https://pubmed.ncbi.nlm.nih.gov/33611446/)]
5. Park L, Schopfer D, Zhang N, Shen H, Whooley M. Participation in cardiac rehabilitation among patients with heart failure. *J Card Fail* 2017 May;23(5):427-431 [FREE Full text] [doi: [10.1016/j.cardfail.2017.02.003](https://doi.org/10.1016/j.cardfail.2017.02.003)] [Medline: [28232047](https://pubmed.ncbi.nlm.nih.gov/28232047/)]
6. Neubeck L, Freedman S, Clark A, Briffa T, Bauman A, Redfern J. Participating in cardiac rehabilitation: a systematic review and meta-synthesis of qualitative data. *Eur J Prev Cardiol* 2012 Jun;19(3):494-503. [doi: [10.1177/1741826711409326](https://doi.org/10.1177/1741826711409326)] [Medline: [22779092](https://pubmed.ncbi.nlm.nih.gov/22779092/)]
7. Turk-Adawi K, Sarrafzadegan N, Grace S. Global availability of cardiac rehabilitation. *Nat Rev Cardiol* 2014 Oct;11(10):586-596 [FREE Full text] [doi: [10.1038/nrcardio.2014.98](https://doi.org/10.1038/nrcardio.2014.98)] [Medline: [25027487](https://pubmed.ncbi.nlm.nih.gov/25027487/)]
8. Marinus N, Vigorito C, Giallauria F, Haenen L, Jansegers T, Dendale P, et al. Frailty is highly prevalent in specific cardiovascular diseases and females, but significantly worsens prognosis in all affected patients: a systematic review. *Ageing Res Rev* 2021 Mar;66:101233. [doi: [10.1016/j.arr.2020.101233](https://doi.org/10.1016/j.arr.2020.101233)] [Medline: [33333322](https://pubmed.ncbi.nlm.nih.gov/33333322/)]
9. Vigorito C, Abreu A, Ambrosetti M, Belardinelli R, Corrà U, Cupples M, et al. Frailty and cardiac rehabilitation: a call to action from the EAPC Cardiac Rehabilitation Section. *Eur J Prev Cardiol* 2017 Apr;24(6):577-590. [doi: [10.1177/2047487316682579](https://doi.org/10.1177/2047487316682579)] [Medline: [27940954](https://pubmed.ncbi.nlm.nih.gov/27940954/)]

10. Howes S, Charles D, Marley J, Pedlow K, McDonough S. Gaming for health: systematic review and meta-analysis of the physical and cognitive effects of active computer gaming in older adults. *Phys Ther* 2017 Dec 01;97(12):1122-1137. [doi: [10.1093/ptj/pzx088](https://doi.org/10.1093/ptj/pzx088)] [Medline: [29077911](https://pubmed.ncbi.nlm.nih.gov/29077911/)]
11. Chao Y, Scherer Y, Montgomery C. Effects of using Nintendo Wii™ exergames in older adults: a review of the literature. *J Aging Health* 2015 Apr;27(3):379-402. [doi: [10.1177/0898264314551171](https://doi.org/10.1177/0898264314551171)] [Medline: [25245519](https://pubmed.ncbi.nlm.nih.gov/25245519/)]
12. Bleakley CM, Charles D, Porter-Armstrong A, McNeill MDJ, McDonough SM, McCormack B. Gaming for health: a systematic review of the physical and cognitive effects of interactive computer games in older adults. *J Appl Gerontol* 2015 Apr;34(3):NP166-NP189. [doi: [10.1177/0733464812470747](https://doi.org/10.1177/0733464812470747)] [Medline: [24652863](https://pubmed.ncbi.nlm.nih.gov/24652863/)]
13. Miller K, Adair B, Pearce A, Said C, Ozanne E, Morris M. Effectiveness and feasibility of virtual reality and gaming system use at home by older adults for enabling physical activity to improve health-related domains: a systematic review. *Age Ageing* 2014 Mar;43(2):188-195. [doi: [10.1093/ageing/aft194](https://doi.org/10.1093/ageing/aft194)] [Medline: [24351549](https://pubmed.ncbi.nlm.nih.gov/24351549/)]
14. Cheek G, Tan D, Low A, Hewitt J. Is Nintendo Wii an effective intervention for individuals with stroke? a systematic review and meta-analysis. *J Am Med Dir Assoc* 2015 Nov 01;16(11):923-932. [doi: [10.1016/j.jamda.2015.06.010](https://doi.org/10.1016/j.jamda.2015.06.010)] [Medline: [26253322](https://pubmed.ncbi.nlm.nih.gov/26253322/)]
15. Balady G, Williams M, Ades P, Bittner V, Comoss P, Foody J, American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology, American Heart Association Council on Cardiovascular Nursing, American Heart Association Council on Epidemiology and Prevention, American Heart Association Council on Nutrition, Physical Activity, and Metabolism, American Association of Cardiovascular and Pulmonary Rehabilitation. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation* 2007 May 22;115(20):2675-2682. [doi: [10.1161/CIRCULATIONAHA.106.180945](https://doi.org/10.1161/CIRCULATIONAHA.106.180945)] [Medline: [17513578](https://pubmed.ncbi.nlm.nih.gov/17513578/)]
16. Bonnechère B, Jansen B, Omelina L, Van Sint Jan S. The use of commercial video games in rehabilitation: a systematic review. *Int J Rehabil Res* 2016 Dec;39(4):277-290. [doi: [10.1097/MRR.0000000000000190](https://doi.org/10.1097/MRR.0000000000000190)] [Medline: [27508968](https://pubmed.ncbi.nlm.nih.gov/27508968/)]
17. Tricco A, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018 Oct 02;169(7):467-473 [FREE Full text] [doi: [10.7326/M18-0850](https://doi.org/10.7326/M18-0850)] [Medline: [30178033](https://pubmed.ncbi.nlm.nih.gov/30178033/)]
18. Alves da Cruz MM, Ricci-Vitor A, Bonini Borges GL, Fernanda da Silva P, Ribeiro F, Marques Vanderlei LC. Acute hemodynamic effects of virtual reality-based therapy in patients of cardiovascular rehabilitation: a cluster randomized crossover trial. *Arch Phys Med Rehabil* 2020 Apr;101(4):642-649. [doi: [10.1016/j.apmr.2019.12.006](https://doi.org/10.1016/j.apmr.2019.12.006)] [Medline: [31926142](https://pubmed.ncbi.nlm.nih.gov/31926142/)]
19. Ruivo J, Karim K, O'Shea R, Oliveira R, Keary L, O'Brien C, et al. In-class active video game supplementation and adherence to cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2017 Jul;37(4):274-278. [doi: [10.1097/HCR.0000000000000224](https://doi.org/10.1097/HCR.0000000000000224)] [Medline: [28350640](https://pubmed.ncbi.nlm.nih.gov/28350640/)]
20. Jaarsma T, Klompstra L, Ben Gal T, Ben Avraham B, Boyne J, Bäck M, et al. Effects of exergaming on exercise capacity in patients with heart failure: results of an international multicentre randomized controlled trial. *Eur J Heart Fail* 2021 Jan;23(1):114-124 [FREE Full text] [doi: [10.1002/ejhf.1754](https://doi.org/10.1002/ejhf.1754)] [Medline: [32167657](https://pubmed.ncbi.nlm.nih.gov/32167657/)]
21. Klompstra L, Jaarsma T, Strömberg A. An in-depth, longitudinal examination of the daily physical activity of a patient with heart failure using a Nintendo Wii at home: a case report. *J Rehabil Med* 2013 Jun;45(6):599-602 [FREE Full text] [doi: [10.2340/16501977-1151](https://doi.org/10.2340/16501977-1151)] [Medline: [23695704](https://pubmed.ncbi.nlm.nih.gov/23695704/)]
22. Klompstra L, Jaarsma T, Strömberg A. Exergaming to increase the exercise capacity and daily physical activity in heart failure patients: a pilot study. *BMC Geriatr* 2014 Nov 18;14:119 [FREE Full text] [doi: [10.1186/1471-2318-14-119](https://doi.org/10.1186/1471-2318-14-119)] [Medline: [25407612](https://pubmed.ncbi.nlm.nih.gov/25407612/)]
23. JCS Joint Working Group. Guidelines for rehabilitation in patients with cardiovascular disease (JCS 2012). *Circ J* 2014;78(8):2022-2093 [FREE Full text] [doi: [10.1253/circj.cj-66-0094](https://doi.org/10.1253/circj.cj-66-0094)] [Medline: [25047729](https://pubmed.ncbi.nlm.nih.gov/25047729/)]
24. Garber C, Blissmer B, Deschenes M, Franklin B, Lamonte M, Lee IM, American College of Sports Medicine. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011 Jul;43(7):1334-1359. [doi: [10.1249/MSS.0b013e318213febf](https://doi.org/10.1249/MSS.0b013e318213febf)] [Medline: [21694556](https://pubmed.ncbi.nlm.nih.gov/21694556/)]
25. Taylor L, Maddison R, Pfaeffli L, Rawstorn J, Gant N, Kerse N. Activity and energy expenditure in older people playing active video games. *Arch Phys Med Rehabil* 2012 Dec;93(12):2281-2286. [doi: [10.1016/j.apmr.2012.03.034](https://doi.org/10.1016/j.apmr.2012.03.034)] [Medline: [22522217](https://pubmed.ncbi.nlm.nih.gov/22522217/)]
26. Hurkmans H, Ribbers G, Streur-Kranenburg M, Stam H, van den Berg-Emons RJ. Energy expenditure in chronic stroke patients playing Wii Sports: a pilot study. *J Neuroeng Rehabil* 2011 Jul 14;8:38 [FREE Full text] [doi: [10.1186/1743-0003-8-38](https://doi.org/10.1186/1743-0003-8-38)] [Medline: [21756315](https://pubmed.ncbi.nlm.nih.gov/21756315/)]
27. Kafri M, Myslinski M, Gade V, Deutsch J. Energy expenditure and exercise intensity of interactive video gaming in individuals poststroke. *Neurorehabil Neural Repair* 2014 Jan;28(1):56-65. [doi: [10.1177/1545968313497100](https://doi.org/10.1177/1545968313497100)] [Medline: [23897906](https://pubmed.ncbi.nlm.nih.gov/23897906/)]

28. McDonall J, Botti M, Redley B, Wood B. Patient participation in a cardiac rehabilitation program. *J Cardiopulm Rehabil Prev* 2013;33(3):185-188. [doi: [10.1097/HCR.0b013e318282551a](https://doi.org/10.1097/HCR.0b013e318282551a)] [Medline: [23403913](#)]
29. Kamiya K, Yamamoto T, Tsuchihashi-Makaya M, Ikegame T, Takahashi T, Sato Y, et al. Nationwide survey of multidisciplinary care and cardiac rehabilitation for patients with heart failure in Japan - an analysis of the AMED-CHF Study. *Circ J* 2019 Jun 25;83(7):1546-1552 [FREE Full text] [doi: [10.1253/circj.CJ-19-0241](https://doi.org/10.1253/circj.CJ-19-0241)] [Medline: [31189753](#)]
30. Kotseva K, Wood D, De Bacquer D, EUROASPIRE investigators. Determinants of participation and risk factor control according to attendance in cardiac rehabilitation programmes in coronary patients in Europe: EUROASPIRE IV survey. *Eur J Prev Cardiol* 2018 Aug;25(12):1242-1251. [doi: [10.1177/2047487318781359](https://doi.org/10.1177/2047487318781359)] [Medline: [29873511](#)]
31. Murray J, Craigs C, Hill K, Honey S, House A. A systematic review of patient reported factors associated with uptake and completion of cardiovascular lifestyle behaviour change. *BMC Cardiovasc Disord* 2012 Dec 08;12:120 [FREE Full text] [doi: [10.1186/1471-2261-12-120](https://doi.org/10.1186/1471-2261-12-120)] [Medline: [23216627](#)]
32. Moradi B, Maleki M, Esmailzadeh M, Abkenar H. Physician-related factors affecting cardiac rehabilitation referral. *J Tehran Heart Cent* 2011 Nov;6(4):187-192 [FREE Full text] [Medline: [23074367](#)]
33. De Vos C, Li X, Van Vlaenderen I, Saka O, Dendale P, Eyssen M, et al. Participating or not in a cardiac rehabilitation programme: factors influencing a patient's decision. *Eur J Prev Cardiol* 2013 Apr;20(2):341-348. [doi: [10.1177/2047487312437057](https://doi.org/10.1177/2047487312437057)] [Medline: [22345682](#)]
34. Thomas R, Beatty A, Beckie T, Brewer L, Brown T, Forman D, et al. Home-based cardiac rehabilitation: a scientific statement from the American Association of Cardiovascular and Pulmonary Rehabilitation, the American Heart Association, and the American College of Cardiology. *J Am Coll Cardiol* 2019 Jul 09;74(1):133-153 [FREE Full text] [doi: [10.1016/j.jacc.2019.03.008](https://doi.org/10.1016/j.jacc.2019.03.008)] [Medline: [31097258](#)]
35. Brouwers R, van Exel HJ, van Hal JMC, Jorstad H, de Kluiver EP, Kraaijenhagen R, Committee for Cardiovascular PreventionCardiac Rehabilitation of the Netherlands Society of Cardiology. Cardiac telerehabilitation as an alternative to centre-based cardiac rehabilitation. *Neth Heart J* 2020 Sep;28(9):443-451 [FREE Full text] [doi: [10.1007/s12471-020-01432-y](https://doi.org/10.1007/s12471-020-01432-y)] [Medline: [32495296](#)]
36. Piotrowicz E, Baranowski R, Bilinska M, Stepnowska M, Piotrowska M, Wójcik A, et al. A new model of home-based telemonitored cardiac rehabilitation in patients with heart failure: effectiveness, quality of life, and adherence. *Eur J Heart Fail* 2010 Feb;12(2):164-171 [FREE Full text] [doi: [10.1093/eurjhf/hfp181](https://doi.org/10.1093/eurjhf/hfp181)] [Medline: [20042423](#)]
37. Anderson L, Sharp G, Norton R, Dalal H, Dean S, Jolly K, et al. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database Syst Rev* 2017 Jun 30;6:CD007130 [FREE Full text] [doi: [10.1002/14651858.CD007130.pub4](https://doi.org/10.1002/14651858.CD007130.pub4)] [Medline: [28665511](#)]
38. Buckingham S, Taylor R, Jolly K, Zawada A, Dean S, Cowie A, et al. Home-based versus centre-based cardiac rehabilitation: abridged Cochrane systematic review and meta-analysis. *Open Heart* 2016;3(2):e000463 [FREE Full text] [doi: [10.1136/openhrt-2016-000463](https://doi.org/10.1136/openhrt-2016-000463)] [Medline: [27738516](#)]
39. Kramer A, Dettmers C, Gruber M. Exergaming with additional postural demands improves balance and gait in patients with multiple sclerosis as much as conventional balance training and leads to high adherence to home-based balance training. *Arch Phys Med Rehabil* 2014 Oct;95(10):1803-1809. [doi: [10.1016/j.apmr.2014.04.020](https://doi.org/10.1016/j.apmr.2014.04.020)] [Medline: [24823959](#)]
40. Lyu M, Chen W, Ding X, Wang J, Pei Z, Zhang B. Development of an EMG-controlled knee exoskeleton to assist home rehabilitation in a game context. *Front Neurobot* 2019;13:67 [FREE Full text] [doi: [10.3389/fnbot.2019.00067](https://doi.org/10.3389/fnbot.2019.00067)] [Medline: [31507400](#)]
41. Ghassemi M, Triandafilou K, Barry A, Stoykov M, Roth E, Mussa-Ivaldi F, et al. Development of an EMG-controlled serious game for rehabilitation. *IEEE Trans Neural Syst Rehabil Eng* 2019 Feb;27(2):283-292 [FREE Full text] [doi: [10.1109/TNSRE.2019.2894102](https://doi.org/10.1109/TNSRE.2019.2894102)] [Medline: [30668478](#)]

Abbreviations

AAU: accelerometer arbitrary unit

AVG: active video game

CVD: cardiovascular disease

PRISMA-ScR: Preferred Reporting Items for Systematic Reviews and Meta-Analyses Extension for Scoping Reviews

6MWT: 6-minute walk test

Edited by N Zary; submitted 12.07.21; peer-reviewed by D Winchester, C Astley; comments to author 13.11.21; revised version received 04.01.22; accepted 16.01.22; published 18.03.22

Please cite as:

Sawa R, Saitoh M, Morisawa T, Takahashi T, Morimoto Y, Kagiya N, Kasai T, Dinesen B, Daida H

The Potential Application of Commercially Available Active Video Games to Cardiac Rehabilitation: Scoping Review

JMIR Serious Games 2022;10(1):e31974

URL: <https://games.jmir.org/2022/1/e31974>

doi: [10.2196/31974](https://doi.org/10.2196/31974)

PMID:

©Ryuichi Sawa, Masakazu Saitoh, Tomoyuki Morisawa, Tetsuya Takahashi, Yuh Morimoto, Nobuyuki Kagiya, Takatoshi Kasai, Birthe Dinesen, Hiroyuki Daida. Originally published in JMIR Serious Games (<https://games.jmir.org>), 18.03.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Serious Games, is properly cited. The complete bibliographic information, a link to the original publication on <https://games.jmir.org>, as well as this copyright and license information must be included.