Aalborg Universitet



Detection of altered pain facilitatory and inhibitory mechanisms in patients with knee osteoarthritis by using a simple bedside tool kit (QuantiPain)

Izumi, Masashi; Hayashi, Yoshihiro; Saito, Ryota; Oda, Shota; Petersen, Kristian Kjær; Arendt-Nielsen, Lars; Ikeuchi, Masahiko

Published in: Pain Reports

DOI (link to publication from Publisher): 10.1097/PR9.000000000000998

Creative Commons License CC BY 4.0

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA):

Izumi, M., Hayashi, Y., Saito, R., Oda, S., Petersen, K. K., Arendt-Nielsen, L., & Ikeuchi, M. (2022). Detection of altered pain facilitatory and inhibitory mechanisms in patients with knee osteoarthritis by using a simple bedside tool kit (QuantiPain). Pain Reports, 7(3), E998. Article e998. https://doi.org/10.1097/PR9.00000000000998

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
 You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: February 07, 2025







Detection of altered pain facilitatory and inhibitory mechanisms in patients with knee osteoarthritis by using a simple bedside tool kit (QuantiPain)

Masashi Izumi^{a,b}, Yoshihiro Hayashi^b, Ryota Saito^c, Shota Oda^b, Kristian Kjær Petersen^d, Lars Arendt-Nielsen^{d,e}, Masahiko Ikeuchi^{a,b}

Abstract

Purpose: Altered pain facilitatory and inhibitory mechanisms have been recognized as an important manifestation in patients with chronic pain, and quantitative sensory testing (QST) can act as a proxy for this process. We have recently developed a simple bedside QST tool kit (*QuantiPain*) for more clinical use. The purpose of this study was to investigate its test–retest reliability and to evaluate its validity compared with the laboratory-based QST protocols in patients with knee osteoarthritis (OA).

Methods: *QuantiPain* consists of 3 items: "pressure algometer" (for pressure pain thresholds [PPTs]), "pinprick" (for temporal summation of pain [TSP]), and "conditioning clamp" (for conditioned pain modulation [CPM]). In experiment-A, intrarater and interrater test–retest reliabilities were investigated in 21 young healthy subjects by using interclass correlation coefficient (ICC). In experiment-B, 40 unilateral painful patients with OA and 40 age-matched, healthy control subjects were included to compare the bedside tool kit against the computerized pressure algometry.

Results: In experiment-A, excellent to moderate intrarater and interrater reliabilities were achieved in PPT and TSP (ICC: 0.60–0.92) while the agreements of CPM were good to poor (ICC: 0.37–0.80). In experiment-B, localized and widespread decrease of PPT, facilitated TSP, and impaired CPM was found by using the bedside tool kit in patients with OA compared with controls (P < 0.05). The data were significantly correlated with the established laboratory-based tools (R = 0.281-0.848, P < 0.05).

Conclusion: *QuantiPain* demonstrated acceptable test-retest reliability and assessment validity with the sensitivity to separate patients with painful OA from controls, which has a potential to create more practical approach for quantifying altered pain mechanisms in clinical settings.

Keywords: Pain, Quantitative sensory testing, Pressure pain threshold, Temporal summation of pain, Conditioned pain modulation, Osteoarthritis

1. Introduction

Pain perception is always subjective and thereby challenging to objectively quantify; however, several biomarkers related to mechanisms, neural activity, and susceptibility offer the possibility of this quantification.⁴⁰ Quantitative sensory testing (QST) is a

psychophysical method and can act as a proxy to investigate the functional states of the somatosensory system by evaluating thresholds or responses to standardized stimuli.³⁴ Quantitative sensory testing has been received much attention with an indirect measure of "pain sensitization" or "pain de-sensitization" in the

PR9 7 (2022) e998

http://dx.doi.org/10.1097/PR9.000000000000998

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Department of Orthopaedic Surgery, Kochi Medical School, Kochi University, Nankoku, Kochi, Japan, ^b Department of Rehabilitation, Kochi Medical School Hospital, Kochi University, Nankoku, Kochi, Japan, ^c Center for Innovative and Translational Medicine, Kochi Medical School, Kochi University, Nankoku, Kochi, Japan, ^d Department of Health Science and Technology, Center for Neuroplasticity and Pain (CNAP), Department of Health Science and Technology, School of Medical Gastroenterology, Mech-Sense, Aalborg University Hospital, Aalborg, Denmark, ^e Department of Medical Gastroenterology, Mech-Sense, Aalborg University Hospital, Aalborg, Denmark

Corresponding author. Address: Department of Orthopaedic Surgery, Kochi Medical School, Kochi University, 185-1 Kohasu, Oko-cho, Nankoku-City, Kochi Pref 783-8505, Japan. Tel.: +81-88-880-2386; fax +81-88-880-2388. E-mail address: izumim@kochi-u.ac.jp (M. Izumi).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.painrpts.com).

Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of The International Association for the Study of Pain. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

peripheral and central pain pathways.^{8,21} Quantitative sensory testing modalities are categorized as static or dynamic procedures that enables to monitor sensory functions regarding different features of transmission and control of pain.⁶

Pressure pain threshold (PPT) has been a widely used static QST for assessing painful musculoskeletal pain disorders.¹⁰ Localized or regional hyperalgesia may indicate sensory dysfunction at the peripheral level, whereas widespread hyperalgesia to remote nonpainful area seems to be a proxy for abnormalities in central nervous system processing.¹⁰ Dynamic QST includes temporal summation of pain (TSP) as a phenomenon in which repeated nociceptive stimuli cause an increase in pain perception²⁴ and conditioned pain modulation (CPM) assessing the endogenous pain modulatory pathways by the "pain inhibits pain" phenomenon.⁴⁹

Altered pain facilitatory and inhibitory mechanisms assessed by QST have been recognized as an important manifestation in patients with chronic pain.³ In addition, QST has been used to predict chronic postoperative pain^{15,28,47} and response to analgesic effects,^{30,48} and a recent review demonstrated that the most frequent reported predictors were PPTs, TSP, and CPM.³² Among multiple chronic pain conditions, osteoarthritis (OA) is one of the most intensively studied diseases, and accumulating evidence suggests that localized and widespread decrease of PPT, facilitated TSP, and impaired CPM are key contributors to chronic pain in subgroups of patients with OA.^{1,4,22,23,39}

Nevertheless, laboratory-based QST tools were complicated, expensive, training necessary, and time consuming,^{17,33} and hence, it has not become a popular pain assessment for routine clinical applications. To solve this problem, we have recently developed a simple bedside tool kit (*QuantiPain*) for evaluating PPT, TSP, and CPM. This study mainly evaluated its (1) test–retest reliability in healthy subjects and (2) validity compared with the laboratory-based QST protocols in patients with knee OA. In addition, we investigated the relationship between the bedside QST data and chronic pain–associated questionnaires that has been receiving attention for contributing pain mechanisms in patients with OA.^{41,51}

2. Materials and methods

This study comprised 2 experiments. First, test–retest reliability of *QuantiPain* assessment was investigated in young healthy subjects (experiment-A). Second, validity of *QuantiPain* assessment was evaluated in healthy subjects and patients with knee OA compared with laboratory-based QST tools (experiment-B).

The study protocol was approved by the Institutional Review Board of Kochi Medical School (No. 31–61). All participants received verbal explanation of this study and provided written informed consent before the investigation. This study was conducted in compliance with the tenets of the Declaration of Helsinki.

2.1. Participants

In experiment-A, 21 healthy, pain-free subjects (10 females, age 27 years [20–38]) were participated. In experiment-B, 40 patients with knee OA (32 females, age 67 years [49–84 years]) suffering for at least 3 months from unilateral knee pain while walking, and age-matched 40 healthy control subjects (17 female, age 61 years [27–84 years]) were recruited. Patients with bilateral knee OA were included only if 1 knee was pain free. In both experiments, subjects and patients who were diagnosed as

having painful musculoskeletal disorders (except knee OA), neurological disorders, psychiatric diseases, skin disorders at examination sites, and taking pain killers 24 hours before the experiment were excluded.

2.2. Experiment-A

QuantiPain consists of 3 items: "pressure algometer" (for pressure pain thresholds, PPTs), "pinprick" (for TSP), and "conditioning clamp" (for CPM) (**Fig. 1A**). The pressure algometer is made by just mounting a 1-cm² plastic probe on the tip of a hand-held analogue mechanical force gauge (available for measurement from 0 to 100 N, IMADA Co, LTD, Aichi, Japan). The custom pinprick (Takei Scientific Instrument Co, LTD, Niigata, Japan) incorporates a 60 g mobile weight that provides identical repeated painful stimuli through the conical tip (3 mm in maximal diameter). The conditioning clamp applies extrasegmental tonic pain stimulus (4.5 kg force applied to approximately 175 mm²). Durability of the conditioning clamp was preliminarily confirmed that the force did not decrease after pinching 200 times.

The participants were carefully familiarized with the methods and then laid in a supine position on the bed and took 5 minutes rest before the start of assessment. Pressure pain threshold was measured by using pressure algometer on tibialis anterior muscle (5 cm distal to the tibial tuberosity) and deltoid muscle (5 cm distal to the acromion) without monitoring of pressure acceleration itself, but it was increased roughly at a speed of 5 N/s, and subjects were instructed to verbally report the point at which pressure changed into pain (Fig. 1B). The force gauge has a peak mode which enables to keep the PPT value accurately. TSP was evaluated by 10 consecutive stimuli (1 second interval between the stimulus) by using pinprick on the same sites of PPT measurement and dorsum of hand (midpoint of the third and fourth metacarpal bone) (Fig. 1C). The TSP effect was calculated as the difference in the pain Visual Analogue Scale (VAS) (mm) between the first and the last stimuli, as previously reported.^{18,31} The VAS was anchored with "no pain" and "worst pain imaginable" at 0 mm and 100 mm, respectively. Conditioned pain modulation was evaluated by using conditioning clamp for applying tonic pain stimulus by pinching the earlobe for 60 seconds¹⁹ (Fig. 1D). When pain VAS of the earlobe became more than 60 mm, PPT was evaluated on contralateral tibialis anterior and deltoid muscle. The CPM effect was calculated as the percent change [(conditioning/baseline \times 100) - 100] and difference [conditioning-baseline] of the PPT, as previously recommended.46 The QST was performed unilaterally, and the test side was randomized in each healthy subject. Each measure except CPM was recorded 3 times, and the average measurement was used for analysis. Regarding each CPM measurement, conditioning pain was applied once and PPTs were recorded twice during the single conditioning period, and the average of the PPT was used for the assessment. All tests were performed at a similar time in the evening.

Two trained experimenters (Y.H. and R.S.) performed QST with an interval of a week. Intrarater and interrater test-retest reliabilities of the PPT, TS, and CPM at each measurement site were analyzed by using interclass correlation coefficient (ICC).

2.3. Experiment-B

Validity of *QuantiPain* assessment was evaluated compared with laboratory-based QST tools on the same day. In patients with OA, bilateral assessments were performed for PPT and TSP, but CPM was evaluated only on the affected side. In healthy controls, the



Figure 1. (A) Overview of the 3 items included in the bedside quantitative sensory testing tool kit, (B) pressure algometer, (C) pinprick, (D) conditioning clamp, and (E) cuff pressure algometry. White arrow: cuff for test stimulus. Black arrow: cuff for conditioning stimulus. The participants continuously rate their pain using a Visual Analogue Scale controller.

assessment side was randomized in each subject. The sequence of QST sessions by using *QuantiPain* or laboratory tools was also randomized. Before QST, the patients with OA were interviewed about the duration of their knee pain and average of the pain VAS (mm) at rest and while walking in the past week. They were also asked to complete chronic pain–associated questionnaires including Central Sensitization Inventory (CSI),²⁰ Hospital Anxiety and Depression Scale (HADS),⁵⁰ and Pain Catastrophizing Scale (PCS).³⁸

QuantiPain was used identically with the experiment-A for PPT (medial joint space of the knee and tibialis anterior muscle), TSP (tibialis anterior muscle), and CPM (tibialis anterior muscle) assessments.

As an established laboratory-based tool, a digital hand-held algometer (SBMEDIC, Hörby, Sweden) mounted with a 1-cm² probe was used for PPT recordings.^{4,45} Pressure was increased gradually at a rate of 30 kPa until the pain threshold was reached and the participants pressed a stop button.

Temporal summation of pain and CPM were measured by using cuff algometry (Cortex, NociTech and Aalborg University, Denmark), which is another all-in-one laboratory-based tool for QST (**Fig. 1E**).^{13,29,37} This consists of a 13-cm-wide tourniquet cuff, a computer-controlled air compressor, and an electronic VAS (Aalborg University, Aalborg, Denmark). The cuff was connected to the compressor and wrapped around the lower leg and was automatically inflated at a rate of 1 kPa/s. The pressure-induced pain intensity was recorded with the electronic 100-mm VAS and sampled at 10 Hz. The subjects were instructed to rate the VAS pain intensity continuously and to press a hand-held pressure release button when the pain was intolerable. This pressure value was defined as pressure tolerance threshold (PTT).

For TSP measurement, the cuff pressure stimuli were applied 10 times with a 1 second interstimulus interval and duration. The applied pressure was equal to the PTT.³⁷ The participants were instructed to continuously rate their pain on a VAS. The TSP effect was calculated as the difference in the VAS between the first and the tenth stimuli.

For CPM measurement, another cuff applied to the contralateral upper arm was promptly inflated to a pressure corresponding to 70% of the PTT as conditioning stimuli.³⁷ The cuff on the ipsilateral lower leg was then inflated at a rate of 1 kPa/s. The participants were instructed to rate the pain on their lower leg. Similar to the *QuantiPain* session, the CPM effect was calculated as the percent change and difference of pressure detection threshold (a pressure value corresponding to the VAS = 10 mm) with and without the conditioning stimuli. Pain VAS for the conditioning tonic pain (*QuantiPain*: earlobe and cuff algometry: upper arm) was also evaluated.

2.4. Statistical analysis

Sample size was determined according to some previous studies. Increasing evidence suggest that a large effect size when comparing QST parameters in patients with OA with healthy subjects⁵ and at least 24 subjects were needed in each group to detect a significant difference based on a large effect size (Cohen d = 0.8) and a significant level at 0.05 and a power of 80%.^{4,15} This was similar to our previously published reliability study on QST profiles.¹¹ Most of the data did not pass Shapiro–Wilk test for normal distribution and were presented in the median and interquartile range. In experiment-A, intrarater and interrater data of the PPT, TSP, and CPM were compared by using the Wilcoxon signed-rank test. Intrarater reliability was

assessed by using ICC (1, k) for data from test 1 and test 2 by a single experimenter (R.S.). Interrater reliability was also assessed by using ICC (3, k) for data from the 2 experimenters. Values less than 0.5, between 0.5 and 0.75, between 0.75 and 0.9, and greater than 0.90 are indicative of poor, moderate, good, and excellent reliability, respectively.¹⁶ In addition, standard error of measurement and smallest real difference were calculated as absolute measure for the reliability of the QST parameters⁴⁴ (see supplementary table 1, available at http://links.lww.com/PR9/A155).

In experiment-B, PPTs were analyzed per each assessment site. The Kruskal–Wallis test was performed on PPT and TSP using the factors of group (control, OA-affected side, and OAcontralateral side). The Mann–Whitney *U* test with the Bonferroni correction was used for post hoc comparisons when the Kruskal–Wallis test showed significant factors.

Comparison of CPM between groups was analyzed by the Mann–Whitney *U* test. The differences of QST tools were analyzed by the Wilcoxon signed-rank test in each group. Furthermore, a ranked distribution analysis of the CPM effects² was added for comparison between the groups and tools, and frequency of antinociceptive (increase PPT) or pronociceptive (decrease PPT) reaction against the conditioning stimulus was evaluated by using the χ^2 test.

Correlations of QST data between *QuantiPain* and the laboratory-based tools were analyzed in patients with OA and control subjects by using the Spearmen correlation coefficient. Moreover, correlations between *QuantiPain* data, knee pain VAS, and chronic pain–associated questionnaires were investigated in patients with OA. All analyses were performed with SPSS version 26.0 software (IBM Corp. Armonk, NY), and P < 0.05 indicated statistical significance.

3. Results

Table 1

3.1. Experiment-A

The median (interquartile range) results of the reliability analyses and ICCs are presented in **Table 1** for intrarater agreement and in **Table 2** for interrater agreement, respectively. The conditioning pain VAS for the CPM assessment was similar between the 2 experimenters (median: 70.0 mm), which seemed relevant to a recent methodology for evoking mechanically induced CPM.¹¹ Regarding intrarater assessment, no significant differences of QST data between tests 1 and 2 were seen and excellent to

FAIN REPUILS

moderate reliability was achieved except CPM which was evaluated by the difference of PPT on deltoid muscle. By contrast, interrater assessment showed significant difference of PPT on both sites and TSP on deltoid muscle, although reliability of these tests was excellent. Agreements of CPM were moderate to poor when it was evaluated by the percent change or difference of PPT on both sites.

3.2. Experiment-B

Demographic data of the patients with knee OA and healthy control subjects are presented in **Table 3**. There was no significant difference of age between groups, but the OA group included more women than the control group. Four patients in the OA group showed radiological bilateral knee OA; however, they had no pain in the contralateral knee.

3.2.1. Quantitative sensory testing measures by different technologies

PPTs recorded on medial joint space of the knee and tibialis anterior muscle were bilaterally lower in patients with OA compared with control subjects. This finding was comparable between using *QuantiPain* (Fig. 2A) and the digital algometer (Fig. 2B), but the difference did not reach statistical significance when using the digital algometer on tibialis anterior muscle. Moreover, excellent correlations were observed between PPTs evaluated by *QuantiPain* and the digital algometer on medial joint space of the knee (affected side: R = 0.848, P < 0.0001, Fig. 2C. and contralateral side: R = 0.815, P < 0.0001, Fig. 2D. and contralateral side: R = 0.807, P < 0.0001).

Enhancement of TSP was observed in both groups; however, it was significantly facilitated in patients with OA compared with control subjects when evaluated by *QuantiPain* (Fig. 3A) and cuff algometry (Fig. 3B). A moderate correlation was seen between TSP evaluated by both systems (affected side: R = 0.447, P < 0.0001, Fig. 3C. and contralateral side: R = 0.284, P = 0.0107).

Conditioned pain modulation was significantly impaired in patients with OA compared with control subjects when evaluated by *QuantiPain* (Figs. 4A and B) and cuff algometry (Figs. 4C and D). A weak but significant correlation was seen between the CPM evaluated by both systems (percent change of PPT: R = 0.281, P = 0.0116, Fig. 4E and difference of PPT: R = 0.3786,

Results of intrarater reliabilit	y analysis.			
	Test 1	Test 2	Wilcoxon (<i>P</i>)	ICC (1, k) [95% CI]
PPT (N)				
Tibialis anterior	51.3 [37.2–60.3]	47.0 [38.0–53.8]	0.677	0.90 [0.76-0.96]
Deltoid	30.5 [24.0–36.7]	33.0 [20.8–37.9]	0.955	0.94 [0.84–0.97]
TSP (mm)				
Tibialis anterior	16.0 [8.0-24.5]	11.0 [6.0-20.0]	0.196	0.84 [0.61-0.93]
Deltoid	15.0 [6.5–27.0]	11.0 [6.0–28.0]	0.265	0.77 [0.43-0.90]
Hand	15.0 [7.5–24.5]	12.0 [6.5–21.5]	0.575	0.60 [0.13-0.83]
СРМ				
Tibialis anterior				
Percent change (%)	17.8 [9.8–26.0]	21.8 [9.6–28.5]	0.677	0.76 [0.41–0.90]
Difference (N)	6.7 [4.5–12.8]	9.0 [5.2–13.7]	0.794	0.80 [0.52-0.92]
Deltoid				
Percent change (%)	32.5 [15.5–50.0]	20.3 [10.7–43.9]	0.313	0.67 [0.20-0.87]
Difference (N)	8.0 [4.3–16.3]	6.0 [4.5-9.7]	0.266	0.39 [-0.47-0.75]

Data from tests 1 and 2 are presented as median [interquartile range]. ICC (1, k) was presented with 95% Cl.

CI, confidence interval; ICC, interclass correlation coefficient; PPT, pressure pain threshold; TSP, temporal summation of pain; CPM, conditioned pain modulation.

Table 2

Results of interrater reliability analysis.

	Experimenter 1	Experimenter 2	Wilcoxon (<i>P</i>)	ICC (3, k) [95% CI]
PPT (N)				
Tibialis anterior	48.4 [39.4–56.3]	55.7 [49.8–71.5]	0.0002	0.92 [0.80-0.97]
Deltoid	31.8 [22.2–37.6]	39.7 [25.5–46.8]	0.0002	0.90 [0.76–0.96]
TSP (mm)				
Tibialis anterior	13.5 [8.0–22.0]	9.0 [3.0–20.0]	0.357	0.86 [0.64-0.94]
Deltoid	14.0 [5.3–21.3]	8.0 [2.0–22.5]	0.038	0.91 [0.78-0.96]
Hand	14.0 [7.5–24.3]	10.0 [5.0–21.0]	0.578	0.71 [0.28–0.88]
CPM				
Tibialis anterior				
Percent change (%)	17.0 [9.2–30.5]	10.4 [8.3–19.1]	0.23	0.61 [0.04-0.84]
Difference (N)	6.8 [4.4–12.5]	7.0 [4.4–13.0]	0.972	0.37 [-0.56-0.74]
Deltoid				
Percent change (%)	27.9 [15.1–38.9]	25.9 [16.1–43.8]	0.876	0.72 [0.31–0.89]
Difference (N)	7.8 [4.8–12.5]	10.0 [5.2–14.6]	0.179	0.45 [-0.36–0.78]

Data from experimenters 1 and 2 are presented as median [interquartile range]. ICC (3, k) was presented with 95% Cl.

Bold indicates significant difference between the experimenters.

CI, confidence interval; ICC, interclass correlation coefficient; PPT, pressure pain threshold; TSP, temporal summation of pain; CPM, conditioned pain modulation.

P = 0.0005, Fig. 4F). The percent change of PPT was significantly greater by using *QuantiPain* compared with cuff algometry in patients with OA (Wilcoxon; P = 0.001), but not in control subjects (Wilcoxon; P = 0.077). Pain VAS for the conditioning tonic pain was significantly greater in *QuantiPain* (earlobe) than cuff algometry (upper arm) in both groups (patients with OA: 80.0[70.0-80.0] mm vs 64.1[48.0-70.0] mm; Wilcoxon; P < 0.000001 and control subjects: 67.0[60.0-80.0] mm vs 62.0 [48.3-70.0] mm; Wilcoxon; P = 0.00028). The conditioning pain VAS was higher in patients with OA than control subjects in *QuantiPain* session (Mann–Whitney, P = 0.005), but this difference was not seen when using cuff algometry (Mann–Whitney, P = 0.972).

The ranked distribution analysis demonstrated that 75% of the patients with OA showed antinociceptive reaction while it was 93% for the control subjects assessed by *QuantiPain* (χ^2 ; *P* = 0.034, **Figs. 5A and B**). By contrast, cuff algometry categorized 43% of the patients with OA and 70% of the control subjects as antinociceptive, respectively (χ^2 ; *P* = 0.013, **Figs. 5C and D**).

3.2.2. Quantitative sensory testing and chronic pain-associated questionnaires

Correlations between *QuantiPain* data, knee pain VAS, and chronic pain–associated questionnaires were displayed in Supplementary table 2 (available at http://links.lww.com/PR9/A155). Ipsilateral PPTs on medial joint space and tibialis anterior muscle were negatively correlated with pain VAS at rest (R = -0.35, -0.33), on walking (R = -0.38, -0.44), and PCS (R = -0.37, -0.37). Ipsilateral TSP was associated with CSI (R = 0.37), whereas CPM was not correlated with all questionnaires. PCS was correlated with CSI (R = 0.53) and HADS anxiety (R = 0.32). CSI was also correlated with HADS anxiety (R = 0.60). Among QST parameters, PPT on tibialis anterior muscle was correlated with TSP (R = -0.38) and CPM (R = 0.48), but no significant correlation was observed between TSP and CPM.

4. Discussion

This study demonstrated that *QuantiPain* had acceptable testretest reliability. In addition, this tool kit successfully detected localized and widespread decrease of PPT, facilitated TSP, and impaired CPM in patients with OA compared with controls, and the data were significantly correlated with laboratory-based QST assessment. Because the tool kit has promising benefits such as being cheap, easy to use, and portable, it seems to have enough potential for clinical application.

4.1. Test-retest reliability of the bedside quantitative sensory testing tool kit

Regarding PPT and TSP, good to excellent intrarater and interrater reliabilities were confirmed except TSP assessed on dorsum of hand being moderate, which were comparable with established pressure algometry and cuff algometry reported in the previous studies.^{12,42} A plausible reason of lower reliability of TSP was the hand being thinner and had more mobile skin with less deep tissues compared with other assessment sites. However, rapid assessment on hand seems to be a great advantage for clinical application.

As recognized, CPM is one of the most unstable QST test partly because of its methodological complexity and variability of test and conditioning stimulus, and hence, intrarater and interrater reliabilities were lower in general.^{26,36} *QuantiPain* showed, however, moderate to good intrarater reliability of CPM effect except the difference of PPT on deltoid, and the ICC (1, k) was comparable with a similar bedside tool reported as 0.67 to 0.72.¹⁹ As for the interrater assessment, moderate reliability was confirmed when evaluated the CPM effect with percent change of PPT, whereas the reliability became poor when evaluated with the difference of PPT. However, the reliability of CPM assessment by *QuantiPain* does not seem to be far less than cuff algometry of which ICC (3, k) was documented as 0.47 to 0.73 in a previous study.¹¹

4.2. Quantitative sensory testing measures by different technologies

Localized and widespread hyperalgesia, facilitated TSP, and impaired CPM were found in patients with OA, which were similar results from laboratory-based tools in this study, and consistent with current understanding of altered pain mechanisms in patients with OA.^{8,21}

Looking at each parameter, PPT measured by our pressure algometer showed excellent correlation with the use of the digital algometer. Although it is not surprising as PPT measurement itself

Table 3
Demographic data of patients with knee osteoarthritis and control
subjects in experiment-B.

Variable	Patients with knee OA	Control subjects
Ν	40	40
Age (y)	69 [58–74]	66 [48–73]
Sex, n (%) Male Female	8 (20) 32 (80)	23 (57) 17 (43)
Pain duration, mo	36.0 [4.5–168.0]	0
VAS at rest, mm	0.0 [0.0–19.3]	0
VAS on walking, mm	48.0 [31.8–67.3]	0
CSI	20.5 [11.3–27.8]	—
HADS Anxiety Depression	3.5 [2.0–6.3] 4.0 [3.0–9.0]	_
PCS	16.0 [8.0–25.0]	—

Data are presented as median [interguartile range].

CSI, Central Sensitization Inventory; HADS, Hospital Anxiety and Depression Scale; OA, osteoarthritis; PCS, Pain Catastrophizing Scale; VAS; Visual Analogue Scale.

was technically almost identical, this finding suggests that roughly increasing pressure at a speed of 5 N/s and verbally indicated thresholds were acceptable measurement for bedside testing.

For TSP, the pinprick stimulation was a different procedure compared with cuff algometry, ie, pinprick mainly stimulated localized superficial tissues while cuff algometry compressed larger area of deep tissues. According to a previous experimental study, TSP was likely to be facilitated by painful stimuli to deep structures rather than superficial tissues.²⁴ Nevertheless, there was a significant moderate correlation between TSP measured by both systems. One possible explanation of this accordance results from structural characteristics of our pinprick device that incorporates a 60 g mobile weight in the body and conical tip. Compared with filaments used in other studies,^{17,34} this device may provide more effective stimuli to superficial and deep structures for facilitating TSP. Because the assessment is extremely easy, this procedure will become a good option for bedside TSP testing.

Impaired CPM in patients with OA was indicated by *QuantiPain* and cuff algometry; however, the correlation of CPM data between both systems was weaker compared with that of PPT and TSP. As mentioned, CPM is a highly variable assessment in general, so the main reason of the discordance was probably derived from the difference of test and conditioning stimulus between the 2 systems. In addition, pain intensity of conditioning stimulus was significantly higher when assessed by *QuantiPain* than cuff algometry in both patients with OA and healthy controls. Because recent studies supported that more painful conditioning stimulus evokes more CPM effect,^{11,25,27} management of the cramping force to earlobe for individual subject might be needed to achieve a more stable result.

The ranked distribution analysis of the CPM provided further interesting information in this study. Frequency of antinociceptive and pronociceptive reaction was significantly different not only between the groups but also between the assessment devises. Because the CPM effects are usually biphasic (ie, antinociceptive and pronociceptive) especially in patients with chronic pain having sensitization, simple analysis using representative values (e.g., average) of the cohorts may be at a risk of overlooking true outcomes. Moreover, it is important that both healthy subjects and patients with OA could individually demonstrate a "scattered," not a "binary" CPM response. In this regard, our new approach would help understanding the characteristics of CPM assessment.



Figure 2. Results of pressure pain threshold (PPT). (A) *QuantiPain* (pressure algometer), (B) laboratory tool (digital algometer), (C) correlation between using both tools measured on TA. **P* < 0.05 vs control, ***P* < 0.01 vs control. MJS; medial joint space, TA; tibialis anterior muscle.



Figure 3. Results of temporal summation of pain (TSP). (A) *QuantiPain* (pinprick), (B) laboratory tool (cuff algometry), and (C) correlation between using both tools. **P* < 0.05 vs control, ***P* < 0.01 vs control.

4.3. Quantitative sensory testing and chronic pain-associated questionnaires

Relationship between QST data and chronic pain-associated questionnaires still remains unclear, but assumed to be weak in recent reports. Walton et al.43 showed that depression, catastrophizing, and kinesiophobia were able to explain small variance in local PPT in people with mechanical neck pain. Coronado et al.⁷ showed that CSI was associated with resilience, anxiety, and negative effects but not with PPT on remote sites in patients with shoulder pain. In patients with knee OA, Gervais-Hupe et al.⁹ reported that CSI was weakly correlated with decreased PPT locally and remotely, and with CPM, but not with TSP. They also mentioned that the CSI is more strongly associated with psychological factors than QST results. Consistent with these reports, this study showed weak correlations between some, but limited QST data and questionnaires (local PPT vs PCS and TSP vs CSI); however, the impact was much smaller than that of PCS vs CSI or CSI vs HADS anxiety. It makes

sense because psychophysical tests and psychological questionnaires are not identical measures, although part of patients with chronic pain has overlapped abnormalities detected by both items.

No significant correlation was observed between TSP and CPM. Temporal summation of pain is often facilitated, and CPM is often impaired in patients with chronic pain compared with healthy subjects.³ However, a recent human experimental study revealed that these 2 dynamic QST assessments were not associated,¹⁴ which was similar to the findings in this study.

4.4. Comparison with other bedside quantitative sensory testing tool kit

Development of easy-to-use bedside QST tool kit has been a hot topic for phenotyping patients with chronic pain. Koulouris et al.¹⁷ developed a bedside equipment for evaluating patients with neuropathic pain. They confirmed moderate test–retest reliability



Figure 4. Results of conditioned pain modulation (CPM). (A) *QuantiPain* (pressure algometer with conditioning clamp) analyzed by percent change of PPT, (B) *QuantiPain* analyzed by difference of PPT, (C) laboratory tool (cuff algometry) analyzed by percent change of PPT, (D) laboratory tool analyzed by difference of PPT, (E) correlation between using both tools analyzed by percent change of PPT, and (F) correlation between using both tools analyzed by difference of PPT. *P < 0.05 vs control, **P < 0.01 vs control. PPT, pressure pain threshold.



Figure 5. Ranked distribution analysis of the CPM effect (presented as percent change of PPT). Positive and negative values show antinociceptive and pronociceptive reaction against the conditioning tonic pain stimulus, respectively. (A) *QuantiPain* (patients with OA), (B) *QuantiPain* (healthy subjects), (C) cuff algometry (patients with OA), and (D) cuff algometry (healthy subjects). CPM, conditioned pain modulation; PPT, pressure pain threshold; OA, osteoarthritis.

of the assessment, and the results were highly correlated with laboratory-based QST variables. Reimer et al.³³ presented another bedside equipment and demonstrated that sensory loss, thermal hyperalgesia, and mechanical hyperalgesia were nominated as bedside cluster assessment. Their new equipment will help stratification of patients with neuropathic pain; however, it seems a bit complicated and concerned that most of the parameters are assessed by static QST with suprathreshold stimulus response.

By contrast, our concept for development of *QuantiPain* is "copying laboratory-based mechanistic QST as simple as possible at bedside," and hence, the algometer was used for analyzing subthreshold stimulus response and detecting PPT. Moreover, TSP and CPM, known as the 2 measure paradigms of dynamic QST, are incorporated by adding quite simple items (pinprick and conditioning clamp) to evaluate overreaction and dysfunction of the central nervous system. This concept would be partly supported by a latest report developing a bedside tool kit for assessing sensitization in patients with chronic OA knee pain.³⁵ They included mechanical pinprick sensitivity, dynamic mechanical allodynia, pressure pain sensitivity, TSP, and CPM for analysis and detected 46% of patients showed signs of sensitization.

4.5. Limitations

There were some limitations to be noted when interpreting the results of this study. First, the number of patients with OA and healthy subjects was relatively small and recruited from single institution because this is a preliminary study investigating test-retest reliability and validity of *QuantiPain*. Second, 80% females were included in patients with OA while only 43% females in control group in experiment 2, which might affect the outcome of comparison between the 2 groups. However, the main purpose of this comparison was to confirm the validity of QuantiPain

compared with laboratory-based QST tools; therefore, the effects of sex difference probably worked equally when using both tools. In addition, correlations of QST variables between both tools were analyzed by using the data from patients with OA and controls together for minimizing the effects of sex difference. Third, *QuantiPain* focused on mechanistic approach of QST and mainly targeted on deep somatic pain. Lack of thermal and light touch stimulus sensitivity might be a disadvantage of *QuantiPain*, especially for neuropathic pain evaluation; however, we prioritized simple protocols than comprehensive sensory testing in this study.

5. Conclusion

The presented, simple bedside tool kit demonstrated acceptable test-retest reliability and assessment validity that would be capable of evaluating painful patients. Although this has not become a complete alternative of laboratory-based tools and further research is warranted for improving reliability, the tool kit has a potential to create more practical approach for quantifying altered pain mechanisms in clinical settings.

Disclosures

M. Izumi received grants from Shionogi & Co., Ltd. during the conduct of this study. The remaining authors have no conflicts of interest to declare.

This study was financially supported by Shionogi & Co, Ltd (Osaka, Japan).

Acknowledgements

This research was conducted as a collaborative project with Kochi University (Nankoku, Kochi, Japan) and Shionogi Inc. (Osaka, Japan). The authors thank all collaborative members of pain biomarker department in Shionogi Inc for practical assistance. Center for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121). Authors' contribution: Study design: M. Izumi, Y. Hayashi, S. Oda, K.K. Petersen, and L. Arendt-Nielsen. Data collection: Y. Hayashi and R. Saito. Data analysis: M. Izumi, Y. Hayashi, R. Saito, S. Oda, and K.K. Petersen. Drafting or revising manuscript: M. Izumi, Y. Hayashi, S. Oda, K.K. Petersen, L. Arendt-Nielsen, and M. Ikeuchi. All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Appendix A. Supplemental digital content

Supplemental digital content associated with this article can be found online at http://links.lww.com/PR9/A155.

Article history:

Received 29 October 2021 Received in revised form 25 January 2022 Accepted 10 February 2022

References

- Arant KR, Katz JN, Neogi T. Quantitative sensory testing: identifying pain characteristics in patients with osteoarthritis. Osteoarthritis Cartilage 2021;20:801–7.
- [2] Arendt-Nielsen L, Larsen JB, Rasmussen S, Krogh M, Borg L, Madeleine P. A novel clinical applicable bed-side tool for assessing conditioning pain modulation: proof-of-concept. Scand J Pain 2020;30:17–31.
- [3] Arendt-Nielsen L, Morlion B, Perrot S, Dahan A, Dickenson A, Kress HG, Wells C, Bouhassira D, Mohr Drewes A. Assessment and manifestation of central sensitisation across different chronic pain conditions. Eur J Pain 2018;22:216–41.
- [4] Arendt-Nielsen L, Nie H, Laursen MB, Laursen BS, Madeleine P, Simonsen OH, Graven-Nielsen T. Sensitization in patients with painful knee osteoarthritis. PAIN 2010;149:573–81.
- [5] Arendt-Nielsen L, Skou ST, Nielsen TA, Petersen KK. Altered central sensitization and pain modulation in the CNS in chronic joint pain. Curr Osteoporos Rep 2015;13:225–34.
- [6] Arendt-Nielsen L, Yarnitsky D. Experimental and clinical applications of quantitative sensory testing applied to skin, muscles and viscera. J Pain 2009;10:556–72.
- [7] Coronado RA, George SZ. The Central Sensitization Inventory and Pain Sensitivity Questionnaire: an exploration of construct validity and associations with widespread pain sensitivity among individuals with shoulder pain. Musculoskelet Sci Pract 2018;36:61–7.
- [8] Fingleton C, Smart K, Moloney N, Fullen BM, Doody C. Pain sensitization in people with knee osteoarthritis: a systematic review and meta-analysis. Osteoarthritis Cartilage 2015;23:1043–56.
- [9] Gervais-Hupé J, Pollice J, Sadi J, Carlesso LC. Validity of the central sensitization inventory with measures of sensitization in people with knee osteoarthritis. Clin Rheumatol 2018;37:3125–32.
- [10] Graven-Nielsen T, Arendt-Nielsen L. Assessment of mechanisms in localized and widespread musculoskeletal pain. Nat Rev Rheumatol 2010;6:599–606.
- [11] Graven-Nielsen T, Izumi M, Petersen KK, Arendt-Nielsen L. Userindependent assessment of conditioning pain modulation by cuff pressure algometry. Eur J Pain 2017;21:552–61.
- [12] Graven-Nielsen T, Vaegter HB, Finocchietti S, Handberg G, Arendt-Nielsen L. Assessment of musculoskeletal pain sensitivity and temporal summation by cuff pressure algometry: a reliability study. PAIN 2015;156: 2193–202.
- [13] Graven-Nielsen T, Wodehouse T, Langford RM, Arendt-Nielsen L, Kidd BL. Normalization of widespread hyperesthesia and facilitated spatial summation of deep-tissue pain in knee osteoarthritis patients after knee replacement. Arthritis Rheum 2012;64:2907–16.
- [14] Holden S, Petersen KK, Arendt-Nielsen L, Graven-Nielsen T. Conditioning pain modulation reduces pain only during the first

stimulation of the temporal summation of pain paradigm in healthy participants. Eur J Pain 2019;23:1390–6.

- [15] Izumi M, Petersen KK, Laursen MB, Arendt-Nielsen L, Graven-Nielsen T. Facilitated temporal summation of pain correlates with clinical pain intensity after hip arthroplasty. PAIN 2017;158:323–32.
- [16] Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. J Chiropr Med 2016;15:155–63.
- [17] Koulouris AE, Edwards RR, Dorado K, Schreiber KL, Lazaridou A, Rajan S, White J, Garcia J, Gibbons C, Freeman R. Reliability and validity of the Boston bedside quantitative sensory testing battery for neuropathic pain. Pain Med 2020;21:2336–47.
- [18] Kurien T, Arendt-Nielsen L, Petersen KK, Graven-Nielsen T, Scammell BE. Preoperative neuropathic pain-like symptoms and central pain mechanisms in knee osteoarthritis predicts poor outcome 6 Months after total knee replacement surgery. J Pain 2018;19:1329–41.
- [19] Larsen JB, Madeleine P, Arendt-Nielsen L. Development of a new bedside-test assessing conditioned pain modulation: a test-retest reliability study. Scand J Pain 2019;19:565–74.
- [20] Mayer TG, Neblett R, Cohen H, Howard KJ, Choi YH, Williams MJ, Perez Y, Gatchel RJ. The development and psychometric validation of the central sensitization inventory. Pain Pract 2012;12:276–85.
- [21] Moore RL, Clifford AM, Moloney N, Doody C, Smart KM, O'Leary H. The relationship between clinical and quantitative measures of pain sensitization in knee osteoarthritis. Clin J Pain 2020;36:336–43.
- [22] Neogi T, Frey-Law L, Scholz J, Niu J, Arendt-Nielsen L, Woolf C, Nevitt M, Bradley L, Felson DT, Study MOM. Sensitivity and sensitisation in relation to pain severity in knee osteoarthritis: trait or state? Ann Rheum Dis 2015; 74:682–8.
- [23] Neogi T, Guermazi A, Roemer F, Nevitt MC, Scholz J, Arendt-Nielsen L, Woolf C, Niu J, Bradley LA, Quinn E, Law LF. Association of joint inflammation with pain sensitization in knee osteoarthritis: the multicenter osteoarthritis study. Arthritis Rheumatol 2016;68:654–61.
- [24] Nie H, Arendt-Nielsen L, Andersen H, Graven-Nielsen T. Temporal summation of pain evoked by mechanical stimulation in deep and superficial tissue. J Pain 2005;6:348–55.
- [25] Nir RR, Granovsky Y, Yarnitsky D, Sprecher E, Granot M. A psychophysical study of endogenous analgesia: the role of the conditioning pain in the induction and magnitude of conditioned pain modulation. Eur J Pain 2011;15:491–7.
- [26] Oono Y, Nie H, Matos RL, Wang K, Arendt-Nielsen L. The inter- and intraindividual variance in descending pain modulation evoked by different conditioning stimuli in healthy men. Scand J Pain 2011;2:162–9.
- [27] Oono Y, Wang K, Svensson P, Arendt-Nielsen L. Conditioned pain modulation evoked by different intensities of mechanical stimuli applied to the craniofacial region in healthy men and women. J Orofac Pain 2011;25: 364–75.
- [28] Petersen KK, Arendt-Nielsen L, Simonsen O, Wilder-Smith O, Laursen MB. Presurgical assessment of temporal summation of pain predicts the development of chronic postoperative pain 12 months after total knee replacement. PAIN 2015;156:55–61.
- [29] Petersen KK, Graven-Nielsen T, Simonsen O, Laursen MB, Arendt-Nielsen L. Preoperative pain mechanisms assessed by cuff algometry are associated with chronic postoperative pain relief after total knee replacement. PAIN 2016;157:1400–6.
- [30] Petersen KK, Olesen AE, Simonsen O, Arendt-Nielsen L. Mechanistic pain profiling as a tool to predict the efficacy of 3-week nonsteroidal antiinflammatory drugs plus paracetamol in patients with painful knee osteoarthritis. PAIN 2019;160:486–92.
- [31] Petersen KK, Simonsen O, Laursen MB, Arendt-Nielsen L. The role of preoperative radiologic severity, sensory testing, and temporal summation on chronic postoperative pain following total knee arthroplasty. Clin J Pain 2018;34:193–7.
- [32] Petersen KK, Vaegter HB, Stubhaug A, Wolff A, Scammell BE, Arendt-Nielsen L, Larsen DB. The predictive value of quantitative sensory testing: a systematic review on chronic postoperative pain and the analgesic effect of pharmacological therapies in patients with chronic pain. PAIN 2021;162:31–44.
- [33] Reimer M, Forstenpointner J, Hartmann A, Otto JC, Vollert J, Gierthmühlen J, Klein T, Hüllemann P, Baron R. Sensory bedside testing: a simple stratification approach for sensory phenotyping. Pain Rep 2020;5:e820.
- [34] Rolke R, Baron R, Maier C, Tölle TR, Treede RD, Beyer A, Binder A, Birbaumer N, Birklein F, Bötefür IC, Braune S, Flor H, Huge V, Klug R, Landwehrmeyer GB, Magerl W, Maihöfner C, Rolko C, Schaub C, Scherens A, Sprenger T, Valet M, Wasserka B. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. PAIN 2006; 123:231–43.

- [35] Sachau J, Otto JC, Kirchhofer V, Larsen JB, Kennes LN, Hüllemann P, Arendt-Nielsen L, Baron R. Development of a bedside tool-kit for assessing sensitization in patients with chronic osteoarthritis knee pain or chronic knee pain after total knee replacement. PAIN 2022;163:308–18.
- [36] Schliessbach J, Lütolf C, Streitberger K, Scaramozzino P, Arendt-Nielsen L, Curatolo M. Reference values of conditioned pain modulation. Scand J Pain 2019;19:279–86.
- [37] Staffe AT, Bech MW, Clemmensen SLK, Nielsen HT, Larsen DB, Petersen KK. Total sleep deprivation increases pain sensitivity, impairs conditioned pain modulation and facilitates temporal summation of pain in healthy participants. PLoS One 2019;14:e0225849.
- [38] Sullivan MJL, Stanish W, Waite H, Sullivan M, Tripp DA. Catastrophizing, pain, and disability in patients with soft-tissue injuries. PAIN 1998;77: 253–60.
- [39] Suokas AK, Walsh DA, McWilliams DF, Condon L, Moreton B, Wylde V, Arendt-Nielsen L, Zhang W. Quantitative sensory testing in painful osteoarthritis: a systematic review and meta-analysis. Osteoarthritis Cartilage 2012;20:1075–85.
- [40] Tracey I, Woolf CJ, Andrews NA. Composite pain biomarker signatures for objective assessment and effective treatment. Neuron 2019;101: 783–800.
- [41] Urquhart DM, Phyomaung PP, Dubowitz J, Fernando S, Wluka AE, Raajmaakers P, Wang Y, Cicuttini FM. Are cognitive and behavioural factors associated with knee pain? A systematic review. Semin Arthritis Rheum 2015;44:445–55.
- [42] Waller R, Straker L, O'Sullivan P, Sterling M, Smith A. Reliability of pressure pain threshold testing in healthy pain free young adults. Scand J Pain 2015;9:38–41.

- [43] Walton DM, Levesque L, Payne M, Schick J. Clinical pressure pain threshold testing in neck pain: comparing protocols, responsiveness, and association with psychological variables. Phys Ther 2014;94:827–37.
- [44] Weir JP. Quantifying test-retest reliability using the intraclass correlation coefficient and the SEM. J Strength Cond Res 2005;19:231–40.
- [45] Wylde V, Sayers A, Lenguerrand E, Gooberman-Hill R, Pyke M, Beswick AD, Dieppe P, Blom AW. Preoperative widespread pain sensitization and chronic pain after hip and knee replacement: a cohort analysis. PAIN 2015;156:47–54.
- [46] Yamitsky D, Bouhassira D, Drewes AM, Fillingim RB, Granot M, Hansson P, Landau R, Marchand S, Matre D, Nilsen KB, Stubhaug A, Treede RD, Wilder-Smith OH. Recommendations on practice of conditioned pain modulation (CPM) testing. Eur J Pain 2015;19:805–6.
- [47] Yarnitsky D, Crispel Y, Eisenberg E, Granovsky Y, Ben-Nun A, Sprecher E, Best LA, Granot M. Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. PAIN 2008; 138:22–8.
- [48] Yamitsky D, Granot M, Nahman-Averbuch H, Khamaisi M, Granovsky Y. Conditioned pain modulation predicts duloxetine efficacy in painful diabetic neuropathy. PAIN 2012;153:1193–8.
- [49] Youssef AM, Macefield VG, Henderson LA. Pain inhibits pain; human brainstem mechanisms. Neuroimage 2016;124:54–62.
- [50] Zigmond AS, Snaith RP. The hospital anxiety and depression scale. Acta Psychiatr Scand 1983;67:361–70.
- [51] Zolio L, Lim KY, McKenzie JE, Yan MK, Estee M, Hussain SM, Cicuttini F, Wluka A. Systematic review and meta-analysis of the prevalence of neuropathic-like pain and/or pain sensitization in people with knee and hip osteoarthritis. Osteoarthritis Cartilage 2021;29:1096–116.