

Development of Neuropathic Post-COVID Pain Symptoms is not Associated with Serological Biomarkers at Hospital Admission in COVID-19 Survivors

A Secondary Analysis

Fernández-de-Las-Peñas, César; Herrero-Montes, Manuel; Ferrer-Pargada, Diego; Izquierdo-Cuervo, Sheila; Palacios-Ceña, Domingo; Arendt-Nielsen, Lars; Torres-Macho, Juan; Parás-Bravo, Paula

Published in:
Pain Medicine

DOI (link to publication from Publisher):
[10.1093/pm/pnac086](https://doi.org/10.1093/pm/pnac086)

Publication date:
2022

Document Version
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Fernández-de-Las-Peñas, C., Herrero-Montes, M., Ferrer-Pargada, D., Izquierdo-Cuervo, S., Palacios-Ceña, D., Arendt-Nielsen, L., Torres-Macho, J., & Parás-Bravo, P. (2022). Development of Neuropathic Post-COVID Pain Symptoms is not Associated with Serological Biomarkers at Hospital Admission in COVID-19 Survivors: A Secondary Analysis. *Pain Medicine*, 23(12), 2092-2094. <https://doi.org/10.1093/pm/pnac086>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from vbn.aau.dk on: December 04, 2025

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Research Letter

Development of Neuropathic Post-COVID Pain Symptoms is not Associated with Serological Biomarkers at Hospital Admission in COVID-19 Survivors: A Secondary Analysis

César Fernández-de-las-Peñas^{1,2} Dr. Med, PhD; Manuel Herrero-Montes^{3,4} PhD; Diego Ferrer-Pargada⁵ MD; Sheila Izquierdo-Cuervo⁵ MD; Domingo Palacios-Ceña¹ PhD; Lars Arendt-Nielsen^{2,6} Dr. Med, PhD; Juan Torres-Macho^{7,8} MD; Paula Parás-Bravo^{3,4} PhD

¹ Department of Physical Therapy, Occupational Therapy, Physical Medicine and Rehabilitation, Universidad Rey Juan Carlos (URJC), Madrid. Spain.

² Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark

³ Departamento de Enfermería, Universidad de Cantabria, Avda. Valdecilla, s/n, 39008 Santander, Spain.

⁴ Instituto de Investigación Sanitaria Valdecilla (IDIVAL), Grupo de Investigación en Enfermería, Calle Cardenal Herrera Oria, s/n, 39011 Santander, Spain.

⁵ Servicio de Neumología, Hospital Universitario Marqués de Valdecilla, Avda. Valdecilla, s/n, 39008 Santander, Spain.

⁶ Department of Medical Gastroenterology, Mech-Sense, Aalborg University Hospital, Aalborg, Denmark

⁷ Department of Internal Medicine, Hospital Universitario Infanta Leonor-Virgen de la Torre, Madrid Spain

⁸ Department of Medicine, School of Medicine, Universidad Complutense de Madrid, Madrid, Spain.

Corresponding / reprint requests author:
César Fernández-de-las-Peñas Telephone number: + 34 91 488 88 84
Facultad de Ciencias de la Salud
Universidad Rey Juan Carlos Fax number: +34 91 488 89 57
Avenida de Atenas s/n
28922 Alcorcón, Madrid, SPAIN
Email: cesar.fernandez@urjc.es

Downloaded from https://academic.oup.com/painmedicine/advance-article/doi/10.1093/pm/pnac086/6604280 by Aalborg University Library user on 13 June 2022

Role of the Funding Source

Proyecto financiado por la convocatoria Next-Val 2021 de la Fundación Instituto de Investigación Marqués de Valdecilla (IDIVAL) and by a grant from the Novo Nordisk Foundation 0067235. The sponsors had no role in the design, collection, management, analysis, or interpretation of the data, draft, review, or approval of the manuscript or its content. The authors were responsible for the decision to submit the manuscript for publication, and the sponsor did not participate in this decision.

Declaration of interests

No conflict of interest is declared by any of the authors

Development of Neuropathic Post-COVID Pain Symptoms is not Associated with Serological Biomarkers at Hospital Admission in COVID-19 Survivors: A Secondary Analysis

Dear Editor

Introduction

Pain symptoms have been found to be present as a post-COVID sequelae in up to 18% of subjects who had survived to the Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) virus [1]. Although post-COVID pain of musculoskeletal origin is the most commonly reported type of pain [2], neuropathic pain is also described as potential post-COVID sequelae. Oguz-Akarsu et al reported that 25% of patients with post-COVID pain has neuropathic symptoms; however, they collected self-reported pain symptoms throughout a telephonic interview [3]. A similar prevalence of neuropathic symptoms (24.6%) has been recently found using a validated the Self-Report Leeds Assessment of Neuropathic Symptoms (S-LANSS) questionnaire by our research group [4].

Identification of potential factors associated with the development of post-COVID neuropathic pain could help for identifying individuals at a higher risk of developing post-COVID pain and, hence, timely interventions and information. Serological biomarkers at the acute phase of COVID-19 infection could be a potential risk factor contributing to the development of long COVID. There is preliminary evidence supporting that neuropathic post-COVID pain can be associated with serum levels of neurofilament light chain (NFL) as a potential biomarker [5]. We present here a secondary analysis of a previous cohort study investigating the prevalence of neuropathic symptoms in previously hospitalized COVID-19 survivors exhibiting “de novo” post-COVID pain [4]. The aim of the current secondary analysis was to investigate the association between serological biomarkers at hospital admission with the development of neuropathic post-COVID symptoms.

Methods

A secondary analysis of a previous observational cross-sectional cohort study was conducted [4]. Briefly, patients hospitalized during the first wave of the pandemic at an urban hospital in Spain due to SARS-CoV-2 infection attending to a specific post-COVID unit from 1st June to 31st October 2021 were invited to participate. They were included if reported pain as their primary post-COVID symptom and did not present a pre-existing history of pain symptoms or any medical comorbidity explaining the presence of pain as previously described [4]. The Institutional Ethic Committee of INDIVAL Cantabria (code 2020.416) approved the study. All participants provided their informed consent.

As previously described in detail [4], participants completed the following self-reported questionnaires: S-LANSS for assessing the presence of neuropathic symptoms [6], the Hospital Anxiety and Depression Scale for the presence of anxiety/depressive levels, the 11-item Tampa Scale for Kinesiophobia for the presence of fear of movement, and the Pain Catastrophizing Scale. In this secondary analysis, we used the cut-off score of ≥ 12 points on the total score of the S-LANSS (range 0 to 24) for determining the presence of neuropathic symptoms [6].

We obtained the following serological biomarkers collected at hospital admission from hospital medical records: glucose, creatinine, aspartate transaminase (AST), alanine transaminase (ALT), lactate dehydrogenase (LDH), creatine kinase (CK), albumin, ferritin, leucocyte count, lymphocyte count, eosinophil count, hemoglobin, platelet count, erythrocyte sedimentation rate (ESR), fibrinogen, and D-dimer.

Data analysis was conducted with STATA 16.1 program (StataCorp. 2019. Stata Statistical Software: Release 16. TX: StataCorp LP. USA). Student t-tests were conducted to compare serological biomarkers mean values between COVID-19 survivors with and without neuropathic post-COVID symptoms. A $P < 0.05$ was considered significant.

Results

Details of the recruitment process and demographics from the sample can be found elsewhere [4]. From 77 individuals initially evaluated, serological biomarkers data were obtained from 67 (87%), which were included in this analysis. Participants were assessed a mean of 6.0±0.8 months after hospitalization. Eighteen (26.7%) exhibited neuropathic post-COVID symptoms (S-LANSS score≥12/24 points). No significant differences in any serological biomarker at hospital admission were observed between individuals with and without neuropathic post-COVID pain symptoms (**Table**).

Discussion

This secondary analysis found no association between serological biomarkers at the acute phase of SARS-CoV-2 infection (hospital admission) and the development of neuropathic post-COVID symptoms 6 months after infection in previously hospitalized COVID-19 survivors. Our results agree with previous data also reporting that laboratory biomarkers obtained at hospital admission are not related to other post-COVID symptoms e.g., fatigue [7]. Similarly, the association between laboratory biomarkers at hospital admission and musculoskeletal post-COVID pain one year after infection is irrelevant [8]

The highly expression of Angiotensin Converting Enzyme-2 (ACE2) receptors within nervous system cells such as neurons and microglia of the spinal cord could explain the neuro-invasive potential of the SARS-CoV-2 virus explaining the presence of neuropathic symptoms in COVID-19 survivors [9]. Since several serological biomarkers analyzed, e.g., higher D-dimer concentration, lower platelet count, increased blood glucose, have been associated with severe COVID-19 [10], our results would suggest that severity of infection is not associated with the development of neuropathic post-COVID symptoms.

Some limitations should be considered. First, current data can be only applicable to previously hospitalized COVID-19 survivors with mild-to-moderate severity. Further, the sample size could be considered small, and it is probably that the lack of association in some comparisons were due to type II error. This study could be used for further sample calculation in future studies. Second, we did not include individuals with pre-existing pain symptoms. Third, as some specific inflammatory biomarkers, e.g., cytokines or C Reactive Protein, were not analyzed, these may exhibit stronger predictive strengths for development of neuropathic post-COVID pain. Finally, we determined the presence of neuropathic pain features based on a patient reported outcome measure (PROM) such as the S-LANSS. The inclusion of objective measures, e.g., electromyography, quantitative sensory testing or skin punch biopsies, could help to confirm or refute the presence of a neuropathic cause of pain symptoms in this population.

In conclusion, serological biomarkers at hospital admission were not associated with the development of neuropathic post-COVID symptoms in previously hospitalized COVID-19 survivors.

References

1. Fernández-de-las-Peñas C, Navarro-Santana M, Plaza-Manzano G, Palacios-Ceña, Arendt-Nielsen L. Time course prevalence of Post-COVID pain symptoms of musculoskeletal origin in patients who had survived to SARS-CoV-2 infection: A systematic review and meta-analysis. Pain 2021 Sep 23.doi: 10.1097/j.pain.0000000000002496.

2. D'Souza RS, Kilgore AE, D'Souza S. Manifestations of pain during the COVID-19 pandemic portrayed on social media: A cross-sectional study. Pain Med. 2022; 23: 229-233.

3. Oguz-Akarsu E, Gullu G, Kilic E, Dinç Y, Ursavas A, Yilmaz E, Zarifoglu M, Karli N; Pandemic Study Team. Insight into pain syndromes in acute phase of mild-to-moderate COVID-19: Frequency, clinical characteristics, and associated factors. Eur J Pain. 2022; 26: 492-504.

4. Herrero-Montes M, Fernández-de-las-Peñas C, Ferrer-Pargada D, Tello-Mena S, Cancela-Cilleruelo I, Rodríguez-Jiménez J, Palacios-Ceña D, Parás-Bravo P. Prevalence of neuropathic component in post-COVID pain symptoms in previously hospitalized COVID-19 survivors. Int J Clin Pract 2022; 3532917.

5. Magdy R, Eid RA, Fathy W, Abdel-Aziz MM, Ibrahim RI, Yehia A, Sheemy MS, Hussein M. Characteristics and risk factors of persistent neuropathic pain in recovered COVID-19 patients. Pain Med. 2022; 23: 774-781

6. Bennett MI, Smith BH, Torrance N, Potter J. The S-LANSS score for identifying pain of predominantly neuropathic origin: validation for use in clinical and postal research. J Pain. 2005; 6: 149-58.

7. Mandal S, Barnett J, Brill SE, Brown JS, Denny EK, Hare SS, Heightman M, Hillman TE, Jacob J, Jarvis HC, Lipman MCI, Naidu SB, Nair A, Porter JC, Tomlinson GS, Hurst JR; ARC Study Group. 'Long-COVID': a cross-sectional study of persisting symptoms, biomarker and imaging abnormalities following hospitalisation for COVID-19. *Thorax*. 2021; 76: 396-398.
8. Fernández-de-las-Peñas C, Ryan-Murua P, de-la-Llave-Rincón AI, Gómez-Mayordomo V, Arendt-Nielsen L, Torres-Macho J. Serological biomarkers of COVID-19 severity at hospital admission are not related to long-term post-COVID pain symptoms in hospitalized COVID-19 survivors. *Pain*. 2022. doi: 10.1097/j.pain.0000000000002608.
9. Torices S, Cabrera R, Stangis M, Naranjo O, Fattakhov N, Teglas T, Adesse D, Toborek M. Expression of SARS-CoV-2-related receptors in cells of the neurovascular unit: implications for HIV-1 infection. *J Neuroinflammation*. 2021; 18: 167.
10. Samprathi M, Jayashree M. Biomarkers in COVID-19: An up-to-date review. *Front Pediatr*. 2021; 8: 607647

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Role of the Funding Source

Proyecto financiado por la convocatoria Next-Val 2021 de la Fundación Instituto de Investigación Marqués de Valdecilla (IDIVAL) and by a grant from the Novo Nordisk Foundation 0067235. The sponsors had no role in the design, collection, management, analysis, or interpretation of the data, draft, review, or approval of the manuscript or its content. The authors were responsible for the decision to submit the manuscript for publication, and the sponsor did not participate in this decision.

Acknowledgements

The Center for Neuroplasticity and Pain (CNAP) is supported by the Danish National Research Foundation (DNRF121) and Norvo Nordic Foundation (NNF21OC0067235).

Declaration of interests

No conflict of interest is declared by any of the authors

Author Contributions

All authors contributed to the study concept and design. CFdLP, and MHM conducted literature review and did the statistical analysis. All authors recruited participants and collected data. PPB supervised the study. All authors contributed to interpretation of data. All authors contributed to drafting the paper. All authors revised the text for intellectual content and have read and approved the final version of the manuscript.

Downloaded from <https://academic.oup.com/painmedicine/advance-article/doi/10.1093/pm/pnac086/6604280> by Aalborg University Library user on 13 June 2022

Table: Laboratory biomarkers of COVID-19 patients according to the presence or absence of neuropathic post-COVID pain symptoms at 6 months after hospital discharge

	S-LANSS≥12 points (n=18)	S-LANSS<12 points (n=49)	P value
Glucose (mg/mL)	118.7 (59.0)	110.0 (28.0)	0.453
Creatinine (mg/dL)	0.9 (0.2)	0.9 (0.4)	0.844
Alanine transaminase (ALT, U/L)	24.8 (10.1)	25.4 (11.7)	0.864
Aspartate transaminase (AST, U/L)	22.5 (7.1)	22.7 (6.4)	0.938
Lactate dehydrogenase (LDH, U/L)	213.5 (32.7)	207.1 (45.1)	0.583
Creatine kinase (CK, mg/dL)	1.5 (2.5)	1.0 (0.4)	0.224
Albumin (g/dL)	4.5 (0.2)	4.55 (0.3)	0.424
Ferritin (ng/mL)	147.2 (187.1)	125.9 (119.8)	0.584
Leucocytes (x10⁹/L)	6.9 (1.7)	7.25 (1.5)	0.419
Lymphocytes (x10⁹/L)	3.4 (0.8)	3.1 (0.75)	0.202
Eosinophils (x10⁹/L)	2.5 (1.9)	2.7 (2.7)	0.786
Haemoglobin (g/dL)	13.9 (1.1)	14.2 (1.5)	0.453
Platelets (x10⁹/L)	238.1 (69.5)	251.0 (54.25)	0.427
Erythrocyte sedimentation rate (ESR, mm/h)	16.1 (17.65)	10.2 (9.3)	0.117
Fibrinogen (mg/dL)	432.3 (102.1)	403.85 (80.9)	0.242
D-dimer (ng/mL)	665.5 (879.7)	513.5 (404.5)	0.339

n: number; SD: Standard Deviation